

Global distribution of the effective aerosol hygroscopicity parameter for CCN activation

K. J. Pringle¹, H. Tost¹, A. Pozzer^{1,2}, U. Pöschl¹, and J. Lelieveld^{1,2}

¹Max Planck Institute for Chemistry, Mainz, Germany

²The Cyprus Institute, Energy, Environment and Water Research Centre, Nicosia, Cyprus

1 Introduction

This supplement contains an alternative data-set of simulated κ values in which κ is calculated considering the hydrophilic particles only. This is equivalent to the κ_a parameter as presented and discussed by Gunthe et al. (2009) and Rose et al. (2010). This is different from the values in the main text, which show the κ calculated considering both CCN active and inactive particles. Alternative (κ_a) versions of Tables 1 to 4, and Figures 1 and 2 are shown.

1.1 Comparison to observations - κ_a compared to κ_t

In Figure 3 we show the comparison between modeled κ and observations for the two different methods of calculating κ : (i) only CCN active particles are considered and (ii) CCN active and inactive particles are considered. When only the hydrophilic (CCN active) aerosol are considered, κ is generally overestimated and the correlation is quite poor ($R^2 = 0.34$), this overestimation is reduced when both hydrophilic and hydrophobic modes are considered, and the correlation improves ($R^2 = 0.53$). This comparison supports the findings of Wex et al. (2010) who re-analysed AMS and HTDMA measurements and found that neglecting the hydrophobic aerosols in calculation of κ for CCN calculations leads to an overestimation of CCN concentrations.

Table 1. Simulated global and regional mean κ_a values (and standard deviation (St Dev)) at the surface and at the simulated PBL height under present day conditions. Standard deviation is calculated for the year from 5-hourly average data. Difference columns show the absolute difference between κ_r and κ_a for the region.

Region	Area (10^{13} m ²)	Mean κ_a Surface	St Dev Surface	Mean κ_a PBL height	St Dev PBL height	Surface difference ($\kappa_a - \kappa_r$)	PBL difference ($\kappa_a - \kappa_r$)
Global (Continental)	14.4	0.34	0.21	0.34	0.20	0.07	0.07
Global (Marine)	37.0	0.75	0.23	0.64	0.23	0.04	0.04
N. America	1.61	0.37	0.16	0.36	0.15	0.07	0.07
S. America	1.90	0.24	0.18	0.26	0.16	0.07	0.07
Africa	3.48	0.25	0.15	0.26	0.13	0.10	0.09
Europe	1.14	0.45	0.17	0.42	0.16	0.10	0.10
Asia	3.64	0.28	0.16	0.28	0.14	0.07	0.07
Australia	0.87	0.29	0.19	0.31	0.17	0.09	0.08
N. Atlantic	1.25	0.64	0.16	0.52	0.17	0.05	0.04
Southern Ocean	1.56	0.93	0.09	0.83	0.15	0.01	0.02

Table 2. Comparison between observed and modelled κ_a values. ^A κ calculated from reported aerosol soluble fraction, following Gunthe et al. (2009). Most measurement sites were surface campaigns, with the exception of Shinozuka et al. (2010) and Hudson (2007), which are flight data. For flight data an average altitude of approx. 1500 (m) was assumed.

Site	Region	Reference	κ observed	κ model
1	Amazon	Gunthe et al. (2009)	0.16±0.06	0.16
2	China	Rose et al. (2008)	0.3	0.46
3	Mexico	Shinozuka et al. (2010)	0.2 - 0.3	0.38
4	US West Coast	Shinozuka et al. (2010)	0.18 - 0.47	0.25
5	Puerto Rico	Allan et al. (2008)	0.6±0.2	0.72
6	Antigua	Hudson (2007)	0.87±0.24	0.75
7	Amazon	Vestin et al. (2007) ^A	0.10	0.11
8	Amazon	Zhou et al. (2002) ^A	0.12	0.15
9	Tenerife	Guibert et al. (2003) ^A	0.43	0.67
10	Germany (Feldberg)	Dusek et al. (2006)	0.15 - 0.3	0.41
11	Germany (Munich)	Kandler and Shütz (2007) ^A	0.36	0.37
12	Eastern Mediterranean	Bougiatioti et al. (2009)	0.24	0.49
13	Toronto	Broekhuizen et al. (2006) ^A	0.15 - 0.3	0.38
14	Ontario	Chang et al. (2009)	0.3	0.33

Table 3. Standard deviation (St Dev) of κ values calculated considering variation in i) all dimensions (x, y, t_5 = area and time (t_5 = 5 hourly time intervals), ii) area only (x, y), iii) time only (t_5 = 5 hourly time intervals) and iv) time only (t_m = monthly mean values used). For each standard deviation value the adjacent column (to the right) shows the number of data points used in the calculation (n).

Region	Area (10^{13} m ²)	Mean κ Surface	x, y, t_5		x, y		t_5		t_m	
			St Dev	n	St Dev	n	St Dev	n	St Dev	n
Global (Cont.)	14.4	0.34	0.21	4761936	0.17	2718	0.02	1752	0.02	12
Global (Marine)	37.0	0.75	0.22	9702576	0.18	5538	0.02	1752	0.02	12
N. America	1.61	0.38	0.16	420480	0.05	240	0.09	1752	0.09	12
S. America	1.90	0.24	0.18	367920	0.15	210	0.02	1752	0.01	12
Africa	3.48	0.25	0.15	672768	0.11	384	0.03	1752	0.03	12
Europe	1.14	0.46	0.17	350400	0.09	200	0.06	1752	0.05	12
Asia	3.64	0.28	0.16	1038936	0.08	593	0.03	1752	0.03	12
Australia	0.87	0.30	0.19	173448	0.14	99	0.04	1752	0.02	12
N. Atlantic	1.25	0.63	0.16	264552	0.07	151	0.10	1752	0.09	12
Southern Ocean	1.56	0.93	0.09	346896	0.05	198	0.04	1752	0.04	12

Table 4. Simulated global and regional mean κ_a values (and standard deviation) at the surface and at the simulated PBL height under preindustrial conditions. Difference columns show the absolute difference between κ_r and κ_a for the region.

Region	Area (10^{13} m ²)	Mean κ_a Surface	St. Dev Surface	Mean κ_a PBL height	St. Dev PBL height	Surface	PBL
						difference ($\kappa_a - s\kappa_r$)	difference ($\kappa_a - \kappa_r$)
Global (Continental)	14.4	0.30	0.25	0.30	0.24	0.06	0.06
Global (Marine)	37.0	0.79	0.22	0.68	0.25	0.03	0.05
N. America	1.61	0.32	0.20	0.32	0.19	0.06	0.06
S. America	1.90	0.23	0.18	0.24	0.17	0.06	0.06
Africa	3.48	0.17	0.14	0.18	0.13	0.08	0.07
Europe	1.14	0.41	0.24	0.39	0.23	0.11	0.12
Asia	3.64	0.21	0.18	0.21	0.16	0.06	0.06
Australia	0.87	0.32	0.21	0.33	0.19	0.09	0.08
N. Atlantic	1.25	0.76	0.17	0.62	0.21	0.05	0.05
Southern Ocean	1.56	0.95	0.06	0.87	0.12	0.00	0.02




Figure 1. Annual mean distribution of κ_a at the surface simulated by EMAC.




Figure 2. Annual mean distribution of κ_a at the altitude of the planetary boundary layer.




Figure 3. Comparison of observed and modelled values of κ assuming (red) κ calculated from hydrophilic modes only (as shown in this supplement, κ_a) and (blue) κ calculated considering hydrophobic and hydrophilic modes (κ_t).

References

- Allan, J. D., Baumgardner, D., Raga, G. B., Mayol-Bracero, O. L., Morales-Garca, F., Garca-Garca, F., Montero-Martinez, G., Borrmann, S., Schneider, J., Mertes, S., Walter, S., Gysel, M., Dusek, U., Frank, G. P., and Krmer, M.: Clouds and aerosols in Puerto Rico – a new evaluation, *Atmospheric Chemistry and Physics*, 8, 1293–1309, <http://www.atmos-chem-phys.net/8/1293/2008/>, 2008.
- Bougiatioti, A., Fountoukis, C., Kalivitis, N., Pandis, S. N., Nenes, A., and Mihalopoulos, N.: Cloud condensation nuclei measurements in the marine boundary layer of the Eastern Mediterranean: CCN closure and droplet growth kinetics, *Atmospheric Chemistry and Physics*, 9, 7053–7066, 2009.
- Broekhuizen, K., Chang, R.-W., Leaitch, W. R., Li, S.-M., and Abbatt, J. P. D.: Closure between measured and modeled cloud condensation nuclei (CCN) using size-resolved aerosol compositions in downtown Toronto, *Atmospheric Chemistry and Physics*, 6, 2513–2524, 2006.
- Chang, R. Y.-W., Slowik, J. G., Shantz, N. C., Vlasenko, A., Liggi, J., Sjostedt, S. J., Leaitch, W. R., and Abbatt, J. P. D.: The hygroscopicity parameter (κ) of ambient organic aerosol at a field site subject to biogenic and anthropogenic influences: Relationship to degree of aerosol oxidation, *Atmospheric Chemistry and Physics Discussions*, 9, 25 323–25 360, 2009.
- Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D., Drewnick, F., Hings, S., Jung, D., Borrmann, S., and Andreae, M. O.: Size Matters More Than Chemistry for Cloud-Nucleating Ability of Aerosol Particles, *Science*, 312, 1375–1378, doi:10.1126/science.1125261, 2006.
- Guibert, S., Snider, J. R., and Brenguier, J.-L.: Aerosol activation in marine stratocumulus clouds: 1. Measurement validation for a closure study, *J. Geophys. Res. - Atmos.*, 108(D15), doi:10.1029/2002JD002 678, 2003.
- Gunthe, S. S., King, S. M., Rose, D., Chen, Q., Roldin, P., Farmer, D. K., Jimenez, J. L., Artaxo, P., Andreae, M. O., Martin, S. T., and Pschl, U.: Cloud condensation nuclei in pristine tropical rainforest air of Amazonia: size-resolved measurements and modeling of atmospheric aerosol composition and CCN activity, *Atmospheric Chemistry and Physics*, 9, 7551–7575, <http://www.atmos-chem-phys.net/9/7551/2009/>, 2009.
- Hudson, J. G.: Variability of the relationship between particle size and cloud-nucleating ability, *Geophys. Res. Lett.*, 34, doi:10.1029/2006GL028 850, 2007.
- Kandler, K. and Shütz: Climatology of the Average Water-soluble Volume Fraction of Atmospheric Aerosol, *Atmos. Res.*, 07, 77–92, 2007.
- Rose, D., Gunthe, S. S., Mikhailov, E., Frank, G. P., Dusek, U., Andreae, M. O., and U, P.: Calibration and measurement uncertainties of a continuous-flow cloud condensation nuclei counter (DMT-CCNC): CCN activation of ammonium sulfate and sodium chloride aerosol particles in theory and experiment, *Atmospheric Chemistry and Physics*, 8, 1153–1179, 2008.
- Rose, D., Nowak, A., Ahtert, P., Wiedensohler, A., Hu, M., Shao, M., Zhang, Y., Andreae, M. O., and U., P.: Cloud condensation nuclei in polluted air and biomass burning smoke near the megacity Guangzhou, China Part 1: Size-resolved measurements and implications for the modeling of aerosol particle hygroscopicity and CCN activity, *Atmospheric Chemistry and Physics*, 10, 3365–3383, 2010.
- Shinozuka, Y., Clarke, A. D., DeCarlo, P. F., Jimenez, J. L., Dunlea, E. J., Roberts, G. C., Tomlinson, J. M., Collins, D. R., Howell, S. G., Kapustin, V. N., McNaughton, C. S., and Zhou, J.: Aerosol optical properties relevant to regional remote sensing of CCN activity and links to their organic mass fraction: airborne observations over Central Mexico and the US West Coast during MILAGRO/INTEX-B, *Atmospheric Chemistry and Physics*, 9, 6727–6742, 2010.
- Vestin, A., Rissler, J., Swietlicki, E., Frank, G. P., and Andreae, M. O.: Cloud-nucleating properties of the Amazonian biomass burning aerosol: Cloud condensation nuclei measurements and modeling, *J. Geophys. Res. - Atmos.*, 112, doi:10.1029/2006JD008 104, 2007.
- Wex, H., McFiggans, G., Henning, S., and Stratmann, F.: Influence of the external mixing state of atmospheric aerosol on derived CCN number concentrations, *Geophys. Res. Lett.*, in press, doi:10.1029/2010GL043 337, 2010.
- Zhou, J., Swietlicki, E., Hansson, H. C., and Artaxo, P.: Submicrometer aerosol particle size distribution and hygroscopic growth measured in the Amazon rain forest during the wet season, *Journal of Geophysical Research (Atmospheres)*, p. doi:10.1029/2000JD000203, 2002.