

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Terpenylic acid and related compounds

F. Yasmeen et al.

Terpenylic acid and related compounds: precursors for dimers in secondary organic aerosol from the ozonolysis of α - and β -pinene

F. Yasmeen^{1,2}, R. Vermeylen¹, R. Szmigielski^{1,*}, Y. Iinuma³, O. Böge³,
H. Herrmann³, W. Maenhaut⁴, and M. Claeys¹

¹Department of Pharmaceutical Sciences, University of Antwerp, 2610 Antwerp, Belgium

²Laboratoire de Radiochimie, Sciences Analytiques et Environnement (LRSAE), Institut de Chimie de Nice (CNRS, FR 3037), University of Nice Sophia-Antipolis, Faculty of Sciences, 06108 Nice, France

³Leibniz-Institut für Troposphärenforschung (IfT), 04318 Leipzig, Germany

⁴Department of Analytical Chemistry, Institute for Nuclear Sciences, Ghent University, 9000 Gent, Belgium

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



* present address: Institute of Physical Chemistry of the Polish Academy of Sciences, 01-224, Warsaw, Poland

Received: 7 April 2010 – Accepted: 14 April 2010 – Published: 23 April 2010

Correspondence to: M. Claeys (magda.claeys@ua.ac.be)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

10, 10865–10888, 2010

Terpenylic acid and related compounds

F. Yasmeeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

In the present study, we have characterized the structure of a higher-molecular weight (MW) 358 α - and β -pinene dimeric secondary organic aerosol (SOA) product that received ample attention in previous molecular characterization studies. Based on mass spectrometric evidence for deprotonated molecules formed by electrospray ionization in the negative ion mode, we propose that diaterpenylic acid is a key monomeric unit for dimers of the ester type. It is shown that *cis*-pinic acid is esterified with the hydroxyl-containing diaterpenylic acid which can be explained through acid-catalyzed hydrolysis of the recently elucidated lactone-containing terpenylic acid and/or diaterpenylic acid acetate, both first-generation oxidation products. To a minor extent, higher-MW 358 and 344 diester products are formed containing other terpenoic acids as monomeric units, i.e., diaterpenylic acid instead of *cis*-pinic acid, and diaterebic acid instead of diaterpenylic acid. It is shown that the MW 358 diester and related MW 344 compounds, which can be regarded as processed SOA products, also occur in ambient fine (PM_{2.5}) rural aerosol collected at night during the warm period of the 2006 summer field campaign conducted at K-pusztá, Hungary, a rural site with coniferous vegetation. This indicates that, under ambient conditions, the higher-MW diesters are formed in the particle phase over a longer time-scale than that required for gas-to-particle partitioning of their monomeric precursors.

1 Introduction

Structural proposals have been made in previous work for higher-molecular weight (MW) secondary organic aerosol (SOA) products that participate in new particle formation and growth, including dimers and oligomers. The structural elucidation at the molecular level of such compounds is a complex and analytically challenging task as it is only readily feasible if the structures of the composing monomers are known. Conversely, lack of knowledge about the monomeric units has severely hampered struc-

Terpenylic acid and related compounds

F. Yasmeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tural characterization efforts of higher-MW di- and oligomeric compounds in previous studies.

Considerable attention has been given in previous work to a higher-MW 358 SOA product from α -pinene ozonolysis (Hoffmann et al., 1998; Gao et al., 2004; Iinuma et al., 2004), which was first detected by direct negative ion atmospheric pressure chemical ionization – mass spectrometry and tentatively assigned to a stable adduct between *cis*-pinic (MW 186) and *cis*-norpinic acid (MW 172), both ozonolysis products (Hoffmann et al., 1998). It was shown in the latter study that the MW 358 compound remained intact upon reversed-phase liquid chromatography; however, a straightforward explanation for the formation of a stable dimer between *cis*-pinic and *cis*-norpinic acid was not presented. Following this initial study by Hoffmann et al. (1998), other tentative structures have been proposed for higher-MW α -pinene ozonolysis products (Gao et al., 2004; Iinuma et al., 2004; Docherty et al., 2005). Gao et al. (2004) attributed the MW 358 α -pinene ozonolysis product to a dehydration product formed between the gem-diol forms of two norpinonic acid molecules. Iinuma et al. (2004) reported MW 354 and 370 dimeric products from the ozonolysis of α -pinene that were enhanced in acidic conditions and tentatively assigned the products to reaction products between the the gem-diol forms of two molecules of pinonaldehyde and those of pinonaldehyde and hydroxypinonaldehyde, respectively. Docherty et al. (2005) proposed peroxy-carboxylic acid dimers for the structure of higher-MW SOA products from the ozonolysis of α -pinene in which peroxy-pinic acid and the gem-diol of a keto or aldehydic compound are connected via a peroxy bridge. Furthermore, the MW 358 product was detected as a major tracer in other α -pinene as well as β -pinene ozonolysis SOA characterization studies (Iinuma et al., 2007; Müller et al., 2008). In addition, Tolocka et al. (2004) characterized dimers in α -pinene ozonolysis SOA and suggested that the dimers were most likely formed by aldol and/or gem-diol formation. Müller et al. (2008) reported the presence of abundant MW 358 and 368 products upon α -pinene ozonolysis and provided evidence for esterification of *cis*-pinic acid with hydroxyl-containing terpenic acids as a major formation pathway; the MW 370 product was attributed to a diester

Terpenylic acid and related compounds

F. Yasmeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



formed between *cis*-pinic acid and 10-hydroxypinonic acid but the structure of the MW 358 product remained unresolved.

Esterification together with anhydride formation were shown to be important routes for dimer formation from the ozonolysis of cyclohexene (Hamilton et al., 2006). In addition, esterification was shown to operate in the formation of di-, tri-, and tetramers in α -pinene ozonolysis based on direct mass spectrometric analysis of complex SOA mixtures and accurate mass measurements that revealed repetitive structures and changes in O:C ratios (Reinhardt et al., 2007). Esterification of small hydroxyl carboxylic acids such as hydracrylic, glycolic, lactic, and 2-methylglyceric acid was also found to be a route resulting in isoprene SOA formation (Altieri et al., 2006; 2008; Surratt et al., 2006; Szmigielski et al., 2007; Perri et al., 2009). Furthermore, theoretical calculations by Barsanti and Pankow (2006) show that esterification is a thermodynamically favorable route for SOA formation under atmospheric conditions.

In the present study, we address the structure of a major MW 358 compound and related minor MW 344 and 358 compounds that are formed upon α - and β -pinene ozonolysis and are also present in ambient fine (PM_{2.5}, particulate matter with aerodynamic diameter <2.5 μ m) rural continental aerosol. We propose that diaterpenylic acid, a hydrolysis product of terpenylic acid (MW 172) and/or diaterpenylic acid acetate (MW 232), recently reported α -pinene SOA tracers formed upon both photooxidation and ozonolysis (Claeys et al., 2009), serves together with *cis*-pinic acid as a monomeric building unit for the higher-MW 358 diester products.

2 Experimental

2.1 Aerosol samples

α - and β -pinene SOA for structural characterization was obtained from ozonolysis experiments carried out in the 19 m³ IfT smog chamber. Briefly, the ozonolysis of α - and β -pinene was carried out in the presence of acidic seed particles [0.03/0.06 M

Terpenylic acid and related compounds

F. Yasmeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Terpenylic acid and related compounds

F. Yasmeen et al.

(NH₄)₂SO₄/H₂SO₄]. The relative humidity and temperature of the chamber were approximately 50% and 23 °C for all experiments. The initial concentrations of α -pinene and ozone were 100 ppb and 72 ppb, respectively. For β -pinene ozonolysis, the initial concentrations were 300 ppb for β -pinene and 108 ppb for ozone. No OH scavenger was used in these chamber experiments. The filter samples taken were extracted with methanol (3 x 20 mL), and the combined methanol extracts were concentrated by rotary evaporation to about 1 mL. Subsequently, the concentrated extracts were filtered over a Teflon filter (0.45 μ m), then further evaporated under a nitrogen stream and reconstituted in 200 μ L methanol. For liquid chromatography/mass spectrometry (LC/MS) analysis a part of the samples was diluted with the same volume of water and an aliquot of 5 μ L was injected.

The ambient aerosol samples used in this work were collected from K-pusztá, Hungary, during the BIOSOL (Formation mechanisms, marker compounds, and source apportionment for biogenic atmospheric aerosols) campaign between 22 May and 29 June 2006. Information about the site and the sampling campaign is given in the supplement (Sect. 1). Day- and night-time samples were collected using a high-volume dichotomous sampler providing two size fractions, a fine (PM_{2.5}) and a coarse size fraction (with aerodynamic diameter >2.5 μ m). A quarter or one eighth (depending on the organic carbon amount) of the PM_{2.5} quartz fiber filter was extracted with methanol (3x20 mL), the residue was redissolved in 150 μ L of a methanol:water (1:1, v/v) mixture, and an aliquot of 5 μ L (corresponding to 5.6 μ g organic carbon in the case of the selected nighttime sample of 19 June, Fig. 2) was injected for LC/MS analysis.

2.2 Chemicals

The terpenoic acid standards used in this work were: *cis*-pinic acid (purity >98%; Sigma-Aldrich, St. Louis, MI, USA) and terebic acid (purity >98%; TCI Europe, Zwijndrecht, Belgium). Terpenylic acid and diaterpenylic acid acetate were available from previous studies (Claeys et al., 2009; linuma et al., 2009).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2.3 LC/MS analysis

The LC/MS system comprised a Surveyor Plus system (pump and autosampler), a linear ion trap mass spectrometer (LXQ) equipped with an electrospray ionization (ESI) source, and a data system using Xcalibur version 2.0 software (Thermo Scientific, San Jose, USA). A T3 Atlantis C18 column (3 μm ; 2.1 \times 150 mm) (Waters, Milford, USA) was employed. The mobile phases consisted of acetic acid 0.1% (v/v) (A) and methanol (B). The applied 80-min gradient elution program was as follows: the concentration of eluent B was kept at 3% for 2 min, then increased to 90% in 18 min, kept at 90% for 43 min, then decreased to 3% in 5 min, and kept at 3% for 12 min. The flow rate was 0.2 mL min^{-1} . The linear ion trap was operated under the following conditions: sheath gas flow (nitrogen), 0.75 L min^{-1} ; auxiliary gas flow (nitrogen), 1.5 L min^{-1} ; source voltage, -4.5 kV; capillary temperature, 350 $^{\circ}\text{C}$; and maximum ion injection time, 200 ms. For MS^2 and MS^3 experiments, an isolation width of 2 m/z units and a normalized collision energy level of 35% were applied.

Accurate mass data for the major MW 358 compound present in α - and β -pinene ozonolysis SOA available from a previous study (Iinuma et al., 2007) indicated an elemental composition of $\text{C}_{17}\text{H}_{25}\text{O}_8$ for the deprotonated molecule.

3 Results and discussion

3.1 Structural characterization of MW 172, 186, and 232 terpenoic acids

In this section we deal with the structural characterization of terpenoic acids that are of relevance as monomeric precursors to the dimeric MW 344 and 358 compounds that will be addressed in the next sections as well as with the structural characterization of some minor monomeric/isobaric compounds. Figure 1a and b present chromatographic data [base peak chromatograms (BPCs) and extracted ion chromatograms (EICs)] for SOA from the ozonolysis of α - and β -pinene, respectively, while Fig. 2

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Terpenylic acid and related compounds

F. Yasmeen et al.

presents chromatographic data for a nighttime PM_{2.5} aerosol sample from K-pusztá. It can be seen that the chromatographic data for α - and β -pinene SOA are rather similar. The major peak on the m/z 171 EICs with RT 17.3 min is assigned to the lactone-containing terpenylic acid, a novel terpenoic acid which has only been recently reported (Claeys et al., 2009), based on comparison of its chromatographic and mass spectrometric behaviors with that of a synthesized reference compound. It is noted that terpenylic acid is an isobaric isomer of *cis*-norpinic acid. For comparison reasons (see next section), (–)ESI-MS data for terpenylic acid are provided in Fig. 3 and Scheme 1. It is noted that the product ions at m/z 127 and 83 are also characteristic for diaterpenylic acid acetate (Claeys et al., 2009) and are also expected to be formed from diaterpenylic acid, produced through hydrolysis of terpenylic acid and diaterpenylic acid acetate (see below). It can be seen that there is also a minor peak on the m/z 171 EIC with RT 19.8 min, which is attributed to *cis*-norpinic acid based on detailed interpretation of the (–)ESI-MS data and comparison with LC/MS data reported in the literature (Glasius et al., 2000; Warnke et al., 2006). (–)ESI-MS data for the latter MW 172 compound are given in the supplement (Fig. S1, Scheme S1; Sect. 2, <http://www.atmos-chem-phys-discuss.net/10/10865/2010/acpd-10-10865-2010-supplement.pdf>). It is likely that in previous work terpenylic acid has often been misassigned to *cis*-norpinic acid, owing to the fact that its mass spectrometric fragmentation is consistent with the presence of two carboxyl groups and thus rather misleading.

The m/z 185 EIC obtained for α - and β -pinene ozonolysis SOA (Fig. 1) shows a prominent peak at RT 20.9 min, which is assigned to *cis*-pinic acid, known to be formed through both ozonolysis and OH radical-initiated reactions (e.g., Yu et al., 1999; Glasius et al., 2000; Larsen et al., 2001; Winterhalter et al., 2003; Ng et al., 2007). For comparison reasons (see next section), (–)ESI-MS data for *cis*-pinic acid are provided in Fig. 3 and Scheme 1. The second m/z 185 compound present in β -pinene ozonolysis SOA at RT 18.8 min corresponds to a new and specific β -pinene SOA tracer, which is tentatively identified as a homolog of terpenylic acid, i.e., ho-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Terpenylic acid and related compounds

F. Yasmeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



moterpenylic acid, based on detailed interpretation of (–)ESI-MS data (supplement; Sect. 3: Fig. S2; Scheme S2, <http://www.atmos-chem-phys-discuss.net/10/10865/2010/acpd-10-10865-2010-supplement.pdf>). Comparison of the MS data obtained for *cis*-pinic acid and homoterpenylic acid (Fig. 3, Scheme 1) shows that the isobaric compounds can be differentiated on the basis of the m/z 185→ m/z 141 MS³ product ion spectrum; in the case of *cis*-pinic acid fragmentation of m/z 141 results in m/z 99 through loss of 42 u (propene), while in the case of homoterpenylic acid m/z 141 produces m/z 97 through loss of 44 u (CO₂). It can also be seen that the relative abundance of m/z 123 in the m/z 185 MS² product ion spectrum is the highest in the case of homoterpenylic acid (i.e., 20% versus 2% for *cis*-pinic acid). A possible pathway to homoterpenylic acid involving reaction of β -pinene with OH radicals (which are also generated in the ozonolysis experiments since no OH scavenger was used), similar to that proposed for terpenylic acid (Claeys et al., 2009), is provided in the supplement (Sect. 3: Scheme S3).

The m/z 231 EIC contains four distinct peaks; the first two peaks with RTs 17.3 and 19.8 min are interpreted as acetic acid (present in the LC eluent) non-covalent adducts of the MW 172 compounds (i.e., terpenylic and *cis*-norpinic acid), while the last two peaks correspond to isomeric MW 232 compounds. Of these, the peak with RT 20.6 min is assigned to diaterpenylic acid acetate, which has recently been reported as a SOA tracer for the photooxidation of 1,8-cineole (Iinuma et al., 2009) and for the photooxidation and ozonolysis of α -pinene (Claeys et al., 2009). The last eluting m/z 231 compound with RT 22.1 min corresponds to an isomer of diaterpenylic acid acetate, a new compound which still remains to be elucidated. In this respect, it has been reported by Iinuma et al. (2004, 2005) that three isomers of a MW 232 compound are detected in α -pinene ozonolysis SOA using capillary electrophoresis/(–)ESI-MS. (–)ESI-MS data for diaterpenylic acid acetate and the isomeric MW 232 compound with RT 22.1 min are given in the supplement (Sect. 4: Figs. S3 and S4; Scheme S4).

Comparison of the chromatographic data for α - and β -pinene SOA and ambient samples from K-pusztá shows the presence of all the terpenoic acids mentioned above

Terpenylic acid and related compounds

F. Yasmeen et al.

in all samples. The data obtained for a nighttime sample of the warm period of the campaign are illustrated in Fig. 2. It is noted that the major MW 358 compound (m/z 357) is only detected in the nighttime samplings of the warm period. Hence, an effort was done to structurally characterize the MW 358 compound and related minor MW 358 and 344 compounds.

3.2 Structural characterization of MW 358 dimeric compounds

Figure 4 presents the (–)ESI-MS data for the major MW 358 compound that can be detected at RT 24.7 min in α -pinene ozonolysis SOA. Exactly the same product ion data were obtained for the major MW 358 compound in the ambient nighttime sample (supplement; Sect. 5: Fig. S5, <http://www.atmos-chem-phys-discuss.net/10/10865/2010/acpd-10-10865-2010-supplement.pdf>), while some differences were observed for the major MW 358 compound in β -pinene ozonolysis SOA (Fig. 5) (see below). The major higher-MW 358 compound in α -pinene ozonolysis SOA is assigned to a covalent dimer of the ester type containing diaterpenylic acid and *cis*-pinic acid as monomeric building units, based on mass spectral considerations. The first-order mass spectrum (Fig. 4a) shows in addition to the deprotonated molecule $[M-H]^-$ (m/z 357) a corresponding Na^+ -containing species $[M-2H + Na]^-$ (m/z 377) and an ion at m/z 343 due to co-elution of a homologous MW 344 species (see below). The m/z 357 MS^2 spectrum (Fig. 4, B1) shows two abundant product ions at m/z 185 and 171, which upon further fragmentation (MS^3) reveal a pattern comparable to that observed for deprotonated *cis*-pinic and terpenylic acid, respectively (Fig. 3). However, there are differences in the relative abundances of the product ions, a feature which can be explained by a different internal energy of the first- (MS^2) and second-generation (MS^3) precursor ions.

The formation of the MW 358 compound can be explained by esterification of *cis*-pinic acid with diaterpenylic acid, which can be generated by acid-catalyzed hydrolysis of terpenylic acid and/or diaterpenylic acid acetate (Scheme 2). In this respect, it has recently been demonstrated that esterification is an important route to the formation of higher-MW products in α -pinene ozonolysis SOA and that *cis*-pinic acid is a key

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

monomeric unit (Müller et al., 2008).

Since *cis*-pinic acid contains two carboxyl groups, two isomeric diester structures are possible. In order to determine which positional isomeric structure belongs to the major MW 358 compound, we also recorded data for the minor MW 358 compound present in β -pinene ozonolysis SOA at RT 23.0 min (Fig. 1b). Selected (–)ESI-MS data (i.e., the m/z 357 MS² spectrum) for the minor MW 358 compound are given in Fig. 5d. The higher-order MS^{*n*} spectra were consistent with diaterpenylic and *cis*-pinic monomeric units (not shown). Compared to the m/z 357 MS² spectrum of the major MW 358 compound (Fig. 5, B1) it can be seen that the spectrum for the minor isomer (Fig. 5d) contains an additional product ion at m/z 189. The latter ion is explained by a neutral loss of 188 u corresponding to *cis*-pinic acid, which can only readily occur in the isomer where the ester linkage is at the carboxylmethyl side (Scheme 2). On the basis of these considerations structures (*a1*) and (*a2*) are tentatively attributed to the major and minor MW 358 compounds, respectively. A possible explanation for the favorable ester formation at the carboxyl (attached to the dimethylcyclobutane ring) instead of the carboxylmethyl side of *cis*-pinic acid is that there is a lower conformational flexibility at this side, facilitating ester formation.

With respect to differences observed between the major MW 358 diester in α - and β -ozonolysis SOA, it can be seen that the m/z 357→ m/z 185→ m/z 141 MS⁴ product ion spectrum in the case of β -pinene shows an additional m/z 97 product ion (Fig. 5, C2). This ion is typical for homoterpenylic acid (Fig. S2, <http://www.atmos-chem-phys-discuss.net/10/10865/2010/acpd-10-10865-2010-supplement.pdf>). It can also be noted that the m/z 357→ m/z 185 MS³ product ion spectrum (Fig. 5, C1) is more similar to the m/z 185 MS² spectrum of homoterpenylic acid (i.e., the relative abundance of m/z 123 is 20%). Hence, it cannot be ruled out that the major MW 358 diester in β -ozonolysis SOA contains a major contribution from a dimer formed between diaterpenylic and homoterpenylic acid.

Terpenylic acid and related compounds

F. Yasmeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3 Structural characterization of MW 344 dimeric compounds

The m/z 343 EICs obtained for α - and β -pinene ozonolysis SOA (Fig. 1) show two major peaks with RTs 17.3 and 24.7 min. The first one is attributed to strong non-covalent dimer formation of terpenylic acid upon (–)ESI-MS, a phenomenon that has been addressed in detail in Claeys et al. (2009) and has been attributed to the presence of one free carboxylic acid group that does not participate in intramolecular hydrogen bonding interactions. It is worth noting that such type of dimer formation is not observed for *cis*-norpinic acid (RT 19.8 min), consistent with the unique non-covalent dimer-forming properties of terpenylic acid. The second major m/z 343 compound is due to a MW 344 compound that co-elutes with the major MW 358 compound (Fig. 4a). (–)ESI-MS data and their interpretation are presented in the supplement (Sect. 6: Fig. S6, <http://www.atmos-chem-phys-discuss.net/10/10865/2010/acpd-10-10865-2010-supplement.pdf>). This MW 344 compound is attributed to a homolog of the major MW 358 compound, which can be explained by reaction between diaterebic acid, a homolog of diaterpenylic acid, and *cis*-pinic acid. A close inspection of the (–)ESI-MS data indeed reveals that the precursor of diaterebic acid, terebic acid, is detected at m/z 157 (RT 15.8 min) with an abundance of about 21%, 13.9%, and 85% of that of terpenylic acid for α - and β -pinene SOA, and the ambient sample, respectively. Terebic acid can be regarded as a processed derivative of terpenylic acid; its possible formation through further reaction with OH radicals (aging) and (–)ESI-MS data are provided in the supplement (Sect. 7: Scheme S5, Fig. S7, <http://www.atmos-chem-phys-discuss.net/10/10865/2010/acpd-10-10865-2010-supplement.pdf>). Similar to terpenylic acid, terebic acid was found to exhibit unique non-covalent dimer-forming properties in (–)ESI-MS. Furthermore, the major MW 344 compound could also be detected in the ambient sample (not shown).

It can also be seen that the m/z 343 EICs for α - and β -pinene ozonolysis SOA (Fig. 1) show minor signals at RTs 23.0 and 23.6 min. The data obtained for the peak

Terpenylic acid and related compounds

F. Yasmeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Terpenylic acid and related compounds

F. Yasmeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



at RT 23.0 min were similar to those for the major peak at RT 24.7 min (not shown); on the basis of these data this minor peak is attributed to a homolog of the minor MW 358 compound (i.e., the dimer where the diaterebic acid residue is at the carboxylmethyl side of *cis*-pinic acid). The minor MW 344 compound eluting at RT 23.6 min is assigned to a dimer of the ester type containing solely terpenylic and diaterpenylic acid units (Scheme 2; structure, *a3*). (–)ESI-MS data for the latter compound are presented in the supplement (Sect. 8: Fig. S8, <http://www.atmos-chem-phys-discuss.net/10/10865/2010/acpd-10-10865-2010-supplement.pdf>). Similar data were also recorded for the ambient sample (not shown).

We also evaluated whether the higher-MW 368 product reported by Müller et al. (2008) corresponding to a diester formed between *cis*-pinic acid and 10-hydroxypinonic acid was present in our samples and how its abundance compared with that of the major MW 358 compound; the MW 368 compound could be detected at RT 25.9 min for α - and β -pinene ozonolysis SOA, and the ambient sample, with an abundance of about 54%, 1.5%, and 54% of that of the major MW 358 compound, respectively. Furthermore, we also evaluated whether higher-MW dimers (MW 372) containing *cis*-pinic acid and the hydrolysis product of homoterpenylic acid (i.e., dihomoterpenylic acid) could be detected in β -pinene ozonolysis SOA and the ambient samples considering that homoterpenylic acid is present in these samples, but no evidence could be found for them.

4 Conclusions

The results of this study are consistent with terpenylic acid and related terpenoic acids (i.e., diaterpenylic acetate and terebic acid) serving together with *cis*-pinic acid as monomeric precursors for a major MW 358 dimeric compound as well as for structurally related minor MW 358 and 344 compounds in α -/ β -pinene ozonolysis SOA. Our study lends support to previous studies (Hamilton et al., 2006; Müller et al., 2008) that provided evidence for esterification being an important route to dimer formation

Terpenylic acid and related compounds

F. Yasmeen et al.

in ozonolysis SOA. It is shown in the present study that the dimeric α -/ β -pinene SOA tracers with MWs 358 and 344 could only be detected in the nighttime samplings of the warm period of the 2006 K-pusztá summer field campaign. This suggests that their formation in the ambient atmosphere occurs at a relatively high temperature (daily maximum temperature $>24^{\circ}\text{C}$) and a long time-scale. This behavior contrasts with the fast formation of the MW 358 diester in α - and β -pinene ozonolysis experiments, where the time-scale at which this product is formed is comparable with that of the corresponding monomeric precursors (Müller et al., 2008; Hall et al., unpublished results). Under the above mentioned ambient conditions the higher-MW dimers appear to be formed in the particle phase over a longer time-scale than that required for gas-to-particle partitioning of their monomeric precursors. Further work is warranted to assess the occurrence of the monoterpene SOA diester products in other forested environments and their usefulness as SOA tracers.

Acknowledgements. This study was financially supported by the following organizations: for Universities of Antwerp (UA) and Ghent: Belgian Federal Science Policy Office (BIOSOL project), and Research Foundation – Flanders; and for Leibniz-Institut für Troposphärenforschung: Deutsche Forschungsgemeinschaft.

References

- Altieri, K. E., Carlton, A. G., Lim, H.-J., Turpin, B. J., and Seitzinger, S. P.: Evidence for oligomer formation in clouds: reactions of isoprene oxidation products, *Environ. Sci. Technol.*, 40, 4956–4960, 2006.
- Altieri, K. E., Seitzinger, S. P., Carlton, A. G., Turpin, B. J., Klein, G. C., and Marshall, A. G.: Oligomers formed through in-cloud methylglyoxal reactions: chemical composition, properties, and mechanisms investigated by ultra-high resolution FT-ICR mass spectrometry, *Atmos. Environ.*, 42, 1476–1490, 2008.
- Barsanti, K. C. and Pankow, J. F.: Thermodynamics of the formation of atmospheric organic particulate matter by accretion reactions – Part 3: Carboxylic and dicarboxylic acids, *Atmos. Environ.*, 40, 6676–6686, 2006.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Terpenylic acid and related compounds

F. Yasmeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Claeys, M., Iinuma, Y., Szmigielski, R., Surratt, J. D., Blockhuys, F., Van Alsenoy, C., Böge, O., Sierau, B., Gómez-González, Y., Vermeylen, R., Van der Veken, P., Shahgholi, M., Chan, A. W. H., Herrmann, H., Seinfeld, J. H., and Maenhaut, W.: Terpenylic acid and related compounds form the oxidation of α -pinene: Implications for new particle formation and growth above forest, *Environ. Sci. Technol.*, 43, 6976–6982, 2009; supporting information available online at: <http://www.acs.com>.

Docherty, K. S., Wu, W., Lim, Y. B., and Ziemann, P. J.: Contributions of organic peroxides to secondary aerosol formed from reactions of monoterpenes with O_3 , *Environ. Sci. Technol.*, 39, 4049–4059, 2005.

Gao, S., Ng, N. L., Keywood, M., Varutbangkul, V., Bahreini, R., Nenes, A., He, J., Yoo, K. Y., Beauchamp, J. L., Hodyss, R., Flagan, R. C., and Seinfeld, J. H.: Particle phase acidity and oligomer formation in secondary organic aerosol, *Environ. Sci. Technol.*, 38, 6582–6589, 2004.

Glasius, M., Lahaniati, M., Calogirou, A., Di Bella, D., Jensen, N. R., Hjorth, J., Kotzias, D., and Larsen, B. R.: Carboxylic acids in secondary aerosols from oxidation of cyclic monoterpenes by ozone, *Environ. Sci. Technol.*, 34, 1001–1010, 2000.

Hamilton, J. F., Lewis, A. C., Reynolds, J. C., Carpenter, L. J., and Lubben, A.: Investigating the composition of organic aerosol resulting from cyclohexene ozonolysis: low molecular weight and heterogeneous reaction products, *Atmos. Chem. Phys.*, 6, 4973–4984, 2006, <http://www.atmos-chem-phys.net/6/4973/2006/>.

Hoffmann, T., Bandur, R., Marggraf, U., and Linscheid, M.: Molecular composition of organic aerosols formed in the α -pinene/ O_3 reaction: Implications for new particle formation processes, *J. Geophys. Res.*, 103, 25569–25578, 1998.

Iinuma, Y., Böge, O., Gnauk, T., and Herrmann, H.: Aerosol-chamber study of the α -pinene/ O_3 reaction: influence of particle acidity on aerosol yields and products, *Atmos. Environ.*, 38, 761–773, 2004.

Iinuma, Y., Böge, O., Miao, Y., Sierau, B., Gnauk, T., and Herrmann, H.: Laboratory studies on secondary organic aerosol formation from terpenes, *Faraday Discuss.*, 130, 279–294, 2005.

Iinuma, Y., Müller, C., Berndt, T., Böge, O., Claeys, M., and Herrmann, H.: Evidence for the existence of organosulfates from beta-pinene ozonolysis in ambient secondary organic aerosol, *Environ. Sci. Technol.*, 41, 6678–6683, supporting information available online at <http://www.acs.com>, 2007.

Iinuma, Y., Böge, O., Keywood, M., Gnauk, T., and Herrmann, H.: Diaterebic acid acetate and

Terpenylic acid and related compounds

F. Yasmeen et al.

diaterpenylic acid acetate: Atmospheric tracers for secondary organic aerosol formation from 1,8-cineole oxidation, *Environ. Sci. Technol.*, 43, 280–285, supporting information available online at <http://www.acs.com>, 2009.

Larsen, B. R., Di Bella, D., Glasius, M., Winterhalter, R., Jensen, N. R., and Hjorth, J.: Gas-phase OH oxidation of monoterpenes: gaseous and particulate products, *J. Atmos. Chem.*, 38, 231–276, 2001.

Müller, L., Reinnig, M.-C., Warnke, J., and Hoffmann, T.: Unambiguous identification of esters as oligomers in secondary organic aerosol formed from cyclohexene and cyclohexene/ α -pinene ozonolysis, *Atmos. Chem. Phys.*, 8, 1423–1433, 2008, <http://www.atmos-chem-phys.net/8/1423/2008/>.

Ng, N. L., Chhabra, P. S., Chan, A. W. H., Surratt, J. D., Kroll, J. H., Kwan, A. J., McCabe, D. C., Wennberg, P. O., Sorooshian, A., Murphy, S. M., Dalleska, N. F., Flagan, R. C., and Seinfeld, J. H.: Effect of NO_x level on secondary organic aerosol (SOA) formation from the photooxidation of terpenes, *Atmos. Chem. Phys.*, 7, 5159–5174, 2007, <http://www.atmos-chem-phys.net/7/5159/2007/>.

Perri, M. J., Seitzinger, S., and Turpin, B. J.: Secondary organic aerosol production from aqueous photooxidation of glycolaldehyde: laboratory experiments, *Atmos. Environ.*, 43, 1487–1497, 2009.

Reinhardt, A., Emmenegger, C., Gerrits, B., Panse, C., Dommen, J., Baltensperger, U., Zenobi, R., and Kalberer, M.: Ultrahigh mass resolution and accurate mass measurements as a tool to characterize oligomers in secondary organic aerosols, *Anal. Chem.*, 79, 4074–4082, 2007.

Surratt, J. D., Murphy, S. M., Kroll, J. H., Ng, N. L., Hildebrandt, L., Sorooshian, A., Szmigielski, R., Vermeylen, R., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Chemical composition of secondary organic aerosol formed from the photooxidation of isoprene, *J. Phys. Chem. A*, 110, 9665–9690, 2006.

Szmigielski, R., Surratt, J. D., Vermeylen, R., Szmigielska, K., Kroll, J. H., Ng, N. L., Murphy, S. M., Sorooshian, A., Seinfeld, J. H., and Claeys, M.: Characterization of 2-methylglyceric acid oligomers in secondary organic aerosol from the photooxidation of isoprene using trimethylsilylation and gas chromatography/ion trap mass spectrometry, *J. Mass Spectrom.*, 42, 101–116, 2007.

Tolocka, M. P., Jang, M., Ginter, J. M., Cox, F. J., Kamens, R. M., and Johnston, M. V.: Formation of oligomers in secondary organic aerosol, *Environ. Sci. Technol.*, 38, 1428–1434,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

supporting information available online at <http://www.acs.com>, 2004.

Warnke, J., Bandur, R., and Hoffmann, T.: Capillary-HPLC-ESI-MS/MS method for the determination of acidic products from the oxidation of monoterpenes in atmospheric aerosol samples, *Anal. Bioanal. Chem.*, 385, 34–45, 2006.

5 Winterhalter, R., Van Dingenen, R., Larsen, B. R., Jensen, N. R., and Hjorth, J.: LC-MS analysis of aerosol particles from the oxidation of α -pinene by ozone and OH-radicals, *Atmos. Chem. Phys. Discuss.*, 3, 1–39, 2003, <http://www.atmos-chem-phys-discuss.net/3/1/2003/>.

10 Yu, J., Cocker III, D. R., Griffin, R. J., Flagan, R. C., and Seinfeld, J. H.: Gas-phase ozone oxidation of monoterpenes: Gaseous and particulate products, *J. Atmos. Chem.*, 34, 207–258, 1999.

Terpenylic acid and related compounds

F. Yasmeen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Terpenylic acid and related compounds

F. Yasmeen et al.




Fig. 1. LC/MS chromatographic data [base peak chromatograms (BPCs) and extracted ion chromatograms (EICs)] for **(A)** α -pinene and **(B)** β -pinene ozonolysis SOA. Abbreviation: NL, normalization level.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Terpenylic acid and related compounds

F. Yasmeen et al.




Fig. 2. LC/MS chromatographic data (BPC and EICs) for fine ambient aerosol ($PM_{2.5}$) from K-pusztá, Hungary, collected during the night of 19 June 2006 of the warm period of the 2006 summer field campaign. The peak at RT 22.1 min (not discussed in the main text) on the m/z 185 EIC is tentatively assigned to *cis*-caronic acid, a positional isomer of *cis*-pinic acid, based on comparison of its chromatographic properties with literature data (Larsen et al., 2001).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Terpenylic acid and related compounds

F. Yasmeen et al.




Fig. 3. (–)ESI-MS data for **(A)** terpenylic acid and **(B)** *cis*-pinic acid.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 4. (–)ESI-MS data for the major MW 358 compound (RT 24.7 min; Fig. 1a) present in α -pinene ozonolysis SOA, assigned to a covalent dimer of the ester type: **(A)** first-order mass spectrum; **(B1)** m/z 357 MS^2 spectrum; **(B2)** and **(B3)** m/z 357 \rightarrow m/z 171 MS^n ($n = 3, 4$) spectra; and, **(C1)** and **(C2)** m/z 357 \rightarrow m/z 185 MS^n ($n = 3, 4$) spectra.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Terpenylic acid and related compounds

F. Yasmeen et al.





Fig. 5. (–)ESI-MS data for the major MW 358 compound [RT 24.7 min; Fig. 1(B)] present in β -pinene ozonolysis SOA, assigned to a covalent dimer of the ester type: **(A)** first-order mass spectrum; **(B1)** m/z 357 MS^2 spectrum; **(B2)** and **(B3)** m/z 357 $\rightarrow m/z$ 171 MS^n ($n = 3, 4$) spectra; and, **(C1)** and **(C2)** m/z 357 $\rightarrow m/z$ 185 MS^n ($n = 3, 4$) spectra. **(D)** m/z 357 MS^2 spectrum for the minor MW 358 compound at RT 23.0 min (Fig. 1b).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Terpenylic acid and related compounds

F. Yasmeen et al.



Scheme 1. Proposed fragmentation pathways for deprotonated (a) terpenylic acid, (b) *cis*-pinic acid, and the (c) major and (d) minor MW 358 compounds.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Terpenylic acid and related compounds

F. Yasmeen et al.



Scheme 2. Proposed formation mechanism for the MW 358 and 344 compounds related to terpenylic acid and diaterpenylic acid acetate.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

