

## Corrigendum to

# “The kinetics and mechanism of an aqueous phase isoprene reaction with hydroxyl radical” published in Atmos. Chem. Phys., 11, 7399–7415, 2011

D. Huang, X. Zhang, Z. M. Chen, Y. Zhao, and X. L. Shen

State Key Laboratory of Environmental Simulation and Pollution Control, College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

The following errors need to be corrected.

In section “Explaining the mechanism of MG and GL formation” (p. 7408), “ $\text{CH}_2(\text{OH})\text{C}(\text{OO}\times)(\text{CH}_3)\text{CHO}$ ” should be “ $\text{CH}_2(\text{OH})\text{C}(\text{OO}\cdot)(\text{CH}_3)\text{CHO}$ ”.

Sect. 4 (p. 7410), “ $Y_{\text{MVK}} (18.9 \pm 0.8\%)$  and  $Y_{\text{MACR}} (9.0 \pm 1.1\%)$ ” should be “ $Y_{\text{MVK}} (24.1 \pm 0.8\%)$  and  $Y_{\text{MACR}} (10.9 \pm 1.1\%)$ ”.

In Table 1, the value of  $R_{\text{sur}}$  for scenario 5, “ $1.3 \times 10^{-17}$ ” should be “ $5.3 \times 10^{-17}$ ”.

On p. 7411, the equation “ $R_{\text{sur}} = k_{\text{sur}} \{\text{ISO}\}_v \{\text{OH}\}_v A_s$ ” should be “ $R_{\text{sur}} = k_{\text{sur}} \{\text{ISO}\}_v \{\text{OH}\}_v l A_s$ ”.

The authors wish to apologize for these errors.



Correspondence to: Z. M. Chen  
(zmchen@pku.edu.cn)