

Supplementary material to

Effect of the summer monsoon on aerosols at two measurement stations in Northern India part 2:
physical and optical properties

Hyvärinen, A.-P.¹, Raatikainen, T.^{1*}, Komppula, M.², Mielonen, T.², Sundström, A.-M.³, Brus,
D.^{1,4}, Panwar, T. S.⁵, Hooda, R. K.⁵, Sharma, V. P.⁵, de Leeuw, G.^{1,3}, and Lihavainen, H.¹

[1] Finnish Meteorological Institute, Erik Palménin aukio 1, P.O. Box 503, FI-00101, Helsinki,
Finland

[2] Finnish Meteorological Institute, Yliopistonranta 1F, P.O. Box 1627, FI-70211 Kuopio, Finland.

[3] Department of Physics, University of Helsinki, P.O. Box 64, FIN-00014 Helsinki, Finland

[4] Laboratory of Aerosol Chemistry and Physics, Institute of Chemical Process Fundamentals
Academy of Sciences of the Czech Republic, Rozvojová 135, CZ-165 02 Prague 6, Czech Republic

[5] The Energy and Resources Institute (TERI), Darbari Seth Block, IHC Complex, Lodhi Road,
New Delhi 110 003, India

[*] Currently at: School of Earth & Atmospheric Sciences, Georgia Institute of Technology, 311
Ferst Drive, Atlanta, GA 30332-0340, USA

Correspondence to: A.-P. Hyvärinen (antti.hyvarinen@fmi.fi)




Figure S1. Single scattering albedo, SSA as a function of total particle count, N_{tot} in Mukteshwar.