



## Corrigendum to

# “The HNO<sub>3</sub> forming branch of the HO<sub>2</sub> + NO reaction: pre-industrial-to-present trends in atmospheric species and radiative forcings” published in Atmos. Chem. Phys., 11, 8929–8943, 2011

O. A. Søvde<sup>1,3</sup>, C. R. Hoyle<sup>2,3,\*</sup>, G. Myhre<sup>1</sup>, and I. S. A. Isaksen<sup>1,3</sup>

<sup>1</sup>Center for International Climate and Environmental Research – Oslo (CICERO), Norway

<sup>2</sup>Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

<sup>3</sup>Department of Geosciences, University of Oslo, Oslo, Norway

\* now at: Laboratory of Atmospheric Chemistry, Paul Scherrer Institut, Villigen, Switzerland

Correspondence to: O. A. Søvde (asovde@cicero.uio.no)

Received: 4 May 2011 – Published in Atmos. Chem. Phys. Discuss.: 16 May 2011

Revised: 16 August 2011 – Accepted: 26 August 2011 – Published: 27 August 2012

In Table 6 of the paper “The HNO<sub>3</sub> forming branch of the HO<sub>2</sub> + NO reaction: pre-industrial-to-present trends in atmospheric species and radiative forcings” some of the entries should be positive, as in the ACPD version. The manuscript text, however, uses the correct signs. The corrected table is given on the right side.

**Table 6.** The different components of the radiative forcing due to the change in O<sub>3</sub> since the pre-industrial times.

RF due to tropospheric O <sub>3</sub> change		
Component	REF	R2
Tropospheric O <sub>3</sub> precursors	0.50	0.44
Cl & Br	−0.05	−0.06
Total RF, trop.	0.45	0.38
RF due to stratospheric O <sub>3</sub> change		
Component	REF	R2
Tropospheric O <sub>3</sub> precursors	0.09	0.08
Cl & Br	−0.21	−0.20
Total RF, strat.	−0.12	−0.12
RF due to tropospheric & stratospheric O <sub>3</sub> change		
Component	REF	R2
Tropospheric O <sub>3</sub> precursors	0.59	0.52
Cl & Br	−0.26	−0.26
Total RF, trop. & strat.	0.33	0.26