



Ozone photochemistry in boreal biomass burning plumes

M. Parrington¹, P. I. Palmer¹, A. C. Lewis², J. D. Lee², A. R. Rickard^{3,4}, P. Di Carlo^{5,6}, J. W. Taylor⁷, J. R. Hopkins², S. Punjabi⁸, D. E. Oram⁹, G. Forster⁹, E. Aruffo^{5,6}, S. J. Moller², S. J.-B. Bauguitte¹⁰, J. D. Allan^{7,11}, H. Coe⁷, and R. J. Leigh¹²

¹School of GeoSciences, The University of Edinburgh, Edinburgh, UK

²National Centre for Atmospheric Science, Department of Chemistry, University of York, York, UK

³National Centre for Atmospheric Science, School of Chemistry, University of Leeds, Leeds, UK

⁴now at the National Centre for Atmospheric Science, Department of Chemistry, University of York, York, UK

⁵Center of Excellence CETEMPS, University of L'Aquila, L'Aquila, Italy

⁶Department of Physical and Chemical Sciences, University of L'Aquila, L'Aquila, Italy

⁷School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Manchester, UK

⁸Department of Chemistry, University of York, York, UK

⁹National Centre for Atmospheric Science, School of Environmental Sciences, University of East Anglia, Norwich, UK

¹⁰Facility for Airborne Atmospheric Measurements, Bedford, UK

¹¹National Centre for Atmospheric Science, University of Manchester, Manchester, UK

¹²Department of Physics and Astronomy, University of Leicester, Leicester, UK

Correspondence to: M. Parrington (mark.parrington@ed.ac.uk)

Received: 26 December 2012 – Published in Atmos. Chem. Phys. Discuss.: 17 January 2013

Revised: 24 June 2013 – Accepted: 25 June 2013 – Published: 1 August 2013

Abstract. We present an analysis of ozone (O₃) photochemistry observed by aircraft measurements of boreal biomass burning plumes over eastern Canada in the summer of 2011. Measurements of O₃ and a number of key chemical species associated with O₃ photochemistry, including non-methane hydrocarbons (NMHCs), nitrogen oxides (NO_x) and total nitrogen containing species (NO_y), were made from the UK FAAM BAe-146 research aircraft as part of the “quantifying the impact of BOREal forest fires on Tropospheric oxidants over the Atlantic using Aircraft and Satellites” (BORTAS) experiment between 12 July and 3 August 2011. The location and timing of the aircraft measurements put BORTAS into a unique position to sample biomass burning plumes from the same source region in Northwestern Ontario with a range of ages. We found that O₃ mixing ratios measured in biomass burning plumes were indistinguishable from non-plume measurements, but evaluating them in relationship to measurements of carbon monoxide (CO), total alkyl nitrates (ΣAN) and the surrogate species NO_z (= NO_y – NO_x) revealed that the potential for O₃ production increased with plume age. We used NMHC ratios to estimate photochemi-

cal ages of the observed biomass burning plumes between 0 and 10 days. The BORTAS measurements provided a wide dynamic range of O₃ production in the sampled biomass burning plumes with ΔO₃/ΔCO enhancement ratios increasing from 0.020 ± 0.008 ppbv ppbv⁻¹ in plumes with photochemical ages less than 2 days to 0.55 ± 0.29 ppbv ppbv⁻¹ in plumes with photochemical ages greater than 5 days. We found that the main contributing factor to the variability in the ΔO₃/ΔCO enhancement ratio was ΔCO in plumes with photochemical ages less than 4 days, and that was a transition to ΔO₃ becoming the main contributing factor in plumes with ages greater than 4 days. In comparing O₃ mixing ratios with components of the NO_y budget, we observed that plumes with ages between 2 and 4 days were characterised by high aerosol loading, relative humidity greater than 40 %, and low ozone production efficiency (OPE) of 7.7 ± 3.5 ppbv ppbv⁻¹ relative to ΣAN and 1.6 ± 0.9 ppbv ppbv⁻¹ relative to NO_z. In plumes with ages greater than 4 days, OPE increased to 472 ± 28 ppbv ppbv⁻¹ relative to ΣAN and 155 ± 5 ppbv ppbv⁻¹ relative to NO_z. From the BORTAS measurements we estimated that aged

plumes with low aerosol loading were close to being in photostationary steady state and O_3 production in younger plumes was inhibited by high aerosol loading and greater production of ΣAN relative to O_3 . The BORTAS measurements of O_3 photochemistry in boreal biomass burning plumes were found to be consistent with previous summertime aircraft measurements made over the same region during the Arctic Research of the Composition of the Troposphere (ARCTAS-B) in 2008 and Atmospheric Boundary Layer Experiment (ABLE 3B) in 1990.

1 Introduction

The quantifying the impact of BOREal forest fires on Tropospheric oxidants over the Atlantic using Aircraft and Satellites (BORTAS) experiment was conducted in two phases: July–August 2010 (phase A) and 2011 (phase B) (Palmer et al., 2013). The overarching objective of BORTAS was to better understand the chemical evolution of plumes emitted from wildfires in boreal regions, with a particular emphasis on the net production of tropospheric O_3 and downwind impacts on air quality. The first phase of the experiment (BORTAS-A) comprised of ground-based, ozonesonde, and satellite measurements over eastern Canada in the summer of 2010, the results from which were reported by Parrington et al. (2012). The second phase of the experiment (BORTAS-B) was an aircraft measurement campaign based out of Halifax, Nova Scotia, Canada, between 12 July and 3 August 2011, supported by ground-based, ozonesonde, and satellite measurements. An overview of the BORTAS-B campaign is given by Palmer et al. (2013). In this paper we analyse O_3 photochemistry in the biomass burning plumes sampled by the aircraft during BORTAS-B.

Tropospheric O_3 is a secondary pollutant formed by the photo-oxidation of carbon monoxide (CO) and volatile organic compounds (VOCs) in the presence of nitrogen oxides (NO_x) and transport from the stratosphere. Removal of O_3 from the troposphere is through dry deposition and photochemical destruction. Tropospheric O_3 plays an important role in the contexts of climate and air quality. Emissions of O_3 precursors (i.e. NO_x and VOCs) can be anthropogenic in origin and can also have natural sources such as biogenic processes and biomass burning. A number of previous measurement campaigns have made in situ observations of photochemistry within boreal biomass burning outflow and evaluated the associated O_3 production and loss. The NASA Atmospheric Boundary Layer Experiment (ABLE) included two high-latitude aircraft measurement campaigns in the summers of 1988 (ABLE 3A) and 1990 (ABLE 3B) (Harris et al., 1992, 1994). Analysis of the ABLE 3A measurements, made over northern and western Alaska, found O_3 production in haze layers associated with fires to be inefficient with less than 0.1 molecules of O_3 formed per molecule

of CO (Wofsy et al., 1992). However, the ABLE 3B measurements, made over central and eastern Canada, observed O_3 enhancements of 5–15 ppbv in aged (> 2 days) biomass burning plumes (Anderson et al., 1994). Aircraft measurements made over the North Atlantic Ocean during the Intercontinental Transport of Ozone and Precursors (ITOP) campaign (Lewis et al., 2007), which was part of the wider International Consortium for Atmospheric Research on Transport and Transformation (ICARTT) (Fehsenfeld et al., 2006) in the summer of 2004, sampled biomass burning plumes associated with fires in Alaska. Examples of elevated O_3 were observed in individual plumes with elevated CO during ITOP but there was no single canonical relationship (Lewis et al., 2007). More recently the NASA Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) campaign in the spring and summer of 2008 made measurements of biomass burning plumes in central Canada. Analysis of the ARCTAS aircraft measurements showed little evidence for O_3 formation within the plumes (Alvarado et al., 2010) other than when the plumes were mixed with outflow from urban pollution (Singh et al., 2010). Ground-based measurements of outflow from boreal biomass burning have also been routinely made at the Pico Mountain Observatory in the mid-Atlantic (Lapina et al., 2006; Val Martin et al., 2006; Helmig et al., 2008). Years with high boreal fire activity have been shown to increase O_3 mixing ratios in the free troposphere over this site by up to 10 ppbv (Lapina et al., 2006) with photochemically aged biomass burning plumes influencing tropospheric oxidant chemistry after 1–2 weeks of transport to the region (Helmig et al., 2008).

Model analyses of photochemistry in boreal biomass burning plumes have also been performed to evaluate O_3 production and loss. Mauzerall et al. (1996) performed 1-D model simulations to calculate in situ production and loss of O_3 based on the ABLE 3B measurements in the summer of 1990. They found biomass burning emissions to be a relatively negligible, contributing less than 2 % to the calculated O_3 budget. McKeen et al. (2001) used a 3-D chemistry transport model (CTM) to evaluate the influence of Canadian biomass burning emissions on O_3 production over north-eastern USA using measurements made from the National Oceanographic and Atmospheric Administration (NOAA) WP-3 aircraft in the summer of 1995. O_3 enhancements of between 10 and 30 ppbv throughout the central and eastern US were attributed to the fires, with the model simulations being sensitive to NO_x/CO and VOC/CO emission ratios. The influence of emissions from forest fires in Alaska and Canada on tropospheric O_3 production during summer 2004 was the subject of model studies by Pfister et al. (2006); Real et al. (2007); Cook et al. (2007). Pfister et al. (2006) utilised the MOZART-4 CTM to evaluate O_3 production in the outflow from fires in Alaska during summer of 2004 with measurements made from aircraft during the ICARTT campaign and at the Pico Mountain Observatory. They reported O_3

enhancements relative to CO of 0.25 ppbv ppbv⁻¹, contributing approximately 3% of the Northern Hemispheric tropospheric O₃ budget. Lagrangian model studies of the outflow over the North Atlantic Ocean (Real et al., 2007) reported that although an O₃ enhancement of 17 ppbv was observed after 5 days, net O₃ loss could occur in the plumes due to biomass burning aerosols in the plume reducing the photolysis rates of O₃ and NO₂.

An overview of observed and model estimates of O₃ production in boreal biomass burning plumes was recently presented by Jaffe and Wigder (2012). A common feature of observation and model studies of O₃ production in biomass burning outflow is the use of O₃ enhancement ratios relative to CO ($\Delta O_3/\Delta CO$) to determine O₃ production. Jaffe and Wigder (2012) broadly showed that the $\Delta O_3/\Delta CO$ ratio increases with plume age. In this paper we diagnose O₃ production in biomass burning plumes using in situ aircraft measurements made over eastern Canada in the summer of 2011. We evaluate a number of different metrics, including $\Delta O_3/\Delta CO$ and O₃ production efficiency relative to the NO_y budget. An overview of the aircraft measurements and chemical processes influencing O₃ photochemistry are presented in Sect. 2. Section 3 presents the O₃ distribution over eastern Canada observed by the aircraft over the campaign period. Section 4 presents an analysis of observed plume photochemistry using photochemical age calculations derived from NMHC ratios, O₃ production efficiencies relative to NO_y partitioning, and photostationary steady state calculations. We conclude in Sect. 5.

2 Airborne photochemical measurements during BORTAS-B

In this section we present a brief overview of tropospheric O₃ photochemistry and the relevant measurements made from the BAe-146 aircraft during the BORTAS-B aircraft measurement campaign, which are used in our analysis.

2.1 Plume photochemistry

The simplest model of O₃ photochemistry in the atmosphere can be represented by the rapid oxidation of nitrogen oxides, NO_x, which represents the sum of nitric oxide (NO) and nitrogen dioxide (NO₂), in the presence of sunlight through the following null cycle:



In polluted air masses such as biomass burning plumes, perturbations of this cycle through additional reactions of NO with peroxy radicals (RO₂ or HO₂) produced by the oxidation of VOCs provide additional pathways for converting NO_x into O₃. Figure 1 shows a schematic of the simplified




Fig. 1. Schematic of the atmospheric O₃-NO_x-hydrocarbon chemistry cycle studied by BORTAS.

O₃-NO_x-VOC chemistry in the atmosphere. The chain of reactions leading to O₃ formation is initiated by the photo-oxidation of a VOC (RH) and its rapid subsequent reaction with molecular oxygen to produce alkyl peroxy radicals (RO₂), for example



Under tropospheric conditions, as indicated by the blue arrows in Fig. 1, alkyl peroxy radicals can react with NO₂ via a three body reaction to form peroxy nitrates (RO₂NO₂):



which will decompose back into the constituent RO₂ and NO₂ under the right environmental conditions (i.e. near surface temperatures) but can also be subject to transport away from the source region as they are thermally stable at lower temperatures, such as those in the free troposphere.

Peroxy radicals are also subject to reaction with NO and can follow two different pathways:



Reaction (R6) results in the formation of O₃ molecules via photolysis of NO₂. Comparison of reactions (R6) and (R7) indicates that O₃ production in biomass burning plumes initiated by photo-oxidation of VOCs is in competition with production of alkyl nitrates (RONO₂); evaluating the relationship between the measured O₃ and RONO₂ mixing ratios provides an indication of the potential for O₃ production related to the above mechanism (e.g. Perring et al., 2010).

The potential of the above reactions to produce O₃ in biomass burning plumes is dependent on the availability of nitrogen oxides, NO_x (=NO+NO₂). A surrogate species, NO_y, is used to represent the sum of all reactive and oxidised nitrogen species in an air mass: NO_y=NO_x+ΣRO₂NO₂+ΣRONO₂+HNO₃+HONO+NO₃+2N₂O₅+NO₃⁻, where ΣRO₂NO₂ represents the total peroxy nitrates (ΣPN) and ΣRONO₂ represents the total alkyl nitrates (ΣAN) (e.g. Day et al., 2003).

A second surrogate species, NO_z ($=\text{NO}_y - \text{NO}_x$), can be defined that represents the sum of the photo-oxidation products of NO_x (Olszyna et al., 1994). Since NO_y is representative of all NO_x emissions, NO_z provides a measure of the number of NO_x molecules that have undergone photo-oxidation. Several studies have shown that the O_3 abundance in polluted air masses is linearly related to the abundance of NO_z (e.g., Trainer et al., 1993; Kleinman, 2000). The enhancement of O_3 abundance relative to NO_z (i.e. $\Delta\text{O}_3/\Delta\text{NO}_z$) provides a measure of O_3 production related to photo-oxidation of NO_x molecules, sometimes referred to as the ozone production efficiency (OPE) (Ryerson et al., 1998; Rickard et al., 2002).

2.2 BORTAS measurements

Measurements of the key tracers relevant to the O_3 photochemistry outlined above were made from the BAe-146 aircraft over the course of the BORTAS-B measurement campaign. Full details of the BAe-146 payload are presented in Table 2 of Palmer et al. (2013) and we provide here brief details of the measurements used in this study.

Measurements of O_3 were made using a TECO 49 ultraviolet absorption instrument (Wilson and Birks, 2006) operated by the UK Facility for Airborne Atmospheric Measurements (FAAM) with an average time of 3 Hz, at a precision of 1 ppbv and accuracy of $\pm 5\%$. Measurements used for identification of biomass burning plume air masses included acetonitrile (CH_3CN) and CO. A proton transfer reaction mass spectrometer (PTR-MS) (Murphy et al., 2010) provided measurements of CH_3CN with an average time of 1 s and mean precision of 37 pptv over all BORTAS flights. CO was measured using a vacuum-UV resonance fluorescence instrument (Gerbig et al., 1999) with an average time of 1 s, precision of 1 ppbv, and accuracy of 3%. We determined the aerosol loading within the sampled plume air with measurements of refractory black carbon using a single particle soot photometer (SP2) and measurements of total scattering coefficient at 550 nm using a TSI Inc. 3563 three-wavelength nephelometer (Anderson and Ogren, 1998). The SP2 uses laser-induced incandescence to measure refractory black carbon on a single particle basis (Schwarz et al., 2006); the inlet and instrumental setup on the BAe-146 are described by McMeeking et al. (2010). The statistical uncertainty in BC mass concentration was 5% for an average time of 15 s, and accuracy was within 20% due to possible differences between the instrumental response to biomass burning BC and the calibrant (Laborde et al., 2012). Measurement of NO_2 photolysis rates were made using upper and lower fixed bandwidth radiometers at wavelengths predominantly between 208 and 500 nm, at a frequency of 1 Hz and with accuracy of 10% (Volz-Thomas et al., 1996). Measurements of VOCs used in this study were made using a dual channel gas chromatograph system with flame ionisation detection (GC-FID) (Hopkins et al., 2003, 2011) to analyse

whole air canister samples taken during each flight. Each canister sample was taken over a 30 s period at points of particular interest along the flight track. The precision and accuracy of the GC-FID measurements are species dependent but typical values are 1–5 pptv and 5%, respectively. Measurements of NO and NO_2 were made using a single channel chemiluminescence instrument manufactured by Air Quality Design (AQD) Inc., USA, as described by Lee et al. (2009). The accuracy of the AQD NO and NO_2 measurements was 10 pptv. Measurements of NO_2 , total NO_y , total peroxyacetyl nitrates ($\Sigma\text{PN} = \Sigma\text{RO}_2\text{NO}_2$), and total alkyl nitrates ($\Sigma\text{AN} = \Sigma\text{RONO}_2$) were made using a Thermal Dissociation-Laser Induced Fluorescence (TD-LIF) instrument (Dari-Salisburgo et al., 2009; Di Carlo et al., 2013) with accuracies of 10%, 22%, 34%, and 46%, respectively.

For the analysis presented here we calculated mixing ratios of NO_x , and subsequently NO_z , from the aircraft measurements of NO, NO_2 and NO_y described above. The NO_x mixing ratio was calculated using NO measured by the AQD instrument and NO_2 measured by the TD-LIF instrument. The AQD instrument measures NO_2 indirectly, after it has undergone photo dissociation to form NO in the presence of UV radiation (Lee et al., 2009), whereas the TD-LIF instrument measures NO_2 directly; we use the TD-LIF in the NO_x calculation to reduce uncertainties and possible interferences. NO_z was then calculated by subtracting the AQD/LIF NO_x values from the LIF total NO_y measurements. We calculated average uncertainties of $13 \pm 6\%$ in the range 6–45% for NO_x and $51 \pm 4\%$ in the range 47–84% for NO_z . For both NO_x and NO_z , calculated uncertainties more than 2σ standard deviations above the average occurred for approximately 5% of the plume measurements with the lowest calculated NO_x and NO_z mixing ratios.

To distinguish measurements made in biomass burning plumes from those made in clean air, we use the threshold of 150 pptv of CH_3CN determined by Palmer et al. (2013) from the 99 percentile of measurements made on flight B625. CH_3CN is predominantly emitted from biomass burning and removed from the atmosphere through uptake by the oceans and reaction with OH, with an atmospheric lifetime of approximately 25 days (Bange and Williams, 2000); therefore, CH_3CN is an excellent tracer for identifying plume measurements.

3 Observed O_3 distribution over eastern Canada

In this section we present an overview of the O_3 distribution observed from the BAe-146 aircraft during the BORTAS-B measurement campaign. Figure 2 shows the relative frequency distribution of O_3 mixing ratios, divided into 1 ppbv bins, measured from the BAe-146 during the eleven research flights undertaken 15–31 July 2011 (Table 1). The frequency distribution of the O_3 measurements made both in and out of biomass burning plumes shows two peaks at approximately




Fig. 2. Frequency distribution of O_3 mixing ratios measured from the BAe-146 on all research flights during the BORTAS campaign period (15 to 31 July 2011). Frequencies are calculated for 1 ppbv bins. The black line shows the distribution of all the measured O_3 mixing ratios, the red line shows the distribution of measurements identified as biomass burning plumes using a threshold of CH_3CN mixing ratios greater than 150 pptv, and the blue line shows measurements made in air masses with CH_3CN mixing ratios less than 150 pptv. The number of data points, n , for each distribution are shown in the legend.

25 and 50 ppbv. The peak in O_3 mixing ratio at 25 ppbv is attributable to measurements made at altitudes below 3 km. The O_3 measurements made during BORTAS-B show a similar distribution to that observed over central and eastern Canada during the ABLE 3B measurement campaign in the summer of 1990 (Anderson et al., 1994).

The statistical distribution of O_3 mixing ratios measured on each of the BORTAS campaign flights (Table 1) is shown in Fig. 3a. Each flight shows a wide range of measured O_3 mixing ratios between 20 and 125 ppbv, reflecting the variability in O_3 over the geographical area and altitude ranges covered by the aircraft over the course of the campaign. The mean O_3 mixing ratio measured over the whole campaign was 50 ppbv, which was reduced to 46 ppbv for measurements made in non-plume ($CH_3CN < 150$ pptv) air masses and increased to 57 ppbv for plume air measurements. In general, O_3 mixing ratios were less than the campaign mean of the plume measurements. Figure 3b shows the statistical distribution of the measured O_3 mixing ratios as a function of the aircraft GPS altitude, divided into 1 km wide bins. Using the medians for each altitude bin, the O_3 mixing ratio increased from less than 40 ppbv between 0.5 and 1.5 km to approximately 60 ppbv between 7.5 and 8.5 km. Plume measurements were higher on average than non-plume measurements, apart from the altitude range between 4.5 and 5.5 km, and measurements were typically less than the campaign mean of the plume data below 5.5 km. Three research flights (B621, B629 and B630) measured O_3 mixing ratios that were




Fig. 3. Box and whisker plots showing the statistical distribution of O_3 measurements made from the BAe-146 during BORTAS as a function of (a) research flight (see Table 1 and Palmer et al., 2013), and (b) aircraft GPS altitude. In both plots solid boxes show the range of the 25th to 75th percentiles and whiskers show the range of the 5th to 95th percentiles with outliers shown as plus symbols. Black boxes and whiskers show all measured values, red boxes and whiskers show plume measurements determined from measured CH_3CN mixing ratios in excess of 150 pptv, and blue boxes and whiskers show non-plume measurements (i.e. CH_3CN less than 150 pptv). Dashed lines in each panel indicate the campaign mean O_3 mixing ratio for each group of data points.

on average greater than the campaign mean. These measurements were typically made at higher altitudes, above 5.5 km, as reflected in Fig. 3b. A number of outlying O_3 mixing ratios were made below the 5th percentile and above the 95th percentile of the observed statistics on each flight, reflecting the wide range of altitudes over which the measurements were made on each flight.

4 Observed boreal biomass burning outflow

In this section we evaluate the processes influencing the O_3 distribution in biomass burning plume air sampled by the BAe-146 over the course of the BORTAS campaign. Figure 4 shows the relative distributions of 60 s averaged O_3 and CO mixing ratios measured from the aircraft. Figure 2 shows that there is no discernible difference between O_3 measurements made in the plume air and those in clean air. CO mixing ratios in plumes range from 64 to 930 ppbv. For CO mixing

Table 1. Flight IDs, dates, and location of research flights made by the BAe-146 during the BORTAS aircraft measurement campaign in 2011. The relative locations of each flight are shown in Palmer et al. (2013).

Flight ID	Date (Day of Year)	Location
B620	15 July (196)	Eastern Quebec
B621	18 July (199)	Newfoundland and Labrador
B622	20 July (201)	Southern Quebec
B623	20 July (201)	Quebec City to Halifax
B624	21 July (202)	Atlantic Ocean to east of Island of Newfoundland
B625	24 July (202)	Gulf of St. Lawrence
B626	26 July (207)	Northwestern Ontario
B627	27 July (208)	Thunder Bay ON to Goose Bay NL
B628	28 July (209)	Labrador coast
B629	31 July (212)	Gulf of St. Lawrence
B630	31 July (212)	Nova Scotia and Island of Newfoundland




Fig. 4. Scatter plot of 60 s averaged O_3 versus CO mixing ratios in ppbv measured from the BAe-146 during the BORTAS campaign period. Black symbols show all measured data and red symbols show the data points identified as biomass burning outflow. The number of data points, n , are shown in the legend.

ratios in excess of 300 ppbv, O_3 mixing ratios lie within a range between approximately 40 and 60 ppbv. In the following subsections we examine O_3 production and photochemistry measured in boreal biomass burning plumes.

4.1 Photochemical ageing of biomass burning plumes

We first investigate photochemical processing within biomass burning plumes sampled by the aircraft through calculation of a photochemical age, i.e. the time taken for a particular measured tracer to be removed chemically from an air mass after emission. We calculate photochemical ages based on ratios of non-methane hydrocarbons (NMHCs) (Parrish et al., 2007). The NMHCs used in this calculation are ethane (C_2H_6), propane (C_3H_8) and *n*-butane (C_4H_{10}) measured by GC-FID from whole air canister samples taken over the course of the BORTAS campaign. Figure 5 shows




Fig. 5. Scatter plot of the natural logarithms of the ratios of $[C_4H_{10}]:[C_2H_6]$ and $[C_3H_8]:[C_2H_6]$ measured from whole air canister samples taken by the BAe-146 during the BORTAS campaign period. All measured data are shown as black diamonds with red diamonds showing measurements made in boreal biomass burning plumes. The green square shows the ratios of the initial concentrations of the three alkanes used in the plot, estimated from BORTAS flight B626 over the burning source region in Northwestern Ontario. The solid red bounding lines represent the behaviour of the hydrocarbon ratios: the steeper slope assumes only oxidation of the alkanes with OH (kinetic slope); the shallower slope assumes only mixing of fresh emissions with background air (mixing slope). The dashed red line shows a linear fit to the plume measurements.

the relationship between natural logarithms of the observed $[C_4H_{10}]:[C_2H_6]$ and $[C_3H_8]:[C_2H_6]$ ratios, where $[X]$ represents the concentration of NMHC X. The ratio of initial mixing ratios for each NMHC pair are determined from flight B626 over Northwestern Ontario on 26 July, which was the main source of biomass burning early in the campaign with fires burning 17–19 July. In the absence of detailed combustion information of these fires, we assume that

the higher values of the natural logarithm of the NMHC ratios from flight B626 are representative of the ratios of the freshest emissions of these NMHCs, and are shown as the solid green square in Fig. 5. We also assume that this ratio is representative of the initial mixing ratios of these NMHCs emitted from forest fire emissions across boreal North America, based on the similarity of vegetation cover over the region (Latifovic et al., 2004). The majority of NMHC ratios measured in biomass burning plumes during BORTAS lie to the left and below this point, indicating that the estimated initial concentrations are a suitable reference point for calculating the relative photochemical ageing of the biomass burning plumes sampled during the campaign. Following Parrish et al. (2007), we show bounding limits for oxidation by OH and dilution or mixing of fresh emissions into aged air masses as solid lines in Fig. 5. The steeper of these lines represents ageing of plume air if oxidation of the NMHCs was the only process influencing their concentration. This is sometimes referred to as the “kinetic” line (Parrish et al., 2007; Helmig et al., 2008), and in our study its slope is calculated from the kinetic reaction rate coefficients of the measured NMHCs with OH. Assuming a typical free tropospheric temperature of 273 K and taking reaction rate coefficients from Atkinson and Arey (2003), we calculate a kinetic slope of 2.61. The less steep of the solid lines in Fig. 5 can be considered a “dilution” line and represents dilution of a plume, starting with the same initial mixing ratios through mixing with background air that is sufficiently aged so that only the least reactive NMHC (i.e. C_2H_6) remains at significant concentrations. The mixing causes the ratio of the NMHC ratios to decrease with a slope of unity (Parrish et al., 2007). In general, NMHC ratios calculated from the BORTAS measurements lie between -3 and -1 for $\ln([C_3H_8]/[C_2H_6])$ and between -6 and -2.5 for $\ln([C_4H_{10}]/[C_2H_6])$ and are consistent with previous observations of NMHC ratios (Rudolph and Johnen, 1990; Parrish et al., 1992; Jobson et al., 1994; Parrish et al., 2007; Helmig et al., 2008). These data points lie within the bounds of the dilution and kinetic slopes, indicating that NMHC concentrations in the measured air masses were subject to a combination of oxidation and mixing between fresh emissions and background air. This is also consistent with the previous studies listed above and is not unexpected given that the BORTAS measurements were made between 1000–3000 km downwind of the emission sources. The gradient of the line of best fit to the data is 1.55, further supporting the assumption that the observations were subject to mixing and photochemical processing. Outliers in Fig. 5, i.e. where $\ln([C_3H_8]/[C_2H_6]) > -1$ and $\ln([C_4H_{10}]/[C_2H_6]) > -2$, show measurements representative of different emission sources and are beyond the scope of the work presented here.

We estimate plume ages in days for the plume air masses sampled during the BORTAS campaign using the $\ln([C_3H_8]/[C_2H_6])$ ratio and assuming a constant average

OH concentration of 2×10^6 molecules cm^{-3} . As no OH measurements were made from the BAe-146 during the campaign, this value was chosen to be representative of a northern mid-latitude summertime OH concentration (Spivakovsky et al., 2000). Varying the average OH concentration in the photochemical age calculations from half to double the assumed value would double or halve the age, respectively. The photochemical ages we calculate from the BORTAS measurements using this assumed OH concentration were similar to transport timescales from the emission source determined by back trajectory calculations (not presented here), providing confidence in our assumption. Estimating the photochemical plume age in this way allows us to compare O_3 production in boreal biomass burning plumes measured over the BORTAS campaign to similar measurements used in previous studies.

4.2 O_3 enhancements in biomass burning plumes

Figure 6 shows the relationships between O_3 and CO measured in biomass burning plumes during the BORTAS measurement campaign to the calculated photochemical ages, sampled at the times of the whole air canister samples. O_3 mixing ratios measured in boreal biomass burning plumes, Fig. 6a, show no clear relationship to the calculated photochemical age over the whole BORTAS campaign period. The O_3 measurements made on flight B626 are distinguished from those made on the other research flights by mixing ratios less than 30 ppbv. It should be noted that the measurements made in biomass burning plumes on flight B626 were made at lower altitudes (below 1.5 km) than on other flights. Measurements with higher CH_3CN mixing ratios typically show moderate O_3 mixing ratios between 40 and 70 ppbv with photochemical ages between 1.5 and 4.5 days. The measurements made with lower CH_3CN exhibit a wider range of O_3 mixing ratios, from 40 to 110 ppbv, and photochemical ages, from 0 to 10 days. The CO mixing ratios measured in boreal biomass burning plumes, Fig. 6b, typically show higher values (between 200 and 1100 ppbv) at photochemical ages of between 2 and 4 days and are coincident with the higher values of CH_3CN . These measurements were made at aircraft altitudes below 3 km during flights B622–B624 on 20 and 21 July in plumes from the fires in Northwestern Ontario with approximate transport times of 2–3 days, calculated by Lagrangian backward trajectories (not shown). Measurements of CO mixing ratios less than 200 ppbv have a wider range of photochemical ages, between 0 and 10 days. On flight B626, CO mixing ratios were measured between 150 and 400 ppbv with photochemical ages from 0 to 8.5 days.

The relationship between the measured O_3 and CO mixing ratios in boreal biomass burning plumes is shown in Fig. 6c for the data sampled at the times of the whole air canister samples. The distribution is similar to that shown by the red symbols in Fig. 4 for the 60 s averaged




Fig. 6. Relationship between O₃ and CO mixing ratios measured from the BAe-146 and sampled at the WAS measurement times, and photochemical ages calculated from the $\ln(\text{C}_3\text{H}_8/\text{C}_2\text{H}_6)$ ratio in boreal biomass burning plumes throughout the BORTAS-B measurement campaign and assuming an OH concentration of 2×10^6 molecules cm^{-3} . (a) and (b) show the measured O₃ and CO mixing ratios, respectively, in units of ppbv as a function of the calculated photochemical age in days. (c) shows the scatter of O₃ versus CO with straight lines showing linear fits to each population of data points. (d) shows the enhancement of O₃ relative to CO ($\Delta\text{O}_3/\Delta\text{CO}$) as a function of photochemical age. In all plots the data are filtered according to their relative abundance of CH₃CN, with black diamonds representing CH₃CN mixing ratios between 150 and 300 pptv and red diamonds representing CH₃CN mixing ratios in excess of 300 pptv. Measurements made on flight B626 over the fire region in Northwestern Ontario are treated separately and are represented by green circles.


data. The three populations of data points defined for flight B626, and moderate and high CH₃CN, show clear distinctions from one another. Straight lines fitted to each population and their slopes provide an indication of O₃ produced since the fire emissions of CO and other precursors at the source, i.e. $\Delta\text{O}_3/\Delta\text{CO}$ (Parrish et al., 1993; Pfister et al., 2006). Measurements of fresh biomass burning air masses were made during flight B626, indicated by a slope of 0.009 ± 0.001 ppbv ppbv⁻¹. Of the measurements made on the other BORTAS research flights, those with the highest CH₃CN mixing ratios were made in moderately fresh plume air masses with photochemical ages of 2–4 days. The gradient of the straight line fitted to these data points is 0.016 ± 0.001 ppbv ppbv⁻¹, suggesting a slight increase in O₃ production downwind of the emission source. Measurements with moderate CH₃CN are typically more aged (photochemical ages > 4 days) and indicate increased O₃ production relative to the other two populations, as reflected in the gradient of the straight line fitted to these data points of

1.292 ± 0.007 ppbv ppbv⁻¹. Figure 6d shows enhancement ratios of O₃ relative to CO as a function of photochemical age. Enhancements of O₃ and CO (i.e. ΔO_3 and ΔCO) are calculated relative to background values determined from the distribution of clean air measurements defined by CH₃CN mixing ratios less than 100 pptv and CO mixing ratios less than 100 ppbv. Average O₃ and CO mixing ratios in the clean air measurements were 42.0 ± 14.3 ppbv in the range 21.5–124.8 ppbv, and 87.8 ± 6.6 ppbv in the range 70.1–99.9 ppbv, respectively. We calculate background O₃ and CO mixing ratios, using the 25th percentile of the distribution of the clean air measurements, to be 25 and 81 ppbv, respectively.

The $\Delta\text{O}_3/\Delta\text{CO}$ ratios for the three populations of data show a similar trend to the gradients calculated for the O₃ to CO relationship in Fig. 6c, with the value of $\Delta\text{O}_3/\Delta\text{CO}$ increasing from the measurements made on flight B626 through the high CH₃CN measurements to the moderate CH₃CN measurements. The measurements made on flight B626 have $\Delta\text{O}_3/\Delta\text{CO}$ ratios between approximately

Table 2. Observed $\Delta\text{O}_3/\Delta\text{CO}$ in biomass burning plumes sampled during BORTAS filtered by photochemical age, following Jaffe and Wigder (2012).

Plume category	Range of $\Delta\text{O}_3/\Delta\text{CO}$ (ppbv ppbv ⁻¹) (no. of measurements)	Mean (ppbv ppbv ⁻¹)	Median (ppbv ppbv ⁻¹)	Standard deviation (ppbv ppbv ⁻¹)
≤ 2 days ¹	0.009–0.029 ($n = 6$)	0.020	0.027	0.009
2–5 days ²	0.02–0.69 ($n = 100$)	0.13	0.12	0.10
2–5 days ³	0.02–1.76 ($n = 134$)	0.27	0.15	0.31
≥ 5 days ³	0.24–2.55 ($n = 52$)	0.81	0.63	0.52
≥ 5 days ⁴	0.24–1.89 ($n = 50$)	0.75	0.63	0.41

¹ Flight B626;² $[\text{CH}_3\text{CN}] > 300$ pptv;³ $[\text{CH}_3\text{CN}] > 150$ pptv;⁴ $\Delta\text{O}_3/\Delta\text{CO} < 2$ ppbv ppbv⁻¹.**Fig. 7.** Scatter plot of \log_{10} of the difference between O_3 and CO enhancements in biomass burning plumes relative to background values (i.e. ΔO_3 and ΔCO) as a function of the $\Delta\text{O}_3/\Delta\text{CO}$ enhancement ratio. The data are filtered based on photochemical ages calculated from the $\ln(\text{C}_3\text{H}_8/\text{C}_2\text{H}_6)$ ratio. The horizontal and vertical dotted lines show the zero difference and unit ratio values, respectively.

0.015 and 0.05 ppbv ppbv⁻¹ over a range of photochemical ages between 0 and 8 days. The $\Delta\text{O}_3/\Delta\text{CO}$ ratio decreases from 0.03 to 0.01 ppbv ppbv⁻¹ between 0 and 2 days of photochemical ageing, with the lowest values associated with higher CO mixing ratios in excess of 250 ppbv, increasing from approximately 150 ppbv (Fig. 6b), and increases from 0.03 to 0.05 ppbv ppbv⁻¹ between 5 and 8 days of photochemical ageing. The high CH_3CN measurements show a wide range of $\Delta\text{O}_3/\Delta\text{CO}$ ratios, from 0.02 to 0.3 ppbv ppbv⁻¹, between 2 and 4 days of photochemical ageing. A few high CH_3CN measurements with photochemical ages between 4 and 8 days have higher $\Delta\text{O}_3/\Delta\text{CO}$ ratios from 0.2 to 0.5 ppbv ppbv⁻¹. In general, the moderate CH_3CN measurements have photochemical ages greater than 4 days and exhibit higher $\Delta\text{O}_3/\Delta\text{CO}$ ratios between

0.1 and 1.0 ppbv ppbv⁻¹. Some measurements with moderate CH_3CN have photochemical ages less than 4 days, although this does not appear to significantly affect the $\Delta\text{O}_3/\Delta\text{CO}$ ratio in these air masses. The measurements made with less than 2 days photochemical ageing and $\Delta\text{O}_3/\Delta\text{CO}$ ratios greater than 0.07 ppbv ppbv⁻¹ may be attributable to a fire source with different initial NMHC mixing ratios than those used in the photochemical age calculation or could be aged air masses that have experienced mixing with air of different origin which has influenced the NMHC abundance. Further analysis of these data points is beyond the scope of the analysis presented here.

Figure 7 shows that the main contributing factor to the variability in the $\Delta\text{O}_3/\Delta\text{CO}$ ratios observed in biomass burning plumes changes as a function of the plume photochemical age. As reflected in Fig. 4a and b, variability in ΔO_3 increases as a function of $\Delta\text{O}_3/\Delta\text{CO}$ while variability in ΔCO decreases to a point where the main contributing factor changes from ΔCO to ΔO_3 . For air masses with photochemical ages of less than 4 days, ΔO_3 is generally less than ΔCO (i.e. $\log_{10}(\Delta\text{O}_3 - \Delta\text{CO}) < 0$ and $\Delta\text{O}_3/\Delta\text{CO} < 1$) and the variability in $\Delta\text{O}_3/\Delta\text{CO}$ is controlled by ΔCO . At photochemical ages greater than 4 days, the variability in $\Delta\text{O}_3/\Delta\text{CO}$ is controlled by a combination of ΔCO , for $\Delta\text{O}_3/\Delta\text{CO}$ between 0.1 and 1.0, and ΔO_3 , for $\Delta\text{O}_3/\Delta\text{CO}$ greater than 1.0.

A review of O_3 production in the outflow from wildfires, including boreal regions, was recently presented by Jaffe and Wigder (2012). Table 2 presents statistics of $\Delta\text{O}_3/\Delta\text{CO}$ ratios in boreal biomass burning plumes sampled during BORTAS, arranged by photochemical age following Jaffe and Wigder (2012). The BORTAS results show that the $\Delta\text{O}_3/\Delta\text{CO}$ ratio increases with plume age in both the range and average (mean and median) values, indicating that O_3 production is enhanced in plumes that have undergone more photochemical processing as they are transported away from the source region. For plume ages less than 2 days, the BORTAS $\Delta\text{O}_3/\Delta\text{CO}$ ratios are within the range of ARCTAS measurements for the same category (Alvarado et al., 2010) and the mean BORTAS value of 0.020 ± 0.008 ppbv ppbv⁻¹

is comparable to the mean of $0.018 \text{ ppbv ppbv}^{-1}$ calculated by Jaffe and Wigder (2012) from studies by Alvarado et al. (2010), Goode et al. (2000), Singh et al. (2010) and DeBell et al. (2004). For plume ages between 2 and 5 days the range of BORTAS $\Delta\text{O}_3/\Delta\text{CO}$ ratios are comparable to the ranges determined for boreal biomass burning plumes from eastern Canada (Mauzerall et al., 1996) and Siberia (Tanimoto et al., 2008). The mean ratio of $0.13 \pm 0.10 \text{ ppbv ppbv}^{-1}$ for the high CH_3CN measurements is comparable to the average ratio of $0.15 \text{ ppbv ppbv}^{-1}$ calculated by Jaffe and Wigder (2012) for this plume category. For the moderate CH_3CN measurements the mean ratio increases to $0.27 \pm 0.31 \text{ ppbv ppbv}^{-1}$, which is also comparable to the mean, although with a much wider range of values (ratio reported by (Jaffe and Wigder, 2012)). For plume ages greater than 5 days, the BORTAS $\Delta\text{O}_3/\Delta\text{CO}$ ratios are typically within the range of values presented by Jaffe and Wigder (2012). The mean $\Delta\text{O}_3/\Delta\text{CO}$ ratio of $0.81 \pm 0.52 \text{ ppbv ppbv}^{-1}$ for BORTAS is higher than the mean values of approximately $0.25 \text{ ppbv ppbv}^{-1}$ for North American boreal biomass burning plumes (Pfister et al., 2006; Val Martin et al., 2006) and is more comparable to similarly aged plumes originating in Siberia (Bertschi and Jaffe, 2005; Honrath et al., 2004). The observed upper-limit of $2.55 \text{ ppbv ppbv}^{-1}$ in the range of $\Delta\text{O}_3/\Delta\text{CO}$ ratios for these measurements is much higher than for the data presented by Jaffe and Wigder (2012). Restricting this upper-limit to 2 ppbv removes two data points from the BORTAS statistics, reducing the mean value from $0.81 \pm 0.57 \text{ ppbv ppbv}^{-1}$ to $0.75 \pm 0.41 \text{ ppbv ppbv}^{-1}$. The observed increase in $\Delta\text{O}_3/\Delta\text{CO}$ ratios calculated from the BORTAS plume measurements with longer photochemical ages and moderate CH_3CN reflects plume dilution and efficient O_3 production at low NO_x mixing ratios (Bertschi and Jaffe, 2005; Jaffe and Wigder, 2012).

4.3 O_3 photochemistry in boreal biomass burning outflow

The $\Delta\text{O}_3/\Delta\text{CO}$ ratios presented above assume that enhancements in plume air over background values provide an indication of O_3 production in biomass burning plumes and that variations in the $\Delta\text{O}_3/\Delta\text{CO}$ ratio are due to variability in ΔO_3 . However, this ratio provides limited information on the photochemistry occurring in the plume air masses. We now evaluate O_3 mixing ratios measured in boreal biomass burning plumes during the BORTAS campaign against measurements of other chemical species associated with O_3 photochemistry.

4.3.1 Ozone production efficiency

A useful concept for understanding O_3 formation in polluted air masses is the ozone production efficiency (OPE) which is defined as the number of O_3 molecules produced

per molecule of NO_x that is oxidised (e.g. Trainer et al., 1993; Olszyna et al., 1994; Ryerson et al., 1998). We evaluate OPE in the plumes sampled during the BORTAS campaign through comparison of the measured O_3 mixing ratios to measurements of alkyl nitrates and the surrogate species NO_z . Photochemical production of O_3 in polluted air masses is dependent on the interaction between VOC and NO_x chemistry as shown in Fig. 1 and Reactions (R4)–(R6)/(R7). Thus, comparing the products of Reactions (R6) and (R7) provides an indication of the potential for O_3 formation in an air mass based on the relative concentrations of alkyl nitrates (RONO_2 in Reactions (R6) and (R7) and referred to as ΣAN from hereon in) and the level of oxidant ($\text{O}_x = \text{O}_3 + \text{NO}_2$). Similarly, the surrogate species NO_z ($= \text{NO}_y - \text{NO}_x$) represents the photo-oxidation products of reactive nitrogen species (i.e. NO_x) and evaluating O_x in relation to NO_z will provide an indication of the potential for O_3 formation in an air mass.

Figures 8a and b show the distribution of O_x relative to ΣAN and NO_z measured in boreal biomass burning plumes during BORTAS, respectively. In both cases three main populations of data points can be clearly defined with a distinctive “L-shape” distribution for measurements made on all BORTAS research flights, not including flight B626 (i.e., O_x mixing ratios in excess of 30 ppbv). Higher values of ΣAN and NO_z generally correspond to lower values of O_x , and lower values of ΣAN and NO_z generally correspond to higher values of O_x . The main factor that distinguishes between the two branches of the “L-shape” pattern is the aerosol loading of the measured air mass. High aerosol loading in the plumes is determined using measured black carbon mass concentration greater than $0.1 \mu\text{g cm}^{-3}$ and nephelometer measurements of the total scatter coefficient at 550 nm greater than $1.1 \times 10^{-5} \text{ m}^{-1}$ (corresponding to a black carbon mass concentration of $0.1 \mu\text{g cm}^{-3}$, and used to fill in gaps in the black carbon measurements). Although the vast majority of scattering is due to nonrefractory aerosols, the black carbon mass concentration and total scatter coefficient measurements are strongly correlated ($r = 0.92$) in the biomass burning plumes sampled during BORTAS, suggesting that these are reliable indicators of plume aerosol loading. The highest mixing ratios of ΣAN ($> 0.5 \text{ ppbv}$) and NO_z ($> 2 \text{ ppbv}$) were measured at night-time and daytime, respectively, in air masses with high aerosol loading and corresponding O_x mixing ratios between approximately 40 and 60 ppbv . Daytime ΣAN and night-time NO_z measurements along the horizontal branch of the “L-shape” distribution were typically made in air masses with high aerosol loading with mixing ratios ranging between 0.05 and 0.2 ppbv , respectively. Measurements made in plume air masses with low aerosol loading (black carbon mass concentration $< 0.1 \mu\text{g cm}^{-3}$) lie along the vertical branch of the “L-shaped” distributions in Fig. 8 and generally have lower ΣAN ($< 0.2 \text{ ppbv}$) and NO_z ($< 0.8 \text{ ppbv}$) mixing ratios and a wider range of O_x mixing ratios between 40 and 90 ppbv .




Fig. 8. Relationship between (a) O_x ($O_3 + NO_2$) and alkyl nitrates (ΣAN), and (b) O_x and NO_z ($NO_y - NO_x$) measured in boreal biomass burning plumes from the BAe-146 over the course of the BORTAS measurement campaign. Night-time and daytime measurements are represented by open and closed diamonds, respectively. Black symbols represent measurement with black carbon mass concentrations less than $0.1 \mu g cm^{-3}$ and red symbols with black carbon mass concentrations greater than $0.1 \mu g cm^{-3}$. Measurements made on flight B626 over the fire region in Northwestern Ontario are treated separately and are represented by closed green circles.

It should be noted that measurements with lower O_x mixing ratios (40–60 ppbv) show a mixture of both high and low aerosol loading and O_x mixing ratios greater than 60 ppbv were made only in air masses with low aerosol loading. The third population of data points, with O_x mixing ratios less than 30 ppbv corresponds to measurements made during flight B626 over Northwestern Ontario and show a narrower range of O_x (25–30 ppbv) for ΣAN between 0.05 and 0.4 ppbv and NO_z between 0.1 and 0.5 ppbv.

We estimate OPE in biomass burning plumes relative to ΣAN and NO_z from the gradients of straight lines fitted to each population of data points in Fig. 8 using a non-linear least squares method. This is a useful diagnostic for giving a general indication of the potential for O_3 formation but it should be noted that calculating OPE in this way

represents an upper limit due to potential loss of O_x and NO_y species from the plume air due to wet and dry deposition (e.g. Wang et al., 1996; Rickard et al., 2002). For relatively fresh biomass burning plumes measured on flight B626 the slope of the fit is 1.6 ± 15.6 ppbv ppbv $^{-1}$ relative to ΣAN and 0.07 ± 13.5 ppbv ppbv $^{-1}$ relative to NO_z (i.e., 1.6 O_x molecules are produced relative to each molecule of ΣAN and 0.07 O_x molecules per molecule of NO_z). The gradient of the fit increases to 7.7 ± 3.5 ppbv ppbv $^{-1}$ relative to ΣAN and 1.6 ± 0.9 ppbv ppbv $^{-1}$ relative to NO_z for the plume measurements made with high aerosol loading and to 472 ± 28 ppbv ppbv $^{-1}$ relative to ΣAN and 155 ± 5 ppbv ppbv $^{-1}$ relative to NO_z for the plume measurements made with low aerosol loading. The change in the relationship between O_x and ΣAN with plume photochemical age observed during BORTAS is consistent with that observed in pollution outflow from Mexico City by Perring et al. (2010) in which slopes of the lines of best fit to aircraft measurements in the outflow increased from approximately 16 ppbv ppbv $^{-1}$ to 89 ppbv ppbv $^{-1}$ as the plume age increased from less than 10 h to between 40 and 50 h. Perring et al. (2010) explained that increased O_3 production with plume age was due to an increase in non-nitrate producing O_3 precursors associated with depleted primary VOCs and enhanced secondary OVOCs. The high OPE relative to ΣAN calculated from the BORTAS measurements of aged plumes with low aerosol loading are consistent with slopes for remote areas of the troposphere such as the remote Pacific (Perring et al., 2010). The OPE values relative to NO_z calculated from the BORTAS measurements represent the extremes of typical OPE values previously reported in the literature and summarised in Table 1 of Rickard et al. (2002). The OPE of 1.6 ppbv ppbv $^{-1}$ relative to NO_z in the BORTAS plume measurements with high aerosol loading is similar to the OPE observed in urban and power station pollution plumes in the eastern USA (Chin et al., 1994; Ryerson et al., 1998; Daum et al., 2000; Sillman, 2000) whereas the OPE of 155 ppbv ppbv $^{-1}$ in the plume measurements with low aerosol loading is more comparable to the OPE observed in aged air masses of continental origin over the western Pacific Ocean (Davis et al., 1996).

An alternative method for calculating OPE from the observed O_x and NO_z mixing ratios is to calculate enhancements of each observed value relative to a background value (i.e. $OPE = \Delta O_x / \Delta NO_z$) similar to the method described for $\Delta O_3 / \Delta CO$ in the previous section. Figure 9 shows the distribution of OPE using $\Delta O_x / \Delta NO_z$ ratios calculated from the BORTAS biomass burning plume measurements as a function of NO_x . Background mixing ratios of O_x and NO_z were determined, in the same way as the background O_3 and CO values described in the previous section, to be 30 ppbv and 0.1 ppbv, respectively. The measurements are filtered for daytime/night-time and high/low aerosol loading as in Fig. 8. Measurements made in plume air with high aerosol loading typically have OPE less than 70 ppbv ppbv $^{-1}$ with




Fig. 9. Relationship between the enhancement ratio of O_x to NO_z ($\Delta O_x/\Delta NO_z$) in units of $ppbv\ ppbv^{-1}$ and NO_x mixing ratio in units of $ppbv$ measured in boreal biomass burning plumes from the BAe-146 over the course of the BORTAS measurement campaign. Symbols are the same as those described for Fig. 8.

NO_x mixing ratios ranging from 0.05 and 0.6 $ppbv$. Measurements with low aerosol loading have higher OPE between 50 and 400 $ppbv\ ppbv^{-1}$ and a narrower range of NO_x mixing ratios between 0 and 0.25 $ppbv$. The NO_x mixing ratios in this branch of the distribution shows a clear distinction between higher ($> 0.1\ ppbv$) and lower ($< 0.1\ ppbv$) NO_x mixing ratios with higher values corresponding to the night-time measurements. The measurements made on flight B626 over Northwestern Ontario have low OPE less than 30 $ppbv\ ppbv^{-1}$ over a similar range of NO_x mixing ratios as the high aerosol measurements. The changes in OPE relative to NO_x for the two populations of data points defined by the aerosol loading indicates that plume measurements were potentially made under two different chemical regimes. For the measurements made on flight B626 and those made during daytime with high aerosol loading, the large range of NO_x mixing ratios and small change in OPE possibly indicate of a more VOC-limited regime. For the low aerosol loading measurements, the much narrower range of NO_x mixing ratio and larger change in OPE possibly indicating a more NO_x -limited regime.

Figure 10 shows the distributions of relative humidity (RH) and aircraft altitude relative to OPE partitioned for night-time/daytime and high/low aerosol loading, as described above. Figure 10a shows that the measurements made with OPE less than 70 $ppbv\ ppbv^{-1}$ are typically characterised by higher relative humidity ($> 40\%$) than the measurements with OPE greater than 70 $ppbv\ ppbv^{-1}$ (RH typically $< 40\%$). The relationship between RH and OPE shows no distinction between daytime and night-time measurements. Figure 10b shows the vertical distribution of RH measured in boreal biomass burning plumes throughout the campaign. In general, measurements with low aerosol loading




Fig. 10. Relationship between (a) relative humidity in percent and the ratio of the $\Delta O_x/\Delta NO_z$ ratio in units of $ppbv\ ppbv^{-1}$, and (b) relative humidity and aircraft GPS altitude in km measured in boreal biomass burning plumes over the course of the BORTAS measurement campaign. Symbols are the same as those described for Fig. 8.

and low RH were made at higher altitudes ($> 3\ km$) during both daytime and night-time. Interestingly, measurements with high aerosol loading and high RH but low OPE were not solely made at lower altitudes but across the entire vertical range of the measurements, from 0.5 to 7 km. In general, plume measurements with low aerosol loading and low RH had corresponding CO, CH_3CN and VOC mixing ratios significantly higher than their respective background values. This, in combination with relatively long photochemical lifetimes of some of these species (approximately 40 days for CO and longer lived VOCs such as C_2H_6), suggests that photochemistry drives the increased OPE in these plumes. Low aerosol loading and low RH may be explained by these more aged air masses having experienced precipitation events during transport from the emission source region. Analysis of ground-based remote sensing measurements made from Halifax, NS, during the BORTAS campaign period identified a biomass burning plume with relatively low fine mode aerosol optical depth correlated with enhanced CO total columns,

which was attributable to precipitation along the plume trajectory (Griffin et al., 2013; Franklin et al., 2013).

The aerosol loading of the measured air masses provides the distinction between the two branches of the “L-shaped” pattern in OPE relative to both ΣAN and NO_z . Suppressed O_3 formation in biomass burning plume air masses with high aerosol loading is perhaps not so unexpected, as increased aerosol optical depth would be expected to attenuate the actinic flux and photolysis of NO_2 to form O_3 . However, this may not completely account for the suppressed O_3 formation as recent laboratory studies have highlighted potential loss of O_3 and NO_y species through heterogeneous uptake by aerosol in biomass burning plumes (Longfellow et al., 2000; Grassian, 2001). Furthermore, light-induced heterogeneous photochemical O_3 loss on the surface of biomass burning aerosols that increases with relative humidity has also been observed in laboratory and field measurements and shown to influence regional tropospheric O_3 distributions (Konovalov et al., 2012). Figure 11 shows an example of the relative change in NO_2 photolysis rate, j_{NO_2} , measured from the BAe-146 in plume and non-plume air masses along a section of BORTAS flight B624 over the North Atlantic Ocean on 21 July 2011 (see Table 1 and Fig. 9 in Palmer et al., 2013). This flight made five interceptions of a biomass burning plume with significantly elevated CO, CH_3CN and black carbon at altitudes between 1.7 and 3.2 km, and provides a useful dataset for evaluating the potential impact of enhanced black carbon aerosol on j_{NO_2} in biomass burning smoke plumes. These measurements were made above a planetary boundary layer between approximately 0.5 and 1.5 km estimated from temperature profiles over Sable Island and wind profiler measurements near Halifax (not shown). During the first two interceptions – prior to approximately 202.75 day and at 202.77 day of 2011 – the j_{NO_2} measurements show no clear deviation from the non-plume measurements. On the final three interceptions, at approximately 202.80, 202.85 and 202.89 days of the same year, the photolysis rate is reduced by a factor of approximately 2 relative to the rate measured in clear air. Although j_{NO_2} shows a reduction in the plume measurements, it is not reduced completely to zero, which potentially indicates that the reduced OPE in these measurements could be due to combination of the processes outlined above. Detailed analysis of these processes, which requires numerical model calculations to quantify the change in j_{NO_2} and its influence on O_3 photochemistry, is beyond the scope of the analysis presented here, and will be investigated through model studies that will be the subject of future papers.

4.3.2 Photostationary steady state and instantaneous O_3 production

To further evaluate O_3 photochemistry in the biomass burning plumes sampled during the BORTAS campaign we use a photostationary steady state calculation to quantify the relative contributions of NO_x and other peroxy radicals to the O_3




Fig. 11. Section of BORTAS flight B624 on 21 July 2011 showing (a) aircraft GPS altitude in km; (b) mixing ratios CO in ppbv and CH_3CN in pptv and black carbon mass concentration in units of $10^3 \mu\text{g cm}^{-3}$; and (c) NO_2 photolysis rate, j_{NO_2} , in units of s^{-1} . Solid symbols in (c) indicate measurements of CH_3CN mixing ratios greater than 150 pptv to indicate plume measurements.

production efficiency. From Reactions (R2) and (R3) and assuming photostationary steady state (i.e. the loss of O_3 via Reaction (R2) is equal to the production of O_3 via Reaction (R3)), the following ratio can be defined:

$$\Phi = \frac{j_{\text{NO}_2}[\text{NO}_2]}{k_{\text{NO}+\text{O}_3}[\text{NO}][\text{O}_3]} \quad (1)$$

where Φ is known as the photostationary ratio, $[\text{X}]$ represents the concentration of species X, j_{NO_2} is the photolysis rate of NO_2 in units of s^{-1} , and $k_{\text{NO}+\text{O}_3}$ is the kinetic rate coefficient for the reaction between NO and O_3 in units of molecules $\text{cm}^{-3} \text{s}^{-1}$. The value of $k_{\text{NO}+\text{O}_3}$ was calculated along the aircraft flight track using the recommended IUPAC definition of $1.4 \times 10^{-12} \exp(-1310/T)$ (Atkinson et al., 2004) where T is the air temperature measured from the aircraft. The photostationary ratio would have a value of unity if chemical cycling between O_3 and NO_x was the only process occurring (Leighton, 1961; Cantrell et al., 2003; Griffin et al., 2007). Deviations of this ratio from unity therefore give an indication of the influence of other chemical processes, for example, interaction between NO and families of organic peroxy radicals (RO_x) and hydro peroxy radicals (HO_x); we




Fig. 12. Relationship between O_3 photostationary state and the NO_y budget measured in boreal biomass burning plumes over the course of the BORTAS campaign. (a) and (b) show the calculated photostationary ratio, Φ , versus NO_x mixing ratios (ppbv) and the ratio of NO_x to NO_y respectively. (c) shows the relationship between NO_x/NO_y and NO_z . (d) shows calculated total peroxy radical mixing ratios, HO_xRO_x , required for a photostationary ratio of unity versus NO_x mixing ratio. Symbols are the same as those described for Fig. 4.

generalise these families of peroxy radicals as HO_xRO_x following Cantrell et al. (2003).

Figure 12 shows plots of photostationary steady state calculations made with the BORTAS aircraft measurements in biomass burning plumes over the course of the campaign. The photostationary ratio defined above is calculated from the measured values of each parameter in Eq. (1). Figure 12a shows the photostationary ratio versus NO_x mixing ratio with symbols denoting measurements made with high/low aerosol, as defined above. The calculated ratios for all the plume data points show a wide range of values between 0 and 10. Ratios for the measurements made with low aerosol loading generally lie in a narrower range from 0 to 2, suggesting that the measurements are closer to photostationary steady state. The measurements made with high aerosol generally have photostationary ratios greater than 2 with a wide range of values extending as high as 10. Figure 12b also shows the calculated photostationary ratio but as a function of the ratio NO_x/NO_y . Note that the expression defining NO_z (i.e. $NO_z = NO_y - NO_x$) can be rewritten as $NO_z = NO_y(1 - NO_x/NO_y)$, in which the term $1 - (NO_x/NO_y)$ can be considered as a measure of the photochemical age of the

air mass (i.e. the fraction of the initial NO_x emitted that has been converted into photooxidation products, as NO_y represents total NO_x emissions) (Olszyna et al., 1994). Values of NO_x/NO_y closer to unity are therefore representative of fresher emissions because there has been less photochemical processing of NO_x . In general, the BORTAS plume measurements have a range of NO_x/NO_y values between 0.0 and 0.4. The data points in Fig. 12 with NO_x/NO_y values greater than 0.4 are measurements made on flight B626 over Northwestern Ontario, indicating that fresher air masses were encountered closer to one of the main source regions. The measure of air mass photochemical age from the ratio of NO_x/NO_y is shown relative to the surrogate species NO_z in Fig. 12c. Measurements with NO_z mixing ratios less than 1 ppbv were made in both fresh and aged plumes, with the fresher plumes ($NO_x/NO_y > 0.4$) all having higher aerosol loading. Measurements with NO_z mixing ratios greater than 1 ppbv were all made in aged plumes ($NO_x/NO_y < 0.2$).

Using the measurements used to calculate the photostationary ratio, and including an additional term for the reaction of NO with peroxy radicals in Eq. (1) (Cantrell et al., 2003; Griffin et al., 2007), a photostationary ratio $\Phi_{HO_xRO_x}$

can be defined:

$$\Phi_{\text{HO}_x\text{RO}_x} = \frac{j_{\text{NO}_2}[\text{NO}_2]}{k_{\text{NO}+\text{O}_3}[\text{NO}][\text{O}_3] + k_{\text{NO}+\text{HO}_2}[\text{NO}][\text{HO}_x\text{RO}_x]} \quad (2)$$

where $k_{\text{NO}+\text{HO}_2}$ is the kinetic rate coefficient for the reaction between NO and HO₂, calculated along the aircraft flight track using the recommended IUPAC definition of $3.6^{-12} \exp(270/T)$ (Atkinson et al., 2004). Assuming photo-stationary steady state, we estimate mixing ratios of HO_xRO_x required to bring the photostationary ratio to unity by rearranging Eq. (2):

$$[\text{HO}_x\text{RO}_x] = \frac{j_{\text{NO}_2}[\text{NO}_2]}{k_{\text{NO}+\text{HO}_2}[\text{NO}]} - \frac{k_{\text{NO}+\text{O}_3}[\text{O}_3]}{k_{\text{NO}+\text{HO}_2}} \quad (3)$$

Figure 12d shows the HO_xRO_x mixing ratios estimated from the BORTAS plume measurements using Eq. (3) versus the NO_x/NO_y ratio. The measurements that were closer to unity in Fig. 12a and b (i.e. night-time or with low aerosol loading) generally show the lowest HO_xRO_x mixing ratios with values below 0.1 ppbv. For the plume measurements with high aerosol loading, the estimated HO_xRO_x mixing ratios are generally greater than 0.1 ppbv and extend up to approximately 0.4 ppbv. In the fresher plumes (i.e. NO_x/NO_y > 0.4), the HO_xRO_x mixing ratios are estimated to be just over 0.1 ppbv, with higher mixing ratios generally associated with more aged plumes. The calculated HO_xRO_x mixing ratios are consistent with values between 0 and 0.35 ppbv calculated from measurements at a ground-based rural site in the northeastern USA (Griffin et al., 2007).

4.3.3 Comparison of O₃ photochemistry from BORTAS and previous campaigns

We put the BORTAS plume measurements into context through comparisons with previous aircraft measurement campaigns with a focus over the central and eastern Canada, and sampled outflow from boreal biomass burning. Measurements from two different campaigns are used for this contextual analysis: the NASA ARCTAS-B measurement campaign during the summer of 2008 (Jacob et al., 2010); and the NASA ABLE 3B measurement campaign during the summer of 1990 (Harriss et al., 1994).

Figure 13 shows a comparison of O₃ production diagnostics (i.e. $\log_{10}(\Delta\text{O}_3 - \Delta\text{CO})$ vs. $\Delta\text{O}_3/\Delta\text{CO}$ and O_x vs. NO_z) derived from aircraft measurements made in biomass burning plumes during ARCTAS-B between 29 June and 13 July 2008 (Fig. 13a and b) and ABLE 3B between 6 July and 15 August 1990 (Fig. 13c and d). For the ARCTAS-B measurements we use 60 s averaged measurements of O₃, NO, NO₂ and NO_y made by chemiluminescence (Weinheimer et al., 1994), CO made by Tunable Diode Laser Absorption Spectroscopy (TDLAS) (Sachse et al., 1987), CH₃CN made by PTR-MS (Wisthaler et al., 2002), and black carbon from SP2 (Moteki and Kondo, 2007; Kondo et al., 2011). For comparison against the BORTAS measurements,

the ARCTAS-B data are filtered to cover a similar ranges of altitude (0.5–8.0 km), longitude (110–50° W), and latitude (40–60° N). Plume measurements are determined as for the BORTAS measurements, with CH₃CN mixing ratios in excess of 150 pptv, and high aerosol loading is determined for black carbon mass concentrations greater than 0.1 μg cm⁻³. For the ABLE 3B measurements we use 90 s averaged measurements of O₃, NO, NO₂ and NO_y, CO, and carbon tetrachloride (C₂Cl₄) (Blake et al., 1994; Harriss et al., 1994; Sandholm et al., 1994; Wofsy et al., 1994). The ABLE 3B deployment was very similar to that of BORTAS, with measurements extending from Northwest Territories and northern Manitoba across Ontario and Quebec to the east coast of Labrador. No CH₃CN measurements were made during ABLE 3B and plume measurements are determined from where measurements of C₂Cl₄ mixing ratios, a tracer of anthropogenic emissions (Wofsy et al., 1994), were below 20 pptv and CO mixing ratios were greater than 150 ppbv. The aerosol loading in the plume measurements are determined from concentrations of total fine (0.195–3.12 μm) and coarse (0.5–8.0 μm) particles greater than 20 cm⁻³.

The ARCTAS-B measurements shown in Fig. 13a, b show similar distributions to the BORTAS measurements shown in Figs. 7 and 8b, respectively. The variability in the ARCTAS-B $\Delta\text{O}_3/\Delta\text{CO}$ ratios is controlled by ΔO_3 in air masses with photochemical ages greater than 4 days, although, as with the BORTAS data, a wide range of $\Delta\text{O}_3/\Delta\text{CO}$ ratios were observed in air masses with these ages. The relationship between O_x and NO_z in the ARCTAS-B data also shows a distinctive “L-shaped” distribution, with relatively higher NO_z (greater than 1 ppbv) and lower O_x (less than 60 ppbv) mixing ratios in plume measurements with high aerosol loading, and lower NO_z/higher O_x associated with low aerosol loading. Slopes of straight lines fitted to the ARCTAS-B data give OPEs relative to NO_z of 4.6 ± 31.7 ppbv ppbv⁻¹ for the high aerosol measurements and 107.8 ± 5.0 ppbv ppbv⁻¹ for the low aerosol measurements. The increased high aerosol OPE and decreased low aerosol OPE compared against the BORTAS measurements further reflect that ARCTAS-B sampled fresher, near-field, biomass burning plumes. The ABLE 3B measurement campaign made considerably less measurements in biomass burning plumes than BORTAS and ARCTAS-B. Figure 13c shows that the ABLE 3B measurements that are attributed to biomass burning plumes, as described above, had $\Delta\text{O}_3/\Delta\text{CO}$ ratios between 0.1 and 1.0 ppbv ppbv⁻¹, indicating that ΔO_3 was always less than ΔCO in the ABLE 3B measurements. In general, the ABLE 3B measurements were shown to have relative high aerosol loading and the OPE relative to NO_z, Fig. 13d, was calculated to be 5.4 ± 49.2 ppbv ppbv⁻¹, consistent with the high aerosol OPE values calculated for BORTAS and ARCTAS-B.

Figure 14 shows the relationship between O₃ mixing ratio and OPE in boreal biomass burning plumes for the BORTAS, ARCTAS-B, and ABLE 3B measurement campaigns.




Fig. 13. Relationships between (a) ΔO_3 and ΔCO , following Fig. 7, and (b) O_x and NO_z , following Fig. 8b, from aircraft measurements made in biomass burning plumes during ARCTAS-B in the summer of 2008, and (c) ΔO_3 and ΔCO and (d) O_x and NO_z from aircraft measurements made in biomass burning plumes during ABLE 3B in the summer of 1990.




Fig. 14. Relationship between O_3 mixing ratio and OPE in boreal biomass burning plumes calculated from aircraft measurements made by the BORTAS (black diamonds), ARCTAS-B (red upward triangles), and ABLE 3B (green downward triangles) campaigns over eastern Canada.

The measurements with low OPE, less than $80 \text{ ppbv ppbv}^{-1}$, show a range of O_3 mixing ratios between 25 and 80 ppbv and show clear distinctions between the measurements made during the three campaigns. The lowest O_3 mixing ratios (25–30 ppbv) were measured on BORTAS flight B626, and the ARCTAS-B measurements typically lie between these values and O_3 mixing ratios measured on the other BORTAS flights, which range from approximately 40 to 55 ppbv. The ABLE 3B measurements at these lower OPEs have higher O_3 mixing ratios, 55–80 ppbv. Higher O_3 mixing ratios, 50–80 ppbv, are observed for OPEs between 80 and $180 \text{ ppbv ppbv}^{-1}$ for the ARCTAS-B measurements, and greater than $100 \text{ ppbv ppbv}^{-1}$ for the BORTAS measurements corresponding to aged plumes with NO_x/NO_y ratios $< 0.2 \text{ ppbv ppbv}^{-1}$ (see Fig. 12c) and low aerosol loading.

5 Conclusions

We have presented an analysis of O_3 photochemistry from in situ aircraft measurements made in boreal biomass burning plumes from a synthesis of aircraft measurements made

over eastern Canada during July and August 2011 as part of the BORTAS measurement campaign. We evaluated the tropospheric O₃ distribution observed over the course of the measurement campaign and the potential for quantifying the photochemical processes that influence O₃ production and loss within boreal biomass burning plumes. The range of the measured O₃ mixing ratios was typically between 45 and 70 ppbv, based on the lower and upper quartiles of the measurements on each research flight. Over the campaign, as a whole, there was no distinguishable difference between O₃ measurements made within plume and those made in clean air. At the highest values of CO (> 300 ppbv) we observed a relatively narrow range of O₃ mixing ratios between approximately 40 and 60 ppbv. We evaluated photochemical processes influencing the observed O₃ distribution using a number of different measurements made from the aircraft throughout the campaign. First, photochemical ageing of the plume measurements was determined using canister samples of different alkanes and their relationship to O₃ enhancements relative to CO. Second, the relative contributions of photochemical cycling of NO_x and peroxy radicals was analysed assuming photochemical steady state.

We calculated photochemical ages of the measured plumes using ratios of different alkanes assuming a common sink through oxidation with OH (Parrish et al., 2007) and found values that were consistent with previous measurements of biomass burning plumes. We used these ratios to estimate that plumes sampled during the campaign ranged in age from 0 to 10 days, indicating that the BORTAS measurements sampled a wider range of plume ages over previous measurement campaigns due to fires in Northwestern Ontario. When biomass burning plume measurements were evaluated relative to these estimated ages, we found that the highest measured values of CO and CH₃CN occurred at ages of 2–3 days, and distances of approximately 1000–3000 km from the expected emissions sources, reflecting the transport time of the plume to where it was intercepted by the aircraft. We calculated O₃ production in boreal biomass burning plumes, using $\Delta\text{O}_3/\Delta\text{CO}$ enhancement ratios, to be 0.020 ± 0.009 ppbv ppbv⁻¹ for plume ages less than 2 days, which increased to 0.13 ± 0.10 ppbv ppbv⁻¹ for plume ages between 2 and 5 days, and to 0.75 ± 0.41 ppbv ppbv⁻¹ for plume ages greater than 5 days. The $\Delta\text{O}_3/\Delta\text{CO}$ enhancement ratio indicated that net O₃ production in boreal biomass burning plumes is a function of photochemical age, consistent with a recent review of O₃ production in boreal biomass burning outflow (Jaffe and Wigder, 2012). The $\Delta\text{O}_3/\Delta\text{CO}$ ratios calculated for the BORTAS measurements were controlled by ΔO_3 after 4 days and a combination of ΔO_3 and ΔCO prior to that.

We performed a detailed analysis of the photochemical processes influencing the O₃ distribution in boreal biomass burning plumes using measurements of total alkyl nitrates (ΣAN), NO_x and NO_y made from the aircraft. Through comparing plume O₃ mixing ratios with ΣAN and the

surrogate species NO_z, we found that O₃ production was suppressed in relatively fresh plumes (photochemical ages less than 4 days) with between 7.7 ± 3.5 ppbv of O_x produced per ppbv of ΣAN and 1.6 ± 0.9 ppbv of O_x produced per ppbv of NO_x that was oxidised. In aged plumes the OPE increased significantly to 472 ± 28 ppbv O_x per ppbv ΣAN and 155 ± 5 ppbv O_x per ppbv NO_z. We found that the plume OPE was strongly correlated with the aerosol loading of the air mass, with higher aerosol loading (black carbon mass concentration > 0.1 $\mu\text{g cm}^{-3}$) generally associated with younger plumes and lower enhancement ratios of $\Delta\text{O}_x/\Delta\text{NO}_z$, and lower aerosol loading (black carbon mass concentration < 0.1 $\mu\text{g cm}^{-3}$) in plumes that are more aged. When compared against NO_x, the $\Delta\text{O}_x/\Delta\text{NO}_z$ ratios indicated that the BORTAS plume measurements were made in different chemical regimes: a more VOC-limited regime for flight B626 and the high aerosol loading measurements, and a more NO_x-limited regime for the low aerosol loading measurements. Furthermore, we found that measurements of plume air with high aerosol loading and relatively low OPE were made at high relative humidity (> 40 %) throughout the vertical extent of the aircraft measurements, and the low aerosol plumes with relatively high OPE were made at low relative humidity and generally higher altitude (typically above 4 km).

When the BORTAS measurements were compared to similar measurements made in boreal biomass burning plumes over eastern Canada by the ARCTAS-B and ABLE 3B campaigns, we found that OPE increased with plume ages calculated by NMHC ratios. In all campaign measurements we found that plumes with high aerosol loading typically had low OPE, less than 80 ppbv ppbv⁻¹, and aged plumes were typically associated with low aerosol loading and relatively high OPE. We showed that on one of the BORTAS flights, which made multiple interceptions of a plume with high aerosol loading, j_{NO_2} was reduced by a factor of approximately 2 within the plume, suggesting that increased optical depth of the plume attenuated the actinic flux and subsequent photolysis of NO₂ to form O₃. However, as j_{NO_2} was not reduced completely to zero, the relatively low OPE in these plumes could also be due to other in-plume processes such as potential loss of O₃ and NO_y species through heterogeneous uptake by aerosol in biomass burning plumes (Longfellow et al., 2000; Grassian, 2001) or light-induced heterogeneous photochemical O₃ loss on the surface of biomass burning aerosols that increases with relative humidity (Konovalov et al., 2012).

Other potential factors not considered in the analysis presented here could have influenced the tropospheric O₃ distribution observed over the course of the BORTAS campaign. Some of the BORTAS measurements were potentially made downwind of anthropogenic emissions sources, such as the northeastern US and the Alberta tar sands, and while great care was taken to avoid pollution outflow from these sources in the deployment of the BORTAS aircraft campaign they

could have provided additional sources of VOCs and NO_x (from PAN decomposition) and influenced O₃ production over the BORTAS region. Measurements were also made downwind of lightning activity across North America and could have provided an additional source of NO_x in the free troposphere. While O₃ enhancements in boreal biomass burning plumes with low aerosol loading and low relative humidity could have been influenced by mixing of the plumes with air of stratospheric origin, the highest O₃ mixing ratios measured during BORTAS were generally less than 100 ppbv and lower than typical stratospheric O₃ mixing ratios. Empirical, observation-based methods provide a powerful tool for diagnosing O₃ photochemistry and NO_y speciation associated with biomass burning. The observational analysis presented here is being investigated further using detailed chemical modelling and will be the subject of future publications using the BORTAS data.

Acknowledgements. The BORTAS project is funded by the Natural Environment Research Council (NERC) under grant NE/F017391/1. Airborne data were obtained using the BAe-146-301 Atmospheric Research Aircraft (ARA) operated by Directflight Ltd (DFL) and managed by the Facility for Airborne Atmospheric Measurements (FAAM), which is a joint entity of NERC and the UK Meteorological Office. We acknowledge the great efforts provided by the staff of FAAM, DFL, Avalon Aero Ltd and Captains Alan Foster and Charlie Whittaker and First Officer Ian Ramsay-Rae. The ARCTAS-B 60-s merges were obtained from the POLARCAT/ARCTAS archive at the NASA Langley Research Center Atmospheric Science Data Center (LaRC DACC). The ABLE 3B 90-s merges were obtained from the NASA Global Tropospheric Experiment data archive at the LaRC DACC. We acknowledge Andrew Weinheimer, Armin Wisthaler, Glenn Diskin, and Yutaka Kondo for their respective measurements of NO_x/NO_y/O₃, CH₃CN, CO, and black carbon made during the ARCTAS-B measurement campaign as well as the science teams of ARCTAS and ABLE 3B for use of the measurements taken during each campaign.

Edited by: B. N. Duncan

References

- Alvarado, M. J., Logan, J. A., Mao, J., Apel, E., Riemer, D., Blake, D., Cohen, R. C., Min, K.-E., Perring, A. E., Browne, E. C., Wooldridge, P. J., Diskin, G. S., Sachse, G. W., Fuelberg, H., Sessions, W. R., Harrigan, D. L., Huey, G., Liao, J., Case-Hanks, A., Jimenez, J. L., Cubison, M. J., Vay, S. A., Weinheimer, A. J., Knapp, D. J., Montzka, D. D., Flocke, F. M., Pollack, I. B., Wennberg, P. O., Kurten, A., Crounse, J., Clair, J. M. St., Wisthaler, A., Mikoviny, T., Yantosca, R. M., Carouge, C. C., and Le Sager, P.: Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: an integrated analysis of aircraft and satellite observations, *Atmos. Chem. Phys.*, 10, 9739–9760, doi:10.5194/acp-10-9739-2010, 2010.
- Anderson, B. E., Gregory, G. L., Barrick, J. D. W., Collins Jr., J. E., Sachse, G. W., Shipham, M. C., and Hudgins, C. H.: Summer-time tropospheric ozone distributions over central and eastern Canada, *J. Geophys. Res.*, 99, 1781–1792, 1994.
- Anderson, T. L. and Ogren, J. A.: Determining aerosol radiative properties using the TSI 3563 integrating nephelometer, *Aerosol Sci. Technol.*, 29, 57–69, 1998.
- Atkinson, R. and Arey, J.: Atmospheric degradation of volatile organic compounds, *Chem. Rev.*, 103, 4605–4638, 2003.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I – gas phase reactions of O_x, HO_x, NO_x and SO_x species, *Atmos. Chem. Phys.*, 4, 1461–1738, doi:10.5194/acp-4-1461-2004, 2004.
- Bange, H. W. and Williams, J.: New directions: Acetonitrile in atmospheric and biogeochemical cycles, *Atmos. Environ.*, 34, 4959–4960, 2000.
- Bertschi, I. T. and Jaffe, D. A.: Long-range transport of ozone, carbon monoxide, and aerosols to the NE Pacific troposphere during the summer of 2003: Observations of smoke plumes from Asian boreal fires, *J. Geophys. Res.*, 110, D05303, doi:10.1029/2004JD005135, 2005.
- Blake, D. R., Smith, Jr., T. W., Chen, T.-Y., Whipple, W. J., and Rowland, F. S.: Effects of biomass burning on summertime non-methane hydrocarbon concentrations in the Canadian wetlands, *J. Geophys. Res.*, 99, 1699–1719, 1994.
- Cantrell, C. A., Mauldin, L., Zondlo, M., Eisele, F., Kosiuch, E., Shetter, R., Lefer, B., Hall, S., Campos, T., Ridley, B., Walega, J., Fried, A., Wert, B., Flocke, F., Weinheimer, A., Hannigan, J., Coffey, M., Atlas, E., Stephens, S., Heikes, B., Snow, J., Blake, D., Blake, N., Katzenstein, A., Lopez, J., Browell, E. V., Dibb, J., Scheuer, E., Seid, G., and Talbot, R.: Steady state free radical budgets and ozone photochemistry during TOPSE, *J. Geophys. Res.*, 108, 8361, doi:10.1029/2002JD002198, 2003.
- Chin, M., Jacob, D. J., and Munger, J. W.: Relationship of ozone and carbon monoxide over North America, *J. Geophys. Res.*, 99, 14565–14573, 1994.
- Cook, P. A., Savage, N. H., Turquety, S., Carver, G. D., O'Connor, F. M., Heckel, A., Stewart, D., Whalley, L. K., Parker, A. E., Schlager, H., Singh, H. B., Avery, M. A., Sachse, G. W., Brune, W., Richter, A., Burrows, J. P., Purvis, R., Lewis, A. C., Reeves, C. E., Monks, P. S., Levine, J. G., and Pyle, J. A.: Forest fire plumes over the North Atlantic: p-TOMCAT model simulations with aircraft and satellite measurements from the ITOP/ICARTT campaign, *J. Geophys. Res.*, 112, D10S43, doi:10.1029/2006JD007563, 2007.
- Dari-Salisburgo, C., Di Carlo, P., Giammaria, F., Kajii, Y., and D'Altorio, A.: Laser induced fluorescence instrument for NO₂ measurements: Observations at a central Italy background site, *Atmos. Environ.*, 43, 970–977, 2009.
- Daum, P. H., Kleinman, L. I., Imre, D., Nunnermacker, L. J., Lee, Y.-N., Springston, S. R., and Newman, L.: Analysis of O₃ formation during a stagnation episode in central Tennessee in summer 1995, *J. Geophys. Res.*, 105, 9107–9119, 2000.
- Davis, D. D., Crawford, J., Chen, G., Chameides, W., Liu, S., Bradshaw, J., Sandholm, S., Sachse, G., Gregory, G., Anderson, B., Barrick, J., Bachmeier, A., Collins, J., Browell, E., Blake, D., Rowland, S., Kondo, Y., Singh, H., Talbot, R., Heikes, B., Merrill, J., Rodriguez, J., and Newell, R. E.: Assessment of ozone photochemistry in the western North Pacific as inferred from

- PEM-West A observations during the fall 1991, *J. Geophys. Res.*, 101, 2111–2134, 1996.
- Day, D. A., Dillon, M. B., Wooldridge, P. J., Thornton, J. A., Rosen, R. S., Wood, E. C., and Cohen, R. C.: On alkyl nitrates, O₃, and the “missing NO_y”, *J. Geophys. Res.*, 108, 4501, doi:10.1029/2003JD003685, 2003.
- DeBell, L. J., Talbot, R. W., Dibb, J. E., Munger, J. W., Fischer, E. V., and Frolking, S. E.: A major regional air pollution event in the northeastern United States caused by extensive forest fires in Quebec, Canada, *J. Geophys. Res.*, 109, D19305, doi:10.1029/2004JD004840, 2004.
- Di Carlo, P., Aruffo, E., Busilacchio, M., Giammaria, F., Darisalisburgo, C., Biancofiore, F., Visconti, G., Lee, J., Moller, S., Reeves, C. E., Bauguitte, S., Forster, G., Jones, R. L., and Ouyang, B.: Aircraft based four-channel thermal dissociation laser induced fluorescence instrument for simultaneous measurements of NO₂, total peroxy nitrate, total alkyl nitrate, and HNO₃, *Atmos. Meas. Tech.*, 6, 971–980, doi:10.5194/amt-6-971-2013, 2013.
- Fehsenfeld, F. C., Ancellet, G., Bates, T. S., Goldstein, A. H., Hardesty, R. M., Honrath, R., Law, K. S., Lewis, A. C., Leaitch, R., McKeen, S., Meagher, J., Parrish, D. D., Pszenny, A. A. P., Russell, P. B., Schlager, H., Seinfeld, J., Talbot, R., and Zbinden, R.: International Consortium for Atmospheric Research on Transport and Transformation (ICARTT): North America to Europe – Overview of the 2004 summer field study, *J. Geophys. Res.*, 111, D23S01, doi:10.1029/2006JD007829, 2006.
- Franklin, J. E., Drummond, J. R., Griffin, D., Pierce, J. R., Waugh, D. L., Palmer, P. I., Parrington, M., Lee, J. D., Lewis, A. C., Rickard, A. R., Taylor, J., Chisholm, L., Duck, T. J., Hopper, J. T., Gibson, M., Curry, K. R., Sakamoto, K. M., Lesins, G., Walker, K. A., Dan, L., and Kliever, J.: A case study of aerosol depletion in a biomass burning plume over Eastern Canada during the 2011 BORTAS field experiment, in preparation, 2013.
- Gerbig, C., Schmitgen, S., Kley, D., and Volz-Thomas, A.: An improved fast-response vacuum-UV resonance fluorescence CO instrument, *J. Geophys. Res.*, 104, 1699–1704, 1999.
- Goode, J. G., Yokelson, R. J., Ward, D. E., Susott, R. A., Babbitt, R. E., Davies, M. A., and Hao, W. M.: Measurements of excess O₃, CO₂, CO, CH₄, C₂H₄, C₂H₂, HCN, NO, NH₃, HCOOH, CH₃COOH, HCHO, and CH₃OH in 1997 Alaskan biomass burning plumes by airborne Fourier transform infrared spectroscopy (AFTIR), *J. Geophys. Res.*, 105, 22147–22166, 2000.
- Grassian, V. H.: Heterogeneous uptake and reaction of nitrogen oxides and volatile organic compounds on the surface of atmospheric particles including oxides, carbonates, soot and mineral dust: Implications for the chemical balance of the troposphere, *Int. Rev. Phys. Chem.*, 20, 467–548, 2001.
- Griffin, D., Walker, K. A., Franklin, J. E., Parrington, M., Whaley, C., Hopper, J., Drummond, J. R., Palmer, P. I., Strong, K., Duck, T. J., Abboud, I., Bernath, P. F., Clerbaux, C., Coheur, P.-F., Curry, K. R., Dan, L., Hyer, E., Kliever, J., Lesins, G., Maurice, M., Saha, A., Tereszchuk, K., and Weaver, D.: Investigation of CO, C₂H₆ and aerosols in a boreal fire plume over eastern Canada during BORTAS 2011 using ground- and satellite-based observations, and model simulations, *Atmos. Chem. Phys. Discuss.*, 13, 11071–11109, doi:10.5194/acpd-13-11071-2013, 2013.
- Griffin, R. J., Beckman, P. J., Talbot, R. W., Sive, B. C., and Varner, R. K.: Deviations from ozone photostationary state during the International Consortium for Atmospheric Research on Transport and Transformation 2004 campaign: Use of measurements and photochemical modeling to assess potential causes, *J. Geophys. Res.*, 112, D10S07, doi:10.1029/2006JD007604, 2007.
- Harriss, R. C., Wofsy, S. C., S., B. D., Shipham, M. C., Jacob, D. J., Hoell Jr., J. M., Bendura, R. J., Drewry, J. W., McNeal, R. J., Navarro, R. L., Gidge, R. N., and Rabine, V. E.: The Arctic Boundary Layer Expedition (ABLE 3A): July–August 1988, *J. Geophys. Res.*, 97, 16383–16394, 1992.
- Harriss, R. C., Wofsy, S. C., Hoell Jr., J. M., Bendura, R. J., Drewry, J. W., McNeal, R. J., Pierce, D., Rabine, V., and Snell, R. L.: The Arctic Boundary Layer Expedition (ABLE 3B): July–August 1990, *J. Geophys. Res.*, 99, 1635, 1994.
- Helmig, D., Tanner, D. M., Honrath, R. E., Owen, R. C., and Parrish, D. D.: Nonmethane hydrocarbons at Pico Mountain, Azores: 1. Oxidation chemistry in the North Atlantic region, *J. Geophys. Res.*, 113, D20S91, doi:10.1029/2007JD008930, 2008.
- Honrath, R. E., Owen, R. C., Val Martin, M., Reid, J. S., Lapina, K., Fialho, P., Dziobak, M. P., Kleissl, J., and Westphal, D. L.: Regional and hemispheric impacts of anthropogenic and biomass burning emissions on summertime CO and O₃ in the North Atlantic lower free troposphere, *J. Geophys. Res.*, 109, D24310, doi:10.1029/2004JD005147, 2004.
- Hopkins, J. R., Lewis, A. C., and Read, K. A.: A two-column method for long-term monitoring of non-methane hydrocarbons (NMHCs) and oxygenated volatile organic compounds (o-VOCs), *J. Environ. Monitor.*, 5, 8–13, 2003.
- Hopkins, J. R., Jones, C. E., and Lewis, A. C.: A dual channel gas chromatograph for atmospheric analysis of volatile organic compounds including oxygenated and monoterpene compounds, *J. Environ. Monitor.*, 13, 2268–2276, 2011.
- Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Emmons, L. K., Ferrare, R. A., Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E., Pederson, J. R., and Fisher, J. A.: The Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) mission: design, execution, and first results, *Atmos. Chem. Phys.*, 10, 5191–5212, doi:10.5194/acp-10-5191-2010, 2010.
- Jaffe, D. A. and Wigder, N. L.: Ozone production from wildfires: A critical review, *Atmos. Environ.*, 51, 1–10, 2012.
- Jobson, B. T., Wu, Z., Niki, H., and Barrie, L. A.: Seasonal trends of isoprene, C₂–C₅ alkanes, and acetylene at a remote boreal site in Canada, *J. Geophys. Res.*, 99, 1589–1599, 1994.
- Kleinman, L. I.: Ozone process insights from field experiments – part II: Observation-based analysis for ozone production, *Atmos. Environ.*, 34, 2023–2033, 2000.
- Kondo, Y., Matsui, H., Moteki, N., Sahu, L., Takegawa, N., Kajino, M., Zhao, Y., Cubison, M. J., Jimenez, J. L., Vay, S., Diskin, G. S., Anderson, B., Wisthaler, A., Mikoviny, T., Fülberg, H. E., Blake, D. R., Huey, G., Weinheimer, A. J., Knapp, D. J., and Brune, W. H.: Emissions of black carbon, organic, and inorganic aerosols from biomass burning in North America and Asia in 2008, *J. Geophys. Res.*, 116, D08204, doi:10.1029/2010JD015152, 2011.
- Kononov, I. B., Beekman, M., D’Anna, B., and George, C.: Significant light induced ozone loss on biomass burning aerosol: Evidence from chemistry-transport modeling based

- on new laboratory studies, *Geophys. Res. Lett.*, 39, L17807, doi:10.1029/2012GL052432, 2012.
- Laborde, M., Mertes, P., Zieger, P., Dommen, J., Baltensperger, U., and Gysel, M.: Sensitivity of the Single Particle Soot Photometer to different black carbon types, *Atmospheric Measurement Techniques*, 5, 1031–1043, 2012.
- Lapina, K., Honrath, R. E., Owen, R. C., Val Martin, M., and Pfister, G.: Evidence of significant large-scale impacts of boreal fires on ozone levels in the midlatitude Northern Hemisphere free troposphere, *Geophys. Res. Lett.*, 33, L10815, doi:10.1029/2006GL025878, 2006.
- Latifovic, R., Zhu, Z.-L., Cihlar, J., Giri, C., and Olthof, I.: Land cover mapping of North and Central America – Global Land Cover 2000, *Remote Sens. Environ.*, 89, 116–127, 2004.
- Lee, J. D., Moller, S. J., Read, K. A., Lewis, A. C., Mendes, L., and Carpenter, L. J.: Year-round measurements of nitrogen oxides and ozone in the tropical North Atlantic marine boundary layer, *J. Geophys. Res.*, 114, D21302, doi:10.1029/2009JD011878, 2009.
- Leighton, P. A.: Photochemistry of air pollution, *Phys. Chem.*, 9, 300 pp., 1961.
- Lewis, A. C., Evans, M. J., Methven, J., Watson, N., Lee, J. D., Hopkins, J. R., Purvis, R. M., Arnold, S. R., McQuaid, J. B., Whalley, L. K., Pilling, M. J., Heard, D. E., Monks, P. S., Parker, A. E., Reeves, C. E., Oram, D. E., Mills, G., Bandy, B. J., Stewart, D., Coe, H., Williams, P., and Crosier, J.: Chemical composition observed over the mid-Atlantic and the detection of pollution signatures far from source regions, *J. Geophys. Res.*, 112, D10S39, doi:10.1029/2006JD007584, 2007.
- Longfellow, C. A., Ravishankara, A. R., and Hanson, D. R.: Reactive and nonreactive uptake on hydrocarbon soot: HNO_3 , O_3 , and N_2O_5 , *J. Geophys. Res.*, 105, 24345–24350, 2000.
- Mauzerall, D. L., Jacob, D. J., Fan, S.-M., Bradshaw, J. D., Gregory, G. L., Sachse, G. W., and Blake, D. R.: Origin of tropospheric ozone at remote high northern latitudes in summer, *J. Geophys. Res.*, 101, 4175–4188, 1996.
- McKeen, S. A., Wotawa, G., Parrish, D. D., Holloway, J. S., Buhr, M. P., Hübler, G., Fehsenfeld, F. C., and Meagher, J. F.: Ozone production from Canadian wildfires during June and July of 1995, *J. Geophys. Res.*, 107, 4192, doi:10.1029/2001JD000697, 2001.
- McMeeking, G. R., Hamburger, T., Liu, D., Flynn, M., Morgan, W. T., Northway, M., Highwood, E. J., Krejci, R., Allan, J. D., Minikin, A., and Coe, H.: Black carbon measurements in the boundary layer over western and northern Europe, *Atmos. Chem. Phys.*, 10, 9393–9414, doi:10.5194/acp-10-9393-2010, 2010.
- Moteki, N. and Kondo, Y.: Effects of mixing state on black carbon measurements by laser-induced incandescence, *Aerosol Sci. Technol.*, 41, 398–417, 2007.
- Murphy, J. G., Oram, D. E., and Reeves, C. E.: Measurements of volatile organic compounds over West Africa, *Atmos. Chem. Phys.*, 10, 5281–5294, doi:10.5194/acp-10-5281-2010, 2010.
- Olszyna, K. J., Bailey, E. M., Simonaitis, R., and Meagher, J. F.: O_3 and NO_y relationships at a rural site, *J. Geophys. Res.*, 99, 14557–14563, 1994.
- Palmer, P. I., Parrington, M., Lee, J. D., Lewis, A. C., Rickard, A. R., Bernath, P. F., Duck, T. J., Waugh, D. L., Tarasick, D. W., Andrews, S., Aruffo, E., Bailey, L. J., Barrett, E., Bauguitte, S. J.-B., Curry, K. R., Di Carlo, P., Chisholm, L., Dan, L., Forster, G., Franklin, J. E., Gibson, M. D., Griffin, D., Helmig, D., Hopkins, J. R., Hopper, J. T., Jenkin, M. E., Kindred, D., Kliever, J., Le Breton, M., Matthiesen, S., Maurice, M., Moller, S., Moore, D. P., Oram, D. E., O’Shea, S. J., Owen, R. C., Pagniello, C. M. L. S., Pawson, S., Percival, C. J., Pierce, J. R., Punjabi, S., Purvis, R. M., Remedios, J. J., Rotermund, K. M., Sakamoto, K. M., da Silva, A. M., Strawbridge, K. B., Strong, K., Taylor, J., Trigwell, R., Tereszchuk, K. A., Walker, K. A., Weaver, D., Whaley, C., and Young, J. C.: Quantifying the impact of Boreal forest fires on Tropospheric oxidants over the Atlantic using Aircraft and Satellites (BORTAS) experiment: design, execution and science overview, *Atmos. Chem. Phys.*, 13, 6239–6261, doi:10.5194/acp-13-6239-2013, 2013.
- Parrington, M., Palmer, P. I., Henze, D. K., Tarasick, D. W., Hyer, E. J., Owen, R. C., Helmig, D., Clerbaux, C., Bowman, K. W., Deeter, M. N., Barratt, E. M., Coheur, P.-F., Hurtmans, D., Jiang, Z., George, M., and Worden, J. R.: The influence of boreal biomass burning emissions on the distribution of tropospheric ozone over North America and the North Atlantic during 2010, *Atmos. Chem. Phys.*, 12, 2077–2098, doi:10.5194/acp-12-2077-2012, 2012.
- Parrish, D. D., Hahn, C. J., Williams, E. J., Norton, R. B., Fehsenfeld, F. C., Singh, H. B., Shetter, J. D., Gandrud, B. W., and Ridley, B. A.: Indications of photochemical histories of Pacific air masses from measurements of atmospheric trace species at Point Arena, California, *J. Geophys. Res.*, 97, 15883–15901, 1992.
- Parrish, D. D., Holloway, J. S., Trainer, M., Murphy, P. C., Forbes, G. L., and Fehsenfeld, F. C.: Export of North American Ozone Pollution to the North Atlantic Ocean, *Science*, 259, 1436–1439, 1993.
- Parrish, D. D., Stohl, A., Forster, C., Atlas, E. L., Blake, D. R., Goldan, P. D., Kuster, W. C., and de Gouw, J. A.: Effects of mixing on evolution of hydrocarbon ratios in the troposphere, *J. Geophys. Res.*, 112, D10S34, doi:10.1029/2006JD007583, 2007.
- Perring, A. E., Bertram, T. H., Farmer, D. K., Wooldridge, P. J., Dibb, J., Blake, N. J., Blake, D. R., Singh, H. B., Fuelberg, H., Diskin, G., Sachse, G., and Cohen, R. C.: The production and persistence of ΣRONO_2 in the Mexico City plume, *Atmos. Chem. Phys.*, 10, 7215–7229, doi:10.5194/acp-10-7215-2010, 2010.
- Pfister, G. G., Emmons, L. K., Hess, P. G., Honrath, R., Lamarque, J.-F., Val Martin, M., Owen, R. C., Avery, M. A., Browell, E. V., Holloway, J. S., Nedelec, P., Purvis, R., Ryerson, T. B., Sachse, G. W., and Schlager, H.: Ozone production from the 2004 North American boreal fires, *J. Geophys. Res.*, 111, D24S07, doi:10.1029/2006JD007695, 2006.
- Real, E., Law, K. S., Weinzierl, B., Fiebig, M., Petzold, A., Wild, O., Methven, J., Arnold, S., Stohl, A., Huntrieser, H., Roiger, A., Schlager, H., Stewart, D., Avery, M., Sachse, G., Browell, E., Ferrare, R., and Blake, D.: Processes influencing ozone levels in Alaskan forest fire plumes during long-range transport over the North Atlantic, *J. Geophys. Res.*, 112, D10S41, doi:10.1029/2006JD007576, 2007.
- Rickard, A. R., Salisbury, G., Monks, P. S., Lewis, A. C., Bauguitte, S., Bandy, B. J., Clemitshaw, K. C., and Penkett, S. A.: Comparison of measured ozone production efficiencies in the marine boundary layer at two European coastal sites under different pollution regimes, *J. Atmos. Chem.*, 43, 107–134, 2002.
- Rudolph, J. and Johnen, F. J.: Measurements of light atmospheric hydrocarbons over the Atlantic in regions of low biological ac-

- tivity, *J. Geophys. Res.*, 95, 20583–20591, 1990.
- Ryerson, T. B., Buhr, M. P., Frost, G. J., Goldan, P. D., Holloway, J. S., Hübler, G., Jobson, B. T., Kuster, W. C., McKee, S. A., Parrish, D. D., Roberts, J. M., Sueper, D. T., Trainer, M., Williams, J., and Fehsenfeld, F. C.: Emissions lifetimes and ozone formation in power plant plumes, *J. Geophys. Res.*, 103, 22569–22583, 1998.
- Sachse, G. W., Hill, G. F., Wade, L. O., and Perry, M. G.: Fast-response, high-precision carbon monoxide sensor using a tunable diode laser absorption technique, *J. Geophys. Res.*, 92, 2071–2081, 1987.
- Sandholm, S., Olson, J., Bradshaw, J., Talbot, R., Singh, H., Gregory, G., Blake, D., Anderson, B., Sachse, G., Barrick, J., Collins, J., Klemm, K., Lefer, B., Klemm, O., Gorzelska, K., Herlth, D., and O'Hara, D.: Summertime partitioning and budget of NO_y compounds in the troposphere over Alaska and Canada: ABLE 3B, *J. Geophys. Res.*, 99, 1837–1861, 1994.
- Schwarz, J. P., Gao, R. S., Fahey, D. W., Thomson, D. S., Watts, L. A., Wilson, J. C., Reeves, J. M., Darbehesti, M., Baumgardner, D. G., Kok, G. L., Chung, S. H., Schulz, M., Hendricks, J., Lauer, A., Kärcher, B., Slowik, J. G., Rosenlof, K. H., Thompson, T. L., Langford, A. O., Loewenstein, M., and Aikin, K. C.: Single-particle measurements of midlatitude black carbon and light-scattering aerosols from the boundary layer to the lower stratosphere, *J. Geophys. Res.*, 111, D16207, doi:10.1029/2006JD007076, 2006.
- Sillman, S.: Ozone production efficiency and loss of NO_x in power plant plumes: Photochemical model and interpretation of measurements in Tennessee, *J. Geophys. Res.*, 105, 9189–9202, 2000.
- Singh, H. B., Anderson, B. E., Brune, W. H., Cai, C., Cohen, R. C., Crawford, J. H., Cubison, M. J., Czech, E. P., Emmons, L., Fuelberg, H. E., Huey, G., Jacob, D. J., Jimenez, J. L., Kaduwela, A., Kondo, Y., Mao, J., Olson, J. R., Sachse, G. W., Vay, S. A., Weinheimer, A., Wennberg, P. O., Wisthaler, A., and the ARCTAS Science Team: Pollution influences on atmospheric composition and chemistry at high northern latitudes: Boreal and California forest fire emissions, *Atmos. Environ.*, 44, 4553–4564, 2010.
- Spivakovsky, C. M., Logan, J. A., Montzka, S. A., Balkanski, Y. J., Foreman-Fowler, M., Jones, D. B. A., Horowitz, L. W., Fusco, A. C., Brenninkmeijer, C. A. M., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Three-dimensional climatological distribution of tropospheric OH: Update and evaluation, *J. Geophys. Res.*, 105, 8931–8980, 2000.
- Tanimoto, H., Matsumoto, K., and Uematsu, M.: Ozone-CO correlations in Siberian wildfire plumes observed at Rishiri Island, *SOLA*, 4, 65–68, doi:10.2151/sola.2008-017, 2008.
- Trainer, M., Parrish, D. D., Buhr, M. P., Norton, R. B., Fehsenfeld, F. C., Anlauf, K. G., Bottenheim, J. W., Tang, Y. Z., Wiebe, H. A., Roberts, J. M., Tanner, R. L., Newman, L., Bowersox, V. C., Meagher, J. F., Olszyna, K. J., Rodgers, M. O., Wang, T., Berresheim, H., Demerjian, K. L., and Roychowdhury, U. K.: Correlation of ozone with NO_y in photochemically aged air, *J. Geophys. Res.*, 98, 2917–2925, 1993.
- Val Martin, M., Honrath, R. E., Owen, R. C., Pfister, G., Fialho, P., and Barata, F.: Significant enhancements of nitrogen oxides, black carbon, and ozone in the North Atlantic lower free troposphere resulting from North American boreal wildfires, *J. Geophys. Res.*, 111, D23S60, doi:10.1029/2006JD007530, 2006.
- Volz-Thomas, A., Lerner, A., Pätz, H.-W., Schulz, M., McKenna, D. S., Schmitt, S., Madronich, S., and Röth, E. P.: Airborne measurements of the photolysis frequency of NO_2 , *J. Geophys. Res.*, 101, 18613–18627, 1996.
- Wang, T., Carroll, M. A., Albercook, G. M., Owens, K. R., Duderstadt, K. A., Markevitch, A. N., Parrish, D. D., Holloway, J. S., Fehsenfeld, F. C., Forbes, G., and Ogren, J.: Ground-based measurements of NO_x and total reactive oxidized nitrogen (NO_y) at Sable Island, Nova Scotia, during the NARE 1993 summer intensive, *J. Geophys. Res.*, 101, 28991–29004, 1996.
- Weinheimer, A. J., Walega, J. G., Ridley, B. A., Gary, B. L., Blake, D. R., Blake, N. J., Rowland, F. S., Sachse, G. W., Anderson, B. E., and Collins, J. E.: Meridional distributions of NO_x , NO_y , and other species in the lower stratosphere and upper troposphere during AASE II, *Geophys. Res. Lett.*, 21, 2583–2586, 1994.
- Wilson, K. L. and Birks, J. W.: Mechanism and elimination of a water vapor interference in the measurement of ozone by UV absorbance, *Environmental Science and Technology*, 40, 6361–6367, 2006.
- Wisthaler, A., Hansel, A., Dickerson, R. R., and Crutzen, P. J.: Organic trace gas measurements by PTR-MS during INDOEX 1999, *J. Geophys. Res.*, 107, 8024, doi:10.1029/2001JD000576, 2002.
- Wofsy, S. C., Sachse, G. W., Gregory, G. L., Blake, D. R., Bradshaw, J. D., Sandholm, S. T., Singh, H. B., Barrick, J. A., Harriss, R. C., Talbot, R. W., Shipham, M. A., Browell, E. V., Jacob, D. J., and Logan, J. A.: Atmospheric chemistry in the Arctic and Subarctic: Influence of natural fires, industrial emissions, and stratospheric inputs, *J. Geophys. Res.*, 97, 16731–16746, 1992.
- Wofsy, S. C., Fan, S.-M., Blake, D. R., Bradshaw, J. D., Sandholm, S. T., Singh, H. B., Sachse, G. W., and Harriss, R. C.: Factors influencing atmospheric composition over subarctic North America during summer, *J. Geophys. Res.*, 99, 1887–1897, 1994.