



Supplement of

Comparison of the HadGEM2 climate-chemistry model against in situ and SCIAMACHY atmospheric methane data

G. D. Hayman et al.

Correspondence to: G. D. Hayman (garr@ceh.ac.uk)

Supplementary material

1 Emission Inventories

1.1 Wetlands and Wetland Emissions

The methane emissions from wetlands used in this study were based on an offline global monthly run of the JULES land-surface model for $0.5^\circ \times 0.5^\circ$ terrestrial grid squares for the period from January 1980 to December 2010, using CRU-NCEP driving meteorological data (Viovy and Ciais, 2009). JULES (Joint UK land earth simulator) is a physically-based model that contains description of water, energy and carbon balances and temperature, moisture and carbon stores (Best et al., 2011; Clark et al., 2011). JULES (v3.1) uses a methane wetland emission parameterization, developed and tested by Gedney et al. (2004) for use at large spatial scales. The wetland parameterization is coupled to the large-scale hydrology scheme of Gedney and Cox (2003), which predicts the distribution of sub-grid scale water table depth and wetland fraction from the overall soil moisture content and the sub-grid scale topography.


Fig. 1 compares maps of the maximum inundation fraction in each grid square for monthly data from January 1993 to December 2007, as derived from (a) the offline JULES run described above and (b) regridding the global inundation extent multi-sensor (GIEMS) product of Prigent et al. (2012). While JULES captures the major wetland areas, it is clear that JULES underestimates the magnitude. The JULES wetlands areas are more extensive spatially, especially over the Amazon, and smoother. Collins et al. (2011) indicated that the greater spatial extent of wetlands over the Amazon was linked to the poor data on the underlying rock topography in this region incorporated into the ancillary topographic index dataset. Changes are being made to the JULES wetland parameterisation and these are currently being evaluated (Gedney, 2014).

In addition to the standard output (denoted *JULES*), a second wetland emission estimate was derived from this JULES offline run by replacing the modelled wetland fraction with the wetland fraction from the regridded GIEMS product (denoted *JULES-GIEMS*). Emission fluxes for the latter could only be derived for the period from January 1993 to December 2007. Fig. 2 shows maps of the annual emissions of methane from wetlands in 2000 derived from the above. The two JULES inventories were separately scaled to give a mean annual emission of 181 Tg CH₄ per annum for the years 1999-2007, the period of the HadGEM2 model runs.

In the *JULES* wetland emission dataset, most of the emissions occur from the tropics, especially from the Amazon. The use of the EO data is to slightly shift the emissions away from the tropics towards the boreal zone. This is more clearly seen in the zonal plots of Fig. 3, which compare the latitudinal distributions of the methane emissions derived from (a) the *JULES* and the *JULES-GIEMS* runs for the year 2000; (b) the inventory of Fung et al. (1991) for wetlands (=wetlands+bogs+swamps+tundra) and for wetlands and rice paddy fields. The JULES methane inventories have more emissions in the tropics compared to the wetlands dataset of Fung et al. (1991).

The EO wetland inundation product does not discriminate between natural wetlands and managed water areas such as rice paddy fields. A correction was made for the rice paddy fields in the EO wetland product using information on the area of cultivation of rice, from both irrigated and rain-fed cultivation (Portmann et al., 2010). The left-hand panel (a) has no correction to the EO wetland fraction. The right-hand panel (b) shows the effect of correcting the EO wetland inundation product for the area of rice paddy fields. The effect of correcting for the methane emissions from rice paddy fields is most noticeable in the latitude band between 10° and 30° N. The corrected wetland emission dataset was used in the HadGEM model run and an additional term of 40 Tg CH₄ per annum was added to the non-wetland methane sources to represent emissions from rice paddy fields.

(a)



(b)





Figure 1: Maps of the maximum inundation fraction in each grid square for monthly data from January 1993 to December 2007, as derived from (a) an offline JULES run using the standard configuration and (b) the Earth Observation Global Inundation Extent Multi-Sensor (GIEMS) product of Prigent et al. (2012).

(a)



(b)




Figure 2: Maps of the annual emissions of methane from wetlands in 2000 from (a) an offline JULES run using the standard configuration and the modelled wetland fraction and (b) the same offline JULES run in which the modelled wetland fraction is replaced with the EO-derived wetland fraction.




Figure 3: The latitudinal distributions of (i) the wetland methane emissions from JULES for the year 2000, denoted *JULES* (blue lines); (ii) the wetland methane emissions from JULES for the year 2000 with the modelled wetland fraction replaced with an EO-derived wetland fraction, denoted *JULES EO* (green lines); (iii) the methane emissions derived from the inventory of Fung et al. (1991) for wetlands (see text) [solid red lines]; (iv) the methane emissions derived from the inventory of Fung et al. (1991) for wetlands and for rice paddy fields [dotted red lines], and (v) the methane emissions derived from the inventory of Fung et al. (1991), as used in the TRANSCOM- CH_4 model intercomparison of Patra et al. (2011). Panel (a) has no correction to the EO wetland fraction. Panel (b) shows the effect of correcting the EO wetland inundation product for the area of rice paddy fields.

Tab. 1 summarises the latitudinal breakdown of the JULES-based global annual methane emissions from wetlands used in this study. The JULES methane inventories have more emissions in the tropics compared to the wetlands dataset of Fung et al. (1991).

Table 1: Latitudinal breakdown of different global annual methane emissions from wetlands for the year 2000.

Emission Dataset	Extratropical SH (90°S–30°S)	Tropical (30°S–30°N)	Extratropical NH (30°N–50°N)	Boreal/Arctic (50°N–90°N)	Global
<i>FUNG</i> Wetlands and Rice paddy fields	5.7	133.2	30.2	90.8	259.9
<i>FUNG</i> Wetlands	5.0	67.1	18.1	90.8	181.0
<i>JULES</i> Wetlands	1.5	167.4	7.7	4.5	181.1
<i>JULES-GIEMS</i> Wetlands: no correction	4.1	132.1	18.3	12.2	166.7
<i>JULES-GIEMS</i> Wetlands: correction	4.9	127.9	17.6	14.8	165.2

1.2 Global Emissions

Year- and month-specific emission datasets were generated for the period from 1997 to 2009 for (a) non-wetland methane sources and (b) the other emitted species in the UKCA standard tropospheric chemistry scheme (CO, NO_x, HCHO, CH₄, C₂H₆, C₃H₈, CH₃CHO, CH₃CHOCH₃ O'Connor et al., 2013). The approach adopted varied depending on the source sector:

- **Anthropogenic:** The emissions from anthropogenic sources were based on the decadal-averaged emission inventories compiled by Lamarque et al. (2010) for the Coupled Carbon Cycle Climate Model Intercomparison Project (C4MIP) undertaken for the fifth IPCC assessment report (AR5). The decadal-averaged inventories were used to derive year-specific emission datasets by scaling the emission totals for the different years and source sectors using sector and species-specific scaling factors based on the annual trends given in various EDGAR time series: (a) version 4.2 for all species (except NMVOCs) and version 4.1 for NMVOCs; (b) v3.2. This approach was also applied to the emissions from aviation (only for oxides of nitrogen) and international shipping.
- **Biomass burning:** Year-specific emission inventories are available from the Global Fire Emissions Database (GFED, v3.1) for the years 1997 to 2009 (van der Werf et al., 2010), on a monthly timestep. The methane emissions were scaled to give the same decadal mean, 25 Tg CH₄ per annum averaged over 1997 to 2005, as used in the study of O'Connor et al. (2013) to evaluate the standard tropospheric chemistry scheme.
- **Other:** Sources such as termites and hydrates for methane and oceanic emissions of CH₄ and other volatile organic compounds were taken from various sources, as described in O'Connor et al. (2013). These datasets contain a single annual cycle, which was assumed to apply for all years.

1.2.1 Methane

Tab. 2 provides a breakdown by source sector of the global annual emissions of methane used in the different HadGEM2 model for the year 2000. Tab. 3 gives the annual global emissions for the different runs from 1997 to 2008.

Table 2: Comparison of the global annual emissions of methane for 2000 (in Tg CH₄ per annum) and their breakdown by major source sector for the HadGEM2 runs using the following wetland emission inventories, *FUNG*, *TRANSCOM-FUNG*, *JULES* and *JULES-GIEMS*.

Emission Sector	<i>FUNG</i> and <i>TRANSCOM-FUNG</i>	<i>JULES</i>	<i>JULES-GIEMS</i>
anthropogenic	299.2	299.2	299.2
shipping	0.4	0.4	0.4
wetlands	181.1	181.2	165.4
biomass burning	15.6	15.6	15.6
termites	20.0	20.0	20.0
hydrates & oceanic emissions	25.0	25.0	25.0
TOTAL	541.3	541.4	525.6

Note: The anthropogenic term includes 40 Tg CH₄ per annum of emissions from rice paddy fields.

Table 3: Time series of the global annual emissions of methane from 1997 to 2008 (in Tg per annum) used in the HadGEM2 runs using the following wetland emission inventories, *FUNG*, *TRANSCOM-FUNG*, *JULES* and *JULES-GIEMS*.

Year	<i>FUNG</i> and <i>TRANSCOM-FUNG</i>	<i>JULES</i>	<i>JULES-GIEMS</i>
1997	578.4	572.6	579.7
1998	556.0	554.1	569.4
1999	543.8	542.1	543.2
2000	541.3	541.4	525.6
2001	541.1	538.2	530.3
2002	551.4	550.8	562.4
2003	548.0	547.4	550.1
2004	546.9	547.6	540.7
2005	549.1	549.8	548.5
2006	554.2	556.8	565.0
2007	547.1	548.6	558.2
2008	542.4	545.7	545.1
2009	542.2	545.9	548.9

Fig. 4-7 show plots of (a) the time series of the annual methane emissions from wetlands and (b) the mean annual cycle of these emissions for all land surfaces and for the 11 terrestrial TRANSCOM regions for the HadGEM2 runs (*FUNG*, *TRANSCOM-FUNG*, *JULES*, *JULES-GIEMS*).

Table 4 shows the contribution of different methane emission sources within a 20° x 20° box centred on two sites: Barrow (Alaska) and Plateau Assy.




Figure 4: Time series of the annual emissions from 1993 to 2007 globally and for the 11 terrestrial TRANSCOM regions for the dataset of Fung et al. (1991). Panel (b) shows the corresponding annual cycles.




Figure 5: Time series of the annual emissions from 1993 to 2007 globally and for the 11 terrestrial TRANSCOM regions for the dataset of Fung et al. (1991). Panel (b) shows the corresponding annual cycles.




Figure 6: Time series of the annual emissions from 1993 to 2007 globally and for the 11 terrestrial TRANSCOM regions for the JULES wetland emission flux. Panel (b) shows the corresponding annual cycles.




Figure 7: Time series of the annual emissions from 1993 to 2007 globally and for the 11 terrestrial TRANSCOM regions for the JULES wetland emission flux driven with the GIEMS EO product. Panel (b) shows the corresponding annual cycles.

Table 4: Contribution of different methane emission sources ($\text{Tg CH}_4 \text{ yr}^{-1}$) within a $20^\circ \times 20^\circ$ box centred on Barrow (Alaska) and Plateau Assy. The percentage contribution is shown in brackets.

(1) Barrow

a) Wetland Emission Inventory: Fung (as used here)

Year	Anthropogenic	Rice	Biomass Burning	Wetlands	Other	Total
1999	0.50 (12.1)	0.00 (0.0)	0.01 (0.3)	3.61 (87.3)	0.01 (0.3)	4.13
2000	0.49 (11.9)	0.00 (0.0)	0.02 (0.5)	3.61 (87.2)	0.01 (0.3)	4.13
2001	0.49 (11.8)	0.00 (0.0)	0.00 (0.0)	3.61 (87.8)	0.01 (0.4)	4.11
2002	0.48 (11.1)	0.00 (0.0)	0.19 (4.4)	3.61 (84.1)	0.01 (0.3)	4.29
2003	0.46 (11.3)	0.00 (0.0)	0.03 (0.6)	3.61 (87.7)	0.01 (0.3)	4.11
2004	0.45 (10.3)	0.00 (0.0)	0.27 (6.1)	3.61 (83.2)	0.01 (0.3)	4.33
2005	0.43 (9.9)	0.00 (0.0)	0.31 (7.2)	3.61 (82.6)	0.01 (0.3)	4.37
2006	0.42 (10.3)	0.00 (0.0)	0.02 (0.4)	3.61 (89.0)	0.01 (0.4)	4.05
2007	0.40 (9.9)	0.00 (0.0)	0.03 (0.8)	3.61 (88.9)	0.01 (0.4)	4.06

b) Wetland Emission Inventory: Fung (as used in the TRANSCOM-CH₄ model intercomparison and scaled to 181 Tg CH₄ yr⁻¹)

Year	Anthropogenic	Rice	Biomass Burning	Wetlands	Other	Total
1999	0.50 (28.1)	0.00 (0.0)	0.01 (0.8)	1.25 (70.3)	0.01 (0.8)	1.77
2000	0.49 (27.7)	0.00 (0.0)	0.02 (1.2)	1.25 (70.2)	0.01 (0.8)	1.77
2001	0.49 (27.8)	0.00 (0.0)	0.00 (0.0)	1.25 (71.4)	0.01 (0.8)	1.74
2002	0.48 (24.8)	0.00 (0.0)	0.19 (9.8)	1.25 (64.6)	0.01 (0.7)	1.93
2003	0.46 (26.5)	0.00 (0.0)	0.03 (1.5)	1.25 (71.2)	0.01 (0.8)	1.75
2004	0.45 (22.7)	0.00 (0.0)	0.27 (13.5)	1.25 (63.1)	0.01 (0.7)	1.97
2005	0.43 (21.5)	0.00 (0.0)	0.31 (15.7)	1.25 (62.1)	0.01 (0.7)	2.01
2006	0.42 (24.6)	0.00 (0.0)	0.02 (1.0)	1.25 (73.6)	0.01 (0.8)	1.69
2007	0.40 (23.8)	0.00 (0.0)	0.03 (1.9)	1.25 (73.5)	0.01 (0.8)	1.69

c) Wetland Emission Inventory: JULES

Year	Anthropogenic	Rice	Biomass Burning	Wetlands	Other	Total
1999	0.50 (90.8)	0.00 (0.0)	0.01 (2.6)	0.02 (4.0)	0.01 (2.6)	0.55
2000	0.49 (87.9)	0.00 (0.0)	0.02 (3.8)	0.03 (5.7)	0.01 (2.6)	0.56
2001	0.49 (90.7)	0.00 (0.0)	0.00 (0.0)	0.04 (6.6)	0.01 (2.7)	0.53
2002	0.48 (69.1)	0.00 (0.0)	0.19 (27.3)	0.01 (1.5)	0.01 (2.1)	0.69
2003	0.46 (88.5)	0.00 (0.0)	0.03 (5.1)	0.02 (3.7)	0.01 (2.7)	0.52
2004	0.45 (61.3)	0.00 (0.0)	0.27 (36.4)	0.00 (0.3)	0.01 (2.0)	0.73
2005	0.43 (56.2)	0.00 (0.0)	0.31 (41.0)	0.01 (0.9)	0.01 (1.9)	0.77
2006	0.42 (90.6)	0.00 (0.0)	0.02 (3.7)	0.01 (2.6)	0.01 (3.1)	0.46
2007	0.40 (88.6)	0.00 (0.0)	0.03 (6.9)	0.01 (1.3)	0.01 (3.2)	0.46

d) Wetland Emission Inventory: JULES-GIEMS

Year	Anthropogenic	Rice	Biomass Burning	Wetlands	Other	Total
1999	0.50 (81.7)	0.00 (0.0)	0.01 (2.3)	0.08 (13.6)	0.01 (2.4)	0.61
2000	0.49 (79.7)	0.00 (0.0)	0.02 (3.5)	0.09 (14.5)	0.01 (2.3)	0.62
2001	0.49 (81.5)	0.00 (0.0)	0.00 (0.0)	0.10 (16.1)	0.01 (2.4)	0.60
2002	0.48 (61.0)	0.00 (0.0)	0.19 (24.1)	0.10 (13.1)	0.01 (1.8)	0.78
2003	0.46 (76.6)	0.00 (0.0)	0.03 (4.4)	0.10 (16.6)	0.01 (2.4)	0.60
2004	0.45 (52.9)	0.00 (0.0)	0.27 (31.4)	0.12 (14.0)	0.01 (1.7)	0.85
2005	0.43 (49.4)	0.00 (0.0)	0.31 (36.1)	0.11 (12.9)	0.01 (1.6)	0.87
2006	0.42 (75.1)	0.00 (0.0)	0.02 (3.1)	0.11 (19.2)	0.01 (2.6)	0.55
2007	0.40 (73.0)	0.00 (0.0)	0.03 (5.7)	0.10 (18.7)	0.01 (2.6)	0.55

85 Table 4 (continued): Contribution of different methane emission sources ($\text{Tg CH}_4 \text{ yr}^{-1}$) within a $20^\circ \times 20^\circ$ box centred on Barrow (Alaska) and Plateau Assy. The percentage contribution is shown in brackets.

(2) Plateau Assy

a) Wetland Emission Inventory: Fung (as used here)

Year	Anthropogenic	Rice	Biomass Burning	Wetlands	Other	Total
1999	3.72 (75.1)	0.14 (2.9)	0.03 (0.7)	0.78 (15.7)	0.28 (5.6)	4.96
2000	3.64 (74.5)	0.14 (3.0)	0.05 (0.9)	0.78 (15.9)	0.28 (5.7)	4.89
2001	3.57 (74.1)	0.14 (3.0)	0.05 (1.0)	0.78 (16.2)	0.28 (5.8)	4.81
2002	3.48 (72.1)	0.14 (3.0)	0.15 (3.0)	0.78 (16.1)	0.28 (5.8)	4.83
2003	3.39 (72.8)	0.14 (3.1)	0.06 (1.3)	0.78 (16.8)	0.28 (6.0)	4.65
2004	3.29 (72.4)	0.14 (3.2)	0.05 (1.2)	0.78 (17.1)	0.28 (6.1)	4.55
2005	3.19 (72.0)	0.14 (3.3)	0.04 (0.9)	0.78 (17.6)	0.28 (6.3)	4.43
2006	3.09 (71.3)	0.14 (3.3)	0.04 (1.0)	0.78 (18.0)	0.28 (6.4)	4.34
2007	3.01 (70.6)	0.14 (3.4)	0.06 (1.3)	0.78 (18.2)	0.28 (6.5)	4.27

b) Wetland Emission Inventory: Fung (as used in the TRANSCOM-CH₄ model intercomparison and scaled to 181 Tg CH₄ yr⁻¹)

Year	Anthropogenic	Rice	Biomass Burning	Wetlands	Other	Total
1999	3.72 (66.3)	0.14 (2.6)	0.03 (0.6)	1.44 (25.6)	0.28 (5.0)	5.61
2000	3.64 (65.7)	0.14 (2.6)	0.05 (0.8)	1.44 (25.9)	0.28 (5.0)	5.55
2001	3.57 (65.2)	0.14 (2.6)	0.05 (0.9)	1.44 (26.3)	0.28 (5.1)	5.47
2002	3.48 (63.5)	0.14 (2.6)	0.15 (2.7)	1.44 (26.2)	0.28 (5.1)	5.49
2003	3.39 (63.8)	0.14 (2.7)	0.06 (1.2)	1.44 (27.1)	0.28 (5.2)	5.31
2004	3.29 (63.2)	0.14 (2.8)	0.05 (1.0)	1.44 (27.6)	0.28 (5.3)	5.20
2005	3.19 (62.7)	0.14 (2.8)	0.04 (0.8)	1.44 (28.2)	0.28 (5.5)	5.08
2006	3.09 (61.9)	0.14 (2.9)	0.04 (0.9)	1.44 (28.8)	0.28 (5.6)	4.99
2007	3.01 (61.2)	0.14 (2.9)	0.06 (1.1)	1.44 (29.1)	0.28 (5.6)	4.93

c) Wetland Emission Inventory: JULES

Year	Anthropogenic	Rice	Biomass Burning	Wetlands	Other	Total
1999	3.72 (83.1)	0.14 (3.2)	0.03 (0.7)	0.30 (6.7)	0.28 (6.2)	4.48
2000	3.64 (82.7)	0.14 (3.3)	0.05 (1.0)	0.29 (6.7)	0.28 (6.3)	4.40
2001	3.57 (82.3)	0.14 (3.3)	0.05 (1.1)	0.30 (6.9)	0.28 (6.4)	4.33
2002	3.48 (80.0)	0.14 (3.3)	0.15 (3.4)	0.30 (6.9)	0.28 (6.4)	4.35
2003	3.39 (81.3)	0.14 (3.5)	0.06 (1.5)	0.29 (7.0)	0.28 (6.7)	4.17
2004	3.29 (80.8)	0.14 (3.5)	0.05 (1.3)	0.30 (7.5)	0.28 (6.8)	4.07
2005	3.19 (80.2)	0.14 (3.6)	0.04 (1.0)	0.32 (8.1)	0.28 (7.0)	3.97
2006	3.09 (80.0)	0.14 (3.7)	0.04 (1.1)	0.31 (8.0)	0.28 (7.2)	3.87
2007	3.01 (79.1)	0.14 (3.8)	0.06 (1.5)	0.32 (8.4)	0.28 (7.3)	3.81

d) Wetland Emission Inventory: JULES-GIEMS

Year	Anthropogenic	Rice	Biomass Burning	Wetlands	Other	Total
1999	3.72 (78.1)	0.14 (3.0)	0.03 (0.7)	0.59 (12.3)	0.28 (5.8)	4.76
2000	3.64 (78.0)	0.14 (3.1)	0.05 (1.0)	0.56 (11.9)	0.28 (6.0)	4.67
2001	3.57 (76.8)	0.14 (3.1)	0.05 (1.0)	0.61 (13.1)	0.28 (6.0)	4.64
2002	3.48 (74.5)	0.14 (3.1)	0.15 (3.1)	0.63 (13.4)	0.28 (5.9)	4.68
2003	3.39 (77.6)	0.14 (3.3)	0.06 (1.4)	0.50 (11.4)	0.28 (6.4)	4.37
2004	3.29 (75.9)	0.14 (3.3)	0.05 (1.3)	0.57 (13.2)	0.28 (6.4)	4.34
2005	3.19 (75.7)	0.14 (3.4)	0.04 (0.9)	0.56 (13.3)	0.28 (6.6)	4.21
2006	3.09 (75.4)	0.14 (3.5)	0.04 (1.0)	0.55 (13.3)	0.28 (6.8)	4.10
2007	3.01 (75.5)	0.14 (3.6)	0.06 (1.4)	0.50 (12.5)	0.28 (7.0)	3.99

90 1.2.2 Other emitted species

Table 5: Breakdown of the global annual emissions by sector and year of those non-methane trace gases that have emissions in the UKCA Standard Troposphere chemistry scheme.

(a) Carbon Monoxide (in Tg CO yr⁻¹)

Year	Anthropogenic	Biomass Burning	Shipping/Aviation	Other	Total
1997	607.5	736.4	1.1	520.0	1864.9
1998	607.5	594.5	1.1	520.0	1723.1
1999	607.5	378.4	1.2	520.0	1507.0
2000	607.5	328.7	1.2	520.0	1457.3
2001	607.5	311.5	1.1	520.0	1440.1
2002	607.5	463.9	1.2	520.0	1592.6
2003	607.5	449.4	1.2	520.0	1578.1
2004	607.5	411.7	1.3	520.0	1540.5
2005	607.5	442.2	1.3	520.0	1571.0
2006	607.5	476.8	1.4	520.0	1605.7
2007	607.5	417.9	1.5	520.0	1546.9

(b) Oxides of Nitrogen (in Tg NO_x as N yr⁻¹)

Year	Anthropogenic	Biomass Burning	Shipping/Aviation	Other	Total
1997	26.1	6.4	6.0	5.9	44.4
1998	26.6	7.1	6.0	5.9	45.6
1999	27.1	4.7	6.4	5.9	44.1
2000	27.7	4.2	6.6	5.9	44.3
2001	28.2	3.9	6.3	5.9	44.2
2002	28.7	5.3	6.5	5.9	46.3
2003	29.4	5.6	6.6	5.9	47.4
2004	29.9	4.9	7.3	5.9	48.0
2005	30.4	5.2	7.3	5.9	48.8
2006	31.0	5.1	7.8	5.9	49.7
2007	31.6	5.0	8.2	5.9	50.7

(c) Formaldehyde (in Tg HCHO yr⁻¹)

Year	Anthropogenic	Biomass Burning	Shipping/Aviation	Other	Total
1997	3.2	8.2	-	-	11.3
1998	3.2	7.8	-	-	11.0
1999	3.2	5.1	-	-	8.2
2000	3.2	4.4	-	-	7.6
2001	3.2	4.1	-	-	7.3
2002	3.2	5.8	-	-	9.0
2003	3.2	5.9	-	-	9.1
2004	3.3	5.4	-	-	8.7
2005	3.3	5.9	-	-	9.3
2006	3.4	5.9	-	-	9.3
2007	3.4	5.8	-	-	9.2

(d) Ethane (in Tg C₂H₆ yr⁻¹)

Year	Anthropogenic	Biomass Burning	Shipping/Aviation	Other	Total
1997	13.9	16.3	0.4	-	30.6
1998	13.9	15.6	0.4	-	29.9
1999	13.9	10.1	0.4	-	24.4
2000	13.9	8.8	0.4	-	23.1
2001	14.0	8.2	0.4	-	22.6
2002	14.0	11.6	0.4	-	26.0
2003	14.2	11.8	0.4	-	26.4
2004	14.5	10.8	0.5	-	25.8
2005	14.6	11.8	0.5	-	26.9
2006	14.9	11.8	0.5	-	27.2
2007	15.1	11.5	0.6	-	27.2

Table 5 (continued): Breakdown of the global annual emissions by sector and year of those non-methane trace gases that have emissions in the UKCA Standard Troposphere chemistry scheme.

(e) Propane (in Tg C₃H₈ yr⁻¹)

Year	Anthropogenic	Biomass Burning	Shipping/Aviation	Other	Total
1997	6.6	8.5	0.7	-	15.7
1998	6.6	8.2	0.7	-	15.4
1999	6.6	5.3	0.7	-	12.5
2000	6.6	4.6	0.7	-	11.9
2001	6.6	4.3	0.7	-	11.6
2002	6.6	6.0	0.7	-	13.3
2003	6.7	6.1	0.7	-	13.6
2004	6.9	5.7	0.8	-	13.3
2005	6.9	6.2	0.8	-	13.9
2006	7.0	6.1	0.9	-	14.0
2007	7.1	6.0	1.0	-	14.1

(f) Acetaldehyde (in Tg CH₃CHO yr⁻¹)

Year	Anthropogenic	Biomass Burning	Shipping/Aviation	Other	Total
1997	-	12.4	-	-	12.4
1998	-	11.9	-	-	11.9
1999	-	7.7	-	-	7.7
2000	-	6.6	-	-	6.6
2001	-	6.2	-	-	6.2
2002	-	8.8	-	-	8.8
2003	-	8.9	-	-	8.9
2004	-	8.2	-	-	8.2
2005	-	9.0	-	-	9.0
2006	-	8.9	-	-	8.9
2007	-	8.7	-	-	8.7

(g) Acetone (in Tg CH₃COCH₃ yr⁻¹)

Year	Anthropogenic	Biomass Burning	Shipping/Aviation	Other	Total
1997	0.7	7.0	-	40.0	47.6
1998	0.7	6.7	-	40.0	47.3
1999	0.7	4.3	-	40.0	45.0
2000	0.7	3.7	-	40.0	44.4
2001	0.7	3.5	-	40.0	44.2
2002	0.7	4.9	-	40.0	45.6
2003	0.7	5.0	-	40.0	45.7
2004	0.7	4.6	-	40.0	45.3
2005	0.7	5.1	-	40.0	45.8
2006	0.7	5.0	-	40.0	45.7
2007	0.7	4.9	-	40.0	45.6

95 Notes to Table 5:

1. For CO, the 'Other' category includes emission terms of (a) 475 Tg CO yr⁻¹ to account for the impact of isoprene emissions from vegetation on atmospheric composition, and (b) 45 Tg yr⁻¹ from oceanic sources.
2. For NO_x, the 'Other' category includes 5.9 Tg NO_x yr⁻¹ from soils. The emissions of NO_x from lightning are calculated interactively (O'Connor et al., 2013).
3. For acetone, the 'Other' category includes 40 Tg yr⁻¹ from vegetation.

100

1.3 Comparison with Estimates from Inverse Modelling

Maps of the global wetland methane emissions are shown for the JULES-based wetland emissions and are compared, as difference maps, with the emissions derived from the inverse modelling
105 study of Bousquet et al. (2011) in Figs. 8 to 13:

- **Fig. 8:** Maps of the global monthly methane emissions from wetlands in 2000 for the *JULES* wetland emission flux.
- **Fig. 9:** Maps of the difference in the global monthly methane emissions from wetlands in 2000 from the *JULES* run and from the inverse modelling study of Bousquet et al. (2011)
110 using the Fung et al. (1991) dataset as the wetland emission prior.
- **Fig. 10:** Maps of the difference in the global monthly methane emissions from wetlands in 2000 from the *JULES* run and from the inverse modelling study of Bousquet et al. (2011) using the Kaplan dataset as the wetland emission prior.
- **Fig. 11:** Maps of the global monthly methane emissions from wetlands in 2000 for the *JULES-GIEMS* wetland emission flux with the modelled wetland fraction replaced with
115 that derived from the GIEMS EO product.
- **Fig. 12:** Maps of the difference in the global monthly methane emissions from wetlands in 2000 from the *JULES-GIEMS* run and from the inverse modelling study of Bousquet et al. (2011) using the Fung et al. (1991) dataset as the wetland emission prior.
- **Fig. 13:** Maps of the difference in the global monthly methane emissions from wetlands in 2000 from the *JULES* run and from the inverse modelling study of Bousquet et al.
120 (2011) using the Kaplan dataset as the wetland emission prior.




Figure 8: Maps of the global monthly wetland methane emissions for 2000 for the JULES wetland emission flux.




Figure 9: Maps of the differences in the global monthly wetland methane emissions for 2000 between the JULES wetland emission flux and the inverse modelling study of Bousquet et al. (2011) using the Fung et al. (1991) dataset as the wetland emission prior.




Figure 10: Maps of the differences in the global monthly wetland methane emissions for 2000 between the JULES wetland emission flux and the inverse modelling study of Bousquet et al. (2011) using the Kaplan dataset as the wetland emission prior.




Figure 11: Maps of the global monthly wetland methane emissions for 2000 for the JULES wetland emission flux driven with the GIEMS EO inundation product.




Figure 12: Maps of the differences in the global monthly wetland methane emissions for 2000 between the JULES wetland emission flux driven with the GIEMS product and the inverse modelling study of Bousquet et al. (2011) using the Fung et al. (1991) dataset as the wetland emission prior.




Figure 13: Maps of the differences in the global monthly wetland methane emissions for 2000 between the JULES wetland emission flux driven with the EO inundation product and the inverse modelling study of Bousquet et al. (2011) using the Kaplan dataset as the wetland emission prior.

2 Comparison with surface and satellite observations

2.1 Surface observations

- 125 The modelled monthly-averaged surface concentrations of atmospheric methane were compared against the monthly-averaged measurements of atmospheric methane dry air mole fractions from the NOAA ESRL GMD Carbon Cycle Cooperative Global Air Sampling Network (Dlugokencky et al., 2012). Fig. 14 shows a map of the locations of the 64 sites used in this work.




Figure 14: Locations of the surface atmospheric methane sampling sites in the NOAA Global Air Sampling Network: ABP - Arembepe, Bahia; ALT - Alert, Nunavut; AMS - Amsterdam Island; ASC - Ascension Island; ASK - Assekrem; AVI - St. Croix, Virgin Islands; AZR - Terceira Island, Azores; BKT - Bukit Kototabang; BME - St. Davids Head, Bermuda; BMW - Tudor Hill, Bermuda; BRW - Barrow, Alaska; BSC - Black Sea, Constanta; CBA - Cold Bay, Alaska; CGO - Cape Grim, Tasmania; CHR - Christmas Island; CMO - Cape Meares, Oregon; CRZ - Crozet Island; EIC - Easter Island; GMI - Mariana Islands; GOZ - Dwejra Point, Gozo; HBA - Halley Station, Antarctica; HPB - Hohenpeissenberg; HUN - Hegyhatsal; ICE - Storhofdi, Vestmannaeyjar; ITN - Grifton, North Carolina; IZO - Tenerife, Canary Islands; KEY - Key Biscayne, Florida; KUM - Cape Kumukahi, Hawaii; KZD - Sary Taukum; KZM - Plateau Assy; LLB - Lac La Biche, Alberta; LLN - Lulin; LMP - Lampedusa; MBC - Mould Bay, Nunavut; MEX - Mex High Altitude Global; MHD - Mace Head, County Galway; MID - Sand Island, Midway; MKN - Mt. Kenya; MLO - Mauna Loa, Hawaii; NMB - Gobabeb; NWR - Niwot Ridge, Colorado; OPW - Olympic Peninsula, Washington; OXK - Ochsenkopf; PAL - Pallas-Sammaltunturi, GAW; PSA - Palmer Station, Antarctica; PTA - Point Arena, California; RPB - Ragged Point; SEY - Mahe Island; SGP - Southern Great Plains, Oklahoma; SHM - Shemya Island, Alaska; SMO - Tutuila; SPO - South Pole, Antarctica; STM - Ocean Station M; SUM - Summit; SYO - Syowa Station, Antarctica; TAP - Tae-ahn Peninsula; TDF - Tierra Del Fuego, Ushuaia; THD - Trinidad Head, California; UTA - Wendover, Utah; UUM - Ulaan Uul; WIS - WIS Station, Negev Desert; WKT - Moody, Texas; WLG - Mt. Waliguan; ZEP - Ny-Alesund, Svalbard.

130 In Figs. 15-18, we show plots of the observed and modelled atmospheric methane concentrations between 2000 and 2010 at 16 of these sites, covering both northern and southern hemisphere locations, for the different model runs: *FUNG*, *JULES* and *JULES-GIEMS*. We present the metrics derived from the statistical analysis of the observed and modelled concentrations in Tab. 6.




Figure 15: Comparison of the surface atmospheric methane mixing ratio (in ppb) as observed (black) and from the HadGEM2 run using the *FUNG* wetland emission inventory (red) at selected sites between 2000 and 2010.




Figure 16: Comparison of the surface atmospheric methane mixing ratio (in ppb) as observed (black) and from the HadGEM2 run using the FUNG wetland emission inventory (red) at selected sites between 2000 and 2010.




Figure 17: Comparison of the surface atmospheric methane mixing ratio (in ppb) as observed (black) and from the HadGEM2 run using the *JULES* wetland emission inventory (red) at selected sites between 2000 and 2010.




Figure 18: Comparison of the surface atmospheric methane mixing ratio (in ppb) as observed (black) and from the HadGEM2 run using the *JULES-GIEMS* wetland emission inventory (red) at selected sites between 2000 and 2010.

Table 6: Statistical outputs from the analysis of the observed and modelled surface methane concentrations at Barrow and the South Pole for the HadGEM2 runs (*FUNG*, *TRANSCOM-FUNG*, *JULES* and *JULES-GIEMS*).

(a) Barrow

Statistic/Metric	<i>FUNG</i>	<i>TRANSCOM-FUNG</i>	<i>JULES</i>	<i>JULES-GIEMS</i>
Number of valid data pairs	119	119	119	119
Linear regression – slope	-12.15	-2.53	1.70	1.68
Linear regression – intercept	24841.0	6685.9	-1308.7	-1262.0
Coefficient of determination (R^2)	0.46	0.29	0.39	0.49
Mean of Observations (in ppb)	1867.8	1867.8	1867.8	1867.8
Mean of Modelled Conc. (in ppb)	2147.1	1969.2	1861.4	1882.1
Mean normalised bias	0.15	0.06	-0.004	0.008
Number of modelled results within a factor of 2 of that observed	1.0	1.0	1.0	1.0
Index of Agreement	0.01	0.09	0.60	0.64
Hit Rate	0.55	0.63	1.00	1.00
Root Mean Square Error (RMSE in ppb)	425.8	136.4	39.2	34.8
Coefficient of Variation in RMSE	0.29	0.07	0.02	0.02

(b) South Pole

Statistic/Metric	<i>FUNG</i>	<i>TRANSCOM-FUNG</i>	<i>JULES</i>	<i>JULES-GIEMS</i>
Number of valid data pairs	119	119	119	119
Linear regression – slope	0.72	1.14	1.07	0.96
Linear regression – intercept	471.7	-228.7	-97.5	79.5
Coefficient of determination (R^2)	0.60	0.74	0.62	0.59
Mean of Observations (in ppb)	1729.2	1729.2	1729.2	1729.2
Mean of Modelled Conc. (in ppb)	1723.8	1740.2	1746.4	1733.1
Mean normalised bias	-0.003	0.01	0.01	0.002
Number of modelled results within a factor of 2 of that observed	1.0	1.0	1.0	1.0
Index of Agreement	0.84	0.79	0.66	0.85
Hit Rate	1.0	1.0	1.0	1.0
Root Mean Square Error (RMSE in ppb)	9.5	13.8	20.0	10.4
Coefficient of Variation in RMSE	0.005	0.008	0.012	0.006

2.2 Satellite observations

- ¹³⁵ The modelled 4-D methane mass mixing ratio fields (longitude, latitude, altitude, time) were converted into 3-D fields (longitude, latitude, time) of the mean atmospheric column methane mixing ratio, using the SCIAMACHY averaging kernels (equation 4 in Schneising et al., 2009).

$$XCH_4^{\text{model}} = \frac{1}{p_0} \sum_l \left[\overline{XCH}_4^l + AK^l \left(XCH_4^{\text{model}\ l} - \overline{XCH}_4^l \right) \right] \Delta p^l \quad (1)$$

¹⁴⁰ where l is the index of the vertical layer, AK^l the averaging kernel, \overline{XCH}_4^l the a-priori mole fraction (1750 ppb below 6km and decreasing above) and $XCH_4^{\text{model}\ l}$ is the modelled mole fraction of layer l . Δp^l is the pressure difference between the upper and lower boundary of layer l and p_0 denotes the surface pressure.

¹⁴⁵ As discussed in the main paper, the modelled atmospheric methane columns underpredicted the columns in the SCIAMACHY dataset. To overcome this limitation, the methane concentration in the upper model layers were replaced initially using data from the thermal infrared channel of the Tropospheric Emission Spectrometer and subsequently with output from a TOM-CAT model run, assimilated with ACE and HALOE data. Fig. 19 presents contour maps of the annual mean atmospheric column measurements of methane as derived from the unconstrained (panel a) and constrained HadGEM2 (panel b) run and from the monthly regridded SCIAMACHY data (v2.3, panel c). The SCIAMACHY and model atmospheric methane columns have been sampled at common, valid space and time points. A land mask was applied to remove all data over the ocean as the SCIAMACHY data are only available over the oceans for the period between 2003 and 2005.

¹⁵⁰ Figure 10 in the main paper shows time series and annual cycles of the area-weighted mean atmospheric column methane mixing ratios between January 2003 and December 2007 from the SCIAMACHY data and the four constrained HadGEM2 runs for all land surface points and for the 11 terrestrial TRANSCOM regions. In Fig. 20, we show similar time series and annual cycle plots using the unconstrained HadGEM2 model outputs. We discuss these figures further in the Discussion in the main paper.




Figure 19: Contour maps of the average atmospheric column methane mixing ratio for 2003-2007 as derived from the unconstrained (panel a) and constrained HadGEM2 (panel b) run and from monthly regridded SCIAMACHY data (v2.3, panel c), sampled at common space and time points.




Figure 20: Time series of the area-weighted average atmospheric column methane mixing ratio from January 2003 to December 2007, as derived from monthly regressed SCIAMACHY data (v2.3) and from the unconstrained HadGEM2 runs using (1) the FUNG (red), (2) TRANSCOM FUNG (magenta), (3) the JULES (blue), and (4) the EDGAR v3.2 (E3.2) anthropogenic methane emission inventories and the EDGAR v3.2 (E3.2) anthropogenic methane emission time series, sampled at co-located space and time points for all land surface points and for the 11 terrestrial TRANSCOM regions (a). (b) shows the corresponding annual cycles. The index of agreement (IOA) is shown for each run (see Sect.3 of the Supplement for the definition of the IOA).

3 Performance metrics

- 160 Evaluation of model performance through statistical metrics focuses on measures that compare a set of N predicted concentrations P_i with their counterpart observed concentrations O_i , where i refers to a given time and/or location. Standard metrics used for air quality model performance evaluation are detailed in numerous papers (e.g., Yu et al., 2006; Dennis et al., 2010) and references therein) and only the ones that are used in our work (main text and Supplement) 165 are reported hereafter. The means of N predictions and observations are defined as

$$P_{mean} = \frac{1}{N} \sum_{i=1}^N P_i \quad \text{and} \quad O_{mean} = \frac{1}{N} \sum_{i=1}^N O_i \quad (2)$$

respectively. The standard deviations of N predictions and observations are defined as

$$\sigma_P = \sqrt{\frac{1}{N} \sum_{i=1}^N (P_i - P_{mean})^2} \quad \text{and} \quad \sigma_O = \sqrt{\frac{1}{N} \sum_{i=1}^N (O_i - O_{mean})^2} \quad (3)$$

respectively. The variables a, b, c, and d are used to calculate the categorical statistics A, B, H, and FAR represent:

- a - all the exceedances that did not occur
- 170 b - all the exceedances that did occur
- c - all the exceedances that were not predicted and not observed, and
- d - all the exceedances that were not predicted but observed, respectively

The following performance metrics were derived:

1. Accuracy (no unit, in %):

$$\text{Accuracy} = \frac{b+c}{a+b+c+d} \times 100$$

2. Bias (no unit):

$$\text{Bias} = \frac{a+b}{b+d}$$

3. Correlation coefficient (no units):

$$\text{Correlation coefficient} = \frac{1}{\sigma_P \sigma_O} \sum_{i=1}^N (P_i - P_{mean})(O_i - O_{mean})$$

4. Fraction of predictions within a factor of 2 of observations (no unit, %):

$$\text{Fraction of Prediction} = \left[\frac{1}{N} \sum_{i=1}^N 1 \mid (0.5 \leq \frac{P_i}{O_i} \leq 2) \right] \times 100$$

5. Fractional Bias (no unit, range [-2,2]):

$$\text{Fractional Bias} = \sum_{i=1}^N (P_i - O_i) / \sum_{i=1}^N (P_i + O_i)/2$$

6. Hit Rate (no unit, in %):

$$\text{Hit Rate} = \frac{b}{b+d} \times 100$$

7. Index of Agreement (no unit, range [0,1]):

$$\text{Index of Agreement} = 1 - \sum_{i=1}^N (P_i - O_i)^2 / \sum_{i=1}^N (|P_i - O_{mean}|^2 + |O_i - O_{mean}|^2)$$

8. Mean Bias (in unit of concentration):

$$\text{Mean Bias} = \frac{1}{N} \sum_{i=1}^N (P_i - O_i)$$

9. Mean Error (in unit of concentration):

$$\text{Mean Error} = \frac{1}{N} \sum_{i=1}^N |P_i - O_i|$$

10. Mean Fractional Bias (no unit, range [-200%, 200%]):

$$\text{Mean Fractional Bias} = \sum_{i=1}^N \frac{(P_i - O_i)}{(P_i + O_i)/2} \times 100$$

11. Mean Fractional Error (no unit, range [0, 200%]):

$$\text{Mean Fractional Error} = \sum_{i=1}^N \frac{|P_i - O_i|}{(P_i + O_i)/2} \times 100$$

12. Mean Normalised Bias (no unit, range [-100%, ∞]):

$$\text{Mean Normalised Bias} = \frac{1}{N} \sum_{i=1}^N \frac{(P_i - O_i)}{O_i} \times 100$$

13. Mean Normalised Error (no unit, in %):

$$\text{Mean Normalised Error} = \frac{1}{N} \sum_{i=1}^N \frac{|P_i - O_i|}{O_i} \times 100$$

14. Normalised Mean Bias (no unit, range [-1, ∞]):

$$\text{Normalised Mean Bias} = \sum_{i=1}^N (P_i - O_i) / \sum_{i=1}^N O_i$$

15. Root Mean Square Error (in unit of concentration):

$$\text{Root Mean Square Error} = \sqrt{\frac{1}{N} \sum_{i=1}^N (P_i - O_i)^2}$$

16. Skill Variance (no units):

$$\text{Skill variance} = \frac{\sigma_P}{\sigma_O}$$

References

- 175 Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description - Part 1: Energy and water fluxes, *Geoscientific Model Development*, 4, 677–699, doi:10.5194/gmd-4-677-2011, <http://www.geosci-model-dev.net/4/677/2011/>, 2011.
- 180 Bousquet, P., Ringeval, B., Pison, I., Dlugokencky, E. J., Brunke, E.-G., Carouge, C., Chevallier, F., Fortems-Cheiney, A., Frankenberg, C., Hauglustaine, D. A., Krummel, P. B., Langenfelds, R. L., Ramonet, M., Schmidt, M., Steele, L. P., Szopa, S., Yver, C., Viovy, N., and Ciais, P.: Source attribution of the changes in atmospheric methane for 2006–2008, *Atmospheric Chemistry and Physics*, 11, 3689–3700, doi:10.5194/acp-11-3689-2011, <http://www.atmos-chem-phys.net/11/3689/2011/>, 2011.
- 185 Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description - Part 2: Carbon fluxes and vegetation dynamics, *Geoscientific Model Development*, 4, 701–722, doi:10.5194/gmd-4-701-2011, <http://www.geosci-model-dev.net/4/701/2011/>, 2011.
- 190 Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System model - HadGEM2, *Geoscientific Model Development*, 4, 1051–1075, doi:10.5194/gmd-4-1051-2011, <http://www.geosci-model-dev.net/4/1051/2011/>, 2011.
- 195 Dennis, R., Fox, T., Fuentes, M., Gilliland, A., Hanna, S., Hogrefe, C., Irwin, J., Rao, S., Scheffe, R., Schere, K., Steyn, D., and Venkatram, A.: A framework for evaluating regional-scale numerical photochemical modeling systems, *Environmental Fluid Mechanics*, 10, 471–489, doi:10.1007/s10652-009-9163-2, <http://dx.doi.org/10.1007/s10652-009-9163-2>, 2010.
- 200 Dlugokencky, E. J., Lang, P. M., Crotwell, A. M., and Masarie, K. A.: Atmospheric Methane Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Co-operative Global Air Sampling Network, 1983–2011, Version: 2012-09-24, <ftp://ftp.cmdl.noaa.gov/ccg/ch4/flask/event/>, 2012.
- 205 Fung, I., John, J., Lerner, J., Matthews, E., Prather, M., Steele, L. P., and Fraser, P. J.: Three-dimensional model synthesis of the global methane cycle, *Journal of Geophysical Research: Atmospheres*, 96, 13 033–13 065, doi:10.1029/91JD01247, <http://dx.doi.org/10.1029/91JD01247>, 1991.
- Gedney, N.: Unpublished results from JULES, 2014.
- 210 Gedney, N. and Cox, P. M.: The Sensitivity of Global Climate Model Simulations to the Representation of Soil Moisture Heterogeneity, *Journal of Hydrometeorology*, 4, 1265–1275, doi:10.1175/1525-7541(2003)004<1265:TSOGCM>2.0.CO;2, [http://dx.doi.org/10.1175/1525-7541\(2003\)004<1265:TSOGCM>2.0.CO;2](http://dx.doi.org/10.1175/1525-7541(2003)004<1265:TSOGCM>2.0.CO;2), 2003.
- 215 Gedney, N., Cox, P. M., and Huntingford, C.: Climate feedback from wetland methane emissions, *Geophysical Research Letters*, 31, L20 503, doi:10.1029/2004GL020919, <http://dx.doi.org/10.1029/2004GL020919>, 2004.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., 220 Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmospheric Chemistry and Physics*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, <http://www.atmos-chem-phys.net/10/7017/2010/>, 2010.
- O'Connor, F. M., Johnson, C. E., Morgenstern, O., Abraham, N. L., Braesicke, P., Dalvi, M., Folberth, G. A., Sanderson, M. G., Telford, P. J., Young, P. J., Zeng, G., Collins, W. J., and Pyle, J. A.: Evaluation of the new UKCA climate-composition model - Part 2: The Troposphere, *Geoscientific Model Development Discussions*, 6, 1743–1857, doi:10.5194/gmdd-6-1743-2013, <http://www.geosci-model-dev-discuss.net/6/1743/2013/>, 2013.
- Patra, P. K., Houweling, S., Krol, M., Bousquet, P., Belikov, D., Bergmann, D., Bian, H., Cameron-Smith, P., Chipperfield, M. P., Corbin, K., Fortems-Cheiney, A., Fraser, A., Gloor, E., Hess, P., Ito, A., Kawa, S. R., Law, R. M., Loh, Z., Maksyutov, S., Meng, L., Palmer, P. I., Prinn, R. G., Rigby, M., Saito, R., and Wilson, C.: TransCom model simulations of CH₄ and related species: linking transport, surface flux and chemical loss with CH₄ variability in the troposphere and lower stratosphere, *Atmospheric Chemistry and Physics*, 11, 12 813–12 837, doi:10.5194/acp-11-12813-2011,

- <http://www.atmos-chem-phys.net/11/12813/2011/>, 2011.
- Portmann, F. T., Siebert, S., and Döll, P.: MIRCA2000 - Global monthly irrigated and rain-fed crop areas around the year 2000: A new high-resolution data set for agricultural and hydrological modeling, *Global Biogeochemical Cycles*, 24, GB1001, doi:10.1029/2008GB003435,
235 <http://dx.doi.org/10.1029/2008GB003435>, 2010.
- Prigent, C., Papa, F., Aires, F., Jimenez, C., Rossow, W. B., and Matthews, E.: Changes in land surface water dynamics since the 1990s and relation to population pressure, *Geophysical Research Letters*, 39, L08 403, doi:10.1029/2012GL051276, <http://dx.doi.org/10.1029/2012GL051276>, 2012.
- Schneising, O., Buchwitz, M., Burrows, J. P., Bovensmann, H., Bergamaschi, P., and Peters, W.:
240 Three years of greenhouse gas column-averaged dry air mole fractions retrieved from satellite - Part 2: Methane, *Atmospheric Chemistry and Physics*, 9, 443–465, doi:10.5194/acp-9-443-2009, <http://www.atmos-chem-phys.net/9/443/2009/>, 2009.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmospheric Chemistry and Physics*, 10, 245 11 707–11 735, doi:10.5194/acp-10-11707-2010, <http://www.atmos-chem-phys.net/10/11707/2010/>, 2010.
- Viovy, N. and Ciais, P.: A combined dataset for ecosystem modelling, available at:
<http://dods.extra.cea.fr/data/p529viov/cruncep/readme.htm> (last access: 10 July 2013), 2009.
- 250 Yu, S., Eder, B., Dennis, R., Chu, S.-H., and Schwartz, S. E.: New unbiased symmetric metrics for evaluation of air quality models, *Atmospheric Science Letters*, 7, 26–34, doi:10.1002/asl.125, <http://dx.doi.org/10.1002/asl.125>, 2006.