

Supplement of Atmos. Chem. Phys., 15, 11537–11555, 2015
<http://www.atmos-chem-phys.net/15/11537/2015/>
doi:10.5194/acp-15-11537-2015-supplement
© Author(s) 2015. CC Attribution 3.0 License.



Atmospheric
Chemistry
and Physics
Open Access
EGU

Supplement of

The importance of Asia as a source of black carbon to the European Arctic during springtime 2013

D. Liu et al.

Correspondence to: D. Liu (dantong.liu@manchester.ac.uk)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.






Fig. S1. (a) The vertical profiles of CO and O₃ (mean $\pm\sigma$) in 400m altitude bins; (b) the CO/O₃ ratio as a function of height; (c) the lowest 5th percentile of CO in each altitude bin, with the black line showing the CO background values as used to calculate ΔCO in this study.


A




B



C



D



E




Fig. S2. In each graph, top panel shows the altitude and FLEXPART FPES fractions for each source region, the plume locations during SLR are marked as Roman numerals along the flight altitude track; the bottom panel shows the time series of BC mass, CO concentration, mass of organic matter and sulphate. (a) to (e) represents the data for different flights. Note that the AMS data was not available for the flight B759 and B760.




Fig. S3. The BC anthropogenic emission inventories in Mar. 2010 from HTAPv2 for the following sectors (a) residential activity, (b) transport, (c) industry, (d) energy and (e) flaring (from ECLIPSE global emission inventory). Note the (b), (c) and (d) use the same legend colour scale.




Fig. S4. A summary of all plumes and profiles. Columns (a) to (e) show the BC mass, ΔCO , BC D_p/D_c , the mass of sulphate and organic matter respectively. The larger markers with error bar show data collected in plumes during straight and level runs and the small markers show data collected during profiles. Each marker shape denotes the flight index and these shapes are applied to all of the plumes and profiles. Note for Clean Air periods, the altitude scale extends to only 3km.




Fig. S5. Average characteristics of sampled air parcel plumes arriving at receptor locations along 12 day backward trajectory pathways, using the identical legends of Fig. 9. A1 is EU plume in the UT (corresponding to plume B761 iv in Table 2); A2 is EU plume in the LT (B761 ii); B1 is NA plume in the UT (B760 i) and B2 is NA plume in the LT (B762 i).

Plume index	Source region	Latitude/Region	BC (ng sm ⁻³)						ΔCO (ppbv)					
			RE	TR	IN	EN	OBB	Fl	RE	TR	IN	EN	OBB	Fl
B759 plumes														
i	AS	30-40°N; Japan, E China	1.70	0.53	1.04	0.01	0.15	0	0.11	0.04	0.09	0.00	0.02	0
ii	AS	30-40°N; Japan, E China	26.58	13.96	18.44	0.09	2.57	0	1.89	1.07	2.22	0.06	0.41	0
iii	AS	40-60°N; N China	24.68	8.61	13.35	0.09	0.39	2.6e-4	1.87	0.73	1.56	0.05	0.06	1.2e-6
iv	AS	30-50°N; N and E China, Japan	8.08	5.43	5.52	0.12	0.51	5.2e-4	0.63	0.65	0.63	0.03	0.07	2.5e-6
v	NA	30-50°N; N US	16.86	37.30	19.96	1.56	3.95	1.2e-4	2.89	6.30	1.51	0.15	0.63	0
vi	AS	20-40°N; S and E China, Japan	60.46	28.20	40.36	0.17	13.03	0	4.15	2.19	4.40	0.12	2.11	0
B760 plumes														
i	NA	20-40°N; S US, Mexico	3.17	11.21	6.54	1.27	2.19	0	0.55	1.58	0.22	0.41	0.32	0
ii	AS	40-60°N; E and N China, Japan	17.06	6.25	8.46	0.08	0.28	1.2e-4	1.34	0.56	1.03	0.03	0.04	0
iii	AS	40-75°N; N China, E Russia	10.36	3.27	3.78	0.10	0.20	6.0e-4	0.80	0.29	0.47	0.02	0.03	2.9e-6
iv	NA	20-40°N; S US, Mexico	0.92	2.17	1.03	0.15	0.20	0	0.13	0.31	0.05	0.02	0.03	0
B761 plumes														
i	EU+NA	30-60°N; E US, N Europe	5.45	11.03	5.28	0.58	2.19	0	0.84	1.77	0.32	0.03	0.34	0
ii	EU	50-80°N; W and N Europe	15.80	14.89	3.56	0.55	2.11	8.8e-4	1.11	1.38	0.33	0.05	0.32	3.1e-6
iii	EU	50-80°N; N Europe	5.55	3.36	0.28	0.06	0.03	1.69e-3	0.25	0.23	0.07	0.02	0.00	6.8e-6
iv	NA	35-60°N; E and S US	6.40	9.7	7.432	0.64	0.18	0	0.72	1.3	0.53	0.14	0.03	0
v	EU	50-80°N; N Europe	51.23	43.29	6.375	2.357	1.11	0	3.2	2.2	0.75	0.19	0.12	0
B762 plumes														
i	NA	40-60°N; E and N US	20.38	34.61	20.18	1.82	0.28	2.15e-3	3.4	6.62	1.68	0.07	0.04	1.2e-6
ii	NA	40-60°N; E and N US	3.86	2.90	1.71	0.11	0.11	3.27e-3	0.34	0.35	0.17	0.02	0.02	1.5e-5
iii	NA	40-60°N; E and N US	19.88	18.85	7.08	1.57	0.30	2.66e-3	1.44	1.57	0.57	0.09	0.05	1.2e-5
iv	EU	45-75°N; N and S Europe	54.33	36.42	4.183	10.86	0.41	8.5e-4	3.38	1.98	0.46	0.18	0.03	0
B763 plumes														
i	AS	30-60°N; N and E China, Japan	72.85	33.94	53.79	0.25	1.07	3.47e-3	5.30	2.53	5.72	0.18	0.16	1.7e-6
ii	NA	35-50°N; E US	0.75	1.43	0.84	0.05	0.01	0	0.11	0.23	0.05	0.00	0.00	0
iii	AS	30-55°N; E and N China, Japan	169.84	66.00	106.33	0.48	1.40	7.4e-4	11.95	4.45	10.11	0.33	0.20	3.6e-6

Table S1. BC and CO source attributions for all of the encountered plumes by FLEXPART FPES, in source sectors: residential (RE), transport (TR), industry (IN), energy (EN) and open biomass burning (OBB). The modelled flaring contribution is deemed to be zero if less than $1e^{-3}$ ng m⁻³ for BC or less than $1e^{-7}$ ppbv for CO.

Source origin	BC/ Δ CO _{measured} (ng sm ⁻³ ppbv ⁻¹)	BC/ Δ CO _{source} (ng sm ⁻³ ppbv ⁻¹)	SF _{BC}
B759 plumes			
AS	2.06±1.55	13.14	0.84±0.12
AS	5.66±3.68	10.94	0.62±0.34
AS	2.17±0.78	11.05	0.80±0.07
AS	3.33±2.09	9.76	0.66±0.21
NA	1.95±1.58	6.93	0.72±0.23
AS	2.05±1.66	10.97	0.81±0.15
B760 plumes			
NA		7.92	
AS	2.61±1.76	10.73	0.76±0.16
AS	2.23±0.72	11.00	0.80±0.07
NA		8.20	
B761 plumes			
EU+NA	2.05±0.69	7.44	0.72±0.09
EU	2.12±0.93	11.55	0.82±0.08
EU	4.15±1.78	16.04	0.74±0.11
NA	0.37±0.28	8.95	0.96±0.03
EU	0.26±0.18	16.16	0.97±0.01
B762 plumes			
NA	0.36±0.27	6.54	0.94±0.04
NA	1.60±1.06	9.64	0.83±0.11
NA	1.21±0.73	12.81	0.91±0.06
EU	0.59±0.63	17.61	0.96±0.03
B763 plumes			
AS	1.75±1.39	11.67	0.85±0.12
NA	1.25±0.70	7.90	0.84±0.09
AS	1.94±1.39	12.73	0.85±0.11

Table S2. Measured and modelled BC/ Δ CO, and derived SF_{BC} for each of the plumes.