



*Supplement of*

## **Depletion of atmospheric gaseous elemental mercury by plant uptake at Mt. Changbai, Northeast China**

**Xuewu Fu et al.**

*Correspondence to:* Xinbin Feng (fengxinbin@vip.skleg.cn)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

1

2

3                   **Contents of the Supporting Information**

4

5       1 Supplementary Text.....	Page S1
6       1.1 Nocturnal boundary layer.....	Page S1
7       1.2 Simulations of atmospheric GEM at Mt. Changbai forest using a box model.....	Page S1
8       1.3 The turbulent diffusivity (Kc).....	Page S2
9       Figure S1.....	Page S3
10      Figure S2.....	Page S4
11      Figure S3.....	Page S5
12      Figure S4.....	Page S6
13      Figure S5.....	Page S7
14      Figure S6.....	Page S8
15      Figure S7.....	Page S9
16      Table S1 .....	Page S10
17      Table S2 .....	Page S11
18      Table S3 .....	Page S12
19      Table S4 .....	Page S13
20      Table S5 .....	Page S14
21      Reference.....	Page S15-S18

22 **1 Supplementary Text**

23 **1.1 Nocturnal boundary layer**

24 The Nocturnal boundary layer (NBL) was calculated by Weather Research and  
25 Forecasting Model (WRF) 3.5 with two-way nested runs. The spatial resolution for  
26 course domain is 30 km with 100×100 grid cells, and for the nested domain is 10 km  
27 with 30×30 grid cells. As our studied site is 100-200 km away from Sea of Japan, we  
28 chose the MYJ scheme for NBL as earlier studies suggested that MYJ scheme was as  
29 first choice for marine atmospheric boundary layer simulations without *a*  
30 *priori* information of atmospheric stability in the region of interest (Huang et al., 2013;  
31 Krogsaeter and Reuder, 2015). For other parameterizations, we selected Kain and  
32 Fritsch cumulus scheme for cumulus parameterization, Lin (Purdue) scheme for  
33 microphysics options RRTM scheme for Radiation Physics Options.

34

35

36 **1.2 Simulations of atmospheric GEM at Mt. Changbai forest using a box model**

37 A box model was applied to estimate the GEM concentration at the height of 24  
38 m agl. Based on the measured characteristics of the GEM depletion events, the  
39 model assumes that vegetative uptake (in terms of dry deposition flux) is the only  
40 pathway for the GEM removal and chemical transformation is not included in the box  
41 modeling. A sensitivity analysis was performed on three parameters: (1) dry  
42 deposition flux (0-10 ng m<sup>-2</sup> h<sup>-1</sup>, the range of measured deposition flux using flux  
43 bags), (2) turbulent diffusivity of the atmosphere (0.1-10 cm s<sup>-1</sup>, typical value under  
44 low wind condition), and (3) a typical nocturnal boundary layer height (100 m agl).  
45 GEM concentration above the stable nocturnal boundary layer was assigned to 1.56  
46 ng m<sup>-3</sup>, the mean observed at 45 m during daytime when the vertical mixing is strong.  
47 The flux of vegetative uptake ( $F_C$ ) and the resulted concentration gradient was  
48 calculated based on the on the algorithm:

$$F_C^{AGM} = - \underbrace{\frac{\kappa \cdot u_*}{\Phi_H(\zeta_1)}}_{K_C} \cdot \frac{\partial C}{\partial \ln(z)} \quad (1)$$

49 where  $k$  is von Kármán constant (~0.41),  $u_*$  is the friction velocity (m s<sup>-1</sup>),  $\psi_H(\zeta_1)$  is  
50 the integrated universal function for sensible heat to correct for deviations from the  
51 ideal logarithmic profile,  $K_C$  term is the turbulent diffusivity (m s<sup>-1</sup>),  $C$  is the  
52 ideal logarithmic profile,  $K_C$  term is the turbulent diffusivity (m s<sup>-1</sup>),  $C$  is the

53 concentration parameter for GEM concentration (ng m<sup>-3</sup>),  $z$  is the height parameter  
54 (m).

55

56 **1.3 The turbulent diffusivity ( $K_c$ )**

57 The flux-gradient approach (Kaimal and Finnigan, 1994) expresses a scalar flux ( $F$ , e.g.  
58 ng m<sup>-2</sup> s<sup>-1</sup>) as the product between the turbulent diffusivity ( $K_C$ , m<sup>2</sup> s<sup>-1</sup>) and a concentration  
59 gradient ( $\partial c / \partial z$ , ng m<sup>-4</sup>) assuming that measurements are made within a vertical layer of  
60 constant flux that forms over homogeneous terrain:

61 
$$F = -K_C \cdot \frac{\partial c}{\partial z} = -\frac{u_* \cdot \kappa \cdot (z-d)}{\phi_H(\varsigma)} \cdot \frac{\partial c}{\partial z} = -\frac{u_* \cdot \kappa}{\phi_H(\varsigma)} \cdot \frac{\partial c}{\partial \ln z} \quad (2),$$

62 where,  $u_*$  is the friction velocity (m s<sup>-1</sup>),  $\kappa$  is the von Kármán constant (taken as 0.4),  
63  $\phi_H(\varsigma)$  is the diabatic influence function for heat (parameterized as a function of  
64  $\varsigma = (z-d)/L$ , where  $L$  is the Obukhov length), whereas  $z$  and  $d$  are the  
65 measurement and (canopy) displacement height (m) respectively. An empirical form of  
66  $\phi_H(\varsigma)$  is  $0.95/\sqrt{(1-11.6\varsigma)}$  and  $0.95 + 7.8\varsigma$  for unstable ( $\varsigma < 0$ ) and stable ( $2 > \varsigma \geq 0$ )  
67 atmospheric conditions respectively (Foken, 2008).

68

69 For flux-gradient measurements made within the roughness sublayer above the canopy  
70 height (i.e.  $h_{canopy} < z < z_*$ ), Eq. 1 is not valid (*underestimates* the magnitude of scalar  
71 flux) and requires further correction following e.g. Garratt (Garratt, 1992) and Simpson et al.  
72 (Simpson et al., 1998):


73 
$$F = -\frac{u_* \cdot \kappa \cdot (z-d)}{\phi_h(\varsigma) \cdot \phi_*(z/z_*)} \cdot \frac{\partial c}{\partial z} \quad (3)$$

74 In contrast to  $\phi_H(\varsigma)$ , the additional correction function  $\phi_*(z/z_*)$  in Eq. 2 is independent of  
75 stability. A common type of parameterization for  $\phi_*$  is  $\exp[-0.7(1 - \frac{z}{z_*})]$  (Garratt, 1992).


76 In-turn, the upper limit of the roughness sublayer ( $z_*$ ) can be estimated by  $2 \cdot h_{canopy} - d$   
77 (Raupach, 1994).

78

79 Figure S1. Map showing the location of Mt. Changbai forest and coal fired power plants in  
80 Northeast Asia.




85 Figure S2. GEM at 4 m agl in the clear spot measured by the Tekran speciation unit, 24 m agl (~3  
86 m above forest canopy, long-term GEM sampling site) and 45 m agl (~24 m above forest  
87 canopy) measured by the Tekran 2537 from 10 to 14 July 2013.




88  
89  
90  
91

92 Figure S3. Time series of soil/air GEM flux and meteorological parameters in Mt. Changbai forest  
93 in July 2013.




94  
95  
96  
97

98 Figure S4. Diurnal trends in vertical gradient of GEM concentrations between the height of 45-24  
99 m, 24-10 m and 10-1m in Mt. Changbai forest from 10 to 15 July 2013.



100  
101  
102  
103

104      Figure S5. Diurnal variations of air temperature (A), wind speed (B), relative humidity (RH, C)  
105      and ozone concentrations (D) in Mt. Changbai forest in leaf-growing season from October 2008 to  
106      December 2014.




107

108

109


110

111      Figure S6. Temporal variation of the turbulent diffusivity at Mt. Changbai forest from 10 to 19  
112      July 2013



113  
114  
115  
116

117 Figure S7. Wind frequency distributions in leaf-growing season and non-leaf-growing season from  
118 Aug 2009 to Jul 2013 (A), monthly mean wind speed from Aug 2009 to Jul 2013 (B), and Leaf  
119 areas index in leaf-growing season during 2003-2005(C) (Shi et al., 2008).



120

121

122 Table S1. Statistical summary of litterfall Hg concentrations and litter mass in Mt. Changbai forest

Litterfall collection site	Time	Species	Concentration (ng g <sup>-1</sup> )	Litter mass (g m <sup>-2</sup> )	Mass-weight concentration (ng g <sup>-1</sup> )
Collector-1	2013-09	<i>Pinus koraiensis</i>	44.5	51.70	
		<i>Acer pseudo-sieboldianum</i>	138.5	24.95	
		<i>Quercus mongolica</i>	170.4	35.60	74.8
		<i>Fraxinus mandshurica</i>	44.5	51.70	
		<i>Tilia amurensis</i>	138.5	24.95	
Collector-1	2013-10	<i>Others</i>	170.4	35.60	
		<i>Acer mono</i>	60.2	1.8	60.2
Collector-2	2013-09	<i>Pinus koraiensis</i>	40.4	6.48	
		<i>Acer pseudo-sieboldianum</i>	46.8	5.38	
		<i>Quercus mongolica</i>	31.2	9.49	
		<i>Acer mono</i>	58.0	4.81	39.7
		<i>Fraxinus mandshurica</i>	34.5	18.54	
Collector-2	2013-10	<i>Tilia amurensis</i>	45.7	52.12	
		<i>Others</i>	36.80	91.16	
		<i>Pinus koraiensis</i>	34.1	4.4	
		<i>Quercus mongolica</i>	30.3	8.8	53.3
Collector-3	2013-09	<i>Acer mono</i>	70.7	5.8	
		<i>Others</i>	81.3	6.6	
		<i>Acer ginnala Maxim</i>	38.1	5.8	
		<i>Pinus koraiensis</i>	29.4	19.0	
		<i>Acer pseudo-sieboldianum</i>	49.6	4.6	
Collector-3	2013-10	<i>Quercus mongolica</i>	32.5	10.8	
		<i>Acer mono</i>	46.9	6.0	31.1
		<i>Fraxinus mandshurica</i>	23.4	1.5	
		<i>Tilia amurensis</i>	40.4	19.6	
		<i>Others</i>	29.8	283.3	
Collector-4	2013-09	<i>Pinus koraiensis</i>	21.7	1.5	
		<i>Quercus mongolica</i>	42.3	3.2	35.8
		<i>Others</i>	31.4	70.6	
Collector-4	2013-10	<i>Acer pseudo-sieboldianum</i>	47.4	2.7	
		<i>Quercus mongolica</i>	26.3	4.1	
		<i>Acer mono</i>	50.9	15.5	34.2
		<i>Tilia amurensis</i>	43.0	11.8	
		<i>Others</i>	32.8	121.1	
Collector-4	2013-10	<i>Pinus koraiensis</i>	40.1	19.4	
		<i>Quercus mongolica</i>	46.0	21.7	64.4
		<i>Others</i>	100.9	23.9	
<b>Overall mean</b>				<b>43.0±29.5</b>	

124 Table S2. Concentration of atmospheric speciated Hg (GEM, PBM, and GOM) and isotopic  
 125 composition of atmospheric GEM at Mt. Changbai forest

Sample ID	Sampling period	GEM conc.	PBM conc.	GOM conc.	$\delta^{202}\text{Hg}$	$\delta^{202}\text{Hg}$	$\Delta^{199}\text{Hg}$	$\Delta^{199}\text{Hg}$	$\Delta^{200}\text{Hg}$	$\Delta^{200}\text{Hg}$	$\Delta^{201}\text{Hg}$	$\Delta^{201}\text{Hg}$
		ng m <sup>-3</sup>	pg m <sup>-3</sup>	pg m <sup>-3</sup>	%o	2σ, %o						
GEM-1	2013/7/8 13:40-2013/7/9 15:30	1.60	2	4	-0.06	0.09	-0.05	0.04	0.01	0.04	-0.01	0.06
GEM-2	2013/7/9 15:30-2013/7/10 15:30	1.46	2	2	0.35	0.09	-0.05	0.04	-0.04	0.04	-0.12	0.06
GEM-3	2013/7/10 15:30-2013/7/11 15:45	1.23	4	2	0.61	0.09	-0.09	0.04	-0.05	0.04	-0.08	0.06
GEM-4	2013/7/11 15:45-2013/7/12 15:50	1.14	5	2	0.80	0.09	-0.08	0.04	-0.03	0.04	-0.06	0.06
GEM-5	2013/7/12 15:50-2013/7/13 16:00	0.91	6	1	0.91	0.09	-0.06	0.04	-0.04	0.04	-0.07	0.06
GEM-6	2013/7/13 16:00-2013/7/14 15:40	1.32	4	2	0.58	0.09	-0.06	0.04	-0.06	0.04	-0.05	0.06
GEM-7	2013/7/14 15:40-2013/7/15 17:10	1.37	2	2	0.58	0.09	-0.08	0.04	-0.05	0.04	-0.04	0.06
GEM-8	2013/7/15 17:15-2013/7/16 17:30	1.24	2	2	0.08	0.09	-0.04	0.04	-0.04	0.04	-0.04	0.06
GEM-9	2013/7/16 17:30-2013/7/17 18:10	1.57	5	1	-0.34	0.09	-0.11	0.04	-0.04	0.04	-0.06	0.06
GEM-10	2013/7/17 18:10-2013/7/18 18:35	1.30	8	1	-0.01	0.09	-0.08	0.04	-0.05	0.04	-0.04	0.06

126

127

128 Table S3. A statistical summary of reported litterfall Hg deposition fluxes and estimated annual  
 129 litterfall Hg deposition of Hg over the world.

Region	Litterfall Hg deposition flux ( $\mu\text{g m}^{-2} \text{yr}^{-1}$ )			Forest area ( $\text{km}^2$ )	Estimated litterfall deposition ( $\text{Mg yr}^{-1}$ )	Reference
	Range	Median	N			
Asia	20.9-220	37.5±76.1	6	5,775,220	217	(Wang et al., 2009; Fu et al., 2010a; Fu et al., 2010b; Zhou et al., 2013)
North America	3.8-30.9	13.9±5.6	47	6,847,010	95	(Lindberg, 1996; Rea et al., 1996; Grigal et al., 2000; St Louis et al., 2001; Sheehan et al., 2006; Demers et al., 2007; Bushey et al., 2008; Fisher and Wolfe, 2012; Juillerat et al., 2012; Risch et al., 2012; Benoit et al., 2013)
Europe (including Russia)	2.7-25.2	14.2±8.9	6	10,156,300	144	(Iverfeldt, 1991; Munthe et al., 1995; Lee et al., 2000; Schwesig and Matzner, 2000; Lindberg et al., 2007; Larssen et al., 2008)
South America	43.0-184	60.0±49.0	9	9,436,410	566	(Roulet et al., 1998; Fostier et al., 2003; Mélières et al., 2003; Silva-Filho et al., 2006; Teixeira et al., 2012)
Africa				6,164,310	159*	Lack of observational data
Oceania				1,951,370	50*	Lack of observational data
<b>Global total</b>					<b>1232</b>	

130 (Estimated litterfall deposition: \* indicates the values were calculated using the global median litterfall Hg deposition flux and the forest  
 131 area in these regions)

132

133

134 Table S4. A statistical summary of reported throughfall Hg deposition fluxes and estimated annual  
 135 throughfall Hg deposition of Hg over the world

Region	Throughfall Hg deposition flux			Forest area (km <sup>2</sup> )	Estimated throughfall deposition (mg yr <sup>-1</sup> )	Reference			
	(µg m <sup>-2</sup> yr <sup>-1</sup> )								
	Range	Median	N						
Asia	10.5-71.3	36.8±29.9	4	5,775,220	213	(Wan et al., 2009; Wang et al., 2009; Fu et al., 2010a; Fu et al., 2010b)			
North America	3.8-30.9	11.8±3.3	10	6,847,010	81	(Lindberg, 1996; Rea et al., 1996; Grigal et al., 2000; St Louis et al., 2001; Sheehan et al., 2006; Choi et al., 2008; Fisher and Wolfe, 2012)			
Europe (including Russia)	6.8-39.0	15.2±10.7	8	10,156,300	154	(Iverfeldt, 1991; Munthe et al., 1995; Lee et al., 2000; Schwesig and Matzner, 2000; Lindberg et al., 2007; Larssen et al., 2008)			
South America	72	72	1	9,436,410	679	(Fostier et al., 2000)			
Africa				6,164,310	160*	Lack of observational data			
Oceania				1,951,370	51*	Lack of observational data			
<b>Global total</b>					<b>1338</b>				

136 (Estimated throughfall deposition: \* indicates the values were calculated using the global median throughfall Hg deposition flux and the  
 137 forest area in these regions)

138

139

140

141 Table S5. A statistical summary of reported forest soil emission fluxes and estimated annual forest  
 142 soil emission fluxes over the world  
 143

Region	Forest soil Hg emission flux			Forest area (km <sup>2</sup> )	Estimated forest soil emission (Mg yr <sup>-1</sup> )	Reference
	Range	Median	N			
Asia	3.3-81.2	30.4±23.5	11	5,775,220	176	(Wang et al., 2006; Fu et al., 2008; Fu et al., 2012; Ma et al., 2013)
North America	-1.3-45.9	9.2±11.4	15	6,847,010	63	(Carpi and Lindberg, 1998; Poissant and Casimir, 1998; Zhang et al., 2001; Nacht and Gustin, 2004; Schroeder et al., 2005; Kuiken et al., 2008a; Kuiken et al., 2008b; Choi and Holsen, 2009)
Europe (including Russia)	-0.1-9.6	2.4±4.3	4	10,156,300	24	(Schroeder et al., 1989; Xiao et al., 1991; Ferrara et al., 1997; Lindberg et al., 1998)
South America	6.0	6.0	1	9,436,410	57	(Carpi et al., 2014)
Africa				6,164,310	47*	Lack of observational data
Oceania				1,951,370	15*	Lack of observational data
<b>Global total</b>					<b>381</b>	

144 (Estimated forest soil emission; \* indicates the values were calculated using the global median forest soil Hg emission flux and the forest  
 145 area in these regions)

146

147

148

149 **Reference:**

- 150 Benoit, J. M., Cato, D. A., Denison, K. C., and Moreira, A. E.: Seasonal Mercury Dynamics in a New  
151 England Vernal Pool, *Wetlands*, 33, 887-894, DOI 10.1007/s13157-013-0447-4, 2013.
- 152 Bushey, J. T., Nallana, A. G., Montesdeoca, M. R., and Driscoll, C. T.: Mercury dynamics of a  
153 northern hardwood canopy, *Atmos Environ*, 42, 6905-6914, DOI  
154 10.1016/j.atmosenv.2008.05.043, 2008.
- 155 Carpi, A., and Lindberg, S. E.: Application of a Teflon (TM) dynamic flux chamber for quantifying  
156 soil mercury flux: Tests and results over background soil, *Atmos Environ*, 32, 873-882, Doi  
157 10.1016/S1352-2310(97)00133-7, 1998.
- 158 Carpi, A., Fostier, A. H., Orta, O. R., dos Santos, J. C., and Gittings, M.: Gaseous mercury emissions  
159 from soil following forest loss and land use changes: Field experiments in the United States and  
160 Brazil, *Atmos Environ*, 96, 423-429, DOI 10.1016/j.atmosenv.2014.08.004, 2014.
- 161 Choi, H. D., Sharac, T. J., and Holsen, T. M.: Mercury deposition in the Adirondacks: A comparison  
162 between precipitation and throughfall, *Atmos Environ*, 42, 1818-1827, DOI  
163 10.1016/j.atmosenv.2007.11.036, 2008.
- 164 Choi, H. D., and Holsen, T. M.: Gaseous mercury fluxes from the forest floor of the Adirondacks,  
165 *Environ Pollut*, 157, 592-600, DOI 10.1016/j.envpol.2008.08.020, 2009.
- 166 Demers, J. D., Driscoll, C. T., Fahey, T. J., and Yavitt, J. B.: Mercury cycling in litter and soil in  
167 different forest types in the Adirondack region, New York, USA, *Ecol Appl*, 17, 1341-1351, Doi  
168 10.1890/06-1697.1, 2007.
- 169 Ferrara, R., Maserati, B. E., Andersson, M., Edner, H., Ragnarson, P., and Svanberg, S.: Mercury  
170 degassing rate from mineralized areas in the Mediterranean basin, *Water Air Soil Poll*, 93, 59-66,  
171 Doi 10.1023/A:1022107205659, 1997.
- 172 Fisher, L. S., and Wolfe, M. H.: Examination of mercury inputs by throughfall and litterfall in the  
173 Great Smoky Mountains National Park, *Atmos Environ*, 47, 554-559, DOI  
174 10.1016/j.atmosenv.2011.10.017, 2012.
- 175 Foken, T.: *Micrometeorology*, Springer-Verlag, Berlin, Heidelberg, 2008.
- 176 Fostier, A. H., Forti, M. C., Guimaraes, J. R. D., Melfi, A. J., Boulet, R., Santo, C. M. E., and Krug, F.  
177 J.: Mercury fluxes in a natural forested Amazonian catchment (Serra do Navio, Amapa State,  
178 Brazil), *Sci Total Environ*, 260, 201-211, Doi 10.1016/S0048-9697(00)00564-7, 2000.
- 179 Fostier, A. H., Cecon, K., and Forti, M. C.: Urban influence on litterfall trace metals fluxes in the  
180 Atlantic forest of Sao Paulo (Brazil), *J Phys Iv*, 107, 491-494, Doi 10.1051/Jp4:20030348, 2003.
- 181 Fu, X., Feng, X., Zhu, W., Rothenberg, S., Yao, H., and Zhang, H.: Elevated atmospheric deposition  
182 and dynamics of mercury in a remote upland forest of southwestern China, *Environ Pollut*, 158,  
183 2324-2333, 10.1016/j.envpol.2010.01.032, 2010a.
- 184 Fu, X. W., Feng, X. B., and Wang, S. F.: Exchange fluxes of Hg between surfaces and atmosphere in  
185 the eastern flank of Mount Gongga, Sichuan province, southwestern China, *J Geophys  
186 Res-Atmos*, 113, Artn D20306  
187 Doi 10.1029/2008jd009814, 2008.
- 188 Fu, X. W., Feng, X., Dong, Z. Q., Yin, R. S., Wang, J. X., Yang, Z. R., and Zhang, H.: Atmospheric  
189 gaseous elemental mercury (GEM) concentrations and mercury depositions at a high-altitude  
190 mountain peak in south China, *Atmos Chem Phys*, 10, 2425-2437, 2010b.
- 191 Fu, X. W., Feng, X. B., Zhang, H., Yu, B., and Chen, L. G.: Mercury emissions from natural surfaces  
192 highly impacted by human activities in Guangzhou province, South China, *Atmos Environ*, 54,

- 193 185-193, DOI 10.1016/j.atmosenv.2012.02.008, 2012.  
194 Garratt, J. R.: The atmospheric boundary layer, Cambridge University Press, Cambridge, UK, 1992.  
195 Grigal, D. F., Kolka, R. K., Fleck, J. A., and Nater, E. A.: Mercury budget of an upland-peatland  
196 watershed, *Biogeochemistry*, 50, 95-109, Doi 10.1023/A:1006322705566, 2000.  
197 Huang, H. Y., Hall, A., and Teixeira, J.: Evaluation of the WRF PBL Parameterizations for Marine  
198 Boundary Layer Clouds: Cumulus and Stratocumulus, *Mon Weather Rev*, 141, 2265-2271,  
199 10.1175/Mwr-D-12-00292.1, 2013.  
200 Iverfeldt, A.: Mercury in forest canopy throughfall water and its relation to atmospheric deposition,  
201 *Water Air Soil Poll*, 56, 553-564, Doi 10.1007/Bf00342299, 1991.  
202 Juillerat, J. I., Ross, D. S., and Bank, M. S.: Mercury in litterfall and upper soil horizons in forested  
203 ecosystems in Vermont, USA, *Environ Toxicol Chem*, 31, 1720-1729, Doi 10.1002/etc.1896,  
204 2012.  
205 Kaimal, J. C., and Finnigan, J. J.: Atmospheric Boundary Layer Flows, Oxford University Press, New  
206 York, Oxford, 1994.  
207 Krogsaeter, O., and Reuder, J.: Validation of boundary layer parameterization schemes in the Weather  
208 Research and Forecasting (WRF) model under the aspect of offshore wind energy  
209 applicationspart II: boundary layer height and atmospheric stability, *Wind Energy*, 18,  
210 1291-1302, 10.1002/we.1765, 2015.  
211 Kuiken, T., Gustin, M., Zhang, H., Lindberg, S., and Sedinger, B.: Mercury emission from terrestrial  
212 background surfaces in the eastern USA. II: Air/surface exchange of mercury within forests  
213 from South Carolina to New England, *Appl Geochem*, 23, 356-368, DOI  
214 10.1016/j.apgeochem.2007.12.007, 2008a.  
215 Kuiken, T., Zhang, H., Gustin, M., and Lindberg, S.: Mercury emission from terrestrial background  
216 surfaces in the eastern USA. Part I: Air/surface exchange of mercury within a southeastern  
217 deciduous forest (Tennessee) over one year, *Appl Geochem*, 23, 345-355, DOI  
218 10.1016/j.apgeociem.2007.12.006, 2008b.  
219 Larssen, T., de Wit, H. A., Wiker, M., and Halse, K.: Mercury budget of a small forested boreal  
220 catchment in southeast Norway, *Sci Total Environ*, 404, 290-296, DOI  
221 10.1016/j.scitotenv.2008.03.013, 2008.  
222 Lee, Y. H., Bishop, K. H., and Munthe, J.: Do concepts about catchment cycling of methylmercury  
223 and mercury in boreal catchments stand the test of time? Six years of atmospheric inputs and  
224 runoff export at Svarthberget, northern Sweden, *Sci Total Environ*, 260, 11-20, Doi  
225 10.1016/S0048-9697(00)00538-6, 2000.  
226 Lindberg, S., Bullock, R., Ebinghaus, R., Engstrom, D., Feng, X. B., Fitzgerald, W., Pirrone, N.,  
227 Prestbo, E., and Seigneur, C.: A synthesis of progress and uncertainties in attributing the sources  
228 of mercury in deposition, *Ambio*, 36, 19-32, 2007.  
229 Lindberg, S. E.: Forests and the global biogeochemical cycle of mercury: The importance of  
230 understanding air/vegetation exchange processes, *Nato Asi* 2, 21, 359-380, 1996.  
231 Lindberg, S. E., Hanson, P. J., Meyers, T. P., and Kim, K. H.: Air/surface exchange of mercury vapor  
232 over forests - The need for a reassessment of continental biogenic emissions, *Atmos Environ*, 32,  
233 895-908, Doi 10.1016/S1352-2310(97)00173-8, 1998.  
234 Mélières, M. A., Pourchet, M., Charles-Dominique, P., and Gaucher, P.: Mercury in canopy leaves of  
235 French Guiana in remote areas, *Sci Total Environ*, 311, 261-267, Doi  
236 10.1016/S0048-9697(03)00142-6, 2003.

- 237 Ma, M., Wang, D. Y., Sun, R. G., Shen, Y. Y., and Huang, L. X.: Gaseous mercury emissions from  
238 subtropical forested and open field soils in a national nature reserve, southwest China, *Atmos*  
239 *Environ.*, 64, 116-123, DOI 10.1016/j.atmosenv.2012.09.038, 2013.
- 240 Munthe, J., Hultberg, H., and Iverfeldt, A.: Mechanisms of Deposition of Methylmercury and  
241 Mercury to Coniferous Forests, *Water Air Soil Poll.*, 80, 363-371, Doi 10.1007/Bf01189686,  
242 1995.
- 243 Nacht, D. M., and Gustin, M. S.: Mercury emissions from background and altered geologic units  
244 throughout Nevada, *Water Air Soil Poll.*, 151, 179-193, Doi  
245 10.1023/B:WATE.0000009907.49577.A8, 2004.
- 246 Poissant, L., and Casimir, A.: Water-air and soil-air exchange rate of total gaseous mercury measured  
247 at background sites, *Atmos Environ.*, 32, 883-893, Doi 10.1016/S1352-2310(97)00132-5, 1998.
- 248 Raupach, M. R.: Simplified Expressions for Vegetation Roughness Length and Zero-Plane  
249 Displacement as Functions of Canopy Height and Area Index, *Bound-Lay Meteorol.*, 71,  
250 211-216, Doi 10.1007/Bf00709229, 1994.
- 251 Rea, A. W., Keeler, G. J., and Scherbatskoy, T.: The deposition of mercury in throughfall and litterfall  
252 in the lake Champlain watershed: A short-term study, *Atmos Environ.*, 30, 3257-3263, Doi  
253 10.1016/1352-2310(96)00087-8, 1996.
- 254 Risch, M. R., DeWild, J. F., Krabbenhoft, D. P., Kolka, R. K., and Zhang, L. M.: Litterfall mercury  
255 dry deposition in the eastern USA, *Environ Pollut.*, 161, 284-290, DOI  
256 10.1016/j.envpol.2011.06.005, 2012.
- 257 Roulet, M., Lucotte, M., Saint-Aubin, A., Tran, S., Rheault, I., Farella, N., Da Silva, E. D.,  
258 Dezencourt, J., Passos, C. J. S., Soares, G. S., Guimaraes, J. R. D., Mergler, D., and Amorim, M.:  
259 The geochemistry of mercury in central Amazonian soils developed on the Alter-do-Chao  
260 formation of the lower Tapajos River Valley, Para state, Brazil, *Sci Total Environ.*, 223, 1-24, Doi  
261 10.1016/S0048-9697(98)00265-4, 1998.
- 262 Schroeder, W. H., Munthe, J., and Lindqvist, O.: Cycling of Mercury between Water, Air, and Soil  
263 Compartments of the Environment, *Water Air Soil Poll.*, 48, 337-347, 1989.
- 264 Schroeder, W. H., Beauchamp, S., Edwards, G., Poissant, L., Rasmussen, P., Tordon, R., Dias, G.,  
265 Kemp, J., Van Heyst, B., and Banic, C. M.: Gaseous mercury emissions from natural sources in  
266 Canadian landscapes, *J Geophys Res-Atmos.*, 110, Artn D18302, Doi 10.1029/2004jd005699,  
267 2005.
- 268 Schwesig, D., and Matzner, E.: Pools and fluxes of mercury and methylmercury in two forested  
269 catchments in Germany, *Sci Total Environ.*, 260, 213-223, Doi 10.1016/S0048-9697(00)00565-9,  
270 2000.
- 271 Sheehan, K. D., Fernandez, I. J., Kahl, J. S., and Amirbahman, A.: Litterfall mercury in two forested  
272 watersheds at Acadia National Park, Maine, USA, *Water Air Soil Poll.*, 170, 249-265, DOI  
273 10.1007/s11270-006-3034-y, 2006.
- 274 Shi, T. T., Guan, D. X., Wang, A. Z., Wu, J. B., Jin, C. J., and Han, S. J.: Comparison of three models  
275 to estimate evapotranspiration for a temperate mixed forest, *Hydrol Process.*, 22, 3431-3443,  
276 10.1002/hyp.6922, 2008.
- 277 Silva-Filho, E. V., Machado, W., Oliveira, R. R., Sella, S. M., and Lacerda, L. D.: Mercury deposition  
278 through litterfall in an Atlantic Forest at Ilha Grande, southeast Brazil, *Chemosphere*, 65,  
279 2477-2484, DOI 10.1016/j.chemosphere.2006.04.053, 2006.
- 280 Simpson, I. J., Thurtell, G. W., Neumann, H. H., Den Hartog, G., and Edwards, G. C.: The validity of

- 281 similarity theory in the roughness sublayer above forests, *Bound-Lay Meteorol*, 87, 69-99, Doi  
282 10.1023/A:1000809902980, 1998.
- 283 St Louis, V. L., Rudd, J. W. M., Kelly, C. A., Hall, B. D., Rolfhus, K. R., Scott, K. J., Lindberg, S. E.,  
284 and Dong, W.: Importance of the forest canopy to fluxes of methyl mercury and total mercury to  
285 boreal ecosystems, *Environmental Science & Technology*, 35, 3089-3098, Doi  
286 10.1021/Es001924p, 2001.
- 287 Teixeira, D. C., Montezuma, R. C., Oliveira, R. R., and Silva, E. V.: Litterfall mercury deposition in  
288 Atlantic forest ecosystem from SE - Brazil, *Environ Pollut*, 164, 11-15, DOI  
289 10.1016/j.envpol.2011.10.032, 2012.
- 290 Wan, Q., Feng, X. B., Lu, J., Zheng, W., Song, X. J., Li, P., Han, S. J., and Xu, H.: Atmospheric  
291 mercury in Changbai Mountain area, northeastern China II. The distribution of reactive gaseous  
292 mercury and particulate mercury and mercury deposition fluxes, *Environ Res*, 109, 721-727,  
293 DOI 10.1016/j.envres.2009.05.006, 2009.
- 294 Wang, D. Y., He, L., Shi, X. J., Wei, S. Q., and Feng, X. B.: Release flux of mercury from different  
295 environmental surfaces in Chongqing, China, *Chemosphere*, 64, 1845-1854, DOI  
296 10.1016/j.chemosphere.2006.01.054, 2006.
- 297 Wang, Z. W., Zhang, X. S., Xiao, J. S., Zhijia, C., and Yu, P. Z.: Mercury fluxes and pools in three  
298 subtropical forested catchments, southwest China, *Environ Pollut*, 157, 801-808, DOI  
299 10.1016/j.envpol.2008.11.018, 2009.
- 300 Xiao, Z. F., Munthe, J., Schroeder, W. H., and Lindqvist, O.: Vertical Fluxes of Volatile Mercury over  
301 Forest Soil and Lake Surfaces in Sweden, *Tellus B*, 43, 267-279, DOI  
302 10.1034/j.1600-0889.1990.t01-1-00009.x-i1, 1991.
- 303 Zhang, H., Lindberg, S. E., Marsik, F. J., and Keeler, G. J.: Mercury air/surface exchange kinetics of  
304 background soils of the Tahquamenon River watershed in the Michigan Upper Peninsula, *Water  
305 Air Soil Poll*, 126, 151-169, Doi 10.1023/A:1005227802306, 2001.
- 306 Zhou, J., Feng, X. B., Liu, H. Y., Zhang, H., Fu, X. W., Bao, Z. D., Wang, X., and Zhang, Y. P.:  
307 Examination of total mercury inputs by precipitation and litterfall in a remote upland forest of  
308 Southwestern China, *Atmos Environ*, 81, 364-372, DOI 10.1016/j.atmosenv.2013.09.010, 2013.  
309