



*Supplement of*

## The magnitude of the snow-sourced reactive nitrogen flux to the boundary layer in the Uintah Basin, Utah, USA

Maria Zatko et al.

Correspondence to: Becky Alexander (beckya@uw.edu)

- acp-16-13837-2016-supplement-title-page.pdf
- acp-2016-320-supplement-version5
  - SupplementalMaterial
    - \* Figure10A.eps
    - \* Figure10B.pdf
    - \* Figure11A.eps
    - \* Figure11B.pdf
    - \* Figure12A.eps
    - \* Figure12B.pdf
    - \* Figure13A.eps
    - \* Figure13B.pdf
    - \* Figure14A.eps
    - \* Figure14B.pdf
    - \* Figure15A.eps
    - \* Figure15B.pdf
    - \* Figure16A.eps
    - \* Figure1A.eps
    - \* Figure1B.pdf
    - \* Figure2A.eps
    - \* Figure2B.eps
    - \* Figure3A.eps
    - \* Figure3B.eps
    - \* Figure4A.eps
    - \* Figure4B.pdf
    - \* Figure5A.eps
    - \* Figure5B.pdf
    - \* Figure6A.eps

- \* Figure6B.pdf
- \* Figure7A.eps
- \* Figure7B.pdf
- \* Figure8A.eps
- \* Figure8B.pdf
- \* Figure9A.eps
- \* Figure9B.pdf
- \* Snow\_SurfaceIrradiance\_ActinicFlux.xlsx
- \* Zatko\_etal\_2016\_SupplementalMaterial.docx

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.