

Supplement of Atmos. Chem. Phys., 16, 14219–14230, 2016
<http://www.atmos-chem-phys.net/16/14219/2016/>
doi:10.5194/acp-16-14219-2016-supplement
© Author(s) 2016. CC Attribution 3.0 License.



Atmospheric
Chemistry
and Physics
Open Access
EGU

Supplement of

Synergetic formation of secondary inorganic and organic aerosol: effect of SO₂ and NH₃ on particle formation and growth

Biwu Chu et al.

Correspondence to: Hong He (honghe@rcees.ac.cn)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

Increase of secondary aerosol formation with SO₂ consumption, particle surface area, and particle volume




Fig. S1 Correlation of secondary aerosol formation with SO₂ consumption, particle surface area, and particle volume.

Estimating concentrations of nitrogen-containing organics (NOC)

Nitrogen-containing organics (NOC) might contain organonitrates, organic ammonium salts, and species with carbon covalently bonded to nitrogen. Organonitrates could be estimated using methods suggested by Farmer et al. (2010), one of which was using NO_x⁺ ratios. The ratio of m/z 30 to m/z 46 for nitrate ($R_{NH_4NO_3}$) is lower than that of organonitrates ($R_{Organonitrates}$) in AMS measurement, so the observed ratio, R_{Obs} , can be used to estimate the fraction of the total nitrate signal due to organonitrates (x)

using the Equation S1:

$$x = \frac{(R_{obs} - R_{NH_4NO_3})(1 + R_{obs})}{(R_{Organonitrates} - R_{NH_4NO_3})(1 + R_{Organonitrates})} \quad (S1)$$

where $R_{NH_4NO_3}$ was measured as 2.35 in the calibration of AMS, and $R_{Organonitrates}$ was assumed to be 15, which was highest value suggested by Farmer et al (2010), to avoid overestimation of organonitrates. The estimated concentrations of organonitrates are shown in Fig. S2.




Fig. S2 The estimated concentrations of organonitrates from NO_x⁺ ratios as a function of SO₂ concentration in photooxidation of toluene/NO_x under (a) NH₃-poor and (b) NH₃-rich conditions

As suggested by Farmer et al. (2010), NH₄⁺ is required to balance with the SO₄²⁻ and NO₃⁻ in the absence of NOC. In this study, the measured concentration of NH₄⁺ was higher than the calculated concentration according to charge balance, especially in the NH₃-rich experiments. This result indicated the presence of NOC with reduced N (Reduced NOC). The NH₃ present may react with the ring opening oxycarboxylic acids from

toluene (Jang and Kamens, 2001) and carbonyl functional group organics (Wang et al., 2010), and result in organic ammonium salt products, imines, imidazole, and so on. The NH_4 balance considering the presence of NOC with reduced N is described by Equation S2:

$$\frac{c_{\text{NH}_4}}{18} - \frac{c_{\text{Reduced NOC}}}{M_{\text{Reduced NOC}}} = 2 \times \frac{c_{\text{SO}_4}}{96} + \left[\frac{c_{\text{NO}_3}}{62} - \frac{c_{\text{Organonitrates}}}{M_{\text{Organonitrates}}} \right] \quad (\text{S2})$$

where c_{NH_4} , c_{SO_4} and c_{NO_3} are the mass concentrations of ammonium and sulfate and nitrate in AMS measurement results, respectively; while $c_{\text{Reduced NOC}}$ and $c_{\text{Organonitrates}}$ are the estimated mass concentrations of NOC with reduced N and organonitrates, respectively. $M_{\text{Reduced NOC}}$ and $M_{\text{Organonitrates}}$ in Equation S2 are the molecular weights of NOC with reduced N and organonitrates, and are assumed to be 100 and 200, respectively. $M_{\text{Reduced NOC}}$ is assumed to have a smaller value than $M_{\text{Organonitrates}}$ to avoid overestimation of NOC with reduced N. Using the estimated concentration of organonitrates according to Equation S1, the estimated concentrations of the NOC with reduced N are calculated and displayed in Fig. S3.




Fig. S3 The estimated concentrations of NOC with reduced N as a function of SO_2 concentration in photooxidation of toluene/ NO_x under (a) NH_3 -poor and (b)

NH₃-rich conditions

The uncertainty in the results in Fig. S2 and Fig. S3 might be large, since a lot of parameters are assumed and the measurement response of AMS to the NOC is still highly uncertain. We analyzed the organic fragments containing N in the AMS measurements. Farmer et al. (2010) suggested the estimation of concentrations of organonitrates using the sum of the signals of the five major C_xH_yO_zN⁺ fragments in AMS measurement results. However, as pointed out by Farmer et al (2010) and other studies (Galloway et al., 2009;Laskin et al., 2014;Liu et al., 2015), NOC with reduced N could also produce these fragments. In this study, we used Equation S3 to provide an estimate of NOC:

$$c_{NOC} = \frac{CH_4NO^+ + C_2H_5NO^+ + C_3H_4NO^+ + CH_2NO_3^+ + CH_2NO_2^+}{R_{C_xH_yO_zN^+}} \quad (S3)$$

where $R_{C_xH_yO_zN^+}$ is the fraction of these five major C_xH_yO_zN⁺ fragments and accounts for the total oxidized N signal of the organonitrates samples, and was measured as 0.045 in the study of Farmer et al. (2010). The estimated concentrations of NOC using Equation S3 are shown in Fig. S4. These results appear to be comparable to the sum of organonitrates in Fig. S2 and NOC with reduced N in Fig. S3. Meanwhile, the concentrations of C_xH_yN⁺ fragments are displayed in Fig. S5. They also have similar change trends with SO₂ concentration as the estimated NOC concentrations.




Fig. S4 The estimated concentrations of NOC from C_xH_yO_zN⁺ fragments as a function of SO₂ concentration in photooxidation of toluene/NO_x under (a) NH₃-poor and (b) NH₃-rich conditions




Fig. S5 The concentrations of C_xH_yN⁺ fragments as a function of SO₂ concentration in photooxidation of toluene/NO_x under (a) NH₃-poor and (b) NH₃-rich conditions

References

Farmer, D. K., Matsunaga, A., Docherty, K. S., Surratt, J. D., Seinfeld, J. H., Ziemann, P. J., and Jimenez, J. L.: Response of an aerosol mass spectrometer to organonitrates and organosulfates and implications for atmospheric chemistry, Proc. Natl. Acad. Sci. USA, 107, 6670-6675, DOI 10.1073/pnas.0912340107, 2010.

Galloway, M. M., Chhabra, P. S., Chan, A. W. H., Surratt, J. D., Flagan, R. C., Seinfeld, J. H., and Keutsch, F. N.: Glyoxal uptake on ammonium sulphate seed aerosol: reaction products and reversibility of uptake under dark and irradiated conditions, *Atmos. Chem. Phys.*, 9, 3331-3345, 2009.

Jang, M. S., and Kamens, R. M.: Characterization of secondary aerosol from the photooxidation of toluene in the presence of NO_x and 1-propene, *Environ. Sci. & Technol.*, 35, 3626-3639, DOI 10.1021/es010676+, 2001.

Laskin, J., Laskin, A., Nizkorodov, S. A., Roach, P., Eckert, P., Gilles, M. K., Wang, B. B., Lee, H. J., and Hu, Q. C.: Molecular Selectivity of Brown Carbon Chromophores, *Environ. Sci. & Technol.*, 48, 12047-12055, 2014.

Liu, Y., Liggio, J., Staebler, R., and Li, S. M.: Reactive uptake of ammonia to secondary organic aerosols: kinetics of organonitrogen formation, *Atmos. Chem. Phys.*, 15, 13569-13584, 10.5194/acp-15-13569-2015, 2015.

Wang, X. F., Gao, S., Yang, X., Chen, H., Chen, J. M., Zhuang, G. S., Surratt, J. D., Chan, M. N., and Seinfeld, J. H.: Evidence for High Molecular Weight Nitrogen-Containing Organic Salts in Urban Aerosols, *Environ. Sci. & Technol.*, 44, 4441-4446, DOI 10.1021/es1001117, 2010.