



*Supplement of*

## **Sensitivity of nitrate aerosols to ammonia emissions and to nitrate chemistry: implications for present and future nitrate optical depth**

**F. Paulot et al.**

*Correspondence to:* F. Paulot (fabien.paulot@noaa.gov)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

Table S1: Heterogeneous reactions and uptake coefficients used in AM3 and AM3N

Aerosol	$\gamma$ (AM3)	$\gamma$ (AM3N)	Reference (AM3N)
$\gamma_{\text{HO}_2}$	all aerosol	1	Mao et al. (2013)
$\gamma_{\text{NO}_2}$	fine aerosol <sup>a</sup>	$10^{-4}$	Wong et al. (2011)
$\gamma_{\text{NO}_3}$	fine aerosol <sup>a</sup>	0.1	Jacob (2000)
$\gamma_{\text{N}_2\text{O}_5}$	fine aerosol <sup>a</sup>	0.1	Macintyre and Evans (2010)
$\gamma_{\text{NO}_3}$	dust <sup>b</sup>	-	Crowley et al. (2010)
$\gamma_{\text{N}_2\text{O}_5}$	dust <sup>b</sup>	-	Crowley et al. (2010)
$\gamma_{\text{SO}_2}$	dust	-	Ullerstam et al. (2002, 2003)
$\gamma_{\text{HNO}_3}$	dust <sup>b</sup>	- $\frac{8\text{RH}}{30(1-\text{RH})(1+7\text{RH})}$ for $\text{RH} < 73\%$ 0.118 for $\text{RH} \geq 73\%$	Vlasenko et al. (2006)

<sup>a</sup> excludes seasalt and dust (including nitrate and sulfate on dust)

<sup>b</sup> if alkalinity is available

Table S2: Present-day budget of  $\text{SO}_4$ ,  $\text{NH}_x$ , and  $\text{NO}_y$  in AM3 and AM3N

	AM3	AM3N	AM3N_fdep_diu	AM3N_diu	AM3N_fdep	AM3N_ns	AM3N_nhet	AM3N_ndust
$\text{SO}_4^a$								
Production (TgS/yr)	37.3	33.1		33.9	33.3	33.8	33.3	34.0
OH	10.4	7.7		7.7	7.7	7.7	8.0	9.1
$\text{H}_2\text{O}_2$	26.7	16.2		15.9	16.0	16.1	16.3	15.9
$\text{O}_3$	0.1	4.5		5.6	4.8	5.2	4.5	5.5
dust	0.0	1.9		1.9	1.9	1.9	1.9	0.0
Loss (TgS/yr)	37.4	33.3		34.0	33.4	33.9	33.4	34.1
Dry deposition	4.7	4.6		4.7	4.6	4.7	4.6	4.0
$\text{SO}_4$	4.7	3.8		3.9	3.8	3.9	3.8	3.9
$\text{SO}_4$ on dust	0.0	0.8		0.8	0.8	0.8	0.8	0.0
Wet deposition	32.7	28.7		29.3	28.8	29.2	28.8	29.4
$\text{SO}_4$	32.7	27.5		28.2	27.6	28.1	27.6	28.3
$\text{SO}_4$ on dust	0.0	1.1		1.1	1.1	1.1	1.1	0.0
Lifetime (days)	4.9	3.8		3.8	3.8	3.8	3.8	4.3
$\text{NH}_x$								
$\text{NH}_3$ emission (TgN/yr)	54.5	54.5		54.5	54.5	54.5	54.5	54.5
Loss (TgN/yr)	54.8	55.0		54.8	54.8	55.0	54.2	55.0
Dry deposition	14.4	23.5		21.3	21.0	23.8	23.2	23.7
Wet deposition	40.4	30.7		32.6	32.9	30.4	30.3	30.5
Gas oxidation	0.0	0.8		0.9	0.9	0.8	0.7	0.7
Lifetime (days)	5.5	2.5		2.7	2.8	2.5	2.5	2.6
$\text{NO}_y$								
NO emission (TgN/yr)	51.4	51.8		51.8	51.7	51.7	51.8	51.8
Loss (TgN/yr)	51.3	51.0		51.2	51.0	51.1	51.0	51.1
Dry deposition	25.4	23.1		25.4	23.1	25.3	23.1	23.4
$\text{HNO}_3$	18.3	10.7		9.4	10.7	9.5	10.7	10.0
$\text{NO}_3$ on dust	0.0	3.4		3.3	3.4	3.3	3.4	0.0
$\text{NH}_4\text{NO}_3$	0.7	0.8		4.4	0.8	4.2	0.8	0.5
Organic nitrogen	3.9	4.0		4.0	4.0	4.0	4.0	4.0
Wet deposition	25.6	27.6		25.5	27.6	25.5	27.6	27.4
$\text{HNO}_3$	23.4	17.8		16.4	17.4	16.7	17.7	18.2
$\text{NO}_3$ on dust	0.0	3.7		3.7	3.7	3.7	3.7	0.0
$\text{NH}_4\text{NO}_3$	0.5	3.5		2.9	3.9	2.5	3.6	2.6
Organic nitrogen	1.7	2.6		2.6	2.6	2.6	2.6	2.6
Lifetime (days)	22.7	13.4		13.3	13.4	13.3	13.4	13.6

<sup>a</sup>  $\text{SO}_2$  emissions are  $74.0 \text{ TgS a}^{-1}$  including  $16.0 \text{ TgS a}^{-1}$

from DMS oxidation

Table S3: Summary of model evaluation<sup>a</sup>

	AM3	AM3N	AM3N_fdep_diu	AM3N_diu	AM3N_fdep	AM3N_ns	AM3N_nhet	AM3N_ndust
$\text{SO}_4$								
Aerosol								
US	0.07 (0.81)	-0.11 (0.89)	-0.06 (0.89)	-0.11 (0.89)	-0.05 (0.89)	-0.05 (0.89)	-0.06 (0.89)	-0.11 (0.89)
Europe	-0.43 (0.24)	-0.22 (0.62)	-0.13 (0.64)	-0.22 (0.62)	-0.13 (0.64)	-0.20 (0.67)	-0.13 (0.64)	-0.24 (0.62)
Wet deposition								
US	0.00 (0.42)	-0.07 (0.59)	-0.08 (0.57)	-0.07 (0.58)	-0.07 (0.58)	-0.08 (0.58)	-0.08 (0.58)	-0.07 (0.58)
Europe	-0.18 (0.53)	-0.32 (0.57)	-0.32 (0.53)	-0.32 (0.57)	-0.31 (0.57)	-0.32 (0.58)	-0.31 (0.57)	-0.32 (0.55)
$\text{NO}_3$								
Aerosol								
US	-0.61 (0.64)	1.03 (0.64)	0.17 (0.65)	0.99 (0.64)	0.16 (0.64)	1.38 (0.61)	0.42 (0.62)	1.06 (0.64)
Europe	-0.78 (0.62)	0.32 (0.62)	-0.30 (0.58)	0.29 (0.62)	-0.30 (0.58)	0.31 (0.61)	-0.13 (0.50)	0.17 (0.62)
Gas+Aerosol								
Europe	-0.18 (0.61)	0.17 (0.75)	-0.29 (0.57)	0.16 (0.75)	-0.29 (0.57)	0.17 (0.74)	-0.13 (0.62)	0.08 (0.76)
Wet deposition								
US	0.14 (0.33)	0.23 (0.52)	0.11 (0.54)	0.23 (0.52)	0.11 (0.54)	0.21 (0.52)	0.17 (0.54)	0.27 (0.54)
Europe	-0.32 (0.57)	-0.29 (0.54)	-0.39 (0.54)	-0.29 (0.54)	-0.39 (0.55)	-0.28 (0.58)	-0.32 (0.52)	-0.27 (0.56)
$\text{NH}_3$								
Gas								
US	-0.75 (0.50)	-0.10 (0.54)	-0.22 (0.53)	-0.29 (0.54)	-0.04 (0.53)	-0.12 (0.50)	-0.06 (0.53)	-0.11 (0.53)
Europe	-0.65 (0.48)	0.23 (0.54)	0.17 (0.50)	0.04 (0.53)	0.36 (0.52)	0.25 (0.46)	0.31 (0.53)	0.21 (0.54)
Gas+aerosol								
Europe	0.69 (0.66)	0.18 (0.64)	0.02 (0.64)	0.07 (0.63)	0.12 (0.64)	0.19 (0.60)	0.14 (0.64)	0.19 (0.64)
Wet deposition								
US	-0.20 (0.50)	-0.20 (0.69)	-0.15 (0.69)	-0.14 (0.69)	-0.20 (0.68)	-0.18 (0.55)	-0.20 (0.69)	-0.20 (0.69)
Europe	-0.23 (0.52)	-0.36 (0.58)	-0.32 (0.58)	-0.31 (0.58)	-0.36 (0.58)	-0.36 (0.49)	-0.36 (0.58)	-0.35 (0.58)
AOD								
MODIS								
World	0.09 (0.57)	-0.08 (0.68)	-0.08 (0.68)	-0.07 (0.68)	-0.08 (0.68)	-0.07 (0.67)	-0.08 (0.68)	-0.06 (0.69)
high $\text{NO}_3$	-0.15 (0.83)	0.11 (0.87)	0.09 (0.87)	0.14 (0.86)	0.06 (0.87)	0.13 (0.86)	0.06 (0.87)	0.12 (0.87)
high $\text{SO}_4$	0.57 (0.83)	0.06 (0.87)	0.06 (0.87)	0.08 (0.87)	0.04 (0.88)	0.06 (0.87)	0.04 (0.88)	0.09 (0.88)
MISR								
World	-0.03 (0.53)	-0.16 (0.59)	-0.16 (0.59)	-0.15 (0.58)	-0.17 (0.59)	-0.16 (0.58)	-0.17 (0.59)	-0.14 (0.59)
high $\text{NO}_3$	-0.12 (0.84)	0.21 (0.87)	0.18 (0.87)	0.24 (0.86)	0.16 (0.87)	0.23 (0.86)	0.16 (0.88)	0.24 (0.87)
high $\text{SO}_4$	0.54 (0.86)	0.12 (0.88)	0.12 (0.88)	0.14 (0.87)	0.10 (0.88)	0.12 (0.87)	0.10 (0.88)	0.15 (0.88)
AERONET								
World	-0.03 (0.72)	-0.10 (0.82)	-0.11 (0.82)	-0.08 (0.82)	-0.13 (0.82)	-0.09 (0.82)	-0.12 (0.83)	-0.07 (0.82)
high $\text{NO}_3$	-0.50 (0.87)	-0.01 (0.76)	-0.07 (0.70)	0.02 (0.72)	-0.08 (0.73)	0.03 (0.79)	-0.09 (0.74)	0.03 (0.75)
high $\text{SO}_4$	0.33 (0.47)	-0.10 (0.74)	-0.10 (0.71)	-0.07 (0.72)	-0.12 (0.73)	-0.10 (0.75)	-0.13 (0.72)	-0.05 (0.74)




Figure S1: Mass extinction of  $(\text{NH}_4)_2\text{SO}_4$  (dotted line) and  $\text{NH}_4\text{NO}_3$  (solid line) at 550nm. The dash line shows the relative error between  $(\text{NH}_4)_2\text{SO}_4$  and  $\text{NH}_4\text{NO}_3$  mass extinctions. Refractive index and hygroscopic growth for  $\text{NH}_4\text{NO}_3$  are take from Gosse et al. (1997) and Tang (1996) respectively.




Figure S2: Contribution of  $\text{SO}_4$ ,  $\text{NO}_3$ , OC, and dust to the simulated dry extinction at Bondville in AM3N\_fdep\_diu.




Figure S3: Observed (black dots) and simulated temperature over Bondville. The dots show the average for each altitude and the bars show the 25<sup>th</sup> and 75<sup>th</sup> percentiles. The model is sampled within one hour of the twice daily observations.




Figure S4: Observed (black dots) and simulated relative humidity over Bondville. The dots show the average for the each altitude and the bars show the 25<sup>th</sup> and 75<sup>th</sup> percentiles. The model is sampled within one hour of the twice daily observations.




Figure S5: Observed (black dots) and simulated (solid line from AM3N) occurrence of RH exceeding 90% over Bondville from 2008 to 2010. The model is sampled within one hour of the bidaily observations.




Figure S6: Comparison between observed and simulated  $\text{SO}_4^{2-}$  for AM3 and AM3N\_fdep\_diu. The ratio between model and observation is shown for each site and each season for AM3 (left panel) and AM3N\_fdep\_diu (middle panel). Observations are averaged on a monthly basis over the 2006–2012 period. Scatterplot of seasonal means is shown on the right for AM3 (black) and AM3N\_fdep\_diu (purple). Note the log scale for the x and y axis.




Figure S7: Same as Fig. S6 but for  $\text{NO}_3^-$  aerosol




Figure S8: Same as Fig. S6 but for  $\text{NH}_3$




Figure S9: Same as Fig. S6 but for  $\text{SO}_4^{2-}$  concentrations in rain from NADP




Figure S10: Same as Fig. S6 but for  $\text{NO}_3^-$  concentration in rain from NADP




Figure S11: Same as Fig. S6 but for  $\text{NH}_4^+$  concentration in rain from NADP




Figure S12: Same as Fig. S6 but for  $\text{SO}_4^{2-}$  surface observations from EMEP.




Figure S13: Same as Fig. S6 but for surface  $\text{NO}_3^-$  from EMEP.




Figure S14: Same as Fig. S6 but for NO<sub>y</sub> from EMEP.




Figure S15: Same as Fig. S6 but for  $\text{NH}_x$  from EMEP.




Figure S16: Same as Fig. S6 but for  $\text{SO}_4^{2-}$  concentration in precipitated water from EMEP




Figure S17: Same as Fig. S6 but for  $\text{NH}_4^+$  concentration in precipitated water from EMEP




Figure S18: Same as Fig. S6 but for  $\text{NO}_3^-$  concentration in precipitated water from EMEP




Figure S19: Same as Fig. S6 but for AOD from AERONET




Figure S20: Average diurnal profile of  $\text{NO}_3^-$  at the YRK SEARCH site ( $33.93^\circ\text{N}$ ,  $274.95^\circ\text{E}$ , 395m asl) in 2008. Simulated nitrate diurnal profiles from AM3N, AM3N\_diu, and AM3N\_fdep\_diu are shown in blue, red, and green respectively. Observations are shown in black

## References

- Crowley, J. N., Ammann, M., Cox, R. A., Hynes, R. G., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume V – heterogeneous reactions on solid substrates, *Atmos. Chem. Phys.*, 10, 9059–9223, doi:10.5194/acp-10-9059-2010, 2010.
- Gosse, S. F., Wang, M., Labrie, D., and Chylek, P.: Imaginary part of the refractive index of sulfates and nitrates in the 07–26- $\mu\text{m}$  spectral region, *Appl. Opt.*, 36, 3622, doi:10.1364/AO.36.003622, 1997.
- Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, 34, 2131–2159, doi:10.1016/S1352-2310(99)00462-8, 2000.
- Macintyre, H. L. and Evans, M. J.: Sensitivity of a global model to the uptake of  $\text{N}_2\text{O}_5$  by tropospheric aerosol, *Atmos. Chem. Phys.*, 10, 7409–7414, doi:10.5194/acp-10-7409-2010, 2010.
- Mao, J., Fan, S., Jacob, D. J., and Travis, K. R.: Radical loss in the atmosphere from Cu-Fe redox coupling in aerosols, *Atmos. Chem. Phys.*, 13, 509–519, doi:10.5194/acp-13-509-2013, 2013.
- Tang, I. N.: Chemical and size effects of hygroscopic aerosols on light scattering coefficients, *J. Geophys. Res. Atmos.*, 101, 19 245–19 250, doi:10.1029/96JD03003, 1996.
- Ullerstam, M., Vogt, R., Langer, S., and Ljungström, E.: The kinetics and mechanism of  $\text{SO}_2$  oxidation by  $\text{O}_3$  on mineral dust, *Phys. Chem. Chem. Phys.*, 4, 4694–4699, doi:10.1039/B203529B, 2002.
- Ullerstam, M., Johnson, M. S., Vogt, R., and Ljungström, E.: DRIFTS and Knudsen cell study of the heterogeneous reactivity of  $\text{SO}_2$  and  $\text{NO}_2$  on mineral dust, *Atmos. Chem. Phys.*, 3, 2043–2051, doi:10.5194/acp-3-2043-2003, 2003.
- Vlasenko, A., Sjogren, S., Weingartner, E., Stemmler, K., Gäggeler, H. W., and Ammann, M.: Effect of humidity on nitric acid uptake to mineral dust aerosol particles, *Atmos. Chem. Phys.*, 6, 2147–2160, doi:10.5194/acp-6-2147-2006, 2006.
- Wong, K. W., Oh, H.-J., Lefer, B. L., Rappenglück, B., and Stutz, J.: Vertical profiles of nitrous acid in the nocturnal urban atmosphere of Houston, TX, *Atmos. Chem. Phys.*, 11, 3595–3609, doi:10.5194/acp-11-3595-2011, 2011.