



Supplement of

Multidecadal variations of the effects of the Quasi-Biennial Oscillation on the climate system

Stefan Brönnimann et al.

Correspondence to: Stefan Brönnimann (stefan.broennimann@giub.unibe.ch)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.



Fig. S1. (left) Registering balloon measurements on Lake Victoria during Berson's East Africa expedition. (right) Stratospheric zonal wind in the profiles from Berson's expedition (note that two ascents were made on 30 Aug 1908). Also indicated are corresponding winds from the reconstructed QBO ("REC").



Fig. S2. Profiles of zonal wind from Batavia, 1910-1911. The QBO phase in the top line refers to the reconstructed QBO at 90 hPa.



Fig. S3. Schematic of the analysis periods and data sets used in this study. Red dotted lines indicate the main analysis periods, black arrow indicate that the data sets reach further back than indicated on the *x*-axis.



Fig. S4. Composite fields (shading) and climatologies (contours) since 1979 in (top) 29CRv2c and (bottom) ERA-20C, Jan.-Mar., for easterly minus westerly QBO phases (Nov.-Dec.) for zonal averages of (a) zonal wind (contours: 20 to 60 m/s in steps of 10 m/s), (b) temperature (contours: 200 to 300 K in steps of 10 K) and (c) GPH. Yellow dashed lines indicate significant difference (p = 0.05).



Fig. S5. Composite fields for easterly minus westerly QBO phases of 200 hPa zonal wind, sea-level pressure, SAT, and snow cover in 20CR. Dashed lines indicate differences that are significant at the 95% confidence level (t-test).



Fig. S6. Composite fields for easterly minus westerly QBO phases for 100 hPa geopotential height in Jan.-Mar., 1908-1957 in reconstructions, 20CRv2c, and ERA-20C. Contours (spacing is 200 gpm) indicate climatology.



Fig. S7. Composite fields for easterly minus westerly QBO phases for snow depth in ERA-20C for (left) 1908-1957, (middle) 1958-2010 and (right) 1908-2010.



Fig. S8. Composite fields of (left) 100 hPa geopotential height and (right) sea-level pressure for easterly minus westerly QBO phases from the FUPSOL simulations (top to bottom: F13, F14, F23, F24) for Jan.-Mar. Hatching denotes differences that are significant at the 95% confidence level (t-test).



Fig. S9. Composite fields of (left) 2 m temperature and (right) snow depth for easterly minus westerly QBO phases from the FUPSOL simulations (top to bottom: F13, F14, F23, F24) for Jan.-Mar. Hatching denotes differences that are significant at the 95% confidence level (t-test).



Fig. S10. Composite fields of 500 hPa omega for easterly minus westerly QBO phases from the FUPSOL simulations (top to bottom: F13, F14, F23, F24) for (left) Dec.-Feb and (right) Jun.-Aug. Hatching denotes differences that are significant at the 95% confidence level (t-test).



Fig. S11. Same as Fig. S10 but for sea-surface temperatures.



Fig. S12. Composite fields for easterly minus westerly QBO phases for (left) 500 hPa vertical velocity, 10 m zonal wind, SST, and 200 hPa zonal wind for boreal winter in 20CRv2c. Dashed lines indicate differences that are significant at the 95% confidence level (t-test).