



Supplement of

**Profiling of aerosol microphysical properties at several
EARLINET/AERONET sites during the July 2012
ChArMEx/EMEP campaign**

María José Granados-Muñoz et al.

Correspondence to: María José Granados Muñoz (mamunoz@jpl.nasa.gov)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

Supplementary material:




Figure S1. Dust mass concentration profiles forecast by BSC-DREAM8b-v2, DREAM8-NMME, NMMB/BSC-Dust and COSMO-MUSCAT at Athens in the period 9 July 2012-11 July 2012. Only those data coincident with AERONET inversion retrievals (± 3 hours) has been presented here.




Figure S2. Dust mass concentration profiles forecast by BSC-DREAM8b-v2, DREAM8-NMME, DREAMABOL and NMMB/BSC-Dust at Barcelona in the period 9 July 2012-11 July 2012. Only those data coincident with AERONET inversion retrievals (± 3 hours) has been presented here.




Figure S3. Dust mass concentration profiles forecast by BSC-DREAM8b-v2, DREAM8-NMME, NMMB/BSC-Dust and COSMO-MUSCAT at Bucharest in the period 9 July 2012-11 July 2012. Only those data coincident with AERONET inversion retrievals (± 3 hours) has been presented here.




Figure S4. Dust mass concentration profiles forecast by BSC-DREAM8b-v2, DREAM8-NMME, NMMB/BSC-Dust and COSMO-MUSCAT at Évora in the period 9 July 2012-11 July 2012. Only those data coincident with AERONET inversion retrievals (± 3 hours) has been presented here.




Figure S5. Correlation coefficient R between simultaneous LIRIC and modelled time series (solid black lines) for every altitude level. Dashed lines represent R values obtained for the modelled time series and LIRIC time series with a time shift of plus (red line) and minus (blue line) 3 hours.