

Supplement of Atmos. Chem. Phys., 16, 7725–7741, 2016
<http://www.atmos-chem-phys.net/16/7725/2016/>
doi:10.5194/acp-16-7725-2016-supplement
© Author(s) 2016. CC Attribution 3.0 License.



Atmospheric
Chemistry
and Physics
Open Access
The EGU logo features the letters 'EGU' in a bold, sans-serif font, with a circular arrow graphic to the left of the 'E'.

Supplement of

Understanding isoprene photooxidation using observations and modeling over a subtropical forest in the southeastern US

Luping Su et al.

Correspondence to: John E. Mak (john.mak@stonybrook.edu)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

1 **1 Algorithms used in ToFIND**

2 **1.1 Peak shape fitting**

3 The peak generated by the PTR-TOF-MS is asymmetric and can be mathematically
4 approximated by peak-shaped functions. Here we choose an empirically transformed
5 Gaussian function (ETG) model to describe a single peak produced by the instrument (Li,
6 1997). The ETG is based on the decomposition of Gaussian function into leading and trailing
7 edge functions. The two functions are weakly combined and subsequently modified, making it
8 very suitable for approximating asymmetric peak shapes. The ETG is given by:

9
$$f(t) = \frac{H}{\{1 + \lambda_L \exp[k_L(t_L - t)]\}^\alpha + \{1 + \lambda_T \exp[k_T(t - t_T)]\}^\beta - 1} \quad (\text{S1})$$

10 where $f(t)$ denotes peak function; t denotes time-of-flight; H is related to peak height; λ_L and
11 λ_T are pre-exponential parameters; k_L and k_T are the parameters related to the speeds of
12 the rise and fall of the leading and trailing edges, respectively; t_L and t_T are the inflection
13 times of the leading and trailing edges, respectively, and are fixed values; α and β are the
14 parameters to further modify the shapes of the lading and trailing edges, respectively. The
15 ETG features in a weak link between the leading and trailing parts of a single peak, thus is
16 very flexible and suitable for fitting asymmetric peaks. The values of t_L and t_T do not need
17 to be accurate, here we use the time-of-flight at the full width at half maximum (FWHM) on
18 the left and right side of a single peak to represent those two values, respectively. There are a
19 total of seven parameters in each ETG (H , λ_L , λ_T , k_L , k_T , α , and β), and the rate of
20 convergence is fast. Levenberg-Marquardt algorithm is used to solve the non-linear least-
21 squares curve fitting problem. To achieve a sufficiently high speed of the curve fitting,
22 MINPACK-1 Least Squares Fitting Library in C
23 (<http://www.physics.wisc.edu/~craigm/idl/cmpfit.html>) is used and integrated into the
24 ToFIND toolbox through MATLAB-C interface (<http://www.imm.dtu.dk/~guuru/>).

25 **1.2 Time-of-flight to m/z conversion**

26 In the field free area of the TOF-MS, the time-of-flight of ions show slow drift due to the
27 change in the length of the ions' flight path caused by temperature variations in this area. To
28 gain a high mass accuracy, continuous mass scale correction is performed on the raw data.
29 Four well identified peaks: three internal ions ($\text{H}_2^{18}\text{OH}^+$, $m/z=21.022$, NO^+ , $m/z=29.997$,

1 and H216OH218OH+, m/z=39.033), and one externally added compound (diiodomethane,
2 fragmented at ion CH2I+, m/z=140.920) are used for mass scale correction peaks. The time-
3 of-flight t and mass scale m/z are related to each other through:

4 $t = a\sqrt{m/z} + b$ (S2)

5 For each raw HDF5 data file, every 3 of the 300 spectra are averaged to produce 100 spectra.
6 For each of the 100 spectra, the peak shape fitting algorithm described above is applied to
7 find the peak centers (t) of the four mass scale correction peaks. Since the exact m/z values
8 for each peak are known, the values of a and b in Eqn. A2 are calculated through linear least-
9 squares method. The relative mass error (RME) is less than 10 parts per million (ppm) over
10 the spectrum after mass scale calibration.

11 **1.3 Peak detection**

12 After time-of-flight to m/z conversion, the 300 mass scale corrected spectra are aligned with
13 each other, and averaged to produce one single spectrum. A target mass list is predefined
14 which includes a series exact m/z of interested VOCs compounds (e.g., methanol, acetone,
15 isoprene, MVK+MACR, and monoterpenes). For one specific target peak, there may be other
16 peaks co-existing in the same nominal m/z, a peak searching routine is used to find all the
17 peaks including the target peak within a certain range of one nominal m/z. The mass list is
18 then updated and used for peak shape fitting (see next section).

19 **1.4 Signal integration**

20 A few issues exist when performing signal integration on each individual peak: (1), due to the
21 low counts rate in each individual 1 Hz spectrum, the peak shape fitting algorithm often fails
22 to converge or generate desirable fitting results when applied to a single spectrum, (2),
23 interference exists between two neighbouring peaks when they overlap with each other due to
24 the limited mass resolution of the PTR-TOF-MS. To improve the accuracy of the peak fitting
25 result, the updated peak list is used and the peak shape fitting algorithm is applied to the
26 averaged spectrum described above to produce a peak function. Based on this, the FWHM for
27 each peak is recalculated and used to define the peak integration boundaries. For the
28 asymmetrical Gaussian-shaped peak which is typical for the PTR-TOF-MS, the integration
29 boundaries on the left and right sides are processed separately. With the updated FWHM
30 values for all the peaks, the ratios of the portion before the peak center to the portion after the

1 peak center are calculated and averaged (r). The integration boundary on the left side is
 2 defined as $a \times \text{FWHM} \times r$ to the left of the peak center. The integration boundary on the right
 3 side is defined as $a \times \text{FWHM} \times (1-r)$ to the right of the peak center. Add a figure to describe the
 4 procedure. The factor a is set to 4 for single peak and 3 for multiple peaks. The factor value is
 5 reduced for multiple peaks to lower the interference of neighbouring peaks.

6 **2 Conversion rate of ISOPOOH to C₄H₇O⁺ in PTR-TOF-MS**

7 To quantify the conversion rate of ISOPOOH to C₄H₇O⁺ when using PTR-TOF-MS, an
 8 experiment was carried out by using 1,2-ISOPPOOH standard during the Focused Isoprene
 9 eXperiment at the California Institute of Technology (FIXCIT) campaign (Nguyen et al.,
 10 2014). The same PTR-TOF-MS was used to perform the measurements. There were three
 11 steps (Figure S0): 1, sampling from ISOPOOH-free air; 2, sampling from Teflon bag filled
 12 with ISOPOOH standard under room temperature through a 1/8 inch O.D. PFA tubing (length
 13 ~40 cm) at ~25 °C; 3, same setup as 2 except that the PFA tubing is submerged in cold trap at
 14 -40 °C. The decreasing of m/z71.049 signal in step 3 is due to the condensing of ISOPOOH
 15 onto the PFA tubing wall surface under low temperature as a result of its low vapor pressure.

16 Signals at peak m/z=71.049 are integrated using method described above. To ensure the
 17 robustness of the results, two independent methods are used to convert the integrated signal
 18 (in unit of cps) at peak m/z=71.049 to its mixing ratio (in unit of ppbv).

19 **2.1 Using reaction rate coefficient**

$$20 \quad \text{VOC[ppbv]} = 10^9 \times \frac{\text{VOC[molec. cm}^{-3}\text{]}}{\text{air[molec. cm}^{-3}\text{]}} \quad (\text{S3})$$

$$21 \quad \text{VOC[molec. cm}^{-3}\text{]} = \frac{1}{kt} \times \frac{\text{VOCH}^+[\text{cps}]}{\text{H}_3\text{O}^+[\text{cps}]} \times \sqrt{\frac{\text{m/z}_{\text{H}_3\text{O}^+}}{\text{m/z}_{\text{VOCH}^+}}} \times \frac{\text{TR}_{\text{H}_3\text{O}^+}}{\text{TR}_{\text{VOCH}^+}} \quad (\text{S4})$$

$$22 \quad t = \frac{l^2}{\mu_0 U_{drift}} \times \frac{T_0}{T_{drift}} \times \frac{P_{drift}}{P_0} \quad (\text{S5})$$

$$23 \quad \text{air[molec. cm}^{-3}\text{]} = \frac{N_A}{22400} \times \frac{T_0}{T_{drift}} \times \frac{P_{drift}}{P_0} \quad (\text{S6})$$

24 the meaning of the symbols are listed as follows,

25

Symbol	Value	Unit	Description
VOC[ppbv]	-	ppbv	VOC mixing ratio
VOC[molec. cm ⁻³]	-	molec. cm ⁻³	VOC number density
air[molec. cm ⁻³]	-	molec. cm ⁻³	air number density
VOCH ⁺ [cps]	-	cps	measured VOCH ⁺ count rates
H ₃ O ⁺ [cps]	-	cps	measured primary ion count rates
μ_0	2.8	cm ² V ⁻¹ s ⁻¹	reduced mobility
p_0	1013.25	mbar	air pressure at standard condition
p_{drift}	2.20	mbar	drift tube pressure
T_0	273.15	K	temperature at standard condition
T_{drift}	332.6	K	drift tube temperature
N_A	6.022×10^{23}	1	Avogadro number
l	9.3	cm	length of the reaction chamber
$TR_{H_3O^+}$	0.67	1	transmission factor of H ₃ O ⁺
TR_{VOCH^+}	1.00	1	transmission factor of VOCH ⁺
k	3.2×10^{-9} ^a	cm ³ s ⁻¹	reaction rate coefficient between VOC and H ₃ O ⁺

1 ^aData obtained from Cappellin et al. (2012).

2 2.2 Using sensitivity

$$3 \quad VOC[\text{ppbv}] = \frac{VOCH^+[\text{ncps}]}{\text{Sensitivity}_{VOCH^+}[\text{ncps ppbv}^{-1}]} \quad (\text{S7})$$

$$4 \quad VOCH^+[\text{ncps}] = \frac{VOCH^+[\text{cps}] \times 10^6}{H_3O^+[\text{cps}]} \times \sqrt{\frac{m/z_{H_3O^+}}{m/z_{VOCH^+}}} \quad (\text{S8})$$

5 where $\text{Sensitivity}_{C4H_7O^+} = 26.83 \text{ ncps ppbv}^{-1}$ by using standard gas calibration.

1 **2.3 Conversion rate**

2 The mixing ratios (mean \pm 1 standard deviation) of m/z71.049 during experiment step 2
3 (Figure S0) are calculated to be 36.20 ± 1.62 ppbv and 34.89 ± 1.54 ppbv by using reaction
4 rate coefficient and sensitivity, respectively. The difference of mixing ratios between the two
5 methods is within 4%. The mixing ratio of m/z71.049 obtained from the sensitivity method is
6 used to calculate the conversion rate of ISOPOOH. The mixing ratio of 1,2-ISOPOOH in the
7 Teflon bag is quantified to be 250 ppbv ($\pm 40\%$ uncertainty) (Nguyen et al., 2014) (A. Teng,
8 personal communication). The conversion rate of ISOPOOH to $C_4H_7O^+$ is estimated to be
9 14% (+14%/-6%).

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27




Figure S0. The mixing ratios of m/z71.049 during an ISOPOOH cold trap experiment. 1, sampling from ISOPOOH-free air; 2, sampling from Teflon bag filled with ISOPOOH standard under room temperature through a 1/8 inch OD PFA tubing (length ~40 cm) at ~25 °C; 3, same setup as 2 except that the PFA tubing is submerged in cold trap at -40 °C.

1

2 Table S1. Other observations during the 2013 SAS campaign used in this study. Days used
 3 are during June 5, 6, 8, 10-13 unless otherwise specified in the table.

Parameter	Platform	Instrument/technique	Uncertainty	Days used	Institute
VOC	Airborne (NCAR C-130)	Proton Transfer Reaction Mass Spectrometer (PTR-MS)	20%	June 12	NCAR, NOAA
ISOPN	SEARCH site	chemical ionization mass spectrometer (CIMS)	25%	June 5, 6, 8, 10-13	Purdue University
O ₃ , NO _x	Airborne (NCAR C-130)	chemiluminescence	10% for NO, 15% for NO ₂ , 5% for O ₃ 5.5% for NO, 15.0% for NO ₂ , 6.1% for O ₃	June 12	NCAR
O ₃ , NO _x	SEARCH tower	ThermoScientific 49i, 42i	32%	June 5, 6, 8, 10-13	ARA
OH	SEARCH tower	laser induced fluorescence (LIF)	15%	June 5, 6, 8, 10-13	Pennsylvania State University
HCHO	SEARCH tower	Fiber - Laser Induced Fluorescence		June 14	University of Wisconsin–Madison
Surface heat flux	AABC tower	Integrated Surface Flux System (ISFS)		June 10-13	NCAR
Potential temperature	AABC tower	ISFS		June 10-13	NCAR
specific humidity	Airborne (NCAR C-130)	onboard sensor		June 12	NCAR
Potential temperature	Airborne (NCAR C-130)	onboard sensor		June 12	NCAR
specific humidity	SEARCH	Ceilometer, Integrated Sounding System (ISS)	13.0% for ceilometer	June 10-13	ARA, NCAR

4

5

6

7

8

9

10

11

1
2

3 Table S2. Chemical species in the complex scheme. See more details at
4 http://wiki.seas.harvard.edu/geos-chem/index.php/New_isoprene_scheme

Species	Formula	Note
A3O2	CH3CH2CH2OO	primary RO2 from C3H8
ACET	CH3C(O)CH3	acetone
ACTA	CH3C(O)OH	acetic acid
ALD2	CH3CHO	acetaldehyde
ALK4	RH	≥C4 alkanes
ATO2	CH3C(O)CH2O2	RO2 from acetone
ATO OH	CH3C(O)CH2OOH	ATO2 peroxide
B3O2	CH3CH(OO)CH3	secondary RO2 from C3H8
C2H6	C2H6	ethane
C3H8	C3H8	propane
CH2O	CH2O	formaldehyde
CH4	CH4	methane
CO	CO	carbon monoxide
CO2	CO2	carbon dioxide
DHMOB	HOCH2C(CH3)(OH)C(=O)CHO	See Paulot et al., ACP (2009)
DIBOO		Dibble peroxy radical
EOH	C2H5OH	ethanol
ETHLN	CHOCH2ONO2	Ethanal nitrate
ETO2	CH3CH2OO	ethylperoxy radical
ETP	CH3CH2OOH	ethylhydroperoxide
GLYC	HOCH2CHO	glycoaldehyde (hydroxyacetaldehyde)
GLYX	CHOCHO	glyoxal
H2	H2	hydrogen atom
H2O	H2O	water vapor
H2O2	H2O2	hydrogen peroxide
HAC	HOCH2C(O)CH3	hydroxyacetone
HCOOH	HCOOH	formic acid
HC5	HOCH2CH=C(CH3)CHO	Hydroxycarbonyl with 5C
HC5OO		Peroxy radical from HC5 (old IAO2?)
HNO2	HONO	nitrous acid
HNO3	HNO3	nitric acid
HNO4	HNO4	pernitric acid
HO2	HO2	hydroperoxyl radical
IALD	HOCH2C(CH3)=CHCHO	hydroxy carbonyl alkenes from isoprene
IAP	HOCH2C(CH3)(OOH)CH(OH)CHO	peroxide from IAO2
IEPOX		Isoprene epoxide
IEPOXOO		RO2 from IEPOX
INO2	O2NOCH2C(OO)(CH3)CH=CH2	RO2 from ISOP+NO3

INPN	O2NOCH2C(OOH)(CH3)CH=CH2	peroxide from INO2
ISN1		nighttime isoprene nitrate
ISNOOA		peroxy radical from ISN1
ISNOOB		peroxy radical from ISN1
ISNOHOO		peroxy radical from ISN1
ISNP	HOCH2C(OOH)(CH3)CH(ONO2)CH2OH	peroxide from ISOPNBO2 and ISOPNDO2
ISOP	CH2=C(CH3)CH=CH2	isoprene
ISOPNB	C5H9NO4	Isoprene nitrate Beta
ISOPND	C5H9NO4	Isoprene nitrate Delta
KO2	RO2 from >3 ketones	RO2 from >3 ketones
M		for three body reactions
MACR	CH2=C(CH3)CHO	methacrolein
MACRN	HOCH2C(ONO2)(CH3)CHO	Nitrate from MVK
MAN2	HOCH2C(ONO2)(CH3)CHO	RO2 from MACR+NO3
MAO3	CH2=C(CH3)C(O)OO	peroxyacetyl from MVK and MACR
MAOP	CH2=C(CH3)C(O)OOH	peroxide from MAO3
MAOPO2	CH2OH-CHOO*CH3C(O)OOH	Peroxy radical from MAOP (addition on the double bond)
MAP	CH3C(O)OOH	peroxyacetic acid
MCO3	CH3C(O)OO	peroxyacetyl radical
MEK	RC(O)R	>3 ketones
MGLY	CH3COCHO	methylglyoxal
MNO3	CH3ONO2	methylnitrate
MOBA	HOC(=O)C(CH3)=CHCHO	5C acid from isoprene
MOBAOO		RO2 from MOBA
MO2	CH3O2	methylperoxy radical
MOH	CH3OH	methanol
MP	CH3OOH	methylhydroperoxide
MRO2	HOCH2C(OO)(CH3)CHO	RO2 from MACR+OH
MRP	HOCH2C(OOH)(CH3)CHO	peroxide from MRO2
MVK	CH2=CHC(=O)CH3	methylvinylketone
MVKN	HOCH2CH(ONO2)C(=O)CH3	Nitrate from MACR
N2	N2	nitrogen
N2O	N2O	nitrous oxide
N2O5	N2O5	dinitrogen pentoxide
NH2	NH2	ammonia radical
NH3	NH3	ammonia
NO	NO	nitric oxide
NO2	NO2	nitrogen dioxide
NO3	NO3	nitrate radical
O2	O2	molecular oxygen
O2CH2OH	O2CH2OH	produced by CH2O+HO2
O3	O3	ozone
OH	OH	hydroxyl radical
PAN	CH3C(O)OONO2	peroxyacetyl nitrate
PMN	CH2=C(CH3)C(O)OONO2	peroxymethacryloyl nitrate (MPAN)

PO2	HOCH2CH(OO)CH3	RO2 from isoprene
PP	HOCH2CH(OOH)CH3	peroxide from PO2
PPN	CH3CH2C(O)OONO2	peroxypropionyl nitrate
PRN1	O2NOCH2CH(OO)CH3	RO2 from propene + NO3
PRPE	C3H6	\geq C4 alkenes
PRPN	O2NOCH2CH(OOH)CH3	peroxide from PRN1
PROPN	CH3C(=O)CH2ONO2	Propanone nitrate
PYAC	CH3COCOOH	Pyruvic acid
R4N1	RO2 from R4N2	RO2 from R4N2
R4N2	RO2NO	\geq C4 alkyl nitrates
R4O2	RO2 from ALK4	RO2 from ALK4
R4P	CH3CH2CH2CH2OOH	peroxide from R4O2
RA3P	CH3CH2CH2OOH	peroxide from A3O2
RB3P	CH3CH(OOH)CH3	peroxide from B3O2
RCHO	CH3CH2CHO	>C2 aldehydes
RCO3	CH3CH2C(O)OO	peroxypropionyl radical
RCOOH	C2H5C(O)OH	>C2 organic acids
RIO1	HOCH2C(OO)(CH3)CH=CHOH	RO2 from isoprene oxidation products
RIO2	HOCH2C(OO)(CH3)CH=CH2	RO2 from isoprene (named as ISOPPO2 in the literature)
RIP	HOCH2C(OOH)(CH3)CH=CH2	peroxide from RIO2 (named as ISOPPOOH in the literature)
ROH	C3H7OH	>C2 alcohols
RP	CH3CH2C(O)OOH	peroxide from RCO3
VRO2	HOCH2CH(OO)C(O)CH3	RO2 from MVK+OH
VRP	HOCH2CH(OOH)C(O)CH3	peroxide from VRO2

1
2
3
4
5
6
7
8
9
10
11
12

1

2

3 Table S3. The complex chemical scheme used in MXLCH.

Reaction	Reaction rate
O + O2 + M -> O3 + M	6E-34*(300/T)^2.4
O + O3 -> 2O2	8.00E-12*exp(-2060/T)
O1D + N2 -> O + N2	2.10E-11*exp(115/T)
O1D + O2 -> O + O2	3.20E-11*exp(70/T)
O1D + H2O -> 2OH	2.20E-10
H2 + O1D -> HO2 + OH	1.10E-10
H2 + OH -> H2O + HO2	5.50E-12*exp(-2000/T)
O + OH -> HO2 + O2	2.20E-11*exp(120/T)
HO2 + O -> OH + O2	3.00E-11*exp(200/T)
OH + O3 -> HO2 + O2	1.70E-12*exp(-940/T)
HO2 + O3 -> OH + 2O2	1.00E-14*exp(-490/T) (2.3E-13*exp(600/T)+1.7E-33*[M]*exp(1000/T))*(1+1.4E-
2HO2 -> H2O2 + O2	21*[H2O]*exp(2200/T))
H2O2 + OH -> H2O + HO2	2.90E-12*exp(-160/T)
OH + HO2 -> H2O + O2	4.80E-11*exp(250/T)
2OH -> H2O + O	4.20E-12*exp(-240/T)
2OH -> H2O2	6.29E-12
N2O + O1D -> N2 + O2	4.90E-11
N2O + O1D -> 2NO	6.70E-11
NO + HO2 -> NO2 + OH	3.50E-12*exp(250/T)
NO + O3 -> NO2 + O2	3.00E-12*exp(-1500.0/T)
NO2 + O -> NO + O2	5.60E-12*exp(180/T)
NO2 + O3 -> NO3 + O2	1.20E-13*exp(-2450/T)
NO3 + HO2 -> OH + NO2	2.30E-12*exp(170/T)
NO2 + NO3 -> N2O5	1.16E-12
N2O5 -> NO2 + NO3	4.70E-02
NO2 + OH -> HNO3	3.50E-12*exp(340/T)
HNO3 + OH -> NO3	1.59E-13
NO3 + NO -> 2NO2	1.50E-11*exp(170/T)
NO2 + HO2 -> HO2NO2	1.37E-12
HO2NO2 + OH -> H2O + NO2 + O2	1.30E-12*exp(380/T)
HO2NO2 -> HO2 + NO2	8.50E-02
CH4 + OH -> MO2 + H2O	2.45E-12*exp(-1775/T)
CH4 + O1D -> .75MO2 + .75OH + .25CH2O + .4HO2 + .05H2	1.50E-10
MO2 + NO -> CH2O + NO2 + HO2	2.80E-12*exp(300/T)
2MO2 -> 2CH2O + 2HO2	5.00E-13*exp(-424/T)
2MO2 -> CH2O + MOH	1.90E-14*exp(706/T)
MO2 + HO2 -> MP + O2	4.10E-13*exp(750/T)
MP + OH -> .7MO2 + .3OH + .3CH2O + H2O	3.80E-12*exp(200/T)

CH2O + OH -> CO + H2O + HO2	9.00E-12
CO + OH -> (CO2) + HO2	2.40E-13
MOH + OH -> HO2 + CH2O	7.30E-12*exp(-620/T)
ISOP + OH -> RIO2	3.1E-11*exp(350/T)
MACR + OH -> 0.53MAO3 + 0.47MRO2	8.0E-12*exp(380/T)
MVK + OH -> VRO2	2.6E-12*exp(610/T)
PMN + OH -> HAC + CO + NO2	2.90E-11
GLYC + OH -> 0.732CH2O + (0.361CO2) + 0.505CO + 0.227OH + 0.773HO2 + 0.134GLYX + 0.134HCOOH	6.54E-12
GLYC + OH -> HCOOH + OH + CO	1.45E-12
GLYX + OH -> HO2 + 2CO	3.1E-12*exp(340/T)
HAC + OH -> MGLY + HO2	1.81E-12*exp(305/T)
HAC + OH -> 0.5HCOOH + OH + 0.5ACTA + (0.5CO2) + 0.5CO + 0.5MO2	3.43E-13*exp(305/T)
PRPN + OH -> 0.209PRN1 + 0.791OH + 0.791PROPNN	8.78E-12*exp(200/T)
ETP + OH -> 0.64OH + 0.36ETO2 + 0.60ALD2	5.18E-12*exp(200/T)
RA3P + OH -> 0.64OH + 0.36A3O2 + 0.64RCHO	5.18E-12*exp(200/T)
RB3P + OH -> 0.791OH + 0.209B3O2 + 0.791ACET	8.78E-12*exp(200/T)
R4P + OH -> 0.791OH + 0.209R4O2 + 0.791RCHO	8.78E-12*exp(200/T)
RP + OH -> RCO3	6.13E-13*exp(200/T)
PP + OH -> 0.791OH + 0.209PO2 + 0.791HAC	8.78E-12*exp(200/T)
RIP + OH -> 0.387RIO2 + 0.613OH + 0.613HC5	4.75E-12*exp(200/T)
RIP + OH -> OH + IEPOX	1.9E-11*exp(390/T)
IEPOX + OH -> IEPOXOO	5.78E-11*exp(-400/T)
IAP + OH -> 0.654OH + 0.654DHMOB + 0.346HC5OO	5.31E-12*exp(200/T)
VRP + OH -> 0.791OH + 0.791MEK + 0.209VRO2	8.78E-12*exp(200/T)
MRP + OH -> MRO2	1.84E-12*exp(200/T)
MRP + OH -> (CO2) + HAC + OH	4.40E-12*exp(380/T)
MAOP + OH -> MAO3	6.13E-13*exp(200/T)
MAOP + OH -> MAOP02	3.60E-12*exp(380/T)
OH + MAP -> 1.0MCO3	6.13E-13*exp(200/T)
HC5 + OH -> HC5OO	3.35E-11*exp(380/T)
ISOPND + OH -> ISOPNDO2	2.64E-11*exp(380/T)
ISOPNB + OH -> ISOPNBO2	3.61E-12*exp(380/T)
ISNP + OH -> 0.612OH + 0.612R4N1 + 0.193ISOPNBO2 + 0.193ISOPNDO2	4.75E-12*exp(200/T)
MVKN + OH -> 0.650HCOOH + NO3 + 0.650MGLY + 0.350CH2O + 0.350PYAC	1.5E-12*exp(380/T)
MACRN + OH -> 1.0MACRNO2	1.39E-11*exp(380/T)
DHMOB + OH -> 1.5CO + 1.0HO2 + 0.5HAC + 0.5MEK	2.52E-11*exp(410/T)
MOBA + OH -> MOBAOO	2.79E-11*exp(380/T)
ETHLN + OH -> CH2O + (CO2) + NO2	1.00E-11
PROPNN + OH -> NO2 + MGLY	1.00E-15
ATOOH + OH -> ATO2 + H2O	2.66E-12*exp(200/T)
ATOOH + OH -> MGLY + OH + H2O	1.14E-12*exp(200/T)
R4N2 + OH -> R4N1 + H2O	1.60E-12
RIO2 + NO -> 0.883NO2 + 0.783HO2 + 0.660CH2O + 0.400MVK + 0.260MACR + 0.070ISOPND + 0.047ISOPNB + 0.123HC5 +	2.70E-12*exp(350/T) ^a

0.1DIBOO

VRO2 + NO -> 0.88NO2 + 0.35HO2 + 0.35CH2O + 0.53MCO3 + 0.53GLYC + 0.35MGLY + 0.12MVKN	2.70E-12*exp(350/T)
MRO2 + NO -> 0.85NO2 + 0.85HO2 + 0.122MGLY + 0.728HAC + 0.728CO + 0.122CH2O + 0.15MACRN	2.70E-12*exp(350/T)
MAN2 + NO -> 1.5NO2 + 0.5CH2O + 0.5MGLY + 0.5PROPNN + 0.5CO + 0.5OH	2.70E-12*exp(350/T)
IEPOXOO + NO -> 0.725HAC + 0.275GLYC + 0.275GLYX + 0.275MGLY + 0.125OH + 0.825HO2 + (0.200CO2) + 0.375CH2O + 0.074HCOOH + 0.251CO + NO2	2.70E-12*exp(350/T)
MAOPO2 + NO -> 1.0HAC + (1.0CO2) + 1.0OH + 1.0NO2	2.48E-12
MAOPO2 + NO -> 1.0HNO3	2.24E-13
HC5OO + NO -> NO2 + 0.216GLYX + 0.234MGLY + 0.234GLYC + 0.216HAC + 0.290DHMOB + 0.170MOBA + 0.090RCHO + HO2 + 0.090CO	2.48E-12
HC5OO + NO -> HNO3	3.42E-13
ISOPNDO2 + NO -> 0.070MACRN + 0.310HCOOH + 0.440HAC + 0.130ETHLN + 0.650CH2O + 1.34NO2 + 0.150GLYC + 0.310NO3 + 0.150PROPNN + 0.340MEK + 0.350HO2	2.48E-12
ISOPNDO2 + NO -> HNO3	3.42E-13
ISOPNBO2 + NO -> 0.6GLYC + 0.6HAC + 0.4CH2O + 1.6NO2 + 0.26MACRN + 0.4HO2 + 0.14MVKN	2.48E-12
ISOPNBO2 + NO -> HNO3	3.42E-13
MACRNO2 + NO -> 0.08ACTA + 0.08CH2O + 0.15NO3 + 0.07HCOOH + 0.070MGLY + 0.850HAC + 0.85NO2 + (0.93CO2) + 1.0NO2	2.7E-12*exp(350/T)
DIBOO + NO -> HO2 + NO2 + 0.520GLYC + 0.520MGLY + 0.480HAC + 0.480GLYX	2.48E-12
DIBOO + NO -> HNO3	3.42E-13
MOBAOO + NO -> RCHO + (CO2) + HO2 + NO2	2.48E-12
MOBAOO + NO -> HNO3	3.42E-13
MAN2 + NO -> 1.5NO2 + 0.5CH2O + 0.5MGLY + 0.5PROPNN + 0.5CO + 0.5OH	2.70E-12*exp(350/T)
MCO3 + NO -> MO2 + NO2 + (CO2)	8.10E-12*exp(270/T)
RCO3 + NO -> NO2 + ETO2	6.70E-12*exp(340/T)
MAO3 + NO -> NO2 + 0.5CH2O + 0.5CO + (CO2) + 0.5MO2 + 0.5MCO3	6.70E-12*exp(340/T)
ATO2 + NO -> 0.96NO2 + 0.960CH2O + 0.960MCO3 + 0.04R4N2	2.80E-12*exp(300/T)
RIO2 + HO2 -> 0.88RIP + 0.12OH + 0.047MACR + 0.073MVK + 0.12HO2 + 0.12CH2O	2.06E-13*exp(1300/T)
VRO2 + HO2 -> 0.1VRP + 0.68OH + 0.578GLYC + 0.578MCO3 + 0.187MEK + 0.102HO2 + 0.102CH2O + 0.102MGLY + 0.033RCHO	1.82E-13*exp(1300/T)
MRO2 + HO2 -> 1.0MRP	1.82E-13*exp(1300/T)
MAN2 + HO2 -> 0.075PROPNN + 0.075CO + 0.075HO2 + 0.075MGLY + 0.075CH2O + 0.075NO2 + 0.15OH + 0.85ISNP	1.82E-13*exp(1300/T)
IEPOXOO + HO2 -> 0.725HAC + 0.275GLYC + 0.275GLYX + 0.275MGLY + 1.125OH + 0.825HO2 + (0.200CO2) + 0.375CH2O + 0.074HCOOH + 0.251CO	2.06E-13*exp(1300/T)
DIBOO + HO2 -> 0.15HO2 + 0.15OH + 0.078GLYC + 0.078MGLY + 0.072HAC + 0.072GLYX + 0.85R4P	2.06E-13*exp(1300/T)
MAOPO2 + HO2 -> 1.0HAC + (1.0CO2) + 2.0OH	1.82E-13*exp(1300/T)

HC5OO + HO2 -> 0.1IAP + 0.9OH + 0.9MGLY + 0.9GLYC + 0.9HO2	2.06E-13*exp(1300/T)
ISOPNDO2 + HO2 -> 0.5OH + 0.035MACRN + 0.155HCOOH + 0.22HAC + 0.065ETHLN + 0.325CH2O + 0.170NO2 + 0.075GLYC + 0.155NO3 + 0.075PROPN + 0.170MEK + 0.175HO2 + 0.5ISNP	2.06E-13*exp(1300/T)
ISOPNBO2 + HO2 -> 0.3GLYC + 0.3HAC + 0.2CH2O + 0.13MACRN + 0.07MVKN + 0.3NO2 + 0.2HO2 + 0.5OH + 0.5ISNP	2.06E-13*exp(1300/T)
MACRNO2 + HO2 -> 0.08ACTA + 0.08CH2O + 0.15NO3 + 0.07HCOOH + 0.07MGLY + 0.85HAC + 0.85NO2 + (0.93CO2) + 1.0OH	1.82E-13*exp(1300/T)
MOBAOO + HO2 -> 0.15OH + 0.15HO2 + 0.15RCHO + (0.15CO2) + 0.85R4P	2.06E-13*exp(1300/T)
MCO3 + HO2 -> 0.15ACTA + 0.15O3 + 0.44OH + 0.44MO2 + 0.41MAP	5.20E-13*exp(980/T)
RCO3 + HO2 -> 0.410RP + 0.150RCOOH + 0.150O3 + 0.440OH + 0.440ETO2	4.30E-13*exp(1040/T)
ATO2 + HO2 -> 0.15MCO3 + 0.15OH + 0.15CH2O + 0.85ATO OH	8.60E-13*exp(700/T)
KO2 + HO2 -> 0.15OH + 0.15ALD2 + 0.15MCO3 + 0.85ATO OH	1.82E-13*exp(1300/T)
MAO3 + HO2 -> 0.44OH + 0.15O3 + 0.59CH2O + 0.39MO2 + 0.41MAOP + 0.39CO	4.30E-13*exp(1040/T)
RIO2 + MO2 -> 1.1HO2 + 1.22CH2O + 0.280MVK + 0.180MACR + 0.3HC5 + 0.24MOH + 0.24ROH	8.37E-14
HC5OO + MO2 -> 0.50HO2 + 0.33CO + 0.09H2 + 0.18HAC + 0.13GLYC + 0.29MGLY + 0.25MEK + 0.95CH2O + 0.25MOH + 0.25ROH + 0.5HO2	8.37E-14
MRO2 + MO2 -> 0.595HAC + 0.255MGLY + 0.595CO + 1.255CH2O + 1.7HO2 + 0.150ROH	8.37E-14
VRO2 + MO2 -> 0.14HO2 + 0.14CH2O + 0.36MCO3 + 0.36GLYC + 0.14MGLY + 0.25MEK + 0.75CH2O + 0.25MOH + 0.25ROH + 0.5HO2	8.37E-14
MAN2 + MO2 -> 0.375PROPN + 0.375CO + 0.375HO2 + 0.375MGLY + 0.375CH2O + 0.375NO2 + 0.250CH2O + 0.250R4N2	8.37E-14
MAOPO2 + MO2 -> 0.7HAC + (0.7CO2) + 0.7OH + 1.0CH2O + 0.7HO2 + 0.3ROH	8.37E-14
2RIO2 -> 1.28HO2 + 0.92CH2O + 0.56MVK + 0.36MACR + 0.48ROH + 0.5HC5	1.54E-13
2MAOPO2 -> 2.0HAC + (2.0CO2) + 2.0OH	8.37E-14
MCO3 + MO2 -> CH2O + MO2 + HO2	1.80E-12*exp(500/T)
MCO3 + MO2 -> ACTA + CH2O	2.00E-13*exp(500/T)
RCO3 + MO2 -> CH2O + HO2 + ETO2	1.68E-12*exp(500/T)
RCO3 + MO2 -> RCOOH + CH2O	1.87E-13*exp(500/T)
MAO3 + MO2 -> CH2O + HO2 + CH2O + MCO3	1.68E-12*exp(500/T)
MAO3 + MO2 -> RCOOH + CH2O	1.87E-13*exp(500/T)
MAOPO2 + MCO3 -> 1.0HAC + (2.0CO2) + OH + MO2	1.68E-12*exp(500/T)
MAOPO2 + MCO3 -> 1.0ACTA + 1.0MEK	1.87E-13*exp(500/T)
R4O2 + MCO3 -> MO2 + 0.32ACET + 0.19MEK + 0.27HO2 + 0.32ALD2 + 0.13RCHO + 0.05A3O2 + 0.18B3O2 + 0.32ETO2	1.68E-12*exp(500/T)
R4O2 + MCO3 -> 1.0ACTA + 1.0MEK	1.87E-13*exp(500/T)
ATO2 + MCO3 -> MCO3 + CH2O + MO2	1.68E-12*exp(500/T)
ATO2 + MCO3 -> MGLY + ACTA	1.87E-13*exp(500/T)
HC5OO + MCO3 -> 0.216GLYX + 0.234MGLY + 0.234GLYC + 0.216HAC + 0.29DHMOB + 0.17MOBA + 0.09RCHO + HO2 +	1.68E-12*exp(500/T)

0.09CO + MO2	
HC5OO + MCO3 -> MEK + ACTA	1.87E-13*exp(500/T)
VRO2 + MCO3 -> 0.4HO2 + 0.4CH2O + 0.6MCO3 + 0.6GLYC + 0.4MGLY + 1.0MO2	1.68E-12*exp(500/T)
VRO2 + MCO3 -> MEK + ACTA	1.87E-13*exp(500/T)
MRO2 + MCO3 -> 0.850HO2 + 0.143MGLY + 0.857HAC + 0.857CO + 0.143CH2O + 1.0MO2	1.68E-12*exp(500/T)
MRO2 + MCO3 -> MEK + ACTA	1.87E-13*exp(500/T)
MAN2 + MCO3 -> 0.5PROPN + 0.5CO + 0.5HO2 + 0.5MGLY + 0.5CH2O + 0.5NO2 + (CO2) + MO2	1.68E-12*exp(500/T)
MAN2 + MCO3 -> RCHO + ACTA + NO2	1.87E-13*exp(500/T)
RIO2 + MCO3 -> 0.887HO2 + 0.747CH2O + 0.453MVK + 0.294MACR + 0.140HC5 + 0.113DIBOO + (CO2) + MO2	1.68E-12*exp(500/T)
RIO2 + MCO3 -> MEK + ACTA	1.87E-13*exp(500/T)
MCO3 + MCO3 -> 2MO2	2.50E-12*exp(500/T)
RCO3 + MCO3 -> MO2 + ETO2	2.50E-12*exp(500/T)
MAO3 + MCO3 -> MO2 + MCO3 + CH2O	2.50E-12*exp(500/T)
MCO3 + NO2 -> PAN	8.59E-12
PAN -> MCO3 + NO2	5.73E-04
RCO3 + NO2 -> PPN	7.40E-12
PPN -> RCO3 + NO2	4.44E-04
MAO3 + NO2 -> PMN	7.40E-12
PMN -> MAO3 + NO2	4.44E-04
MACRNO2 + NO2 -> PMNN	7.40E-12
PMNN -> MACRNO2 + NO2	4.44E-04
ISOP + O3 -> 0.244MVK + 0.325MACR + 0.845CH2O + 0.110H2O2 + 0.522CO + 0.204HCOOH + 0.199MCO3 + 0.026HO2 + 0.270OH + 0.128PRPE + 0.051MO2	1.00E-14*exp(-1970/T)
MVK + O3 -> 0.202OH + 0.202HO2 + 0.352HCOOH + 0.535CO + 0.050ALD2 + 0.950MGLY + 0.050CH2O	8.50E-16*exp(-1520/T)
MACR + O3 -> 0.261OH + 0.202HO2 + 0.326HCOOH + 0.569CO + 0.880MGLY + 0.120CH2O	1.40E-15*exp(-2100/T)
HC5 + O3 -> 0.6MGLY + 0.1OH + 0.12CH2O + 0.28GLYC + 0.3O3 + 0.4CO + 0.2H2 + 0.2HAC + 0.2HCOOH	6.16E-15*exp(-1814/T)
ISOPNB + O3 -> 0.610MVKN + 0.390MACRN + 0.27OH + CH2O	1.06E-16
ISOPND + O3 -> 0.5PROPN + 0.5ETHLN + 0.27OH + 0.5GLYC + 0.5HAC	5.30E-17
MOBA + O3 -> OH + HO2 + (CO2) + MEK	2.00E-17
PMN + O3 -> NO2 + 0.6CH2O + HO2	8.20E-18
RIO2 -> 2.0HO2 + 1.0CH2O + 0.5MGLY + 0.5GLYC + 0.5GLYX + 0.500HAC + 1.0OH	4.07E+08*exp(-7694/T)
MRO2 -> 1.0CO + 1.0HAC + 1.0OH	2.90E+07*exp(-5297/T)
C10H16 + OH -> TERPO2	1.20E-11*exp(444/T)
C10H16 + O3 -> .7OH + MVK + MACR + HO2	1.00E-15*exp(-732/T)
C10H16 + NO3 -> TERPO2 + NO2	1.20E-12*exp(490/T)
TERPO2 + NO -> .1ACET + MVK + MACR + NO2	4.20E-12*exp(180/T)
TERPO2 + HO2 -> TERPOOH	7.50E-13*exp(700/T)
TERPOOH + OH -> TERPO2	3.80E-12*exp(200/T)
O3 -> O2 + O1D	2.80E-04*exp(-1.75/cos(x))
O3 -> O2 + O	6.76E-04*exp(-0.30/cos(x))

H2O2 -> 2OH	2.10E-05*exp(-0.86/cos(x))
NO2 -> NO + O	1.71E-02*exp(-0.55/cos(x))
NO3 -> NO + O2	3.03E-02*exp(-0.22/cos(x))
NO3 -> NO2 + O	2.49E-01*exp(-0.25/cos(x))
N2O5 -> NO3 + NO + O	1.02E-06*exp(-5.50/cos(x))
N2O5 -> NO3 + NO2	1.25E-04*exp(-0.85/cos(x))
HNO3 -> OH + NO2	2.8E-06*exp(-1.20/cos(x))
HO2NO2 -> HO2 + NO2	2.05E-05*exp(-1.20/cos(x))
CH2O -> 2HO2 + CO	1.05E-04*exp(-1.00/cos(x))
CH2O -> H2 + CO	1.25E-04*exp(-0.80/cos(x))
MP -> CH2O + HO2 + OH	1.65E-05*exp(-0.88/cos(x))
ALD2 -> MO2 + HO2 + CO	2.95E-05*exp(-1.40/cos(x))
ALD2 -> CH4 + CO	1.25E-08*exp(-6.00/cos(x))
PAN -> 0.6MCO3 + 0.6NO2 + 0.4MO2 + 0.4NO3	2.5E-06*exp(-1.00/cos(x))
RCHO -> ETO2 + HO2 + CO	7.35E-05*exp(-1.10/cos(x))
ACET -> MCO3 + MO2	3.76E-06*exp(-1.80/cos(x))
MEK -> 0.85MCO3 + 0.85ETO2 + 0.15MO2 + 0.15RCO3	3.05E-05*exp(-1.20/cos(x))
GLYC -> CH2O + 2.0HO2 + CO	2.20E-05*exp(-1.20/cos(x))
GLYX -> 2.0CO + 2.0HO2	1.16E-04*exp(-0.45/cos(x))
MGLY -> MCO3 + CO + HO2	2.60E-04*exp(-0.60/cos(x))
MVK -> PRPE + CO	6.72E-06*exp(-0.77/cos(x))
MVK -> MCO3 + CH2O + CO + HO2	2.24E-06*exp(-0.77/cos(x))
MVK -> MO2 + MAO3	2.24E-06*exp(-0.77/cos(x))
MACR -> MAO3 + HO2	5.85E-06*exp(-0.60/cos(x))
MACR -> CO + HO2 + CH2O + MCO3	5.85E-06*exp(-0.60/cos(x))
HAC -> MCO3 + CH2O + HO2	4.05E-06*exp(-1.10/cos(x))
INPN -> OH + HO2 + RCHO + NO2	1.65E-05*exp(-0.88/cos(x))
PRPN -> OH + HO2 + RCHO + NO2	1.65E-05*exp(-0.88/cos(x))
ETP -> OH + HO2 + ALD2	1.65E-05*exp(-0.88/cos(x))
RA3P -> OH + HO2 + RCHO	1.65E-05*exp(-0.88/cos(x))
RB3P -> OH + HO2 + ACET	1.65E-05*exp(-0.88/cos(x))
R4P -> OH + HO2 + RCHO	1.65E-05*exp(-0.88/cos(x))
PP -> OH + HO2 + ALD2 + CH2O	1.65E-05*exp(-0.88/cos(x))
RP -> OH + HO2 + ALD2	1.65E-05*exp(-0.88/cos(x))
RIP -> OH + HO2 + 0.710CH2O + 0.425MVK + 0.285MACR + 0.29HC5	1.65E-05*exp(-0.88/cos(x))
IAP -> OH + HO2 + 0.67CO + 0.190H2 + 0.36HAC + 0.26GLYC + 0.580MGLY	1.65E-05*exp(-0.88/cos(x))
ISNP -> OH + HO2 + RCHO + NO2	1.65E-05*exp(-0.88/cos(x))
VRP -> OH + 0.3HO2 + 0.3CH2O + 0.7MCO3 + 0.7GLYC + 0.3MGLY	1.65E-05*exp(-0.88/cos(x))
MRP -> OH + HO2 + HAC + CO + CH2O	1.65E-05*exp(-0.88/cos(x))
MAOP -> OH + CH2O + MCO3	1.65E-05*exp(-0.88/cos(x))
R4N2 -> NO2 + 0.320ACET + 0.190MEK + 0.180MO2 + 0.270HO2 + 0.320ALD2 + 0.130RCHO + 0.050A3O2 + 0.180B3O2 + 0.320ETO2	3.94E-06*exp(-1.20/cos(x))
MAP -> OH + MO2	1.65E-05*exp(-0.88/cos(x))
MACRN -> NO2 + HAC + MGLY + 0.5CH2O + HO2 + 0.5CO	2.50E-06*exp(-1.00/cos(x))
MVKN -> GLYC + NO2 + MCO3	2.50E-06*exp(-1.00/cos(x))
ISOPNB -> HC5 + NO2 + HO2	2.50E-06*exp(-1.00/cos(x))

ISOPND -> HC5 + NO2 + HO2	2.50E-06*exp(-1.00/cos(x))
PROPNN -> CH2O + NO2 + CO + MO2	2.50E-06*exp(-1.00/cos(x))
ATOOH -> OH + CH2O + MCO3	1.65E-05*exp(-0.88/cos(x))

1

2 ^aThe stoichiometric coefficient is in accordance with 12% ISOPN yield.

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24


1
 2
 3 Table S4. The relationship between solar zenith angle (sza) and photolysis rates ($j =$
 4 $a * \exp(b / \cos(sza))$), which are obtained from the TUV radiation model. Adjusted R^2 indicates
 5 the adjusted correlation coefficient for each curve fitting result. The last two columns show
 6 the photolysis rates obtained from TUV simulations and NCAR C-130 observations at 10:37
 7 CST (which corresponds to sza of 0.32), respectively. J-value observations onboard NCAR C-
 8 130 over the AABC and SEARCH sites are available on 06/14/2013. The j-values collected
 9 below pressure altitude of 1000 m while over the two ground sites are averaged and used here
 10 (sampling time ranges between 10:02 CST and 11:11 CST with middle sampling time at
 11 10:37 CST). NA indicates no observations were available for the specific reaction.

Number	Reaction	a	b	Adjusted R^2	TUV (s-1)	C-130 (s-1)
R_01	O3 -> O2 + O(1D)	2.80E-4	-1.75	0.996	4.47E-05	3.74E-05
R_02	O3 -> O2 + O(3P)	6.76E-4	-0.30	0.996	4.92E-04	NA
R_03	H2O2 -> 2 OH	2.10E-5	-0.86	0.993	8.42E-06	7.96E-06
R_04	NO2 -> NO + O(3P)	1.88E-2	-0.50	0.999	1.11E-02	9.48E-03
R_05	NO3 -> NO + O2	3.03E-2	-0.22	0.998	2.40E-02	NA
R_06	NO3 -> NO2 + O(3P)	2.49E-1	-0.25	0.999	1.91E-01	NA
R_07	N2O5 -> NO3 + NO + O(3P)	1.02E-6	-5.50	0.998	3.03E-09	NA
R_08	N2O5 -> NO3 + NO2	1.25E-4	-0.85	0.994	5.09E-05	4.73E-05
R_09	HNO2 -> OH + NO	3.85E-3	-0.50	0.995	2.26E-03	1.54E-03
R_10	HNO3 -> OH + NO2	2.80E-6	-1.20	0.993	7.93E-07	6.82E-07
R_11	HNO4 -> HO2 + NO2	2.05E-5	-1.20	0.995	5.80E-06	7.22E-06
R_12	CH2O -> H + HCO	1.05E-4	-1.00	0.996	3.68E-05	3.53E-05
R_13	CH2O -> H2 + CO	1.25E-4	-0.80	0.990	5.37E-05	5.38E-05
R_14	CH3CHO -> CH3 + HCO	2.95E-5	-1.40	0.996	6.74E-06	5.79E-06
R_15	CH3CHO -> CH4 + CO	1.25E-8	-6.00	0.992	2.16E-11	NA
R_16	C2H5CHO -> C2H5 + HCO	7.35E-5	-1.10	0.988	2.28E-05	1.94E-05
R_17	CH2(OH)CHO -> Products	2.20E-5	-1.20	0.995	6.23E-06	NA
R_18	CH2=C(CH3)CHO -> Products	1.17E-5	-0.60	0.996	6.21E-06	NA
R_19	CH3COCH3 -> CH3CO + CH3	3.76E-6	-1.80	0.992	5.61E-07	1.20E-06
R_20	CH3COCHCH2 -> Products	1.12E-5	-0.77	0.991	4.92E-06	NA
R_21	CH3COCH2CH3 -> CH3CO + CH2CH3	3.05E-5	-1.20	0.994	8.52E-06	7.26E-06
R_22	CH2(OH)COCH3 -> CH3CO + CH2(OH)	4.05E-6	-1.10	0.990	1.26E-06	NA
R_23	CH3OOH -> CH3O + OH	1.65E-5	-0.88	0.991	6.44E-06	NA
R_24	CH3CO(OOH) -> Products	2.74E-6	-1.00	0.994	9.68E-07	NA
R_25	CH3ONO2 -> CH3O + NO2	3.94E-6	-1.20	0.994	1.12E-06	NA
R_26	CH3CO(OONO2) -> CH3CO(OO) + NO2	1.51E-6	-1.00	0.992	5.26E-07	5.54E-07
R_27	CH3CO(OONO2) -> CH3CO(O) +	1.00E-6	-1.00	0.993	3.51E-07	2.37E-07

NO3						
R_28	CH3CH2CO(OONO2) -> CH3CH2CO(OO) + NO2	2.50E-6	-1.00	0.992	8.68E-07	NA
R_29	CHOCHO -> HCO + HCO	1.16E-4	-0.45	0.998	7.22E-05	7.59E-05
R_30	CH3COCHO -> CH3CO + HCO	2.60E-4	-0.60	0.996	1.38E-04	1.45E-04


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

1
2
3
4




5
6 Figure S1. Vertical profiles of potential temperature and specific humidity collected from the
7 sounding system at the SEARCH site on (a) 2013-6-4, (b) 2013-6-11, (c) 2013-6-12, and (d)
8 2013-6-13. Time series shown in the legend are in CST. Data shown here are representative
9 of the other sampling days in this study.
10
11
12
13
14

1
2
3
4
5
6




7
8 Figure S2. Top panel, photosynthetically active radiation (PAR) measured at 32 m on the
9 AABC flux tower. Bottom panel, ozone mixing ratios measured at the SEARCH site. x-axis
10 show sampling time in month-day. Thin blue line indicates daily measurements. Bold red line
11 indicates a one-day diurnal profile obtained by averaging the data between 2013-6-1 and
12 2013-6-14, which is then duplicated for each day to show the deviation of daily profile from
13 the averaged profile.
14
15
16
17

1
2
3
4
5
6
7
8




9
10 Figure S3. Diurnal evolution of (a) surface sensible heat flux, (b) surface latent heat flux. The
11 blue lines indicate the averaged data obtained from the AABC tower over the low cloud cover
12 days (June 10-13). The shaded areas indicate 1 standard deviation of the observations. The red
13 line indicates the fitted curve, which is prescribed to MXLCH.
14
15
16
17
18
19
20

1
2
3
4
5
6
7




8
9 Figure S4. Summary of the horizontal wind data during 06:00-16:30 CST of the selected days
10 (June 5, 6, 8, 10-13, 2013). The wind data were collected on top of the AABC flux tower at
11 44 m height above the ground level.
12
13
14
15
16

1
2
3
4
5
6
7
8




9
10 Figure S5. Diurnal evolution of (a) isoprene flux, (b) monoterpenes fluxes. The circles
11 indicate averaged flux during within the time bin (30 min). Error bars indicate 1 standard
12 deviation. The red line indicates the fitted curve, which is prescribed to MXLCH. The
13 isoprene peak flux value prescribed in MXLCH is on the lower bound of the observations.
14
15
16
17
18
19

1
2
3
4
5
6
7
8



9
10 Figure S6. Comparison of TUV outputs with observation for photolysis reaction $O_3 \rightarrow O_2 +$
11 $O(1D)$. Error bars indicate 1 standard deviation. See Table S4 for curve fitting parameters and
12 methods of processing the observational data.
13
14
15

1
2
3
4
5



6
7
8
9
10
11
12
13

Figure S7. Summary of the sensitivities for (a) isoprene, (b) MVK+MACR, and (c) monoterpenes obtained from standard gas calibrations during the whole campaign period.

1

2 **References**

- 3 Cappellin, L., Karl, T., Probst, M., Ismailova, O., Winkler, P. M., Soukoulis, C., Aprea, E.,
4 Mark, T. D., Gasperi, F., and Biasioli, F.: On Quantitative Determination of Volatile Organic
5 Compound Concentrations Using Proton Transfer Reaction Time-of-Flight Mass
6 Spectrometry, Environ Sci Technol, 46, 2283-2290, 2012.
- 7 Li, J. W.: Development and evaluation of flexible empirical peak functions for processing
8 chromatographic peaks, Anal Chem, 69, 4452-4462, Doi 10.1021/Ac970481d, 1997.
- 9 Nguyen, T. B., Crounse, J. D., Schwantes, R. H., Teng, A. P., Bates, K. H., Zhang, X., St
10 Clair, J. M., Brune, W. H., Tyndall, G. S., Keutsch, F. N., Seinfeld, J. H., and Wennberg, P.
11 O.: Overview of the Focused Isoprene eXperiment at the California Institute of Technology
12 (FIXCIT): mechanistic chamber studies on the oxidation of biogenic compounds, Atmos
13 Chem Phys, 14, 13531-13549, 10.5194/acp-14-13531-2014, 2014.

14