



Corrigendum to
**“Influence of enhanced Asian NO_x emissions on ozone
in the upper troposphere and lower stratosphere
in chemistry–climate model simulations” published in Atmos.
Chem. Phys., 17, 1297–1311, 2017**

Chaitri Roy¹, Suvarna Fadnavis¹, Rolf Müller², D. C. Ayantika¹, Felix Ploeger², and Alexandru Rap³

¹Indian Institute of Tropical Meteorology, Pune, India

²Forschungszentrum Jülich GmbH, IEK7, Jülich, Germany

³School of Earth and Environment, University of Leeds, Leeds, UK

Correspondence to: Suvarna Fadnavis (suvarna@tropmet.res.in)

Published: 26 July 2017

We discovered an error in Fig. 7: the correct unit for the Fig. 7 colour scale should be “(K day⁻¹ × 10⁻³)” instead of “(K day⁻¹)”. The corrected figure with caption is shown below.




Figure 7. Latitude–pressure distribution of ozone heating rate changes ($\text{K day}^{-1} \times 10^{-3}$) for the (a) Ind38 (averaged over $70\text{--}100^\circ \text{E}$), (b) Ind73 (averaged $70\text{--}100^\circ \text{E}$), and (c) Chin73 (averaged over $70\text{--}100^\circ \text{E}$) perturbed simulations, compared to the CTRL simulation.