



Dust deposition and ambient PM₁₀ concentration in northwest China: spatial and temporal variability

Xiao-Xiao Zhang^{1,2}, Brenton Sharratt³, Xi Chen¹, Zi-Fa Wang², Lian-You Liu⁴, Yu-Hong Guo², Jie Li², Huan-Sheng Chen², and Wen-Yi Yang²

¹State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi, 830011, China

²State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China

³USDA-ARS, 215 Johnson Hall, Washington State University, Pullman, WA 99164, USA

⁴Key Laboratory of Environmental Change and Natural Disaster, Ministry of Education, Beijing Normal University, Beijing, 100875, China

Correspondence to: Xiao-Xiao Zhang (zhangxx@ms.xjb.ac.cn)

Received: 15 June 2016 – Published in Atmos. Chem. Phys. Discuss.: 23 June 2016

Revised: 10 January 2017 – Accepted: 15 January 2017 – Published: 3 February 2017

Abstract. Eolian dust transport and deposition are important geophysical processes which influence global biogeochemical cycles. Currently, reliable deposition data are scarce in central and east Asia. Located at the boundary of central and east Asia, Xinjiang Province of northwestern China has long played a strategic role in cultural and economic trade between Asia and Europe. In this paper, we investigated the spatial distribution and temporal variation in dust deposition and ambient PM₁₀ (particulate matter in aerodynamic diameter $\leq 10\ \mu\text{m}$) concentration from 2000 to 2013 in Xinjiang Province. This variation was assessed using environmental monitoring records from 14 stations in the province. Over the 14 years, annual average dust deposition across stations in the province ranged from 255.7 to 421.4 t km⁻². Annual dust deposition was greater in southern Xinjiang (663.6 t km⁻²) than northern (147.8 t km⁻²) and eastern Xinjiang (194.9 t km⁻²). Annual average PM₁₀ concentration across stations in the province varied from 100 to 196 $\mu\text{g m}^{-3}$ and was 70, 115 and 239 $\mu\text{g m}^{-3}$ in northern, eastern and southern Xinjiang, respectively. The highest annual dust deposition (1394.1 t km⁻²) and ambient PM₁₀ concentration (352 $\mu\text{g m}^{-3}$) were observed in Hotan, which is located in southern Xinjiang and at the southern boundary of the Taklamakan Desert. Dust deposition was more intense during the spring and summer than other seasons. PM₁₀ was the main air pollutant that significantly influenced regional

air quality. Annual average dust deposition increased logarithmically with ambient PM₁₀ concentration ($R^2 \geq 0.81$). While the annual average dust storm frequency remained unchanged from 2000 to 2013, there was a positive relationship between dust storm days and dust deposition and PM₁₀ concentration across stations. This study suggests that sand storms are a major factor affecting the temporal variability and spatial distribution of dust deposition in northwest China.

1 Introduction

Airborne dust generated by eolian activity is an environmental concern in central and east Asia (Huang et al., 2011; Chen et al., 2014). Historically, eolian activity and airborne dust influenced civilization along the ancient Silk Road which connected Asia and Europe (Zhang, 1984; Dong et al., 2012; Groll et al., 2013). Today, airborne dust is recognized as a factor affecting global radiation and warming (Stanhill, 2005; Carslaw et al., 2013; IPCC, 2013; Huang et al., 2009; Chen et al., 2013; Huang et al., 2014) and air quality in distant lands (Tsoar and Pye, 1987; Xu et al., 2007; Uno et al., 2009; Li et al., 2012). Deposition of airborne dust also plays a significant role in soil formation and biological diversity in arid and semi-arid regions (Simonson, 1995; Lin and Feng,

Table 1. Observations of dust deposition in desertification regions.

Continent	Location	Period	Dust deposition t km ⁻² yr ⁻¹	Citation
N. America	Kansas, USA	1964–1966	53.5–62.1	USDA (1968)
	New Mexico, USA	1962–1972	9.3–125.8	Gile and Grossman (1979)
	Arizona, USA	1972–1973	54	Péwé (1981)
Europe	Spain	2002–2003	17–79	Menéndez et al. (2007)
Africa	Nigeria	1976–1979	137–181	McTainsh and Walker (1982)
	Niger	1985	164–212	Drees et al. (1993)
	Libya	2000–2001	420	O'Hara et al. (2006)
Oceania	Australia	2000–2001	5–100	Cattle et al. (2009)
Asia	Israel	1968–1973	57–217	Yaalon and Ganor (1975)
	Kuwait	1982	2600	Khalaf and Al-Hashash (1983)
	Saudi Arabia	1991–1992	4704	Modaihash (1997)
	Lanzhou, China	1988–1991	108	Derbyshire et al. (1998)
	Loess Plateau, China	2003–2004	133	Liu et al. (2004)
	Urumqi, China	1981–2004	284.5	Zhang et al. (2010)
	Iran	2008–2009	72–120	Saeid et al. (2012)
	Uzbekistan	2003–2010	8365	Groll et al. (2013)

Table 2. Dust deposition and PM₁₀ concentrations at 14 stations in Xinjiang.

No.	Station	Region ¹	Latitude	Longitude	Population ² (million)	Annual dust deposition (t km ⁻²)	Annual PM ₁₀ concentration (µg m ⁻³)
1	Urumqi	NJ	43.832° N	87.616° E	2.26	229.4	141
2	Changji	NJ	44.017° N	87.308° E	0.36	295.7	76
3	Shihezi	NJ	44.306° N	86.080° E	0.62	107.7	61
4	Bole	NJ	44.900° N	82.071° E	0.27	133	48
5	Karamay	NJ	45.580° N	84.889° E	0.29	81.1	54
6	Tacheng	NJ	46.691° N	82.952° E	0.17	84.9	39
7	Yining	NJ	43.912° N	81.329° E	0.53	142.7	78
8	Kuytun	NJ	44.426° N	84.903° E	0.30	108.1	66
9	Hami	EJ	42.818° N	93.514° E	0.48	209.8	84
10	Turpan	EJ	42.957° N	89.179° E	0.28	180.1	145
11	Korla	SJ	41.727° N	86.174° E	0.57	231.8	131
12	Hotan	SJ	37.113° N	79.922° E	0.33	1394.1	352
13	Kashgar	SJ	39.471° N	75.989° E	0.57	516.9	236
14	Aksu	SJ	41.170° N	80.230° E	0.51	511.5	238

¹ Xinjiang Province was classified into three regions: northern Xinjiang (NJ), eastern Xinjiang (EJ) and southern Xinjiang (SJ).

² Population in 2013 as reported by the Xinjiang Statistical Bureau.

2015; Varga et al., 2016). An understanding of atmospheric dust sources, emissions and deposition is therefore essential to improve our knowledge of dust impact on regional air quality.

Dust in the atmosphere and its subsequent deposition are vital indicators of eolian activity and environmental quality. Deposition has been measured directly at only a few sites; therefore, reliable dust deposition data are lacking around the world (Pye, 1987; Mahowald et al., 1999, 2009; Prospero, 1999; Zhang et al., 2010; Huneus et al., 2011; Shao et al., 2011). Annual dust deposition ranges from 10 to 200 t km⁻²

on continents and 1–2 orders of magnitude lower over oceans (Pye, 1987; Duce et al., 1991; Ginoux et al., 2001, 2012). It is estimated that the annual average dust deposition rate in desert areas ranged between 14 and 2100 t km⁻² (Zhang et al., 1997). Observations of dust deposition have been made over deserts with an enhanced awareness of its significance. Table 1 lists observations of dust deposition in desert regions and other regions of the world prone to eolian activity. According to these observations (Table 1), dust deposition is high in Asia with an annual deposition of 8365 t km⁻² in the Aral Sea Basin (Wake and Mayewski, 1994; Groll et




Figure 1. Location of Xinjiang Province in China (gray area outlined on inset map). Dust deposition and concentrations were measured at stations signified by small triangles. Land use types are identified across the province according to Wang et al. (2005).

al., 2013). Recent investigations suggest that the intensity of dust deposition is closely related to weather. For example, dust deposition during extreme high winds can be 10 to 25 times higher than the annual average (Liu et al., 2004; Zhang et al., 2010; Goudie, 2014). The observations on dust deposition are ordinarily scattered and discontinuous. The limited observational data restricted our understanding of dust fluxes between the atmosphere and land surface; thus numerical simulations are needed to evaluate dust fluxes and the rate of global dust deposition. Ginoux et al. (2001) simulated dust deposition at 16 sites around the world and predicted the annual global dust deposition was approximately 1842 megatons. Shao et al. (2011) estimated that over 2000 megatons of dust is emitted from the Earth's surface into the atmosphere annually. Zheng et al. (2016) estimated that annual average global dust deposition was approximately 1161 megatons. However, uncertainties remain in estimating the dust deposition budget of the Earth system because of the lack of observational data and inaccuracies of parameters in numerical simulations (Ginoux et al., 2001, 2012; Huneus et al., 2011; Shao et al., 2011; J. Zhang et al., 2014). Observation of worldwide dust deposition is urgently needed to assess biogeochemical cycle of dust on Earth.

Located in east Asia and at the boundary of central Asia, Xinjiang Province of northwestern China has long played a strategic role in cultural and economic trade between Asia and Europe. Xinjiang Province experiences severe sand and

dust storms and is highly susceptible to desertification (Chen, 2010). Xinjiang Province is one of two major source regions of atmospheric dust in China, the other region being western Inner Mongolia (Xuan, 1999; Xuan et al., 2000). Long-range transport of dust from the region strongly affects air quality in east Asia (Derbyshire et al., 1998; Uno et al., 2009). In fact, dust from the region can be transported across the Pacific Ocean and thus impact air quality in North America (Husar et al., 2001; Osada et al., 2014). Indeed, particulate matter associated with dust transport can severely deteriorate air quality (Sharratt and Lauer, 2006; Shoemaker et al., 2013). Over the past decades, many observations have been made of processes that govern dust emissions, transport and deposition in Asia (Shao et al., 2011; Groll et al., 2013). Little is known, however, concerning dust deposition and concentrations in Xinjiang Province. In fact, temporal and spatial variations in dust deposition and concentration have not been characterized despite the importance of dust transport from the region. To improve our understanding of the fate and transport of airborne dust in central and east Asia, there is a need for continuous and long-term records of dust deposition and concentration. The purpose of this study is to characterize the spatiotemporal distribution of dust deposition and particulate matter concentration in Xinjiang Province. This characterization will strengthen our comprehension of aerosol transport in east Asia and provide aerosol data for modeling dust transport in global desertification regions.




Figure 2. Annual average dust deposition reported at 14 stations in Xinjiang Province from 2000 to 2013. Land use types across the province are identified according to Wang et al. (2005).

2 Methods

2.1 Study area

Xinjiang Province is located in northwest China and is the largest inland province which covers an area of more than 1.6 million km² (Fig. 1). The Taklamakan and Gurbantunggut deserts are located in the province. The Taklamakan Desert, located in the southern region, is the world's largest shifting-sand-dune desert. The Gurbantunggut Desert, located in the northern region, is the largest fixed-dune desert in China. The province is in part characterized by extreme aridity and eolian desertification. The average annual precipitation varies from more than 700 mm in high-altitude forests and mountains to less than 50 mm in the deserts. Annual potential evaporation can exceed 2000 mm in desert regions (Chen, 2010). Sand and dust storms occur throughout the year but are most common in spring. In this study, the province was divided by latitude and longitude into three regions, those being northern Xinjiang, eastern Xinjiang and southern Xinjiang (Table 2).

2.2 Experimental data

Dust deposition and PM₁₀ concentration were measured at environmental monitoring stations maintained by the Xinjiang Environmental Protection Administration, a division of the Ministry of Environmental Protection (MEP) in China. Data collected at 14 stations (Fig. 1 and Table 2) were used

in this study and represent a spatial distribution within this region.

Dust deposition was determined by the gravimetric method and documented at monthly intervals. Glass cylinders were used to monitor dust deposition. Three cylinders (replicates) were installed to monitor dust deposition at each station. The cylinders (0.15 m in diameter, 0.3 m tall and open at the top) were partly filled with an ethylene glycol (C₂H₆O₂)–water solution prior to deployment. The solution enabled trapping of dust in a liquid medium at temperatures below 0 °C and also minimized evaporation from the cylinder. Cylinders were mounted vertically on a tower at approximate 10 m above ground. The mass of dust collected by the cylinders was determined after washing the contents out of the cylinders and oven-drying the contents at 105 °C. Dust deposition rate was calculated as the mass of dust per unit area per unit time and expressed in units of t km⁻² month⁻¹ (MEP, 1994). Monthly and yearly dust deposition data were available through the MEP for the 14 stations from 2000 to 2013.

Ambient PM₁₀ concentration was measured with high-volume samplers designed to collect particulate matter by filtration. The samplers were installed at 1.5 m above the ground and equipped with fiberglass filters for trapping PM₁₀. PM₁₀ concentration was determined based upon gravimetric filter analysis and flow rate of each sampler. Daily PM₁₀ concentration data were obtained by the arithmetic mean of four samplers, with the sampling time being > 18 h for each sampler. PM₁₀ was expressed in units




Figure 3. Annual average dust deposition in Xinjiang Province from 2000 to 2013. Dust deposition in northern, eastern and southern Xinjiang is the average deposition at 8, 2 and 4 stations, respectively.

of $\mu\text{g m}^{-3}$ (MEP, 2011). Annual PM₁₀ data were available through the MEP for the 14 stations (Xinjiang Statistical Bureau, 2014).

Daily meteorological data – including dust days, surface wind speed and precipitation – were collected from the China Meteorological Administration. A dust day was defined by visibility according to World Meteorological Organization (WMO) protocol; days on which visibility was < 10 km at any observation time throughout the day constituted a dust day. The WMO further classifies dust days as dust in suspension, blowing dust and dust storms (<http://www.wmo.int/pages/prog/www/WMOCodes.html>; Shao and Dong, 2006). Dust in suspension constitutes days on which dust is emitted at the station at the time of observation and visibility is < 10 km, blowing dust constitutes days on which dust or sand is emitted at the station and visibility is 1–10 km, and dust storms constitutes days on which dust or sand is emitted at the station and visibility is < 1 km. Observations of visibility and wind characteristics at each station were taken at 3 h intervals throughout the day.

Daily air pollution index (API) data were obtained from air quality monitoring statistics published by the MEP (<http://datacenter.mep.gov.cn>). These data were used to illustrate the impact of airborne dust versus other air pollutants on air quality. The API is calculated according to the daily concentration of three main air pollutants (USEPA, 2006; Wang et al., 2013), namely PM₁₀, SO₂ and NO₂. The API is calculated as

$$\text{API} = \max(\text{API}_i), \quad (1)$$

$$\text{API}_i = \frac{\text{API}_u - \text{API}_L}{C_u - C_L} \times (C_i - C_L) + \text{API}_L, \quad (2)$$




Figure 4. Monthly average dust deposition in Xinjiang Province from 2000 to 2013. Dust deposition in northern, eastern and southern Xinjiang is the average deposition at 8, 2 and 4 stations, respectively.

where API_i is the index for pollutant i (i.e., PM₁₀, SO₂ and NO₂), API_u and API_L are the upper and lower limits of the index for a specific category of air quality (i.e., excellent, slight, moderate, moderately severe and severe), C_i is the observed concentration of pollutant, and C_u and C_L are the upper and lower limits of the pollutant for a specific category of air quality.

Information regarding the determination of the API index can be obtained from the MEP (MEP, 2012a, b). Based on the API, air quality was classified as excellent with an API of 0 to 50, slight pollution with an API of 50 to 100, moderate pollution with an API of 100 to 200, moderately severe pollution with an API of 200 to 300 and severe pollution with an API of 300 to 500. For the purpose of this study, we used only API data collected in 2010 since annual deposition and PM₁₀ concentration appeared to typify that which occurred between 2000 and 2013 in eastern, northern and southern Xinjiang Province.

Temporal trends in dust deposition, PM₁₀ concentration and dust days were evaluated by testing the significance of the slope estimate using a t test at a probability level (P value) of 0.05.

3 Results and discussion

3.1 Dust deposition

Detailed information on dust deposition during 2000–2013 was obtained from 14 environmental monitoring stations (Table 2). Annual average dust deposition across all stations in Xinjiang Province was 301.9 t km^{-2} . The highest annual deposition occurred in Hotan and Kashgar in southern Xinjiang, while the lowest deposition occurred in Kara-




Figure 5. Annual average PM₁₀ concentration reported for 14 stations in Xinjiang Province from 2000 to 2013. Land use types are identified across the province according to Wang et al. (2005).

may in northern Xinjiang. Based upon spatial characteristics in annual dust deposition, deposition increased from north to south across the province (Fig. 2). The annual average dust deposition was 147.8, 194.9 and 663.6 t km⁻² in northern, eastern and southern Xinjiang, respectively. Generally, the origin of mineral dust could be attributed to both natural and anthropogenic sources (Miller-Schulze et al., 2015). Although dust deposition was relatively low (< 150 t km⁻²) for the majority of stations in northern Xinjiang Province, dust deposition was at least 50 % higher for stations within the industrial belt on the northern slope of the Tianshan Mountains. This industrial belt includes Changji and Urumqi. High dust deposition in the industrial belt was due to local industry, coal burning and vehicle exhaust (Matinmin and Meixner, 2011; X. X. Zhang et al., 2014). Therefore, the mixing of the anthropogenic aerosol with transported desert dust contributed to deposition in Changji and Urumqi (Li, et al., 2008).

Figures 3 and 4 depict the temporal variation in dust deposition from 2000 to 2013. The highest annual deposition occurred in 2012 in southern Xinjiang, 2002 and 2012 in eastern Xinjiang, and 2001 in northern Xinjiang. Over the 14-year period, dust deposition varied with time across Xinjiang Province. The slope estimate of the relation between average dust deposition and year (-6.4 ± 0.1 t km⁻² yr⁻¹) was significant at $P = 0.05$. This trend was most apparent in northern Xinjiang (slope estimate was -5.6 ± 0.1 t km⁻² yr⁻¹) and least apparent in southern Xinjiang (slope estimate was -1.9 ± 0.1 t km⁻² yr⁻¹). High dust deposition occurred in spring in eastern and northern Xinjiang and in spring

and summer in southern Xinjiang (Fig. 4). Dust deposition peaked in April in eastern and northern Xinjiang and in May in southern Xinjiang. This corresponds to the onset of wind erosion caused by intensifying zonal flow and rising air temperatures before the arrival of the summer monsoon (Song et al., 2016). The maximum monthly average dust deposition was 97.5 t km⁻² in southern Xinjiang, which was 6.9 and 8 times more than the deposition in northern and eastern Xinjiang, respectively. These results suggest that dust deposition in south Xinjiang is of similar magnitude to deposition that occurs in the Middle East and Sahel regions (Khalaf and Al-Hashash, 1983; McTainsh and Walker, 1982; O'Hara et al., 2006).

3.2 PM₁₀ concentration

The annual average PM₁₀ concentration in Xinjiang was 125 µg m⁻³ based upon data collected at 14 stations from 2000 to 2013. Ten stations (71 %) in our study had an annual average PM₁₀ concentration above the People's Republic of China Class II residential standard of 70 µg m⁻³. The highest annual average PM₁₀ concentration (352 µg m⁻³) occurred in Hotan in southern Xinjiang, while the lowest average PM₁₀ concentration (46 µg m⁻³) occurred in Tacheng in northern Xinjiang. The annual average PM₁₀ concentration appeared to increase from northern to southern regions (Fig. 5). Annual average PM₁₀ concentration in Xinjiang ranged from 100 to 196 µg m⁻³ (Fig. 6) across years. The annual average PM₁₀ concentration was 70, 115 and




Figure 6. Annual average PM₁₀ concentration in Xinjiang Province from 2000 to 2013.




Figure 7. Relationship between annual dust deposition and PM₁₀ concentration in Xinjiang Province. Each point represents data averaged across 2000 to 2013 at one station.

239 $\mu\text{g m}^{-3}$ in northern, eastern and southern Xinjiang, respectively. The high annual concentration in southern Xinjiang is of the same magnitude as found in other desertification regions of the world such as south Asia, the Middle East and the western Sahel (WHO, 2016a). These high concentrations of suspended particulates, especially finer particulate, may influence the health of sensitive populations who are susceptible to respiratory illness (Goudie, 2014).

Over the period of record (2000–2013), there was a trend of decreasing PM₁₀ concentration in Xinjiang Province. The slope estimate of the relation between annual PM₁₀ concentration and year ($-4.2 \pm 0.1 \mu\text{g m}^{-3} \text{yr}^{-1}$) was significant at $P = 0.05$. This trend was most apparent in southern Xinjiang (slope estimate was $-11.8 \pm 0.1 \mu\text{g m}^{-3} \text{yr}^{-1}$). However, PM₁₀ concentration appeared to increase with time in eastern and northern Xinjiang (slope estimates were

1.3 ± 0.1 and $3 \pm 0.1 \mu\text{g m}^{-3} \text{yr}^{-1}$, respectively). The slope estimate, however, was not statistically different from zero and indicated no apparent trend in PM₁₀ concentration with time in northern Xinjiang. A decrease in both dust deposition and PM₁₀ concentration over 2000 to 2013 suggests a positive relationship between dust deposition and PM₁₀ concentration. This relationship is supported by data in Fig. 7. Dust particles are delivered back to the surface by both dry and wet deposition (Shao, 2000). In arid and semi-arid regions of central Asia, the deposition process is mainly dominated by dry deposition because of less precipitation, which is comprised of gravitational settling, turbulent diffusion and molecular diffusion (Zhang and Shao, 2014; Xi and Sokolik, 2015). Those physical processes from the air to surface are complex and dependent on dust concentration: the higher the dust concentration, the higher the dust deposition (Slinn and Slinn, 1980; Wesely and Hicks, 2000; Petroff, et al., 2008; J. Zhang et al., 2014). Figure 7 showed that dust deposition significantly increased with high PM₁₀ concentration above 200 $\mu\text{g m}^{-3}$. A logarithmic function fit the data better than a linear function, suggesting that changes in atmospheric PM₁₀ concentration are smaller at higher rates of deposition with a correlation coefficient $R^2 \geq 0.81$. This trend could be due to deposition of larger or more massive particles under more severe dust or sand storms. While PM₁₀ concentration may rise under more severe wind erosion events, the limited supply of PM₁₀ in sand (major soil type in the province) will likely suppress a rise in PM₁₀ concentration in the atmosphere under more severe erosion events. Nevertheless, from 2000 to 2013, the decline in both dust deposition and PM₁₀ concentration across Xinjiang could be due to less frequent or intense dust storms because dust deposition in major cities of northern China was found to be closely related to the frequency of sand and dust storms (Zhang et al., 2010).

3.3 Influence of atmospheric dust deposition on local air quality

Daily ambient air quality has been reported by the MEP since 2000. Airborne dust is one of three pollutants influencing the API; thus the relative contribution of dust to the API was of interest. Accordingly, we made a comparative analysis to identify the impact of airborne dust on air quality in Urumqi and Kuytun in northern Xinjiang, Turpan and Hami in eastern Xinjiang, and Kashgar and Hotan in southern Xinjiang (Fig. 8). In 2010, there were 178, 286, 351, 334, 363 and 360 days on which PM₁₀ was the main constituent of the API in Kuytun, Urumqi, Turpan, Hami, Kashgar and Hotan, respectively (Fig. 8). The PM₁₀ constituent accounted for 48.7, 78.4, 96.2, 91.5, 99.5 and 99.6% of the API in the respective above cities. These data suggest that particulate matter is the main air pollutant in Xinjiang. Severe PM₁₀ pollution (API > 300) occurred mainly in spring, which was closely associated with the seasonality of strong winds and dust storm activity (Li et al., 2004). Stations in southern Xinjiang (Kash-




Figure 8. Daily air pollution index for Kuytun and Urumqi in northern Xinjiang, Hami and Turpan in eastern Xinjiang, and Kashgar and Hotan in southern Xinjiang in 2010. The main air pollutant contributing to the daily API is identified for each station. “Not detected” indicates excellent air quality (API < 50).

gar and Hotan) had higher APIs caused by elevated PM₁₀ concentrations throughout the year. This can be attributed to the violent and persistent eolian activity around the Taklamakan Desert (Pi et al., 2014). Consequently, PM₁₀ is an important pollutant which dominates ambient air quality in Xinjiang.

3.4 Factors influencing dust deposition and PM₁₀ concentration

Many factors influence ambient particulate concentration and dust deposition, but weather appears to be a dominant factor in arid regions (Zhang et al., 1996, 2010). In fact, dust activity is highly correlated with variability in global climate and atmospheric circulation (Gong, et al, 2006; Mao et al.,




Figure 9. Dust day frequency in Xinjiang Province from 2000 to 2013.

2011; Shao et al., 2013). The Eurasian atmospheric circulation greatly affects weather in central and east Asia (Zhang et al., 1997; Kang et al., 2013; Xi and Sokolik, 2015). Dust activities are primarily driven by the strength of cyclones and the Siberian High affecting the study region (Park et al., 2011; Shao et al., 2013). Strong winds associated with this atmospheric circulation cause large amounts of dust to be emitted into the atmosphere. Deserts in central Asia are an important source of atmospheric mineral dust (Miller-Schulze et al., 2015). Under the strong westerly circulation, atmospheric dust can be transported a few hundred kilometers to the east and be deposited through wet scavenging and dry settling (Shao, 2000; Chen et al., 2014). Despite the Taklamakan and Gurbantunggut deserts being local sources of dust in Xinjiang Province, long-range transport of dust from the central Asian Aralkum, Karakum, Caspian and Kyzylkum deserts (Indoitu et al., 2012) could also contribute to the dust deposition and ambient PM₁₀ concentration in neighboring Xinjiang Province. Since the 1980s, the Aralkum Desert in Uzbekistan and Kazakhstan has become one of world’s youngest deserts and a potential source of salt dust in east Asia (Indoitu et al., 2012; Groll et al., 2013; Opp et al., 2016).

Climate also directly influences the atmospheric environment of arid and semi-arid areas (Wei et al., 2004; Zu et al., 2008; Huang et al., 2014). The annual average precipitation in north, east and south Xinjiang is 237, 94 and 87 mm, respectively. Dust emission was negatively correlated with precipitation (Gong et al., 2006). Therefore, the lack of precipitation contributes to dust emissions. In fact, regions with lower precipitation have higher PM₁₀ concentrations and dust deposition in Xinjiang Province. Daily average wind speed in north, east and south Xinjiang is 2.5, 2.2 and 1.8 m s⁻¹, respectively. In contrast to precipitation, regional differences in wind speed fail to account for differences in PM₁₀ concentrations and dust deposition. Dust dis-




Figure 10. Relationship between annual dust deposition and dust day frequency in Xinjiang Province. Each point represents data averaged across 2000 to 2013 at one station.




Figure 11. Relationship between PM₁₀ concentration and dust day frequency in Xinjiang Province. Each point represents data averaged across 2000 to 2013 at one station.

tribution in south Xinjiang (including the Tarim Basin and Taklamakan Desert), however, is strongly affected by wind flow patterns. Eolian transport in the Taklamakan Desert is predominantly from the northeast toward the south (Wang et al., 2014; Rittner et al., 2016). Huang et al. (2014) reported that the Taklamakan Desert is a source of fine dust particles ($\leq 3 \mu\text{m}$ in aerodynamic diameter) which significantly influences East Asia. Strong northeast winds dominate the prevailing wind regime in the eastern Taklamakan Desert; these winds influence air quality in both the eastern and the south-eastern parts of the desert. The western and northern parts of the Taklamakan Desert and Tarim Basin are highly affected by west, northwest and north winds (Sun and Liu, 2006; Zan et al., 2014; Li et al., 2015). Under prevailing winds, dust aerosols are transported from the northern to the southern Taklamakan Desert (e.g., Hotan city) and thereby cause high ambient PM₁₀ concentration and dust deposition.

Spatial differences in dust deposition and PM₁₀ concentration across Xinjiang Province may also be due in part to differences in frequency of dust days in the region. Dust storms normally occurred in all seasons in southern Xinjiang. The magnitude of wind erosion and dust day frequency in southern Xinjiang is nearly twice as large as in northern and eastern Xinjiang (Wang et al., 2006). Figure 9 displays the variation in dust day frequency in Xinjiang Province from 2000 to 2013. The data indicate that the annual average frequency of dust days fluctuated from 15 to 52 days. The frequency of dust days in the southern region ranged from 41 to 133 days, while the frequency of dust days in eastern and northern regions ranged from 2 to 45 days and from 0 to 3 days, respectively, across years. The slope estimate of the relationship between dust days and years (0.11 day yr^{-1}) indicated no apparent trend for an increase or decrease in dust days

from 2000 to 2013. Thus, despite no temporal trend in dust days, we observed a decline in dust deposition and PM₁₀ concentration across years. This decline in dust deposition or PM₁₀ concentration could be due to a decrease in frequency of severe dust days versus frequency of dust days from 2000 to 2013 in the region. We are unaware of any previous study which has examined dust day severity in Xinjiang Province; thus we used data available through the China Meteorological Administration to assess trends in dust day severity. Dust days were characterized according to dust-in-suspension, blowing dust and dust storm events. Although there was no trend in the frequency of blowing dust and dust storm events from 2000 to 2013, there was a trend for fewer dust-in-suspension events from 2000 to 2013 (Fig. S1 in the Supplement). Thus, there appeared to be a close association between frequency of dust-in-suspension events and PM₁₀ concentration and dust deposition. Nevertheless, in examining the relationship between average annual dust days and dust deposition or PM₁₀ concentration across stations, the frequency of dust days was closely related to dust deposition ($R^2 = 0.93$; Fig. 10) and ambient PM₁₀ concentration ($R^2 = 0.89$; Fig. 11). There was a significant increase in dust deposition ($7.91 \text{ t km}^{-2} \text{ day}^{-1}$) and PM₁₀ concentration ($2.06 \mu\text{g m}^{-3} \text{ day}^{-1}$) associated with an increase in dust days.

4 Conclusions

The atmospheric environment of central and east Asia is severely affected by the airborne dust; thus this study was undertaken to quantify dust deposition and ambient PM₁₀ concentration in east Asia. Data collected at 14 environmental

monitoring stations from 2000 to 2013 in Xinjiang Province, China, confirmed that annual average dust deposition ranged from 255.7 to 421.4 t km⁻². Annual average PM₁₀ concentration varied from 100 to 196 μg m⁻³. The highest dust deposition was observed in Hotan in the southern Taklamakan Desert with 1394.1 t km⁻², which is 10 times that in China's Loess Plateau (Liu et al., 2004). The highest ambient PM₁₀ concentration was also observed in Hotan with 352 μg m⁻³, which far exceeds the World Health Organization's long-term exposure standard (WHO, 2016b). These observation results provide concrete evidence on the study area as "dust region" described by Shao et al. (2011) and Ginoux et al. (2012), and they suggest that dust sources in east Asia affect regional air quality and is a potential contributor of global dust.

The spatial distribution and temporal variability in dust deposition and ambient PM₁₀ concentration showed significant variation and a trend for lower deposition and concentration with time. The interannual dynamic of dust deposition varied significantly with seasonality. Spring and summer had the highest dust deposition (1.3 times the average), followed by autumn and winter. The highest intensity of dust deposition was observed in May, followed by April, June and July.

In dust source areas such as Xinjiang, China, windblown sand and dust affect air quality, especially during the spring season. The analysis of the data indicated no trend in frequency of dust days from 2000 to 2013. A positive relationship existed, however, between dust days and dust deposition as well as airborne PM₁₀ concentration across stations. The effect of weather on dust deposition and ambient air quality cannot be expressed by a simple correlation and should not be extrapolated based on the currently limited evidence. This study provides information on the potential spatiotemporal dust deposition and ambient dust aerosol variation in east Asia. Although longer-term datasets are needed to address trends over longer time periods, this work can aid in adjusting model parameters in simulating dry dust deposition or PM₁₀ concentration in desertification regions of east Asia.

5 Data availability

The data used in this study are available upon request by contacting corresponding author (zhangxx@ms.xjb.ac.cn).

The Supplement related to this article is available online at doi:10.5194/acp-17-1699-2017-supplement.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The authors would like to thank anonymous reviewers for their useful comments that contributed to improving the manuscript. This work was supported by the National Natural Science Foundation of China (no. 41301655); the West Light Foundation of the Chinese Academy of Sciences (no. XBBS201104); and the Open Funds (no. LAPC-KF-2013-17) of the State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, China.

Edited by: J. Huang

Reviewed by: three anonymous referees

References

- Carslaw, K. S., Lee, L. A., Reddington, C. L., Pringle, K. J., Rap, A., Forster, P. M., Mann, G. W., Spracklen, D. V., Woodhouse, M. T., Regayre, L. A., and Pierce, J. R.: Large contribution of natural aerosols to uncertainty in indirect forcing, *Nature*, 503, 67–71, doi:10.1038/nature12674, 2013.
- Cattle, S. R., McTainsh, G. H., and Elias, S.: Aeolian dust deposition rates, particle-sizes and contributions to soils along a transect in semi-arid New South Wales, Australia, *Sedimentology*, 56, 765–783, doi:10.1111/j.1365-3091.2008.00996.x, 2009.
- Chen, S., Huang, J., Zhao, C., Qian, Y., Leung, L. R., and Yang, B.: Modeling the transport and radiative forcing of Taklimakan dust over the Tibetan Plateau: A case study in the summer of 2006, *J. Geophys. Res.-Atmos.*, 118, 797–812, doi:10.1002/jgrd.50122, 2013.
- Chen, S., Zhao, C., Qian, Y., Leung, L. R., Huang, J., Huang, Z., Bi, J., Zhang, W., Shi, J., Yang, L., Li, D., and Li, J.: Regional modeling of dust mass balance and radiative forcing over East Asia using WRF-Chem, *Aeolian Res.*, 15, 15–30, doi:10.1016/j.aeolia.2014.02.001, 2014.
- Chen, X.: Physical geography of arid land in China, Science Press, Beijing, 124–155, 2010.
- Derbyshire, E., Meng, X., and Kemp, R.A.: Provenance, transport and characteristics of modern aeolian dust in western Gansu Province, China, and interpretation of the quaternary loess record, *J. Arid Environ.*, 39, 497–516, doi:10.1006/jare.1997.0369, 1998.
- Dong, Z., Lv, P., Qiang, G., Xia, X., Zhao, Y., and Mu G.: Research progress in China's Lop Nur, *Earth-Sci. Rev.*, 111, 142–153, doi:10.1016/j.earscirev.2011.11.003, 2012.
- Drees, L. R., Manu, A., and Wilding, L. P.: Characteristics of aeolian dusts in Niger, West Africa, *Geoderma*, 59, 213–233, doi:10.1016/0016-7061(93)90070-2, 1993.
- Duce, R. A., Liss, P. S., Merrill, J. T., Atlas, E. L., Buat-Menard, P., Hicks, B. B., Miller, J. M., Prospero, J. M., Arimoto, R., Church, T. M., Ellis, W., Galloway, J. N., and Hansen, L.: The atmospheric input of trace species to the world ocean, *Global Biogeochem. Cy.*, 5, 193–259, doi:10.1029/91GB01778, 1991.
- Gile, L. H. and Grossman, R. B.: The desert project soil monograph: soils and landscapes of a desert region astride the Rio Grande Valley near Las Cruces, New Mexico, USDA-SCS, Washington D.C., 1979.
- Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S.: Sources and distributions of dust

- aerosols simulated with the GOCART model, *J. Geophys. Res.*, 106, 20255–20273, doi:10.1029/2000JD000053, 2001.
- Ginoux, P., Prospero, J. M., Gill, T. E., Hsu, N. C., and Zhao, M.: Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products, *Rev. Geophys.*, 50, RG3005, doi:10.1029/2012RG000388, 2012.
- Gong, S. L., Zhang, X. Y., Zhao, T. L., Zhang, X. B., Barrie, L. A., Mckendry, I. G., and Zhao, C. S.: A Simulated climatology of Asian dust aerosol and its trans-Pacific transport, Part II: Inter-annual variability and climate connections, *J. Climate*, 19, 104–122, doi:10.1175/JCLI3606.1, 2006.
- Goudie, A. S.: Desert dust and human health disorders, *Environ. Int.*, 63, 101–113, doi:10.1016/j.envint.2013.10.011, 2014.
- Groll, M., Opp, Chr., and Aslanov, I.: Spatial and temporal distribution of the dust deposition in Central Asia – results from a long term monitoring program, *Aeolian Res.*, 9, 49–62, doi:10.1016/j.aeolia.2012.08.002, 2013.
- Huang, J., Fu, Q., Su, J., Tang, Q., Minnis, P., Hu, Y., Yi, Y., and Zhao, Q.: Taklimakan dust aerosol radiative heating derived from CALIPSO observations using the Fu-Liou radiation model with CERES constraints, *Atmos. Chem. Phys.*, 9, 4011–4021, doi:10.5194/acp-9-4011-2009, 2009.
- Huang, J., Wang, T., Wang, W., Li, Z., and Yan, H.: Climate effects of dust aerosols over East Asian arid and semi-arid regions, *J. Geophys. Res.-Atmos.*, 119, 11398–11416, doi:10.1002/2014JD021796, 2014.
- Huang, X., Oberhänsli, H., von Suchodoletz H., and Sorrel P.: Dust deposition in the Aral Sea: implications for changes in atmospheric circulation in central Asia during the past 2000 years, *Quaternary Sci. Rev.*, 30, 3661–3674, doi:10.1016/j.quascirev.2011.09.011, 2011.
- Huneus, N., Schulz, M., Balkanski, Y., Griesfeller, J., Prospero, J., Kinne, S., Bauer, S., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Fillmore, D., Ghan, S., Ginoux, P., Grini, A., Horowitz, L., Koch, D., Krol, M. C., Landing, W., Liu, X., Mahowald, N., Miller, R., Morcrette, J.-J., Myhre, G., Penner, J., Perlwitz, J., Stier, P., Takemura, T., and Zender, C. S.: Global dust model intercomparison in AeroCom phase I, *Atmos. Chem. Phys.*, 11, 7781–7816, doi:10.5194/acp-11-7781-2011, 2011.
- Husar, R. B., Tratt, D. M., Schichtel, B. A., Falke, S. R., Li, F., Jaffe, D., Gasso, S., Gill, T., Laulainen, N. S., Lu, F., Reheis, M. C., Chun, Y., Westphal, D., Holben, B. N., Gueymard, C., Mckendry, I., Kuring, N., Feldman, G. C., McClain, C., Frouin, R. J., Merrill, J., DuBois, D., Vignola, F., Murayama, T., Nickovic, S., Wilson, W. E., Sassen, K., Sugimoto, N., and Malm, W. C.: The Asian dust events of April 1998, *J. Geophys. Res.*, 106, 18317–18330, doi:10.1029/2000JD900788, 2001.
- Indoitu, R., Orlovsky, L., and Orlovsky, N.: Dust storms in Central Asia: Spatial and temporal variations, *J. Arid Environ.*, 85, 62–70, doi:10.1016/j.jaridenv.2012.03.018, 2012.
- IPCC: Intergovernmental Panel on Climate Change, *Climate change-the physical science basis*, Cambridge University Press, New York, 252–271, 2013.
- Kang, L., Ji, M., Huang, J., and Guan, X.: Impact of Eurasian atmospheric circulation on dust events in spring over northern China, *J. Desert Res.*, 33, 1453–1460, doi:10.7522/j.issn.1000-694X.2013.00211, 2013.
- Khalaf, F. and Al-Hashash, M.: Aeolian sedimentation in the north-western part of the Arabian Gulf, *J. Arid Environ.*, 6, 319–332, 1983.
- Li, F., Zhao, L., Zhang, H., Zhang, T., and Shirato, Y.: Wind erosion and airborne dust deposition in farmland during spring in the Horqin Sandy Land of eastern Inner Mongolia, China, *Soil Till. Res.*, 75, 121–130, doi:10.1016/j.still.2003.08.001, 2004.
- Li, J., Zhuang, G., Huang, K., Lin, Y., Xu, C., and Yu, S.: Characteristics and sources of air-borne particulate in Urumqi, China, the upstream area of Asia dust, *Atmos. Environ.*, 42, 776–787, doi:10.1016/j.atmosenv.2007.09.062, 2008.
- Li, J., Wang, Z., Zhuang, G., Luo, G., Sun, Y., and Wang, Q.: Mixing of Asian mineral dust with anthropogenic pollutants over East Asia: a model case study of a super-duststorm in March 2010, *Atmos. Chem. Phys.*, 12, 7591–7607, doi:10.5194/acp-12-7591-2012, 2012.
- Li, X., Feng, G., Sharratt, B., and Zheng, Z.: Aerodynamic properties of agricultural and natural surfaces in north-western Tarim Basin, *Agr. Forest Meteorol.*, 204, 37–45, doi:10.1016/j.agrformet.2015.01.005, 2015.
- Lin, Y. and Feng, J.: Aeolian dust contribution to the formation of alpine soils at Amdo (Northern Tibetan Plateau), *Geoderma*, 259–260, 104–115, doi:10.1016/j.geoderma.2015.05.012, 2015.
- Liu, L. Y., Shi, P. J., Gao, S. Y., Zou, X. Y., Erdon, H., Yan, P., Li, X. Y., Ta, W. Q., Wang, J. H., and Zhang, C. L.: Dustfall in China's western loess plateau as influenced by dust storm and haze events, *Atmos. Environ.*, 38, 1699–1703, doi:10.1016/j.atmosenv.2004.01.003, 2004.
- Mahowald, N. M., Kohfeld, K. E., Hansson, M., Balkanski, Y., Harrison, S. P., Prentice, I. C., Michael, S., and Rodhe, H.: Dust sources and deposition during the last glacial maximum and current climate: a comparison of model results with paleodata from ice cores and marine sediments, *J. Geophys. Res.*, 104, 15895–916, doi:10.1029/1999JD900084, 1999.
- Mahowald, N. M., Engelstaedter, S., Luo, C., Sealy, A., Artaxo, P., Benitez-Nelson, C., Bonnet, S., Chen, Y., Chuang, P.Y., Cohen, D. D., Dulac, F., Herut, B., Johansen, A. M., Kubilay, N., Losno, R., Maenhaut, W., Paytan, A., Prospero, J. M., Shank, L. M., and Siefert, R. L.: Atmospheric iron deposition: global distribution, variability, and human perturbations, *Annu. Rev. Mar. Sci.*, 1, 245–278, doi:10.1146/annurev.marine.010908.163727, 2009.
- Mamtimin, B. and Meixner, F.: Air pollution and meteorological processes in the growing dryland city of Urumqi (Xinjiang, China), *Sci. Total Environ.*, 409, 1277–1290, doi:10.1016/j.scitotenv.2010.12.010, 2011.
- Mao, R., Ho, C., Shao, Y., Gong, D., and Kim, J.: Influence of Arctic Oscillation on dust activity over northeast Asia, *Atmos. Environ.*, 45, 326–337, doi:10.1016/j.atmosenv.2010.10.020, 2011.
- McTainsh, G. H. and Walker, P.H.: Nature and distribution of Harmattan dust, *Z. Geomorphol.*, 26, 417–435, 1982.
- Menéndez, I., Díaz-Hernández, J. L., and Mangasa, J.: Airborne dust accumulation and soil development in the North-East sector of Gran Canaria (Canary Islands, Spain), *J. Arid Environ.*, 71, 57–81, doi:10.1016/j.jaridenv.2007.03.011, 2007.
- MEP: Ministry of Environmental Protection, China, *Ambient Air Determination of Dustfall Gravimetric Method (GB/T 15265–1994)*, Standards Press of China, Beijing, 1994.
- MEP: Ministry of Environmental Protection, China, *Determination of atmospheric particles PM₁₀ and PM_{2.5} in ambient air by*

- gravimetric method (HJ 618–2011), Standards Press of China, Beijing, 2011.
- MEP: Ministry of Environmental Protection, China, Ambient air quality standards (GB 3095–2012), Standards Press of China, Beijing, 2012a.
- MEP: Ministry of Environmental Protection, China, Technical Regulation on Ambient Air Quality Index (HJ 633–2012), Standards Press of China, Beijing, 2012b.
- Miller-Schulze, J. P., Shafer, M., Schauer, J. J., Heo, J., Solomon, P. A., Lantz, J., Artamonova, M., Chen, B., Imashev, S., Sverdluk, L., Carmichael, G., and DeMinter, J.: Seasonal contribution of mineral dust and other major components to particulate matter at two remote sites in Central Asia, *Atmos. Environ.*, 119, 11–20, doi:10.1016/j.atmosenv.2015.07.011, 2015.
- Modaihash, A. S.: Characteristics and composition of the falling dust sediments on Riyadh city, Saudi Arabia, *J. Arid Environ.*, 36, 211–223, doi:10.1006/jare.1996.0225, 1997.
- O'Hara, S. L., Clarke, M. L., and Elatrash, M. S.: Field measurements of desert dust deposition in Libya, *Atmos. Environ.*, 40, 3881–3897, doi:10.1016/j.atmosenv.2006.02.020, 2006.
- Opp, C., Groll, M., Aslanov, I., Lotz T., and Vereshagina, N.: Aeolian dust deposition in the southern Aral Sea region (Uzbekistan): Ground-based monitoring results from the LUCA project, *Quatern. Int.*, doi:10.1016/j.quaint.2015.12.103, 2016.
- Osada, K., Ura, S., Kagawa, M., Mikami, M., Tanaka, T. Y., Matoba, S., Aoki, K., Shinoda, M., Kurosaki, Y., Hayashi, M., Shimizu, A., and Uematsu, M.: Wet and dry deposition of mineral dust particles in Japan: factors related to temporal variation and spatial distribution, *Atmos. Chem. Phys.*, 14, 1107–1121, doi:10.5194/acp-14-1107-2014, 2014.
- Park, T. W., Ho, C. H., Jeong, S. J., Choi, Y. S., Park, S. K., and Song, C. K.: Different characteristics of cold day and cold surge frequency over East Asia in a global warming situation, *J. Geophys. Res.*, 116, D12118, doi:10.1029/2010JD015369, 2011.
- Péwé, T. L.: Desert dust: origin, characteristics, and effect on man, *Geol. Soc. Am. Bull.*, 186, 1–10, doi:10.1130/SPE186-p1, 1981.
- Petroff, A., Mailliat, A., Amielh, M., and Anselmet, F.: Aerosol dry deposition on vegetative canopies. Part I: Review of present knowledge, *Atmos. Environ.*, 42, 3625–3653, doi:10.1016/j.atmosenv.2007.09.043, 2008.
- Pi, H., Feng, G., Sharratt, B., Li, X., and Zheng, Z.: Validation of SWEEP for contrasting agricultural land use types in the Tarim Basin, *Soil Sci.*, 179, 433–445, doi:10.1097/SS.0000000000000083, 2014.
- Prospero J. M.: Long-range transport of mineral dust in the global atmosphere: impact of African dust on the environment of the southeastern United States, *P. Natl. Acad. Sci. USA*, 96, 3396–3403, doi:10.1073/pnas.96.7.3396, 1999.
- Pye, K.: *Aeolian dust and dust deposits*, Academic Press, London, 5–63, 1987.
- Rittner, M., Vermeesch, P., Carter, A., Bird, A., Stevens, T., Garzanti, E., Andò, S., Vezzoli, G., Dutt, R., Xu, Z., and Lu, H.: The provenance of Taklamakan desert sand, *Earth Planet. Sc. Lett.*, 437, 127–137, doi:10.1016/j.epsl.2015.12.036, 2016.
- Saeid, H., Khademia, H., Angel, F., and Ahmad, L.: Characteristics of dust deposited along a transect between central Iran and the Zagros Mountains, *Catena*, 88, 27–36, doi:10.1016/j.catena.2011.09.002, 2012.
- Shao, Y.: *Physics and modeling of wind erosion*, Dordrecht, Kluwer Academic Publishers, 225–278, 2000.
- Shao, Y. and Dong, C.: A review on East Asian dust storm climate, modelling and monitoring, *Global Planet. Change*, 52, 1–22, doi:10.1016/j.gloplacha.2006.02.011, 2006.
- Shao, Y., Wyrwoll, K., Chappell, A., Huang, J., Lin, Z., McTainsh, G., Mikami, M., Tanaka, T., Wang, X., and Yoon, S.: Dust cycle: an emerging core theme in Earth system science, *Aeolian Res.*, 2, 181–204, doi:10.1016/j.aeolia.2011.02.001, 2011.
- Shao, Y., Kolse, M., and Wyrwoll, K.: Recent global dust trend and connections to climate forcing, *J. Geophys. Res.*, 118, 11107–11118, doi:10.1002/jgrd.50836, 2013.
- Sharratt, B. and Lauer, D.: Particulate matter concentration and air quality affected by windblown dust in the Columbia Plateau, *J. Environ. Qual.*, 35, 2011–2016, doi:10.2134/jeq2006.0212, 2006.
- Shoemaker, J. K., Schrag, D. P., Molina, M. J., and Ramanathan, V.: What role for short-lived climate pollutants in mitigation policy, *Science*, 342, 1323–1324, doi:10.1126/science.1240162, 2013.
- Simonson, R.: Airborne dust and its significance to soil, *Geoderma*, 65, 1–43, doi:10.1016/0016-7061(94)00031-5, 1995.
- Slinn, S. A. and Slinn, W. G. N.: Predictions for particle deposition on natural waters, *Atmos. Environ.*, 14, 1013–1016, doi:10.1016/0004-6981(80)90032-3, 1980.
- Song, H., Zhang, K., Piao, S., and Wan, S.: Spatial and temporal variations of spring dust emissions in northern China over the last 30 years, *Atmos. Environ.*, 126, 117–127, doi:10.1016/j.atmosenv.2015.11.052, 2016.
- Stanhill, G.: Global dimming: a new aspect of climate change, *Weather*, 60, 11–14, doi:10.1256/wea.210.03, 2005.
- Sun, J. and Liu, T.: The age of the Taklimakan Desert, *Science*, 312, 1621, doi:10.1126/science.1124616, 2006.
- Tsoar, H. and Pye, K.: Dust transport and the question of desert loess formation, *Sedimentology*, 34, 139–153, doi:10.1111/j.1365-3091.1987.tb00566.x, 1987.
- Uno, I., Eguchi, K., and Yumimoto, K.: Asian dust transported one full circuit around the globe, *Nat. Geosci.*, 2, 557–560, doi:10.1038/ngeo583, 2009.
- USEPA: Guideline for reporting of daily air quality – the Air Quality Index (AQI), US Environmental Protection Agency, Office of Air Quality Planning and Standards, Research Triangle Park, North Carolina, 2006.
- USDA: Dust deposition and weather, *Weatherwise*, 21, 65–70, doi:10.1080/00431672.1968.9932796, 1968.
- Varga, G., Cserhádi, C., Kovács, J., and Szaljai, Z.: Saharan dust deposition in the Carpathian Basin and its possible effects on interglacial soil formation, *Aeolian Res.*, 22, 1–12, doi:10.1016/j.aeolia.2016.05.004, 2016.
- Wake, C. P. and Mayewski, P. A.: Modern aeolian dust deposition in central Asia, *Tellus*, 46B, 220–233, doi:10.1034/j.1600-0889.1994.t01-2-00005.x, 1994.
- Wang, H., Jia, X., Lib, K., and Wang, H.: External supply of dust in the Taklamakan sand sea, Northwest China, reveals the dust-forming processes of the modern sand sea surface, *Catena*, 119, 104–115, doi:10.1016/j.catena.2014.03.015, 2014.
- Wang, L., Zhang, P., Tan, S., Zhao, X., Cheng, D., Wei, W., Su, J., and Pan, X.: Assessment of urban air quality in China using air pollution indices (APIs), *J. Air Waste Manage. Assoc.*, 63, 170–178, doi:10.1080/10962247.2012.739583, 2013.

- Wang, X., Oenema, O., Hoogmoed, W. B., Perdok, U. D., and Cai, D.: Dust storm erosion and its impact on soil carbon and nitrogen losses in northern China, *Catena*, 66, 221–227, doi:10.1016/j.catena.2006.02.006, 2006.
- Wang, Y. M., Wang, J. H., Qi, Y., and Yan, C. Z.: China 1 : 100000 Desert (Sand land) Map, Environmental and Ecological Science Data Centre for West China (<http://westdc.westgis.ac.cn>), doi:10.3972/westdc.006.2013.db, 2005.
- Wei, W. S., Gao, W. D., Shi, Y. G., and Osamu, A. B. E.: Influence of climate and environment change on dust storms in Xinjiang, China, *J. Arid Land Geogr.*, 27, 137–141, doi:10.13826/j.cnki.cn65-1103/x.2004.02.002, 2004.
- Wesely M. L. and Hicks B. B.: A review of the current status of knowledge on dry deposition, *Atmos. Environ.*, 34, 2261–2282, doi:10.1016/S1352-2310(99)00467-7, 2000.
- WHO: Ambient Air Pollution database, available at: <http://gamapserver.who.int/mapLibrary/app/searchResults.aspx>, last access: May 2016a.
- WHO: Ambient (outdoor) air quality and health, available at: <http://www.who.int/mediacentre/factsheets/fs313/en/>, last access: September 2016b.
- Xi, X. and Sokolik, I. N.: Dust interannual variability and trend in Central Asia from 2000 to 2014 and their climatic linkages, *J. Geophys. Res.*, 120, 12175–12197, doi:10.1002/2015JD024092, 2015.
- Xinjiang Statistical Bureau: Xinjiang Statistical Yearbook 2014, China Statistical Press, Beijing, 60–225, 2014.
- Xu, J., Hou, S., Qin, D., Kang, S., Ren, J., and Ming, J.: Dust storm activity over the Tibetan Plateau recorded by a shallow ice core from the north slope of Mt. Qomolangma (Everest), Tibet-Himal region, *Geophys. Res. Lett.*, 34, L17504, doi:10.1029/2007GL030853, 2007.
- Xuan, J.: Dust emission factors for environment of northern China, *Atmos. Environ.*, 33, 1767–1776, doi:10.1016/S1352-2310(98)00339-2, 1999.
- Xuan, J., Liu, G. L., and Du, K.: Dust emission inventory in northern China, *Atmos. Environ.*, 34, 4565–4570, doi:10.1016/S1352-2310(00)00203-X, 2000.
- Yalon, D. H. and Ganor, E.: Rates of aeolian dust accretion in the Mediterranean and desert fringe environments of Israel, 9th International Sedimentological Congress, Nice, France, 2, 169–174, 1975.
- Zan, J., Fang, X., Appel, E., Yan, M., and Yang, S.: New insights into the magnetic variations of aeolian sands in the Tarim Basin and its paleoclimatic implications, *Phys. Earth Planet. In.*, 229, 82–87, doi:10.1016/j.pepi.2014.01.010, 2014.
- Zhang, D.: Preliminary analysis on climatology of dust deposition in China history, *Sci. Sinica*, 3, 278–288, 1984.
- Zhang, J. and Shao, Y.: A new parameterization of particle dry deposition over rough surfaces, *Atmos. Chem. Phys.*, 14, 12429–12440, doi:10.5194/acp-14-12429-2014, 2014.
- Zhang, J., Shao, Y., and Huang, N.: Measurements of dust deposition velocity in a wind-tunnel experiment, *Atmos. Chem. Phys.*, 14, 8869–8882, doi:10.5194/acp-14-8869-2014, 2014.
- Zhang, X. X., Shi, P. J., Liu, L. Y., Tang, Y., Cao, H. W., Zhang, X. N., Hu, X., Guo, L. L., Lue, Y. L., Qu, Z. Q., Jia, Z. J., and Yang, Y. Y.: Ambient TSP concentration and dustfall in major cities of China: spatial distribution and temporal variability, *Atmos. Environ.*, 44, 1641–1648, doi:10.1016/j.atmosenv.2010.01.035, 2010.
- Zhang, X. X., Chen, X., Guo, Y. H., Wang, Z. F., Liu, L. Y., Paul, C., Li, S. Y., and Pi, H. W.: Ambient TSP concentration and dustfall variation in Urumqi, China, *J. Arid Land*, 6, 668–677, doi:10.1007/s40333-014-0069-6, 2014.
- Zhang, X. Y., Shen, Z. B., Zhang, G. Y., Chen, T., and Liu, H.: Remote mineral aerosols in westerlies and their contributions to the Chinese loess, *Sci. China Ser. D*, 39, 134–143, 1996.
- Zhang, X. Y., Arimoto, R., and An, Z. S.: Dust emission from Chinese desert sources linked to variations in atmospheric circulation, *J. Geophys. Res.*, 102, 28041–28047, doi:10.1029/97JD02300, 1997.
- Zheng, Y., Zhao, T., Che, H., Liu, Y., Han, Y., Liu, C., Xiong, J., Liu, J., and Zhou, Y.: A 20-year simulated climatology of global dust aerosol deposition, *Sci. Total Environ.*, 557–558, 861–868, doi:10.1016/j.scitotenv.2016.03.086, 2016.
- Zu, R., Xue, X., Qiang, M., Yang, B., Qu, J., and Zhang, K.: Characteristics of near-surface wind regimes in the Taklimakan Desert, China, *Geomorphology*, 96, 39–47, doi:10.1016/j.geomorph.2007.07.008, 2008.