

Supplement of Atmos. Chem. Phys., 18, 11581–11597, 2018
<https://doi.org/10.5194/acp-18-11581-2018-supplement>
© Author(s) 2018. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Production of N_2O_5 and ClNO_2 in summer in urban Beijing, China

Wei Zhou et al.

Correspondence to: Yele Sun (sunyele@mail.iap.ac.cn) and Hugh Coe (hugh.coe@manchester.ac.uk)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.




Figure S1. High resolution peak fitting for (a) m/z 235 as $\text{I}\cdot\text{N}_2\text{O}_5^-$, (b) m/z 208 as $\text{I}\cdot\text{ClNO}_2^-$, (c) m/z 210 as $\text{I}\cdot^{37}\text{ClNO}_2^-$, and (d) average high-resolution mass spectrum for one night from the IAP-CIMS measurement.




Figure S2. Comparison of IAP-CIMS raw signals of N_2O_5 with those measured by the Cambridge Broadband Cavity Enhanced Absorption Spectrometer (BBCEAS). The derived sensitivity of IAP-CIMS N_2O_5 is $0.54 \text{ cps ppt}^{-1}$.




Figure S3. Time series of NO and BC during the campaign.




Figure S4. Variations of (a) aerosol surface area (S_a), and (b) aerosol liquid water content (LWC) as a function of RH. (c) The average mixing ratios of and NO_3 , N_2O_5 , NO_2 , O_3 and the nitrate radical production rate $p(\text{NO}_3)$ for four different nights (i.e., P1, P2, P3 and P4).




Figure S5. The 48-h back trajectories arrived at the sampling site during four different nights (i.e., P1, P2, P3 and P4). Note that the trajectories at each night are divided into two periods, i.e., before and after midnight.




Figure S6. The correlations between ClNO_2 and N_2O_5 for four different nights (i.e., P1, P2, P3 and P4). All the correlations are divided into two periods, i.e., before midnight and after midnight.