

	355 nm	532 nm	1064 nm	Reference
Gobi dust				
Vancouver, Canada	–	0.27	–	Cottle et al. (2013)
Dushanbe, Tajikistan	0.18–0.29	0.31–0.35	–	Hofer et al. (2017)
Tsukuba, Japan	–	0.35	0.35	Sugimoto and Lee (2006)
Gwangju, South Korea	–	0.30–0.33	–	Shin et al. (2015)
Arabian dust				
Limassol, Cyprus	–	0.28-0.35	–	Mamouri and Ansmann (2013)
Saharan dust				
Ouarzazate, Morocco	0.240 ± 0.007	0.310 ± 0.020	0.270 ± 0.040	Freudenthaler et al. (2009)
M'Bour, Senegal	–	0.300 ± 0.045	–	Veselovskii et al. (2016)
Evora, Portugal	–	0.280 ± 0.040	–	Preißler et al. (2011)
Munich, Germany	0.290 ± 0.070	0.340 ± 0.020	–	Wiegner et al. (2011)
Barbados	0.252 ± 0.030	0.280 ± 0.020	0.225 ± 0.020	Haarig et al. (2017)
Caribbean	–	0.327 ± 0.018	0.278 ± 0.001	Burton et al. (2015)
Midwest USA	0.246 ± 0.018	0.304 ± 0.005	0.270 ± 0.005	Burton et al. (2015)
Great Basin dust				
Pico de Orizaba	–	0.334 ± 0.018	0.400 ± 0.009	Burton et al. (2015)
Chihuahuan desert	0.243 ± 0.046	0.373 ± 0.014	0.383 ± 0.006	Burton et al. (2015)