

	Wavelength	Wet season (Feb–May)		Dry season (Aug–Nov)		Entire period (2012–2017)	
		Mean ± SD	IQR	Mean ± SD	IQR	Mean ± SD	IQR
Scattering coefficient σ_{sp} (M m^{-1})	450 nm	9 ± 11	(5.1–11)	47 ± 35	(24–64)	31 ± 35	(10–39)
	550 nm	7.5 ± 9.3	(3.8–8.7)	33 ± 25	(17–46)	22 ± 25	(7–28)
	637 nm	6.4 ± 8.9	(3.0–7.4)	26 ± 19	(13–35)	17 ± 19	(6–23)
Absorption coefficient σ_{ap} (M m^{-1})	637 nm	0.68 ± 0.91	(0.17–0.72)	4.0 ± 2.2	(2.4–5.1)	2.1 ± 2.2	(0.43–3.2)
Single scattering albedo ω_0	637 nm	0.93 ± 0.04	(0.91–0.96)	0.87 ± 0.03	(0.84–0.89)	0.89 ± 0.04	(0.86–0.93)
Scattering Ångström exp. ¹ \AA_{sca}		1.29 ± 0.50	(0.98–1.65)	1.71 ± 0.24	(1.53–1.89)	1.54 ± 0.42	(1.32–1.84)
Absorption Ångström exp. ¹ \AA_{abs}		0.91 ± 0.19	(0.80–0.98)	0.94 ± 0.16	(0.84–1.03)	0.93 ± 0.16	(0.83–1.01)
Mass absorption cross section α_{abs} ($\text{m}^2 \text{ g}^{-1}$) ²	637 nm	11.4 ± 1.2	(10.8–12.0)	12.3 ± 1.3 ³	(11.4–13.3) ³	11.9 ± 1.4	(11.0–13.0)