

Supplement of Atmos. Chem. Phys., 18, 13581–13600, 2018
<https://doi.org/10.5194/acp-18-13581-2018-supplement>
© Author(s) 2018. This work is distributed under
the Creative Commons Attribution 4.0 License.



Atmospheric
Chemistry
and Physics
Open Access
EGU

Supplement of

Counteractive effects of regional transport and emission control on the formation of fine particles: a case study during the Hangzhou G20 summit

Ying Ji et al.

Correspondence to: Kan Huang (huangkan@fudan.edu.cn) and Congrui Deng (congruideng@fudan.edu.cn)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.




Figure S1. Temporal evolutions of precipitation during entire study period.




Figure S2. The hourly SOC plotted against temperature and colored with RH during the whole study period.




Figure S3. Time series of hourly concentrations of SNA , Cl⁻, and meteorological parameters (WS, WD, RH, and Radiation) in S4




Figure S4. Concentrations of NO₂ and SO₂ in the morning (0:00-5:00AM LST) of 2 September in urban cities of Shandong.




Figure S5. The illustration of minimum R^2 method to calculate $(OC/EC)_{pri}$ for each stage. The red curve indicates the correlation coefficient (R^2) between OC and EC as a function of assumed $(OC/EC)_{pri}$. The shaded area in tan is the frequency distribution of the OC/EC ratio for the entire OC and EC data set. The green dashed curve is the cumulative frequency curve of OC/EC ratio.

Table S1. The temperature in different areas that the hotspots had covered during the three days before S3 according to the 72h air mass back trajectory in CWT analysis.

Date	Tangshan	Langfan	Cangzhou	Lianyungang	Yangzhou	Wuxi	Huangshan	Xuancheng	Shangrao
8/25	23	25	23	26	28	29	30	29	32
8/26	21	22	21	23	24	27	29	27	32
8/27	22	22	22	23	24	25	25	24	27
8/28	22	24	22	24	24	25	26	24	25
8/29	23	23	23	25	23	25	24	24	25
8/30	23	24	24	24	23	25	24	23	25
8/31	24	26	25	24	24	25	24	25	24

Table S2. A review of SOC/OC ratios in different urban areas.

City	period	Categories	Method	SOC/OC	Reference
Hangzhou, China	Summer	PM _{2.5}	EC-Tracer method	38%	My study
Hangzhou, China	Winter	PM _{2.5}	EC-Tracer method	40%	Li et al. (2016)
Hangzhou, China	Winter, Spring, Summer, Autumn	PM ₁₀	EC-Tracer method	44.7%, 37.1%, 45.8%, 58%	Jiao et al. (2007)
Tianjin, China	Annual average	PM _{2.5}	EC-Tracer method	61.70%	Gu et al. (2010)
Beijing, China	winter, Spring, Summer, Autumn	PM _{2.5}	EC-Tracer method	19%, 27%, 45%, 23%	Lin et al. (2009)
California	Summer	PM _{2.5}	EC-Tracer method	30%-80%	Gray et al. (1986); Pandis et al. (1992); Hildemann et al. (1993)