

Supplement of Atmos. Chem. Phys., 18, 4055–4068, 2018
<https://doi.org/10.5194/acp-18-4055-2018-supplement>
© Author(s) 2018. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Primary and secondary organic aerosols in summer 2016 in Beijing

Rongzhi Tang et al.

Correspondence to: Song Guo (songguo@pku.edu.cn)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Figure captions

Fig. S1 Locations of the sampling sites CP and PKUERS

Fig. S2 Mixing ratios of gaseous pollutants and meteorological conditions at (a) CP and (b) PKUERS, including concentrations of SO₂, O₃, NO, NO₂, temperature, relative humidity, wind speed as well as wind direction. Parameters of the gaseous pollutants and meteorological conditions from 1 to 5 June were not available due to technical reasons.

Fig. S3 Back trajectory clusters during the campaign at (a) CP (b) PKUERS

Fig. S4 PM_{2.5} chemical compositions at the regional site CP (a & c) and urban site PKUERS (b & d)

Fig. S5 Three categories of PM_{2.5} chemical compositions according to back trajectory clustering analysis

Fig. S6 Primary organic matter from different directions of CP and PKUERS

Fig. S7 Daily average concentrations for different classes of primary organic matter, (a) n-alkanes (b) saccharides (c) PAHs (d) hopanes




Fig. S1 Locations of the sampling sites CP and PKUERS




Fig. S2 Mixing ratios of gaseous pollutants and meteorological conditions at (a) CP and (b) PKUERS, including concentrations of SO₂, O₃, NO, NO₂, temperature, relative humidity, wind speed as well as wind direction. Parameters of the gaseous pollutants and meteorological conditions from 1 to 5 June were not available due to technical reasons.




Fig. S3 Back trajectory clusters during the campaign at (a) CP (b) PKUERS




Fig. S4 PM_{2.5} chemical compositions at the regional site CP (a & c) and urban site PKUERS (b & d)




Fig. S5 Three categories of PM_{2.5} chemical compositions according to back trajectory clustering analysis




Fig. S6 Primary organic matter from different directions of CP and PKUERS




Fig. S7 Daily average concentrations for different classes of primary organic matter, (a) n-alkanes (b) saccharides (c) PAHs (d) hopanes.

Table

Table S1 Mixing ratios of gaseous pollutants (SO₂, O₃, NO, NO₂)

Type	Sites	NO	NO ₂	O ₃	SO ₂	CO
Urban	PKUERS (2010) (Zheng et al., 2016a)	7.1	26.0	32.0	4.9	820.0
	PKUERS (2016)	10.5	32.3	48.6	3.9	623.6
	CP(2016)	3.1	19.7	57.7	3.0	492.4

Table S2 Concentrations of tracer compounds for primary sources

	CP			PKUERS		
	Daytime	Nighttime	Diurnal	Daytime	Nighttime	Diurnal
Tracers (ng·m⁻³)						
n-alkanes (C28-C33)						
C-28	2.74±1.18	3.34±2.29	3.10±1.84	3.99±1.93	5.06±2.44	4.54±2.24
C-29	23.12±14.38	22.54±13.38	22.82±13.68	20.49±8.48	24.96±12.69	22.79±10.94
C-30	1.54±0.62	2.19±1.25	1.88±1.04	2.06±0.72	3.25±1.30	2.72±1.23
C-31	5.22±1.98	5.85±3.40	5.54±2.78	5.23±2.31	7.24±3.72	6.23±3.22
C-32	1.67±1.05	1.72±1.11	1.70±1.07	1.91±0.92	2.69±1.71	2.34±1.43
C-33	1.83±0.94	2.49±1.25	2.19±1.15	2.54±1.12	2.74±1.51	2.65±1.32
Levoglucosan						
Levoglucosan	28.86±16.95	75.92±40.91	53.03±39.26	39.09±25.04	79.56±39.97	59.87±38.93
Hopanes						
17 α (H)-22,29,30-trishopane	0.61±0.28	0.65±0.44	0.63±0.36	0.88±0.55	0.93±0.52	0.90±0.53
17 β (H)-21 α (H)-norhopane	1.42±0.48	1.66±0.97	1.55±0.70	1.93±1.24	1.79±0.96	1.86±1.09
17 α (H)-21 β (H)-hopane	0.80±0.32	0.95±0.59	0.88±0.47	1.12±0.64	1.16±0.55	1.14±0.58
PAHs						
benzo(b)fluoranthene	0.64±0.45	2.45±1.68	1.57±1.53	0.53±0.47	1.07±1.00	0.81±0.82
benzo(k)fluoranthene	0.50±0.25	1.42±0.73	0.98±0.72	0.50±0.38	0.69±0.54	0.60±0.47
benzo(e)pyrene	0.38±0.22	1.09±0.67	0.74±0.61	0.59±0.39	0.70±0.40	0.65±0.39
benzo(ghi)perylene	0.71±0.40	2.48±1.63	1.81±1.56	1.15±0.43	1.84±0.80	1.54±0.73
indeno(1,2,3-cd)pyrene	0.76±0.52	2.78±2.24	2.01±2.03	0.79±0.26	2.02±0.97	1.48±0.96

Table S3 Comparison of SOA tracers for the biogenic and anthropogenic sources in different regions

Species	Tracer (ng·m ⁻³)	IITB	CH	YL	CL	RTP	Yufa 2008	PKUERS 2008	PKUERS 2016	CP 2016
Isoprene	I-1	0.4±0.4	0.3±0.2	1.0±1.3	-	26.5±20.9	29.9±19.9	21.0±19.5	1.2±0.7	1.4±1.7
	I-2	-	0.2±0.1	8.3±12.9	11.2±5.8	63.4±19.2	63.6±46.8	42.2±28.2	7.7±4.7	9.9±4.6
	I-3	-	0.5±0.3	20.3±20.3	24.1±13.4	85.7±27.6	121.5±101.4	77.2±60.2	16.8±6.3	10.5±5.7
	∑isoprene	1.9±2.0	1.0±0.6	29.6±34.5	35.3±19.2	175.6±49.7	215.0±160.3	140.4±100.9	25.7±11.7	18.8±12.0
α-pinene	A-1	-	-	8.4±10.7	17.2±6.3	1.8±0.7	9.6±7.2	8.7±7.1	7.0±2.7	9.2±5.5
	A-2	-	-	5.9±4.9	17.1±5.4	16.3±7.3	5.3±2.8	6.8±7.1	4.8±2.8	4.2±2.4
	A-3	-	-	-	-	25.9±13.9	4.3±6.9	4.6±4.0	15.6±8.3	11.8±6.3
	A-4	-	-	4.6±3.6	-	46.1±18.9	7.7±5.5	8.3±5.8	4.8±1.2	4.4±1.3
	A-5	2.0±1.2	-	10.2±7.0	-	53.5±17.9	57.5±27.4	51.7±31.3	16.8±6.3	10.5±5.7
	A-6	-	-	9.7±15.1	-	12.0±1.0	8.3±6.7	9.9±9.1	13.0±4.8	13.6±8.0
	A-7	0.3±0.2	-	8.0±7.5	2.2±1.0	4.7±1.3	6.1±5.5	8.5±10.1	12.4±6.0	13.6±8.0
	PA	0.6±0.3	0.2±0.5	-	4.4±3.5	9.2±3.6	3.4±5.7	3.2±6.4	10.0±4.5	9.5±12.7
	PNA	5.5±3.9	0.1±0.1	-	6.7±1.9	-	21.9±9.2	11.9±6.3	11.9±3.3	15.4±8.1
	∑α-pinene	8.4±5.6	0.3±0.6	46.8±56.3	47.6±18.1	169.5±64.6	124.1±47.8	113.5±63.3	96.4±39.9	92.2±58.0
β-caryophyllene	C-1	0.7±0.3	0.2±0.5	12.5±16.9	1.4±0.4	25.3±4.2	3.6±2.3	5.1±4.5	6.0±2.8	6.1±3.5
Toluene	T-3	0.1±0.1	-	1.7±1.8	8.3±2.8	4.1±1.9	11.7±6.9	13.3±7.7	11.0±3.7	9.7±7.3

I-1 represented 2-methylglyceric acid, I-2 represented 2-Methylthreitol, I-3 represented 2-methylerythritol, A-1 represented 3-Isopropylpentanedioic acid, A-2 represented 3-Acetylpentanedioic acid, A-3 represented 2-Hydroxy-4-isopropyladipic acid, A-4 represented 3-Acetyl hexanedioic acid, A-5 represented 3-Hydroxyglutaric acid, A-6 represented 2-Hydroxy-4,4-dimethylglutaric acid, A-7 represented 3-(2-Hydroxy-ethyl)-2,2-dimethyl-cyclobutane-carboxylic acid, PA represented pinic acid, PNA represented pinonic acid, T-3 represented 2,3-dihydroxy-4-oxopentanoic acid, C-1 represented β-caryophyllinic acid.