

A parameterization of the heterogeneous hydrolysis of N₂O₅ for mass-based aerosol models: improvement of particulate nitrate prediction

Ying Chen^{1,2,a}, Ralf Wolke¹, Liang Ran³, Wolfram Birmili^{1,4}, Gerald Spindler¹, Wolfram Schröder¹, Hang Su^{2,5}, Yafang Cheng^{2,5}, Ina Tegen¹, and Alfred Wiedensohler¹

¹Leibniz-Institute for Tropospheric Research (TROPOS), 04318 Leipzig, Germany

²Multiphase Chemistry Department, Max Planck Institute for Chemistry, 55128 Mainz, Germany

³Key Laboratory of Middle Atmosphere and Global Environment Observation, Institute of

Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China

⁴German Environment Agency, 06844 Dessau-Roßlau, Germany

⁵Institute for Environmental and Climate Research, Jinan University, Guangzhou, China

^anow at: Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

Correspondence: Ying Chen (chen@tropos.de) and Ralf Wolke (wolke@tropos.de)

Received: 3 February 2017 – Discussion started: 16 February 2017 Revised: 28 June 2017 – Accepted: 4 December 2017 – Published: 19 January 2018

Abstract. The heterogeneous hydrolysis of N₂O₅ on the surface of deliquescent aerosol leads to HNO₃ formation and acts as a major sink of NO_x in the atmosphere during nighttime. The reaction constant of this heterogeneous hydrolysis is determined by temperature (T), relative humidity (RH), aerosol particle composition, and the surface area concentration (S). However, these parameters were not comprehensively considered in the parameterization of the heterogeneous hydrolysis of N2O5 in previous mass-based 3-D aerosol modelling studies. In this investigation, we propose a sophisticated parameterization (NewN2O5) of N2O5 heterogeneous hydrolysis with respect to T, RH, aerosol particle compositions, and S based on laboratory experiments. We evaluated closure between NewN2O5 and a state-of-the-art parameterization based on a sectional aerosol treatment. The comparison showed a good linear relationship (R = 0.91) between these two parameterizations. NewN2O5 was incorporated into a 3-D fully online coupled model, COSMO-MUSCAT, with the mass-based aerosol treatment. As a case study, we used the data from the HOPE Melpitz campaign (10-25 September 2013) to validate model performance. Here, we investigated the improvement of nitrate prediction over western and central Europe. The modelled particulate nitrate mass concentrations ([NO₃]) were validated by filter measurements over Germany (Neuglobsow, Schmücke, Zingst, and Melpitz). The modelled $[NO_3^-]$ was significantly overestimated for this period by a factor of 5-19, with the corrected NH₃ emissions (reduced by 50%) and the original parameterization of N2O5 heterogeneous hydrolysis. The NewN2O5 significantly reduces the overestimation of $[NO_3^-]$ by ~ 35 %. Particularly, the overestimation factor was reduced to approximately 1.4 in our case study (12, 17-18 and 25 September 2013) when $[NO_3^-]$ was dominated by local chemical formations. In our case, the suppression of organic coating was negligible over western and central Europe, with an influence on $[NO_3^-]$ of less than 2 % on average and 20 % at the most significant moment. To obtain a significant impact of the organic coating effect, N₂O₅, SOA, and NH₃ need to be present when RH is high and T is low. However, those conditions were rarely fulfilled simultaneously over western and central Europe. Hence, the organic coating effect on the reaction probability of N2O5 may not be as significant as expected over western and central Europe.

1 Introduction

The budget of nitrogen oxides (NO_x) is of fundamental importance for tropospheric chemistry (Ehhalt and Drummond, 1982). The removal of nitrogen from the atmosphere is largely contributed by the formation of HNO₃, which is eventually transferred to particles and deposited (Riemer et al., 2003). In the daytime, HNO₃ is mainly produced via the reaction of OH and NO₂. At night, the major sink of NO_x is heterogeneously hydrolyzed N₂O₅ on the surface of deliquescent aerosol (Jacob, 2000; Brown and Stutz, 2012; Platt et al., 1984; Brown et al., 2004). Given that NO_x is the key precursor of ozone, chemical mechanisms controlling the budget of NO_x also greatly affect ozone and thereby the atmospheric oxidizing capacity on a global scale (Dentener and Crutzen, 1993; Evans and Jacob, 2005).

The reaction probability ($\gamma_{N_2O_5}$) is used to quantify the reaction constant of N₂O₅ hydrolysis ($k_{N_2O_5}$) on the surface of deliquescent aerosol. Several techniques could be employed to measure $\gamma_{N_2O_5}$ on the surface of different aqueous solutions (Hallquist et al., 2000; Robinson et al., 1997; Hu and Abbatt, 1997; Behnke et al., 1997; Fenter et al., 1996; Van Doren et al., 1990; Mozurkewich and Calvert, 1988). It was reported that the typical values for $\gamma_{N_2O_5}$ are on the order of 10^{-2} . Organic coating of the particles may reduce this reaction probability. Anttila et al. (2006) proposed a parameterization (Anttila06) that described the organic coating suppression effect on $\gamma_{N_2O_5}$. Based on laboratory experiments and Anttila06, Gaston et al. (2014) reported that the suppression of $\gamma_{N_2O_5}$ by organic coating is dependent on a range of factors including the O: C ratio, the organic particle mass fraction, and the relative humidity (RH). Bertram and Thornton (2009) parameterized the effect of chloride salts on $\gamma_{N_2O_5}$ as a function of RH. However, the influence of temperature was not considered in their study. Recently, Morgan et al. (2015) demonstrated that this influence of chloride may not be represented properly, and the "high" O: C regime defined in Gaston et al. (2014) was scarcely observed over north-western Europe by airborne regional measurements. Several laboratory studies reported that $\gamma_{N_2O_5}$ substantially varies with temperature, RH, and particle composition (Mentel et al., 1999; Kane et al., 2001; Hallquist et al., 2003; Chang et al., 2011; Brown and Stutz, 2012; Gaston et al., 2014). Davis et al. (2008) developed a parameterization scheme (Davis08) of $\gamma_{N_2O_5}$ for aerosols containing nitrate, sulfate, and ammonium. It was developed on the basis of numerous previous laboratory studies (Badger et al., 2006; Folkers et al., 2003; Hallquist et al., 2003; Folkers, 2002; Kane et al., 2001; Hu and Abbatt, 1997; Mozurkewich and Calvert, 1988) with respect to temperature, RH, and particle compositions.

Several studies have implemented the heterogeneous hydrolysis of N₂O₅ in global and regional chemical transport models in order to investigate its influences on atmospheric chemistry. Dentener and Crutzen (1993) assumed a constant $\gamma_{N_2O_5}$ of 0.1 in a global modelling study, which might be

overestimated. Chang et al. (1987) proposed a scheme to parameterize the $k_{N_2O_5}$ for 3-D models without complex aerosol treatments. Riemer et al. (2003) reported that the scheme of Chang et al. (1987) could only be representative of heavily polluted conditions or in the presence of cloud droplets, with a particle surface area concentration of $2700 \,\mu\text{m}^2 \,\text{cm}^{-3}$. Riemer et al. (2003) proposed a more complex scheme (P1 in the literature) with respect to the particle surface area concentration (S) and $\gamma_{N_2O_5}$ of nitrate and sulfate, which were reported by Mentel et al. (1999) and Wahner et al. (1998). However, the influence of temperature and RH on $\gamma_{N_2O_5}$ was still not considered in the P1 of Riemer et al. (2003). Later, by applying Anttila06 to the P1 of Riemer et al. (2003), Riemer et al. (2009) reported an up to 90 % decrease in particulate nitrate due to organic coating in the regions where both N₂O₅ and secondary organic compounds were built up. Evans and Jacob (2005) proposed a parameterization scheme (EJ05) that has an extensive description of aerosol composition to improve the GEOS-CHEM simulations. EJ05 included $\gamma_{N_2O_5}$ of dust (Bauer et al., 2004), sea salt (Sander et al., 2003), sulfate (Hallquist et al., 2003; Kane et al., 2001), elemental carbon (EC; Sander et al., 2003), and organic carbon (OC; Thornton et al., 2003) and also took into account the dependence of $\gamma_{N_2O_5}$ on RH. However, $\gamma_{N_2O_5}$ of nitrate and its dependence on temperature were not carefully considered in EJ05. Archer-Nicholls et al. (2014) incorporated Bertram and Thornton (2009) into WRF-Chem. Lowe et al. (2015) further took the organic coating effect into account by applying Anttila06 to Bertram and Thornton (2009). However, $\gamma_{N_2O_5}$ with respect to EC, OC, and dust was lacking in Bertram and Thornton (2009). Furthermore, as mentioned above, the reported influence of chloride on $\gamma_{N_2O_5}$ (Bertram and Thornton, 2009) may not be realistic in north-western Europe (Morgan et al., 2015). Recently, Chang et al. (2016) improved the P1 (Riemer et al., 2003) with the Davis08 + Anttila06 scheme and incorporated it into WRF-Chem with a sectional aerosol treatment (MO-SAIC, Zaveri et al., 2008). They validated the improved P1 with the aircraft measurements from the CalNex 2010 campaign. Davis08 + Anttila06 showed a better result than that from the scheme according to Bertram and Thornton (2009) and significantly improved the model performance (Chang et al., 2016). However, the influences of black carbon (BC), sea salt aerosol (SSA), and dust were still missing in the parameterizations according to Chang et al. (2016). The P1 scheme (Riemer et al., 2003; Chang et al., 2016) is very helpful for models with complex aerosol treatments (modal and/or sectional aerosol approach; e.g. WRF-Chem with MOSAIC). However, it cannot be easily adopted in the computationally efficient mass-based aerosol approaches, which are commonly used in atmospheric chemistry transport or climate models, for example EMEP (Simpson et al., 2012) and GEOS-Chem (Walker et al., 2012), or long-term modelling studies (e.g. Bellouin et al., 2011; Hardiman et al., 2017). Riemer et al. (2003) also improved a simplified scheme (P2 in the literature) based on the work of Chang et al. (1987), which is easily adopted in the mass-based aerosol models and is currently used in COSMO–MUSCAT (Consortium for Small-scale Modelling and Multi-Scale Chemistry Aerosol Transport, http://projects.tropos.de/cosmo_muscat; Wolke et al., 2004, 2012) with the mass-based aerosol treatment according to Simpson et al. (2003). However, the P2 still showed a large difference in comparison to the more complete P1 (Riemer et al., 2003). The reasons may be the missing complex considerations of *S* and $\gamma_{N_2O_5}$ in the P2.

To improve the representativeness of the heterogeneous hydrolysis of N_2O_5 in 3-D models with mass-based aerosol treatment, we propose a new parameterization (NewN2O5) with respect to temperature, RH, particle composition, and particle surface area. NewN2O5 was validated by the state-of-the-art parameterization in Chang et al. (2016). We also incorporated NewN2O5 into the 3-D fully online coupled model COSMO–MUSCAT in order to investigate the improvement of particulate nitrate prediction. The measurements of the HOPE campaign (HD(CP)² Observational Prototype Experiment; Macke et al., 2017) at the Melpitz site (12.93° E, 51.53° N; 86 m a.s.l.; a regional background observatory of central Europe) and three other stations of the German federal environmental agency (UBA) over Germany were used to validate the simulation results.

2 Data and methods

2.1 The model system COSMO-MUSCAT

The online coupled chemical transport model COSMO– MUSCAT is qualified for the operation forecast of pollutants and process studies in regional and local areas (Heinold et al., 2011; Renner and Wolke, 2010; Hinneburg et al., 2009; Stern et al., 2008). Two nested domains with 50 vertical layers were used for this model study. The outer domain covers the whole of Europe, with a spatial grid resolution of $14 \text{ km} \times 14 \text{ km}$. The inner domain (N2) covers Germany, the Netherlands, and nearby regions, with a spatial grid resolution of $7 \text{ km} \times 7 \text{ km}$ (Fig. 1). The simulation period was divided into several overlapping short periods, each of which included a 1-day spin-up followed by a 2-day run with meteorology and chemistry coupled. The main features of the model system are described below. More details are given in Wolke et al. (2004, 2012) and Baldauf et al. (2011).

An adequate modelling of dynamics requires an online coupling between the chemical transport model MUSCAT and the meteorological model COSMO. Here, the compressible non-hydrostatic flow in a moist atmosphere is described by the primitive hydro-thermodynamical equations (Steppeler et al., 2003; Doms et al., 2011a). The vertical diffusion is parameterized by a level 2.5 closure scheme, which adopts a prognostic equation for turbulent kinetic energy (Doms et al., 2011b). Moist convection is parameterized according to Tiedtke (1989). A two-stream formulation (Ritter and Geleyn, 1992) is applied for radiative transfer.

Radiative fluxes could be modified by aerosol clouds and tracer gases via absorption, scattering, and emission. The reanalysis data of the German Weather Service DWD derived from the global meteorological model GME (Majewski et al., 2002) were used for initial and boundary conditions. MUSCAT describes the transport, chemical, and removal processes. The gaseous chemistry is represented by RACM-MIM2, which consists of 87 species and more than 200 reactions (Karl et al., 2006; Stockwell et al., 1997). A simplified mass-based approach (similar to the EMEP model; Simpson et al., 2003) is used to represent the aerosol processes with high efficiency. The formation of secondary inorganic particulate matter is through reactions of ammonia with sulfuric acid and nitric acid, which are produced from gaseous precursors SO_2 and NO_x (Hinneburg et al., 2009). The applied particle-gas partitioning depends on temperature and humidity. As in ISORROPIA, under dry and warm conditions the equilibrium shifts toward the gas phase (Nenes et al., 1998). By using the equilibrium approach of Mozurkewich (1993), the partitioning scheme was comparable to Galperin and Sofiev (1998). The extended SORGAM (Schell et al., 2001; Li et al., 2013) is coupled with the massbased aerosol approach to predict the formation of secondary organic aerosol (SOA). Dry deposition is modelled by using the resistance approach following Seinfeld and Pandis (2006) and considering the kinetic viscosity, the atmospheric turbulence state, and the gravitational settling of particles. The resistances for the aerodynamic and quasi-laminar layer are taken from COSMO and analogous to the deposition of water vapour. The parameterization of the wet deposition is dependent upon size-resolved collection efficiency and scavenging (Simpson et al., 2003).

The European anthropogenic emission inventory and the temporal-resolved emission factors are provided by TNO for the AQMEII project (Pouliot et al., 2012; Wolke et al., 2012). The inventory includes the gaseous pollutants (CO, NO_x , SO₂, NH_3 methane, and non-methane volatile organic compounds) and primary emitted particulate matter (PM_{2.5}, PM_{2.5-10}, organic carbon OC, and elemental carbon EC) with a spatial resolution of $0.125^{\circ} \times 0.0625^{\circ}$ (longitude–latitude, about $7 \text{ km} \times 7 \text{ km}$). Note that EC and BC are usually interchangeable in modelling studies (Vignati et al., 2010; Chen et al., 2016a; Nordmann et al., 2014). The emission of NH_3 was reduced by 50 %, since over 90 % of NH₃ emissions in Europe are contributed by agricultural sources (Hertel et al., 2011; Erisman et al., 2008; Reidy et al., 2008), and agriculture emissions of NH₃ are overestimated by $\sim 50 \%$ or even more (Sintermann et al., 2012; Backes et al., 2016). Also, Chen et al. (2016b) adopted the same NH₃ emission inventory in the WRF-Chem model and reported that total NH₃ was overestimated by a factory of ~ 2 at Melpitz during the campaign period. The modelled dust emissions depend on surface wind friction velocities, sur-



Figure 1. Results (domain N2) of the NewN2O5 case of averaged NH_3 mass concentration during 10–25 September 2013. The added emissions of marker tracer (T-NH3) from NH_3 source regions (the Netherlands and southern Germany) are marked by the white bars. The locations of the considered measurement stations are also marked: Neuglobsow, Schmücke, and Zingst are marked by black dots; Melpitz is marked with a red star and its results are discussed in Figs. 2 and 3. The region with the most significant organic coating effect is highlighted by the black polygon and will be analysed together with Fig. 6.

face roughness, soil particle size distribution, and soil moisture (Heinold et al., 2011). Emissions of sea salt depend on salinity and wind speed (Long et al., 2011). Biogenic emissions depend on land use and meteorology in the approach of Steinbrecher et al. (2009) and for "soil NO" in Williams et al. (1992) and Stohl et al. (1996). The Saarikoski et al. (2007) scheme was applied to estimate the biomass burning emissions.

The chemical inactive tracers (T1, T2, and T3) were added into RACM-MIM2 to investigate the chemical fluxes of the selected reactions. T1, T2, and T3 (representing R1, R2, and R3, respectively) were reset to zero every hour in the simulation in order to quantify the chemical fluxes from N₂O₅ to nitrate, avoiding the influence of transport. T1 represents the contribution of N₂O₅ to the chemical formation of nitric acid; T3 and T2 represent the local chemical net formation of particulate nitrate.

$$N_2O_5 + H_2O \xrightarrow{(acrosof)} 2HNO_3 + T1$$
 (R1)

$$NH_3 + HNO_3 \rightarrow NH_4NO_3 + T2$$
 (R2)

$$NH_4NO_3 \rightarrow NH_3 + HNO_3 + T3 \tag{R3}$$

Furthermore, in order to investigate the influence of transport from the NH₃ source regions (the Netherlands and southern Germany) on particulate nitrate, the marker tracer (T-NH3) was emitted into the first layer of regions with high ammonia emissions (white bars in Fig. 1) with an emission rate of $2 \times 10^{-8} \text{ mol m}^{-2} \text{ s}^{-1}$.

2.2 A novel mass-based parameterization of the heterogeneous hydrolysis of N₂O₅

The heterogeneous hydrolysis of N₂O₅ on the surface of deliquescent aerosol is believed to be considerable, since N₂O₅ reacts quite slowly with water vapour (Platt et al., 1984). Many laboratory experiments have shown that $\gamma_{N_2O_5}$ depends principally on particle composition and water content (and so atmospheric RH). Reducing the RH, or adding organics or nitrate to the solutions, suppresses $\gamma_{N_2O_5}$ by an order of magnitude (Chang et al., 2011, and the references therein).

The P2 of Riemer et al. (2003), which was adapted from Chang et al. (1987), was originally incorporated in COSMO–MUSCAT to represent the heterogeneous hydrolysis of N₂O₅ (OldN2O5). Reaction R1 is implemented into chemical transport models as a first-order loss (Riemer et al., 2003). The reaction constant ($k_{N_2O_5}$) is defined as

$$k_{\rm N_2O_5} = \frac{1}{4} \cdot v_{\rm N_2O_5} \cdot S \cdot \gamma_{\rm N_2O_5},\tag{1}$$

where *S* is the surface area concentration of aerosol particles and $v_{N_2O_5}$ is the mean molecular velocity of N₂O₅. Based on the first-order reaction constant, Chang et al. (1987) proposed the following scheme to represent $k_{N_2O_5}$.

$$k_{\rm N_2O_5} = \frac{1}{600 \exp\left(-\left(\frac{\rm RH}{28}\right)^{2.8}\right) + a}$$
(2)

(aaraaal)

RH is the relative humidity in percent, which was used as an indicator for the influence of hygroscopic growth on *S*, $k_{N_2O_5}$ results in min⁻¹, and a = 5 was originally adopted in Chang et al. (1987). According to Riemer et al. (2003), a = 5represents the surface area concentration of 2700 µm² cm⁻³ when RH is higher than 60 %. However, this high surface area concentration can only be expected in areas suffering from severe pollution or where cloud droplets exist. Therefore, we adopted a = 17 in this study as suggested by Riemer et al. (2003). However, the complex considerations of *S* and $\gamma_{N_2O_5}$ are still missing in the OldN2O5. In this study, we propose a sophisticated parameterization to improve the OldN2O5 for mass-based aerosol models, with a full consideration of temperature, RH, aerosol particle compositions, and *S*.

As shown in Eq. (1), $k_{N_2O_5}$ is linearly related to *S* and $\gamma_{N_2O_5}$. We adapted Eq. (2) with factors f_s and $f_{\gamma N_2O_5}$, which represent the impact of *S* and $\gamma_{N_2O_5}$, respectively, as shown in Eq. (3); f_s can be estimated from the particle mass concentration according to Eq. (4) and $f_{\gamma N_2O_5}$ can be estimated from the core-shell model of aerosol particles considering the secondary organic coating effect according to Anttila et al. (2006) and Riemer et al. (2009), as given in Eq. (5). The influence of the O : C ratio on the organic coating effect (Gaston et al., 2014) was not considered here, since the O : C ratio information is not available in COSMO–MUSCAT. Also, the "high" O : C regime defined in Gaston et al. (2014) may not represent the north-western European case compared with airborne measurements (Morgan et al., 2015).

$$k_{\rm N_2O_5} = \frac{1}{600 \exp\left(-\left(\frac{\rm RH}{28}\right)^{2.8}\right) + 17} \cdot f_s \cdot f_{\gamma_{\rm N_2O_5}} \tag{3}$$

$$f_s = (SA_{fine} \cdot PM_{fine} + SA_{coarse} \cdot PM_{coarse}) / S_{ref}$$
(4)

 SA_{fine} and SA_{coarse} represent the specific surface area for fine- and coarse-mode particles in $m^2 g^{-1}$, and PM_{fine} and PM_{coarse} represent the mass concentration of fine- and coarse-mode particles in $\mu g m^{-3}$. A value $11 m^2 g^{-1}$ was used for SA_{fine}, considering recently reported values of 11.9 and $10.2 \text{ m}^2 \text{ g}^{-1}$ from laboratory studies (Okuda, 2013) and measurements in Japanese urban regions (Hatoya et al., 2016). A value of $1.2 \text{ m}^2 \text{ g}^{-1}$ was used for SA_{coarse} (Okuda, 2013). Sref is the reference particle surface area concentration; here, we suggest $S_{\text{ref}} = 600 \,\mu\text{m}^2 \,\text{cm}^{-3}$. Equation (2) will provide a result that is very close to a complex parameterization with a particle surface area concentration of $600 \,\mu\text{m}^2 \,\text{cm}^{-3}$ (Riemer et al., 2003) when a = 17 and RH>60%. Note that a small initial overestimation of particulate nitrate may result in a significant overprediction of nitrate through the integration in models due to a feedback in this scheme. That is, higher nitrate concentrations result in a larger f_s and promise a higher $k_{N_2O_5}$, leading to a higher production of nitrate. In order to avoid the uncertainty of this feedback mechanism and to calculate a reasonable $k_{N_2O_5}$ in this case study, the nitrate mass concentration in Eq. (4) is considered as 1.3 times the sulfate mass concentration based on filter measurements during the HOPE Melpitz campaign.

$$f_{\gamma N_2 O_5} = \left(\gamma_{\text{core}}^{-1} + \gamma_{\text{coating}}^{-1}\right)^{-1} / \gamma_{\text{ref}},\tag{5}$$

where γ_{core} is the N₂O₅ reaction probability with the core of the particle, which can be estimated by Table 1; $\gamma_{coating}$ is the N₂O₅ reaction probability with the secondary organic coating shell of the particle, which can be estimated by Eq. (6) according to Anttila et al. (2006) and Riemer et al. (2009); γ_{ref} is the reference reaction probability. Here, we suggest $\gamma_{ref} = 0.1$, since Eq. (2) is developed on the basis of the assumption $\gamma_{N_2O_5} = 0.1$ (Riemer et al., 2003).

$$\gamma_{\text{coating}} = \frac{4RTH_{N_2O_5}D_{N_2O_5}R_{\text{core}}}{v_{N_2O_5}l_{\text{shell}}R_{\text{particle}}},$$
(6)

where *T* is the temperature, $D_{N_2O_5}$ is the diffusion coefficient of N₂O₅ in the organic coating, $H_{N_2O_5}$ is the Henry's Law constant of N₂O₅ for the organic coating shell, *R* is the universal gas constant, R_{particle} is the radius of the particle, R_{core} is the radius of the core, and l_{shell} is the thickness of the organic coating shell.

The γ_{core} values can be estimated from previous laboratory experiments (Table 1) of inorganic and primary organic compositions (Davis et al., 2008; Evans and Jacob, 2005, and references therein;). Davis et al. (2008) developed a parameterization for N₂O₅ hydrolysis on ammonium–sulfate–nitrate particles with respect to RH and temperature. Evans and Jacob (2005) provided the parameterizations for N₂O₅ hydrolysis on primary organic particles (Thornton et al., 2003), black carbon (Sander et al., 2003), sea salt (Sander et al., 2003), and dust (Bauer et al., 2004); γ_{core} can be derived by a mass-weighted average (Riemer et al., 2003) of each single-component parameterization (Table 1).

2.3 Estimation of reaction probabilities with a sectional aerosol model

The Weather Research and Forecasting–Chemistry model (WRF-Chem V3.5.1) is a fully online coupled regional air quality model. Chang et al. (2016) incorporated several parameterizations for N_2O_5 hydrolysis into a sectional aerosol treatment (MOSAIC; Zaveri et al., 2008) in WRF-Chem. The "Davis" approach from Chang et al. (2016), hereinafter referred to as Ch&Davis, was chosen to be compared with NewN2O5. The reasons for this choice will be discussed in detail in Sect. 3.1.

In order to validate the mass-based NewN2O5 with the sectional-based Ch&Davis, we performed WRF-Chem simulations during the HOPE Melpitz campaign. The same WRF-Chem results were adopted for offline estimating $k_{N_2O_5}$ according to NewN2O5 and Ch&Davis. We followed the physics-relating configuration according to Chen et al. (2016a), which reproduced meteorological conditions

Particle type	Parameterization		Parameters	References
r article type	i aranicunzation		i arameters	and remarks
Core of particle	$\gamma_{\text{core}} = \sum_{i} \gamma_i \cdot \text{ratio}_i$		ratio _i = $\frac{[m_i]}{[m_{core}]}$ [m_{core}] = $\sum_{i}^{[m_i]}$ <i>i</i> : the following particle types <i>i</i> = [ASN, OC, SSA, dust, BC]	γ : reaction probability m: mass [mass]: mass concentration
ASN: (A) Ammonium (S) Sulfate (N) Nitrate	$\gamma_{ASN} = \gamma_{AB}^* \cdot x_{AB} + \gamma_{aq/}^*$ $\gamma_{AB}^* = \min(\gamma_{AB}, 0.08585)$ $\gamma_{aq,AS}^* = \min(\gamma_{aq,AS}, 0.0)$ $\gamma_{aq,AS} = \frac{1}{1 + e^{-\lambda_{aq,AS}}}$ $\gamma_{d,AS}^* = \min(\gamma_{d,AS}, 0.012)$ $\gamma_{AN}^* = \min(\gamma_{AN}, 0.0154)$ $\lambda_{AB} = \beta_{10} + \beta_{11} \cdot RH + \mu$ $\lambda_{aq,AS} = (\beta_{10} + \beta_{20}) + \beta$ $\lambda_{d,AS} = \beta_{d0} + \beta_{d1} \cdot RH + \mu$ $\lambda_{d,AN} = \beta_{30} + \beta_{31} \cdot RH$	$d_{AS} \cdot x_{AS} + \gamma_{AN}^* \cdot x_{AN}$ $i), \gamma_{AB} = \frac{1}{1 + e^{-\lambda_{AB}}}$ i53), $24), \gamma_{d,AS} = \frac{1}{1 + e^{-\lambda_{d,AS}}}$ $, \gamma_{AN} = \frac{1}{1 + e^{-\lambda_{AN}}}$ $\beta_{12} \cdot T_{291}$ $11 \cdot \text{RH} + (\beta_{12} + \beta_{22}) \cdot T_{291}$ $-\beta_{d2} \cdot T_{293}$	$\begin{aligned} x_{AB} &= 1 - (x_{AS} + x_{AN}) \\ x_{AS} &= \max \left(0, \min \left(1 - x_{AN}, \frac{[A]}{[N] + [S]} - 1 \right) \right) \\ x_{AN} &= \frac{[N]}{[N] + [S]} \\ \beta_{10} &= -4.10612 \\ \beta_{11} &= 0.02386 \\ \beta_{12} &= -0.23771 \\ \beta_{20} &= -0.80570 \\ \beta_{22} &= 0.10225 \\ \beta_{30} &= -8.10774 \\ \beta_{31} &= 0.04902 \\ \beta_{d0} &= -6.13376 \\ \beta_{d1} &= 0.03592 \\ \beta_{d2} &= -0.19688 \\ T_{291} &= \max(T - 291, 0) \\ T_{293} &= \max(T - 293, 0) \end{aligned}$	Davis et al. (2008) AB: ammonium bisulfate AS: ammonium sulfate AN: ammonium nitrate A: NH_4^+ S: SO_4^{2-} N: NO_3^- Unit of RH: % Unit of T: K aq: aqueous phase d: dry phase (crystallized) AS crystallizes when RH < 32.8 % and forms a solid phase (Martine et al., 2003)
Organic carbon (Primary OC)	$\gamma_{\rm OC} = \rm RH \times 5.2 \times 10^{-4}$ $\gamma_{\rm OC} = 0.03$	RH < 57% RH ≥ 57%		Evans and Jacob (2005) Thornton et al. (2003)
Sea salt aerosol (SSA)	$\gamma_{\rm SSA} = 0.005$ $\gamma_{\rm SSA} = 0.03$	$\begin{array}{l} RH < 62\% \\ RH \geq 62\% \end{array}$		Evans and Jacob (2005) Sander et al. (2003)
Dust	$\gamma_{\text{Dust}} = 0.01$			Evans and Jacob (2005) Bauer et al. (2004)
Black carbon (BC)	$\gamma_{\rm BC} = 0.005$			Sander et al. (2003)

Table 1. Representation of the reaction probability of aerosol particle core (γ_{core}) for N₂O₅ hydrolysis.

during the HOPE Melpitz campaign well. The sea salt emission (Gong, 2003) was reduced by a factor of 20 in WRF-Chem, considering that Gong (2003) may highly overestimate sea salt emission (Neumann et al., 2016), and thus leads to an overestimation of sea salt by a factor of 20 during the HOPE campaign at Melpitz (Chen et al., 2016b). The configuration of chemical and aerosol treatments followed Chang et al. (2016). The CBMZ (Zaveri and Peter, 1999) mechanism was used to describe gas-phase reactions. MOSAIC (Zaveri et al., 2008) with eight size bins was chosen to represent aerosol properties. Three nested domains (Fig. S1 in the Supplement) with 39 vertical layers were set up for the simulated case, with a resolution of 54, 18, and 6 km, respectively.

In Ch&Davis aerosol liquid water is considered when calculating the particle surface area for each size bin. Details of the sectional-based method for estimating *S* in the Ch&Davis scheme are given by Chang et al. (2016). In the NewN2O5 scheme, the first six bins (with diameter in the range of 40 nm–2.5 µm) are counted as fine mode, and the last two bins (2.5–10 µm) are counted as coarse mode. This definition is identical to COSMO–MUSCAT. In order to be consistent with COSMO–MUSCAT, the organic coating effect is considered for fine particles in NewN2O5, since the maximum effective particle diameter of the Anttila06 scheme is $2 \mu m$ (Anttila et al., 2006). In order to quantify the uncertainty stemming from the different *S* treatments between NewN2O5 (mass based) and Ch&Davis (sectional based), an estimation result according to an adapted NewN2O5 (with sectional-based *S*) will also be discussed in Sect. 3.1.

2.4 Observations

The filter chemical composition measurements of the HOPE Melpitz campaign (10–25 September 2013) and at three UBA stations (Neuglobsow, Schmücke, and Zingst; http: //www.umweltbundesamt.de) were used to validate the modelled results. The observations at the TROPOS research station at Melpitz represent the regional background of central Europe (Spindler et al., 2010, 2012; Brüggemann and Spindler, 1999; Poulain et al., 2011; Birmili et al., 2001). During the HOPE Melpitz campaign, high-volume samplers (DIGITEL DHA-80; Walter Riemer Messtechnik, Germany) with a sampling flow of $\sim 30 \,\mathrm{m^3 \, h^{-1}}$ were used to collect 24 h daily filter samples with 10 µm cut-off inlets. Additionally, 24 h filter sampler measurements with a PM₁₀ inlet at three UBA stations in Germany were collected every third day. The filter material is quartz fibre (Munktell, Grycksbo, Sweden; type MK 360), which allows for the determination of particle mass, water-soluble ions $(SO_4^{2-}, NO_3^{-}, NH_4^{+}, Cl^{-},$ Na⁺, K⁺, Mg²⁺, and Ca²⁺), OC, and EC from one filter. The filters were preheated before sampling for at least 24 h at 105 °C to minimize the blank values of OC. More details about filter measurement are given in Spindler et al. (2013). Near-ground meteorological parameters (e.g. relative humidity, temperature, wind speed, wind direction) were simultaneously measured at Melpitz. More details about the HOPE campaign are given in Macke et al. (2017).

3 Results and discussion

The COSMO-MUSCAT model performance was examined by comparing simulated meteorological fields with the Melpitz near-ground measurements (Fig. 2). Generally, the meteorological conditions during the HOPE Melpitz campaign were well captured by the model, with correlation coefficients (R) of 0.87, 0.85, 0.73, and 0.85 for temperature, RH, 10 m wind speed, and wind direction, respectively. The factors between modelled results and the meteorological measurements were ~ 1 , except for an overestimation of wind speed with a factor of 1.44, possibly due to the vertical resolution of the model. Nevertheless, the temperature and RH, which are the most important meteorological parameters in this study for N₂O₅ heterogeneous hydrolysis during nighttime, were in good agreement with the measurement. Although model simulations slightly underestimated RH during the night-time of 17 and 22 September (Fig. 2b), modelled RH was still higher than 80 % where $k_{N_2O_5}$ is insensitive to RH as shown in Table 1 and Riemer et al. (2003). Therefore, this bias of RH will not lead to a significant uncertainty in nitrate simulation. However, the overestimation of wind speed may favour the transport of ammonia from western Europe (e.g. the Netherlands). This could be a possible reason for the nitrate overprediction in the NewN2O5 case (Fig. 3d), especially during 20–24 September when a western wind was constantly dominant (Fig. 2d).

3.1 Evaluating closure for mass-based NewN2O5 and a sectional approach

In order to confirm that the mass-based NewN2O5 estimates $k_{N_2O_5}$ with a reliable accuracy, we evaluated closure between NewN2O5 and a sectional-based state-of-the-art parameterization (Chang et al., 2016) based on the WRF-Chem (MOSAIC) results. Chang et al. (2016) reported that the Davis + coat (Daivs08 + Anttila06) approach produced the best agreement of $\gamma_{N_2O_5}$ with aircraft observations during the CalNex 2010 campaign, with overestimation by a factor mostly within in a range of 2-8 (Fig. S2b). Without considering OC coating effect (Davis08 only), Ch&Davis still showed a relatively good linear relationship with the observed $\gamma_{N_2O_5}$, which was, however, overestimated with a higher factor ranging about 3-10 (Fig. S2a). Considering the different treatments of OC coating between NewN2O5 (SOA coating only) and Chang et al. (2016) (Davis + coat, primary OC, and SOA), NewN2O5 was validated using the Ch&Davis scheme. This would not significantly influence the comparison results, since the HOPE Melpitz campaign was an OC-low case, with only a \sim 7 % contribution from total OC mass based on filter measurements at Melpitz. Therefore, not much SOA was available for the coating effect in contrast to the OC-high case (contributed about 50-80% to total mass; Fig. 9 in Chang et al., 2016) in the CalNex 2010 campaign. The coating effect exerted a negligible influence at Melpitz, and this point will be discussed in detail in Sect. 3.4. We validated the NewN2O5 scheme by comparing $k_{N_2O_5}$ instead of $\gamma_{N_2O_5}$ because NewN2O5 scheme was developed on the basis of a parameterization to directly calculate $k_{N_2O_5}$ as proposed by Chang et al. (1987) and Riemer et al (2003).

As shown in Fig. 4, the $k_{N_2O_5}$ showed a good linear relationship (R = 0.91) between NewN2O5 and Ch&Davis that is much better than using the OldN2O5 (Fig. 4). Mass-based NewN2O5 estimated lower $k_{N_2O_5}$ than the sectional-based Ch&Davis by a factor of ~ 8. However, Ch&Davis may overestimate the $\gamma_{N_2O_5}$ by a factor of 3–10 (Chang et al, 2016, see also Fig. S2a). Assuming that *S* was correctly given by the WRF-Chem sectional aerosol module, we can expect that Ch&Davis may overestimate $k_{N_2O_5}$ by a factor of 3–10 according to Eq. (1). Therefore, NewN2O5 may provide a $k_{N_2O_5}$ in the range of 0.36–1.2 times the realistic one.

Two important uncertainties need to be kept in mind in this validation. First, the estimation of S is very challenging due to the uncertainties of, for example, particle num-



Figure 2. Comparison between modelled and measured meteorological conditions. (a) Temperature (T); (b) relative humidity (RH); (c) wind speed; (d) wind direction.

ber and mass size distribution, partitioning processes, and secondary formation. In addition, the hygroscopic growth of particles can also be an important source of the uncertainty of S due to the challenge in the estimation of particle liquid water, especially at low RH, even by a complex aerosol treatment (Chang et al., 2016). About a 30 % difference in $k_{N_2O_5}$ between NewN2O5 and Ch&Davis stems from the different treatments of S. As shown in Fig. 4, the factor between NewN2O5 and Ch&Davis was reduced from ~ 8.3 to ~ 5.9 with a slightly increase in R when we adopted the sectional-based S (same as Ch&Davis) in NewN2O5. Second, Ch&Davis was validated by aircraft measurements in an OC-high case during the CalNex 2010 campaign. Therefore, the overestimation factor of Ch&Davis may not be as high as expected in an OC-low case during the HOPE Melpitz campaign. However, SSA, BC, and dust should exert an influence in an OC-low case and should also be considered in a parameterization, as we did in NewN2O5. This can also be a reason for the difference between Ch&Davis and NewN2O5.

3.2 Improvement of the particulate nitrate prediction

In previous evaluation studies (Im et al., 2015; Wolke et al., 2012), the COSMO–MUSCAT model predicted particulate nitrate mass concentrations ($[NO_3^-]$) in fair agreement with the measurements, with an overestimation in the range of 50% on long-term average. This is comparable with other models (Im et al., 2015). However, short periods with strong overestimations of $[NO_3^-]$ were also observed in previous

Table 2. Sensitivity simulation cases.

Case	N ₂ O ₅ parameterization	NH ₃ emission
OldN2O5-FullNH3	P2 of Riemer et al. (2003)	100 %
OldN2O5	P2 of Riemer et al. (2003)	50 % *
NewN2O5	New scheme (this study)	50 % *

* Suggested by Sintermann et al. (2012), Backes et al. (2016), and Chen et al. (2016b).

studies. This seems to be the case for the HOPE Melpitz campaign simulation, during which COSMO–MUSCAT highly overpredicted $[NO_3^-]$ over Germany in this study (Fig. 3), and WRF-Chem in a previous study (Chen et al., 2016b). In order to evaluate the improvement of the NewN2O5 scheme and quantify the influence of NH₃ emission overestimation on the particulate nitrate prediction, three sensitivity simulations were conducted (Table 2).

In this HOPE Melpitz campaign case, the particulate nitrate mass concentrations were overestimated by factors of 23.7, 12, 6.5, and 6.3 for Neuglobsow, Schmücke, Zingst, and Melpitz, respectively (Fig. 3). The modelled NO_x was in line with the observed concentration level at Melpitz and should not be the reason for the overprediction of particulate nitrate (see details in the Supplement Sect. S1 and Fig. S3). Nevertheless, the overestimation of NH₃ emissions might contribute about 20–30 % of the particulate nitrate overprediction compared between the OldN2O5-FullNH3 and



Figure 3. Comparison of particulate nitrate mass concentration between filter measurements and modelled results: (a) Neuglobsow; (b) Schmuecke; (c) Zingst; (d) Melpitz. Modelled concentrations at Melpitz: (e) N_2O_5 ; (f) marker species T1 for chemical reaction R1; (g) marker species for the chemical formation of particulate nitrate (T3-T2); (h) the NH₃ marker tracer (T-NH3) for transport from the Netherlands and southern Germany. The light-red colour bars indicate the results of the OldN2O5-FullNH3 case, the red colour bars indicate the results of the OldN2O5 case, and the blue colour bars indicate the results of the NewN2O5 case. The shaded periods indicate the dominating processes for high concentrations of particulate nitrate: chemical formation (red), transport (blue), and boundary conditions (grey).

OldN2O5 cases. This is in line with previous studies (Renner and Wolke, 2010; Backes et al., 2016). However, even with a 50 % reduction of NH_3 emissions, the particulate nitrate was still highly overestimated over Germany with factors of about 19, 9, 4.5, and 5 for these four stations, respectively. The NewN2O5 scheme would further moderate the

overprediction by another $\sim 35 \%$ (Fig. 3). Correspondingly, the overestimation factors of particulate nitrate were reduced to about 10.7, 6, 2.5, and 3 for the four stations, respectively. The N₂O₅ was almost completely consumed by the heterogeneous reaction at Melpitz in the OldN2O5 case, but not in the NewN2O5 case (Fig. 3e). This is due to a significant de-



Figure 4. Comparison between the sectional-based Ch&Davis ("Davis" of Chang et al., 2016) and mass-based NewN2O5 (mass based, black), NewN2O5 (with a sectional-based particle surface area, blue), and OldN2O5 (red). The results are offline calculated on the basis of the WRF-Chem simulation with a sectional aerosol treatment (MOSAIC).

crease (by an average of more than a factor of 20, see Fig. 4) in the reaction constant of the heterogeneous hydrolysis of N₂O₅ by NewN2O5. However, there must be other reasons that might explain the remaining overestimations in the simulated particulate nitrate mass concentrations. One possible reason can be the underprediction of the coating organic matter budget in the model leading to an overestimation of $\gamma_{N_2O_5}$ (Chang et al., 2016); other possible reasons should be investigated in future studies, such as the deposition process, long-range transport, the formation of nitrogen-containing OC, and neutralization processes.

The improvement of particulate nitrate prediction with NewN2O5 can be more clearly shown associated with the tracers (T1 in Fig. 3f, T3-T2 in Fig. 3g, and T-NH3 in Fig. 3h) and the comparison with Melpitz measurements (Fig. 3d), which were sampled on the filter every day and analysed offline. The overestimation of [NO₃] on 10-11 September (grey shaded period in Fig. 3) stemmed from the uncertainty of boundary conditions in the model. As shown in Fig. S4, an air mass with high $[NO_3^-]$ was transported from the south-western boundary area to Melpitz. The $[NO_3^-]$ at Melpitz was dominated by transport from the Netherlands and southern Germany on 13-14 and 19-24 September (blue shaded period in Fig. 3), as indicated by the high T-NH3 concentration (Fig. 3h) and the negligible local chemical formations (Fig. 3g). In contrast, local chemical formations dominated the $[NO_3^-]$ on 12, 17– 18, and 25 September (red shaded period in Fig. 3). During the red shaded period, T-NH3 was almost zero (Fig. 3h) and the modelled wind speed was less than 4 m s^{-1} on average (Fig. 2c). A much stronger reduction in the overestimation of particulate nitrate occurred during the red shaded period (a factor of ~ 1.4 on average), which was dominated by local chemical formations. This further confirmed the improvement of the heterogeneous hydrolysis of N₂O₅ by NewN2O5. During 15–16 September (without the shaded period in Fig. 3), the contributions from both transport and local chemical formations of particulate nitrate were very limited (Fig. 3f–h), resulting in a very low [NO₃⁻].

3.3 Comparison between NewN2O5 and OldN2O5

The NewN2O5 case improved the particulate nitrate overestimation problem compared with OldN2O5. Meanwhile, the spatial distribution pattern of [NO₃⁻] was similar between these two cases (Fig. 5). Here, we focus on the nighttime period of the HOPE Melpitz campaign, since the N₂O₅ heterogeneous reaction is much more significant during the night than in the daytime. The lowest $[NO_3^-]$ was found over Poland and ocean regions during night-time, and $[NO_3^-]$ was lower than 4 and $3 \mu g m^{-3}$ in the OldN2O5 and NewN2O5 cases, respectively. Moderate [NO₃⁻] was found over central Europe (Melpitz and the surrounding region) at about 6–8 and 4–5.5 μ g m⁻³ in the OldN2O5 and NewN2O5 cases, respectively. The highest $[NO_3^-]$ occurs over the region of the Netherlands and nearby regions at about 9-12 and 6- $8 \,\mu g \,m^{-3}$ in the OldN2O5 and NewN2O5 cases, respectively, due to the high agriculture emission of NH₃ in this region. There was also a remarkably high amount of particulate nitrate over southern Germany of about 8–10 and 5–6.5 μ g m⁻³ in the OldN2O5 and NewN2O5 cases, respectively. In general, the $[NO_3^-]$ was reduced by ~ 35 % over the entire N2 domain (Fig. 5). The most significant reduction of $[NO_3^-]$ is found over the Netherlands and southern Germany where the highest $[NO_3^-]$ (reduced by about 3–4.5 µg m⁻³) was found, followed by the region near Melpitz (reduced by about 2- $3 \,\mu g \, m^{-3}$; Fig. 5c). This is caused by a significant reduction (by more than a factor of 20; see Fig. 4) of $k_{N_2O_5}$, which results from the consideration of particle mass concentration influence on S and comprehensive treatments for $\gamma_{N_2O_5}$. Therefore, the regions with high $[NO_3^-]$ during night-time indicate considerable nitrate formation from the heterogeneous hydrolysis of N₂O₅, during which [NO₃⁻] was reduced by about 3–4.5 μ g m⁻³ (~ 35 %; see Fig. 5) in the new scheme. However, this heterogeneous hydrolysis was negligible over the regions where $[NO_3^-]$ was low during night-time and did not have a relevant contribution to the formation of particulate nitrate. Hence, the improvement of particulate nitrate prediction by NewN2O5 was more significant over the high- $[NO_3^-]$ regions than the low- $[NO_3^-]$ regions.





Figure 5. Horizontal distribution of averaged modelled particulate nitrate mass concentration during night-time on 10–25 September. (a) OldN2O5 case; (b) NewN2O5 case; (c) difference between OldN2O5 and NewN2O5 cases.

3.4 Influence of organic coating on N₂O₅ heterogeneous hydrolysis

The secondary organic coating on particle surfaces may significantly decrease the reaction probability of N2O5 and influence the particulate nitrate concentration. Riemer et al. (2009) reported that organic coating could decrease $[NO_3^-]$ by up to 90 % when both N₂O₅ and secondary organic compounds were built up. The highest reduction over Europe was found over the Netherlands followed by western Germany (both covered by the domain N2) in their study. In addition to N2O5 and secondary organic compounds, the meteorological conditions (e.g. RH and temperature) may also exert an influence on the organic coating effect. In this study, we introduced a parameterization (NewN2O5) for the heterogeneous hydrolysis of N₂O₅ considering meteorological conditions. The influence of the organic coating suppression effect on particulate nitrate prediction was investigated by a comparison between NewN2O5 with and without the SOA coating effect.

At night-time, much higher N_2O_5 concentrations occurred and heterogeneous hydrolysis is more important than during daytime (Jacob, 2000). As shown in Fig. 6a and b, the influence of the organic coating effect was negligible over the domain N2 including the Netherlands and Germany. Even on 24 September at 23:00 CET when changes were most significant, organic coating reduced $[NO_3^-]$ only by about $2-4 \,\mu g \, m^{-3}$ (less than 10–20%) over the black-polygonmarked and nearby regions (Fig. S5). Meanwhile, for nighttime averages during the campaign, organic coating could only reduce $[NO_3^-]$ by less than 0.1 µg m⁻³ or 2 % over the whole domain (Fig. 6). This is because appropriate meteorological conditions, as described in the following, are needed in NewN2O5 for a significant organic coating. In addition to the simultaneous build-up of SOA and N2O5 (Riemer et al., 2009), high NH₃ concentrations and $\gamma_{N_2O_5}$ are also indispensable conditions for a significant organic coating effect. High NH₃ concentrations are necessary for neutralizing the HNO₃ from the heterogeneous hydrolysis of N₂O₅ during the night. High $\gamma_{N_2O_5}$ causes a significant reduction of $\gamma_{N_2O_5}$ by organic coating (Chang et al., 2016; Riemer et al., 2009). Therefore, a large impact should be expected in regions with high RH and low temperature and hence a high $\gamma_{N_2O_5}$. As show in Fig. 6, the most significant organic coating effect (still less than 2% influence on $[NO_3^-]$) could be found over the Netherlands and nearby regions (black polygon). Over this area, these five conditions were fulfilled to some extent: (1) temperature was 13.5–14.5 °C; (2) RH was higher than 75 %; (3) SOA concentration was $\sim 1.6 \,\mu g \, m^{-3}$; (4) N₂O₅ concentration was about $0.4-0.6 \,\mu g \,m^{-3}$; and (5) NH₃ concentration was about $4-6 \,\mu g \, m^{-3}$ (Fig. 1). There was almost no influence of organic coating over the other regions (Fig. 6a and b). These five conditions (not very high temperature; relatively high RH; build-up of SOA, N₂O₅, and NH₃) could



Figure 6. Horizontal distribution of averaged model results during night-time on 10–25 September computed with NewN2O5. (a) Difference in particulate nitrate mass concentration between model cases with and without considering the organic coating effect; (b) difference in particulate nitrate mass concentration in percent between model cases with and without considering the organic coating effect; (c) temperature; (d) RH; (e) SOA mass concentration; (f) N₂O₅ mass concentration. The region with the most significant organic coating effect is highlighted by the black polygon.

not be simultaneously fulfilled over western and central Europe, and therefore the organic coating effect was not very significant.

4 Conclusions

Generally, the COSMO–MUSCAT model predicted particulate nitrate in a reasonable range on long-term average. The results were comparable with other models in previous studies. However during the HOPE Melpitz campaign (10–25 September 2013), particulate nitrate was significantly overestimated by the COSMO–MUSCAT model over Germany, despite a good performance of meteorological simulation. This can be partly ($\sim 35 \%$) attributed to the parameterization of the heterogeneous hydrolysis of N₂O₅ (OldN2O5). A sophisticated mass-based parameterization of the heterogeneous hydrolysis of N₂O₅ (NewN2O5) was proposed in this study, aiming at improving the particulate nitrate prediction in atmospheric modelling. This mass-based NewN2O5 was validated with a state-of-the-art parameterization (Chang et al., 2016), which is based on a sectional aerosol treatment. The validation results showed a good linear relationship (R = 0.91) and indicated that NewN2O5 could estimate the reaction probability of N₂O₅ in a reasonable range, within about 0.36–1.2 times the realistic one.

In order to quantify the improvement of the nitrate prediction by using NewN2O5, sensitivity studies were performed based on more realistic NH₃ emissions, which are reduced by 50%. This correction was based on previous studies that showed NH₃ emissions were overestimated by a factor of \sim 2. The overestimation of NH₃ emissions led to about 20– 30 % overprediction of particulate nitrate over Germany. The horizontal distribution patterns of particulate nitrate were in good agreement between the OldN2O5 and NewN2O5 cases. The OldN2O5 case overestimated particulate nitrate by a factor of 19, 9, 4.5, and 5 for Neuglobsow, Schmücke, Zingst, and Melpitz, respectively. This may be caused by a lacking consideration of particle surface area (S) and complex treatments of $\gamma_{N_2O_5}$. Based on many previous laboratory experiments, the influences of temperature, RH, aerosol particle compositions, and surface area concentration on the heterogeneous reaction constant of N2O5 were considered in NewN2O5. The reaction constant was on average reduced by a factor of more than 20 in NewN2O5. Correspondingly, the overestimation of particulate nitrate was reduced by \sim 35 % for the whole period. Particularly, NewN2O5 significantly improved particulate nitrate prediction, with a factor of ~ 1.4 compared with the filter measurements, when particulate nitrate was dominated by local chemical formations at Melpitz (12, 17–18, and 25 September).

In this study, we additionally investigated how the decrease in $\gamma_{N_2O_5}$ due to organic coating (Anttila et al., 2006) influences the particulate nitrate prediction over western and central Europe. Based on NewN2O5, the simulation results with and without organic coating were analysed. Our results showed a negligible (less than 2% or $0.1 \,\mu g \, m^{-3}$) influence of coating on particulate nitrate over the Netherlands and Germany. Since, in addition to the considerable amounts of N_2O_5 , SOA, and NH₃ that must be present at the same location, appropriate meteorological conditions (relatively high RH and low temperature) are also indispensable for organic coating to exert an impact. This is because low RH and high temperature would lead to a low $\gamma_{N_2O_5}$ value, and therefore no significant organic coating suppression on $\gamma_{N_2O_5}$ would be observed. These conditions were rarely fulfilled simultaneously over western and central Europe; hence, the influence of the organic coating effect on particulate nitrate prediction was negligible in this study.

This study suggests that temperature, RH, particle compositions, and surface area concentration should be taken into account in the parameterization of the heterogeneous hydrolysis of N₂O₅. A sophisticated parameterization is proposed for the mass-based aerosol models. It should be included in model simulations to improve the representativeness of N₂O₅ hydrolysis in the ambient atmosphere. The results also indicate that the organic coating effect on suppressing the heterogeneous hydrolysis of N_2O_5 may not be as significant as expected over Europe.

Data availability. All data needed to evaluate the conclusions in the paper are presented in the paper and/or the Supplement. Additional data related to this paper may be requested from the corresponding authors: wolke@tropos.de and chen@tropos.de.

The Supplement related to this article is available online at https://doi.org/10.5194/acp-18-673-2018-supplement.

Competing interests. The authors declare that they have no conflict of interest.

Special issue statement. This article is part of the special issue "HD(CP)2 Observational Prototype Experiment (ACP/AMT interjournal SI)". It is not associated with a conference.

Acknowledgements. The HOPE campaign was funded by the German Research Ministry under project number 01LK1212 C. We would like to thank TNO (the Netherlands) and the AQMEII project (http://aqmeii.jrc.ec.europa.eu/aqmeii2.htm) for providing the European anthropogenic emission inventory and the German Federal Environmental Agency (UBA) for providing the filter measurements of particle compositions. Furthermore, JSC Jülich and DWD Offenbach supported the work by providing computing time and meteorological data. We would also like to thank Ulrich Pöschl, Hartmut Herrmann, and Andreas Tilgner for the discussion and help.

Edited by: Herman Russchenberg Reviewed by: two anonymous referees

References

- Anttila, T., Kiendler-Scharr, A., Tillmann, R., and Mentel, T. F.: On the Reactive Uptake of Gaseous Compounds by Organic-Coated Aqueous Aerosols:? Theoretical Analysis and Application to the Heterogeneous Hydrolysis of N₂O₅, J. Phys. Chem. A, 110, 10435–10443, https://doi.org/10.1021/jp062403c, 2006.
- Archer-Nicholls, S., Lowe, D., Utembe, S., Allan, J., Zaveri, R. A., Fast, J. D., Hodnebrog, Ø., Denier van der Gon, H., and McFiggans, G.: Gaseous chemistry and aerosol mechanism developments for version 3.5.1 of the online regional model, WRF-Chem, Geosci. Model Dev., 7, 2557–2579, https://doi.org/10.5194/gmd-7-2557-2014, 2014.
- Backes, A. M., Aulinger, A., Bieser, J., Matthias, V., and Quante, M.: Ammonia emissions in Europe, part II: How ammonia emission abatement strategies affect secondary aerosols, Atmos. Environ., 126, 153–161, https://doi.org/10.1016/j.atmosenv.2015.11.039, 2016.

- Badger, C. L., Griffiths, P. T., George, I., Abbatt, J. P. D., and Cox, R. A.: Reactive Uptake of N₂O₅ by Aerosol Particles Containing Mixtures of Humic Acid and Ammonium Sulfate, J. Phys. Chem. A, 110, 6986–6994, https://doi.org/10.1021/jp0562678, 2006.
- Baldauf, M., Seifert, A., Förstner, J., Majewski, D., Raschendorfer, M., and Reinhardt, T.: Operational Convective-Scale Numerical Weather Prediction with the COSMO Model: Description and Sensitivities, Mon. Weather Rev., 139, 3887–3905, https://doi.org/10.1175/mwr-d-10-05013.1, 2011.
- Bauer, S. E., Balkanski, Y., Schulz, M., Hauglustaine, D. A., and Dentener, F.: Global modeling of heterogeneous chemistry on mineral aerosol surfaces: Influence on tropospheric ozone chemistry and comparison to observations, J. Geophys. Res.-Atmos., 109, D02304, https://doi.org/10.1029/2003JD003868, 2004.
- Behnke, W., George, C., Scheer, V., and Zetzsch, C.: Production and decay of ClNO₂ from the reaction of gaseous N₂O₅ with NaCl solution: Bulk and aerosol experiments, J. Geophys. Res.-Atmos., 102, 3795–3804, https://doi.org/10.1029/96JD03057, 1997.
- Bellouin, N., Rae, J., Jones, A., Johnson, C., Haywood, J., and Boucher, O.: Aerosol forcing in the Climate Model Intercomparison Project (CMIP5) simulations by HadGEM2-ES and the role of ammonium nitrate, J. Geophys. Res.-Atmos., 116, D20206, https://doi.org/10.1029/2011JD016074, 2011.
- Bertram, T. H. and Thornton, J. A.: Toward a general parameterization of N_2O_5 reactivity on aqueous particles: the competing effects of particle liquid water, nitrate and chloride, Atmos. Chem. Phys., 9, 8351–8363, https://doi.org/10.5194/acp-9-8351-2009, 2009.
- Birmili, W., Wiedensohler, A., Heintzenberg, J., and Lehmann, K.: Atmospheric particle number size distribution in central Europe: Statistical relations to air masses and meteorology, J. Geophys. Res., 106, 32005–32018, https://doi.org/10.1029/2000JD000220, 2001.
- Brown, S. S. and Stutz, J.: Nighttime radical observations and chemistry, Chem. Soc. Rev., 41, 6405–6447, https://doi.org/10.1039/c2cs35181a, 2012.
- Brown, S. S., Dibb, J. E., Stark, H., Aldener, M., Vozella, M., Whitlow, S., Williams, E. J., Lerner, B. M., Jakoubek, R., Middlebrook, A. M., DeGouw, J. A., Warneke, C., Goldan, P. D., Kuster, W. C., Angevine, W. M., Sueper, D. T., Quinn, P. K., Bates, T. S., Meagher, J. F., Fehsenfeld, F. C., and Ravishankara, A. R.: Nighttime removal of NO_x in the summer marine boundary layer, Geophys. Res. Lett., 31, L07108, https://doi.org/10.1029/2004GL019412, 2004.
- Brüggemann, E. and Spindler, G.: Wet and dry deposition of sulphur at the site Melpitz in East Germany, Water Air Soil Pollut., 109, 81–99, 1999.
- Chang, J. S., Brost, R. A., Isaksen, I. S. A., Madronich, S., Middleton, P., Stockwell, W. R., and Walcek, C. J.: A threedimensional Eulerian acid deposition model: Physical concepts and formulation, J. Geophys. Res.-Atmos., 92, 14681–14700, https://doi.org/10.1029/JD092iD12p14681, 1987.
- Chang, W. L., Bhave, P. V., Brown, S. S., Riemer, N., Stutz, J., and Dabdub, D.: Heterogeneous Atmospheric Chemistry, Ambient Measurements, and Model Calculations of N₂O₅: A Review, Aerosol Sci. Tech., 45, 665–695, https://doi.org/10.1080/02786826.2010.551672, 2011.

- Chang, W. L., Brown, S. S., Stutz, J., Middlebrook, A. M., Bahreini, R., Wagner, N. L., Dubé, W. P., Pollack, I. B., Ryerson, T. B., and Riemer, N.: Evaluating N₂O₅ heterogeneous hydrolysis parameterizations for CalNex 2010, J. Geophys. Res.-Atmos., 121, 5051–5070, https://doi.org/10.1002/2015JD024737, 2016.
- Chen, Y., Cheng, Y.-F., Nordmann, S., Birmili, W., Denier van der Gon, H. A. C., Ma, N., Wolke, R., Wehner, B., Sun, J., Spindler, G., Mu, Q., Pöschl, U., Su, H., and Wiedensohler, A.: Evaluation of the size segregation of elemental carbon (EC) emission in Europe: influence on the simulation of EC long-range transportation, Atmos. Chem. Phys., 16, 1823–1835, https://doi.org/10.5194/acp-16-1823-2016, 2016a.
- Chen, Y., Cheng, Y., Ma, N., Wolke, R., Nordmann, S., Schüttauf, S., Ran, L., Wehner, B., Birmili, W., van der Gon, H. A. C. D., Mu, Q., Barthel, S., Spindler, G., Stieger, B., Müller, K., Zheng, G.-J., Pöschl, U., Su, H., and Wiedensohler, A.: Sea salt emission, transport and influence on sizesegregated nitrate simulation: a case study in northwestern Europe by WRF-Chem, Atmos. Chem. Phys., 16, 12081–12097, https://doi.org/10.5194/acp-16-12081-2016, 2016b.
- Davis, J. M., Bhave, P. V., and Foley, K. M.: Parameterization of N₂O₅ reaction probabilities on the surface of particles containing ammonium, sulfate, and nitrate, Atmos. Chem. Phys., 8, 5295– 5311, https://doi.org/10.5194/acp-8-5295-2008, 2008.
- Dentener, F. J. and Crutzen, P. J.: Reaction of N_2O_5 on tropospheric aerosols: Impact on the global distributions of NO_x , O_3 , and OH, J. Geophys. Res.-Atmos., 98, 7149–7163, https://doi.org/10.1029/92JD02979, 1993.
- Doms, G., Schättler, U., Thomas, S., Gassmann, A., and Baldauf, M.: A Description of the Nonhydrostatic Regional COSMO-Model, in: Part I: Dynamics and Numerics, Deutscher Wetterdienst, Offenbach, available at: http://www.cosmo-model.org (3 December 2017), 2011a.
- Doms, G., Forstner, J., Heise, E., Herzog, H.-J., Raschendorfer, M., Reinhardt, T., Ritter, B., Schrodin, R., Schulz, J. P., and Vogel, G.: A Description of the Nonhydrostatic Regional COSMO-Model, in: Part II: Physical Parametrizations, Deutscher Wetterdienst, Offenbach, available at: http://www.cosmo-model.org, 2011b.
- Ehhalt, D. and Drummond, J. W.: The tropospheric cycle of NO_x : chemistry of the unpolluted and polluted troposphere, edited by: Georgii, W. J., Proc. NATO Adv. Study Inst., Corfu, Greece, 28 September–10 October 1981, Reidel, Dordrecht, 219–251, 1982.
- Erisman, J. W., Bleeker, A., Hensen, A., and Vermeulen, A.: Agricultural air quality in Europe and the future perspectives, Atmos. Environ., 42, 3209–3217, https://doi.org/10.1016/j.atmosenv.2007.04.004, 2008.
- Evans, M. J. and Jacob, D. J.: Impact of new laboratory studies of N₂O₅ hydrolysis on global model budgets of tropospheric nitrogen oxides, ozone, and OH, Geophys. Res. Lett., 32, L09813, https://doi.org/10.1029/2005GL022469, 2005.
- Fenter, F. F., Caloz, F., and Rossi, M. J.: Heterogeneous Kinetics of N₂O₅ Uptake on Salt, with a Systematic Study of the Role of Surface Presentation (for N₂O₅ and HNO₃), J. Phys. Chem., 100, 1008–1019, https://doi.org/10.1021/jp9503829, 1996.
- Folkers, M.: Bestimmung der Reaktionswahrscheinlichkeit von N₂O₅ an troposphärisch relevanten Aerosolen, PhD thesis, Universität zu Köln, 2002.

Y. Chen et al.: A parameterization of the heterogeneous hydrolysis of N₂O₅

- Folkers, M., Mentel, T. F., and Wahner, A.: Influence of an organic coating on the reactivity of aqueous aerosols probed by the heterogeneous hydrolysis of N₂O₅, Geophys. Res. Lett., 30, 1644, https://doi.org/10.1029/2003GL017168, 2003.
- Galperin, M. V. and Sofiev, M. A.: The long-range transport of ammonia and ammonium in the Northern Hemisphere, Atmos. Environ., 32, 373–380, https://doi.org/10.1016/S1352-2310(97)00045-9, 1998.
- Gaston, C. J., Thornton, J. A., and Ng, N. L.: Reactive uptake of N₂O₅ to internally mixed inorganic and organic particles: the role of organic carbon oxidation state and inferred organic phase separations, Atmos. Chem. Phys., 14, 5693–5707, https://doi.org/10.5194/acp-14-5693-2014, 2014.
- Gong, S. L.: A parameterization of sea-salt aerosol source function for sub- and super-micron particles, Global Biogeochem. Cy., 17, 1097, https://doi.org/10.1029/2003GB002079, 2003.
- Hallquist, M., Stewart, D. J., Baker, J., and Cox, R. A.: Hydrolysis of N₂O₅ on Submicron Sulfuric Acid Aerosols, J. Phys. Chem. A, 104, 3984–3990, https://doi.org/10.1021/jp9939625, 2000.
- Hallquist, M., Stewart, D. J., Stephenson, S. K., and Anthony Cox, R.: Hydrolysis of N₂O₅ on sub-micron sulfate aerosols, Phys. Chem. Chem. Phys., 5, 3453–3463, https://doi.org/10.1039/B301827J, 2003.
- Hardiman, S. C., Butchart, N., O'Connor, F. M., and Rumbold, S. T.: The Met Office HadGEM3–ES chemistry– climate model: evaluation of stratospheric dynamics and its impact on ozone, Geosci. Model Dev., 10, 1209–1232, https://doi.org/10.5194/gmd-10-1209-2017, 2017.
- Hatoya, K., Okuda, T., Funato, K., and Inoue, K.: Online Measurement of the Surface Area Concentration of Aerosols in Yokohama, Japan, using the Diffusion Charging Method, Asian Journal of Atmos. Environ., 10, 1–12, https://doi.org/10.5572/ajae.2016.10.1.001, 2016.
- Heinold, B., Tegen, I. N. A., Schepanski, K., Tesche, M., Esselborn, M., Freudenthaler, V., Gross, S., Kandler, K., Knippertz, P., MÜLler, D., Schladitz, A., Toledano, C., Weinzierl, B., Ansmann, A., Althausen, D., Müller, T., Petzold, A., and Wiedensohler, A.: Regional modelling of Saharan dust and biomass-burning smoke, Tellus B, 63, 781–799, https://doi.org/10.1111/j.1600-0889.2011.00570.x, 2011.
- Hertel, O., Reis, S., Skjøth, C. A., Bleeker, A., Harrison, R., Cape, J. N., Fowler, D., Skiba, U., Simpson, D., Jickells, T., Baker, A., Kulmala, M., Gyldenkaerne, S., Sørensen, L. L., and Erisman, J. W.: Nitrogen processes in the atmosphere, in: The European Nitrogen Assessment Sources, Effects and Policy Perspectives, edited by: Sutton, M. A., Howard, C. M., Erisman, J. W., Billen, G., Grennfelt, P., van Grinsven, H., and Grizzetti, B., 177–207, Cambridge University Press, Cambridge, UK, available at: http://www.cambridge.org/ena (3 December 2017), 2011.
- Hinneburg, D., Renner, E., and Wolke, R.: Formation of secondary inorganic aerosols by power plant emissions exhausted through cooling towers in Saxony, Environ. Sci. Pollut. R., 16, 25–35, https://doi.org/10.1007/s11356-008-0081-5, 2009.
- Hu, J. H. and Abbatt, J. P. D.: Reaction Probabilities for N₂O₅ Hydrolysis on Sulfuric Acid and Ammonium Sulfate Aerosols at Room Temperature, J. Phys. Chem. A, 101, 871–878, https://doi.org/10.1021/jp9627436, 1997.
- Im, U., Bianconi, R., Solazzo, E., Kioutsioukis, I., Badia, A., Balzarini, A., Baró, R., Bellasio, R., Brunner, D., Chemel, C.,

Curci, G., Denier van der Gon, H., Flemming, J., Forkel, R., Giordano, L., Jiménez-Guerrero, P., Hirtl, M., Hodzic, A., Honzak, L., Jorba, O., Knote, C., Makar, P. A., Manders-Groot, A., Neal, L., Pérez, J. L., Pirovano, G., Pouliot, G., San Jose, R., Savage, N., Schroder, W., Sokhi, R. S., Syrakov, D., Torian, A., Tuccella, P., Wang, K., Werhahn, J., Wolke, R., Zabkar, R., Zhang, Y., Zhang, J., Hogrefe, C., and Galmarini, S.: Evaluation of operational online-coupled regional air quality models over Europe and North America in the context of AQMEII phase 2. Part II: Particulate matter, Atmos. Environ., 115, 421–441, https://doi.org/10.1016/j.atmosenv.2014.08.072, 2015.

- Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, Atmos. Environ., 34, 2131–2159, 2000.
- Kane, S. M., Caloz, F., and Leu, M.-T.: Heterogeneous Uptake of Gaseous N₂O₅ by (NH₄)₂SO₄, NH₄HSO₄, and H₂SO₄ Aerosols, J. Phys. Chem. A, 105, 6465–6470, https://doi.org/10.1021/jp010490x, 2001.
- Karl, M., Dorn, H.-P., Holland, F., Koppmann, R., Poppe, D., Rupp, L., Schaub, A., and Wahner, A.: Product study of the reaction of OH radicals with isoprene in the atmosphere simulation chamber SAPHIR, J. Atmos. Chem., 55, 167–187, https://doi.org/10.1007/s10874-006-9034-x, 2006.
- Li, Y. P., Elbern, H., Lu, K. D., Friese, E., Kiendler-Scharr, A., Mentel, Th. F., Wang, X. S., Wahner, A., and Zhang, Y. H.: Updated aerosol module and its application to simulate secondary organic aerosols during IMPACT campaign May 2008, Atmos. Chem. Phys., 13, 6289–6304, https://doi.org/10.5194/acp-13-6289-2013, 2013.
- Long, M. S., Keene, W. C., Kieber, D. J., Erickson, D. J., and Maring, H.: A sea-state based source function for sizeand composition-resolved marine aerosol production, Atmos. Chem. Phys., 11, 1203–1216, https://doi.org/10.5194/acp-11-1203-2011, 2011.
- Lowe, D., Archer-Nicholls, S., Morgan, W., Allan, J., Utembe, S., Ouyang, B., Aruffo, E., Le Breton, M., Zaveri, R. A., Di Carlo, P., Percival, C., Coe, H., Jones, R., and McFiggans, G.: WRF-Chem model predictions of the regional impacts of N₂O₅ heterogeneous processes on night-time chemistry over north-western Europe, Atmos. Chem. Phys., 15, 1385–1409, https://doi.org/10.5194/acp-15-1385-2015, 2015.
- Macke, A., Seifert, P., Baars, H., Barthlott, C., Beekmans, C., Behrendt, A., Bohn, B., Brueck, M., Bühl, J., Crewell, S., Damian, T., Deneke, H., Düsing, S., Foth, A., Di Girolamo, P., Hammann, E., Heinze, R., Hirsikko, A., Kalisch, J., Kalthoff, N., Kinne, S., Kohler, M., Löhnert, U., Madhavan, B. L., Maurer, V., Muppa, S. K., Schween, J., Serikov, I., Siebert, H., Simmer, C., Späth, F., Steinke, S., Träumner, K., Trömel, S., Wehner, B., Wieser, A., Wulfmeyer, V., and Xie, X.: The HD(CP)2 Observational Prototype Experiment (HOPE) – an overview, Atmos. Chem. Phys., 17, 4887–4914, https://doi.org/10.5194/acp-17-4887-2017, 2017.
- Majewski, D., Liermann, D., Prohl, P., Ritter, B., Buchhold, M., Hanisch, T., Paul, G., Wergen, W., and Baumgardner, J.: The Operational Global Icosahedral–Hexagonal Gridpoint Model GME: Description and High-Resolution Tests, Mon. Weather Rev., 130, 319–338, https://doi.org/10.1175/1520-0493(2002)130<0319:TOGIHG> 2.0.CO;2, 2002.
- Mentel, T., F., Sohn, M., and Wahner, A.: Nitrate effect in the heterogeneous hydrolysis of dinitrogen pentoxide on aque-

ous aerosols, Phys. Chem. Chem. Phys., 1, 5451–5457, https://doi.org/10.1039/A905338G, 1999.

- Morgan, W. T., Ouyang, B., Allan, J. D., Aruffo, E., Di Carlo, P., Kennedy, O. J., Lowe, D., Flynn, M. J., Rosenberg, P. D., Williams, P. I., Jones, R., McFiggans, G. B., and Coe, H.: Influence of aerosol chemical composition on N₂O₅ uptake: airborne regional measurements in northwestern Europe, Atmos. Chem. Phys., 15, 973–990, https://doi.org/10.5194/acp-15-973-2015, 2015.
- Mozurkewich, M.: The dissociation constant of ammonium nitrate and its dependence on temperature, relative humidity and particle size, Atmos. Environ., 27A, 261–270, 1993.
- Mozurkewich, M. and Calvert, J. G.: Reaction probability of N₂O₅ on aqueous aerosols, J. Geophys. Res.-Atmos., 93, 15889–15896, https://doi.org/10.1029/JD093iD12p15889, 1988.
- Nenes, A., Pandis, S. N., and Pilinis, C.: ISORROPIA: A New Thermodynamic Equilibrium Model for Multiphase Multicomponent Inorganic Aerosols, Aquat. Geochem., 4, 123–152, https://doi.org/10.1023/a:1009604003981, 1998.
- Nordmann, S., Cheng, Y. F., Carmichael, G. R., Yu, M., Denier van der Gon, H. A. C., Zhang, Q., Saide, P. E., Pöschl, U., Su, H., Birmili, W., and Wiedensohler, A.: Atmospheric black carbon and warming effects influenced by the source and absorption enhancement in central Europe, Atmos. Chem. Phys., 14, 12683– 12699, https://doi.org/10.5194/acp-14-12683-2014, 2014.
- Neumann, D., Matthias, V., Bieser, J., Aulinger, A., and Quante, M.: A comparison of sea salt emission parameterizations in northwestern Europe using a chemistry transport model setup, Atmos. Chem. Phys., 16, 9905–9933, https://doi.org/10.5194/acp-16-9905-2016, 2016.
- Okuda, T.: Measurement of the specific surface area and particle size distribution of atmospheric aerosol reference materials, Atmos. Environ., 75, 1–5, https://doi.org/10.1016/j.atmosenv.2013.04.033, 2013.
- Platt, U. F., Winer, A. M., Biermann, H. W., Atkinson, R., and Pitts, J. N.: Measurement of nitrate radical concentrations in continental air, Environ. Sci. Technol., 18, 365–369, https://doi.org/10.1021/es00123a015, 1984.
- Poulain, L., Spindler, G., Birmili, W., Plass-Dülmer, C., Wiedensohler, A., and Herrmann, H.: Seasonal and diurnal variations of particulate nitrate and organic matter at the IfT research station Melpitz, Atmos. Chem. Phys., 11, 12579–12599, https://doi.org/10.5194/acp-11-12579-2011, 2011.
- Pouliot, G., Pierce, T., Denier van der Gon, H., Schaap, M., Moran, M., and Nopmongcol, U.: Comparing emission inventories and model-ready emission datasets between Europe and North America for the AQMEII project, Atmos. Environ., 53, 4– 14, https://doi.org/10.1016/j.atmosenv.2011.12.041, 2012.
- Reidy, B., Dämmgen, U., Döhler, H., Eurich-Menden, B., van Evert, F. K., Hutchings, N. J., Luesink, H. H., Menzi, H., Misselbrook, T. H., Monteny, G. J., and Webb, J.: Comparison of models used for national agricultural ammonia emission inventories in Europe: Liquid manure systems, Atmos. Environ., 42, 3452–3464, https://doi.org/10.1016/j.atmosenv.2007.04.009, 2008.
- Renner, E. and Wolke, R.: Modelling the formation and atmospheric transport of secondary inorganic aerosols with special attention to regions with high ammonia emissions, Atmos. Environ., 44, 1904–1912, https://doi.org/10.1016/j.atmosenv.2010.02.018, 2010.

- Riemer, N., Vogel, H., Vogel, B., Schell, B., Ackermann, I., Kessler, C., and Hass, H.: Impact of the heterogeneous hydrolysis of N_2O_5 on chemistry and nitrate aerosol formation in the lower troposphere under photosmog conditions, J. Geophys. Res.-Atmos., 108, 4144, https://doi.org/10.1029/2002JD002436, 2003.
- Riemer, N., Vogel, H., Vogel, B., Anttila, T., Kiendler-Scharr, A., and Mentel, T. F.: Relative importance of organic coatings for the heterogeneous hydrolysis of N₂O₅ during summer in Europe, J. Geophys. Res.-Atmos., 114, D17307, https://doi.org/10.1029/2008JD011369, 2009.
- Ritter, B. and Geleyn, J.-F.: A Comprehensive Radiation Scheme for Numerical Weather Prediction Models with Potential Applications in Climate Simulations, Mon. Weather Rev., 120, 303–325, https://doi.org/10.1175/1520-0493(1992)120<0303:acrsfn>2.0.co;2, 1992.
- Robinson, G. N., Worsnop, D. R., Jayne, J. T., Kolb, C. E., and Davidovits, P.: Heterogeneous uptake of CIONO₂ and N₂O₅ by sulfuric acid solutions, J. Geophys. Res.-Atmos., 102, 3583–3601, https://doi.org/10.1029/96JD03457, 1997.
- Saarikoski, S., Sillanpää, M., Sofiev, M., Timonen, H., Saarnio, K., Teinilä, K., Karppinen, A., Kukkonen, J., and Hillamo, R.: Chemical composition of aerosols during a major biomass burning episode over northern Europe in spring 2006: Experimental and modelling assessments, Atmos. Environ., 41, 3577–3589, https://doi.org/10.1016/j.atmosenv.2006.12.053, 2007.
- Sander, S. P., Friedl, R. R., Ravishankara, A. R., M., G. D., Kolb, C. E., Kurylo, M. J., Molina, M. J., Moortgat, G. K., and Finlayson-Pitts, B. J.: Chemical kinetics and photochemical data for use in atmospheric studies, evaluation number 14, JPL Publ., 02-25, 334 pp., 2003.
- Schell, B., Ackermann, I. J., Hass, H., Binkowski, F. S., and Ebel, A.: Modeling the formation of secondary organic aerosol within a comprehensive air quality model system, J. Geophys. Res.-Atmos., 106, 28275–28293, https://doi.org/10.1029/2001JD000384, 2001.
- Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics: From Air Pollution to Climate Change, John Wiley & Sons, New York, 2nd Edn., 2006.
- Simpson, D., Fagerli, H., Jonson, J. E., Tsyro, S., and Wind, P.: Transboundary ACIDIFICATION, Eutrophication and Ground Level Ozone in Europe. PART I: Unified EMEP Model Description, EMEP/MSC-W: EMEP Report, available at: http://emep.int/publ/reports/2003/emep_report_1_part1_ 2003.pdf (3 December 2017), 2003.
- Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L. D., Fagerli, H., Flechard, C. R., Hayman, G. D., Gauss, M., Jonson, J. E., Jenkin, M. E., Nyíri, A., Richter, C., Semeena, V. S., Tsyro, S., Tuovinen, J.-P., Valdebenito, Á., and Wind, P.: The EMEP MSC-W chemical transport model technical description, Atmos. Chem. Phys., 12, 7825–7865, https://doi.org/10.5194/acp-12-7825-2012, 2012.
- Sintermann, J., Neftel, A., Ammann, C., Háni, C., Hensen, A., Loubet, B., and Flechard, C. R.: Are ammonia emissions from field-applied slurry substantially over-estimated in European emission inventories?, Biogeosciences, 9, 1611–1632, https://doi.org/10.5194/bg-9-1611-2012, 2012.
- Spindler, G., Brüggemann, E., Gnauk, T., Grüner, A., Müller, K., and Herrmann, H.: A four-year size-segregated characterization

Y. Chen et al.: A parameterization of the heterogeneous hydrolysis of N₂O₅

study of particles PM_{10} , $PM_{2.5}$ and PM_1 depending on air mass origin at Melpitz, Atmos. Environ., 44, 164–173, 2010.

- Spindler, G., Gnauk, T., Grüner, A., Iinuma, Y., Müller, K., Scheinhardt, S., and Herrmann, H.: Size-segregated characterization of PM₁₀ at the EMEP site Melpitz (Germany) using a five-stage impactor: a six year study, J. Atmos. Chem., 69, 127–157, https://doi.org/10.1007/s10874-012-9233-6, 2012.
- Spindler, G., Grüner, A., Müller, K., Schlimper, S., and Herrmann, H.: Long-term size-segregated particle (PM₁₀, PM_{2.5}, PM₁) characterization study at Melpitz – influence of air mass inflow, weather conditions and season, J. Atmos. Chem., 70, 165–195, https://doi.org/10.1007/s10874-013-9263-8, 2013.
- Steppeler, J., Doms, G., Schättler, U., Bitzer, H. W., Gassmann, A., Damrath, U., and Gregoric, G.: Meso-gamma scale forecasts using the nonhydrostatic model LM, Meteorol. Atmos. Phys., 82, 75–96, https://doi.org/10.1007/s00703-001-0592-9, 2003.
- Stern, R., Builtjes, P., Schaap, M., Timmermans, R., Vautard, R., Hodzic, A., Memmesheimer, M., Feldmann, H., Renner, E., Wolke, R., and Kerschbaumer, A.: A model inter-comparison study focussing on episodes with elevated PM10 concentrations, Atmos. Environ., 42, 4567–4588, https://doi.org/10.1016/j.atmosenv.2008.01.068, 2008.
- Steinbrecher, R., Smiatek, G., Köble, R., Seufert, G., Theloke, J., Hauff, K., Ciccioli, P., Vautard, R., and Curci, G.: Intra- and inter-annual variability of VOC emissions from natural and semi-natural vegetation in Europe and neighbouring countries, Atmos. Environ., 43, 1380–1391, https://doi.org/10.1016/j.atmosenv.2008.09.072, 2009.
- Stockwell, W. R., Kirchner, F., Kuhn, M., and Seefeld, S.: A new mechanism for regional atmospheric chemistry modeling, J. Geophys. Res.-Atmos., 102, 25847–25879, https://doi.org/10.1029/97JD00849, 1997.
- Stohl, A., Williams, E., Wotawa, G., and Kromp-Kolb, H.: A European inventory of soil nitric oxide emissions and the effect of these emissions on the photochemical formation of ozone, Atmos. Environ., 30, 3741–3755, https://doi.org/10.1016/1352-2310(96)00104-5, 1996.
- Thornton, J. A., Braban, C. F., and Abbatt, J. P. D.: N₂O₅ hydrolysis on sub-micron organic aerosols: the effect of relative humidity, particle phase, and particle size, Phys. Chem. Chem. Phys., 5, 4593–4603, https://doi.org/10.1039/B307498F, 2003.
- Tiedtke, M.: A Comprehensive Mass Flux Scheme for Cumulus Parameterization in Large-Scale Models, Mon. Weather Rev., 117, 1779–1800, https://doi.org/10.1175/1520-0493(1989)117<1779:acmfsf>2.0.co;2, 1989.

- Van Doren, J. M., Watson, L. R., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Temperature dependence of the uptake coefficients of nitric acid, hydrochloric acid and nitrogen oxide (N₂O₅) by water droplets, J. Phys. Chem., 94, 3265–3269, https://doi.org/10.1021/j100371a009, 1990.
- Vignati, E., Karl, M., Krol, M., Wilson, J., Stier, P., and Cavalli, F.: Sources of uncertainties in modelling black carbon at the global scale, Atmos. Chem. Phys., 10, 2595–2611, https://doi.org/10.5194/acp-10-2595-2010, 2010.
- Walker, J. M., Philip, S., Martin, R. V., and Seinfeld, J. H.: Simulation of nitrate, sulfate, and ammonium aerosols over the United States, Atmos. Chem. Phys., 12, 11213–11227, https://doi.org/10.5194/acp-12-11213-2012, 2012.
- Wahner, A., Mentel, T. F., Sohn, M., and Stier, J.: Heterogeneous reaction of N_2O_5 on sodium nitrate aerosol, J. Geophys. Res.-Atmos., 103, 31103–31112, https://doi.org/10.1029/1998JD100022, 1998.
- Williams, E. J., Guenther, A., and Fehsenfeldi, F. C.: An inventory of nitric oxide emissions from soils in the United States, J. Geophys. Res.-Atmos., 97, 7511–7519, https://doi.org/10.1029/92JD00412, 1992.
- Wolke, R., Knoth, O., Hellmuth, O., Schröder, W., and Renner, E.: The parallel model system LM-MUSCAT for chemistrytransport simulations: Coupling scheme, parallelization and application, edited by: Joubert, G. R., Nagel, W. E., Peters, F. J., and Walter, W. V., Parallel Computing: Software Technology, Algorithms, Architectures, and Applications, Elsevier, Amsterdam, The Netherlands, 13, 363–370, https://doi.org/10.1016/S0927-5452(04)80048-0, 2004.
- Wolke, R., Schröder, W., Schrödner, R., and Renner, E.: Influence of grid resolution and meteorological forcing on simulated European air quality: A sensitivity study with the modeling system COSMO–MUSCAT, Atmos. Environ., 53, 110–130, https://doi.org/10.1016/j.atmosenv.2012.02.085, 2012.
- Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, J. Geophys. Res., 104, 30387–30415, 1999.
- Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), J. Geophys. Res.-Atmos., 113, D13204, https://doi.org/10.1029/2007JD008782, 2008.