

Supplement of Atmos. Chem. Phys., 19, 10319–10334, 2019
<https://doi.org/10.5194/acp-19-10319-2019-supplement>
© Author(s) 2019. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Distinctions in source regions and formation mechanisms of secondary aerosol in Beijing from summer to winter

Jing Duan et al.

Correspondence to: Ru-Jin Huang (rujin.huang@ieecas.cn)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Table S1. Summary of mass concentrations of aerosol species, gaseous pollutants and meteorological parameters during entire study and different seasons. PM₁ chemical composition during clean days (C), M-polluted days (M) and H-polluted days (H) in different seasons are also shown.

Species	Entire study	Late Summer			Autumn				Early Winter			
		Entire	C	M	Entire	C	M	H	Entire	C	M	H
Aerosol species ($\mu\text{g m}^{-3}$)												
PM ₁	44.7	21.6	15.6	46.9	43.3	9.3	54.2	110.5	64.3	8.1	43.5	109.7
OA	22.4	13.8	10.5	28.1	21.2	5.8	25.5	50.8	29.6	4.5	20.4	49.4
HOA	2.2	1.8	1.3	3.9	2.1	0.4	2.5	6.1	3.3	0.2	1.2	6.4
COA	3.9	1.4	1.5	2.0	4.3	1.9	4.6	8.1	4.7	1.1	4.3	6.9
CCOA	3.1	0.7	0.5	1.7	2.0	0.6	2.3	4.6	7.7	1.1	5.3	12.8
LSOA	9.0	3.2	2.2	6.7	9.2	1.5	10.2	25.4	12.1	1.5	7.5	20.7
RSOA	4.2	6.6	5.0	13.8	3.8	1.5	5.9	6.6	1.8	0.6	2.0	2.5
SO ₄	5.6	3.1	1.9	8.0	4.8	0.8	7.6	11.0	9.6	1.5	6.5	16.5
NO ₃	8.7	1.3	0.9	3.3	9.5	0.6	12.5	27.6	12.9	0.7	8.7	23.0
NH ₄	3.8	1.5	0.9	3.8	3.5	0.5	3.7	11.0	6.4	0.7	4.4	9.9
Chl	0.8	0.2	0.05	0.5	0.8	0.1	0.7	2.2	1.8	0.2	1.3	3.3
BC	3.4	1.7	1.4	3.3	3.5	1.5	4.3	7.7	3.9	0.4	2.2	7.7
Gaseous pollutants												
SO ₂ (ppb)	5.0	3.0	2.7	3.5	4.4	3.0	4.4	6.2	7.9	3.7	5.2	11.7
CO (ppm)	0.9	0.5	0.6	0.4	0.7	0.3	0.8	1.2	1.7	0.2	0.9	3.0
NO (ppb)	19.1	4.7	3.7	4.2	18.1	8.9	15.5	28.2	32.7	4.3	18.9	55.9
NO ₂ (ppb)	29.9	18.7	14.6	23.1	32.0	22.3	32.3	48.7	34.0	11.9	25.5	49.1
O ₃ (ppb)	18.0	35.4	31.5	54.1	17.5	27.2	16.1	20.8	5.7	19.7	4.9	2.7
Meteorological parameters												
RH (%)	72.5	71.1	71.4	64.5	69.6	56.7	82.3	75.9	73.9	34.3	79.4	82.7
T (°C)	12.7	23.9	24.2	27.0	14.4	17.9	14.9	14.5	1.5	-3.0	1.3	2.8
WS (m s ⁻¹)	0.95	0.88	0.89	0.93	0.97	1.23	0.63	0.55	1.26	3.60	1.12	0.63




Figure S1. PMF profiles of OA sources for 4, 5 and 6 factor solutions.




Figure S2. ME-2 profiles of OA sources. The HOA profile is from that of Ng et al. (2011b), and the COA profile is from 4-factor PMF result. The others are unconstrained factors.




Figure S3. Mass fractions of PM₁ species and OA factors in later summer (a), autumn (b) and early winter (c).




Figure S4. Diurnal cycles of PM₁ species and OA factors during late summer (a, d), autumn (b, e) and early winter (c, f).




Figure S5. Bivariate polar plots of LSOA (a) and RSOA (b) as functions of wind direction and wind speed (m s^{-1}) during the entire study period.




Figure S6. Bivariate polar plots of O_x as functions of wind direction and wind speed (m s^{-1}) during late summer.