



## ***Corrigendum to*** **“Rate enhancement in collisions of sulfuric acid molecules due to long-range intermolecular forces” published in Atmos. Chem. Phys., 19, 13355–13366, 2019**

**Roope Halonen<sup>1</sup>, Evgeni Zapadinsky<sup>1</sup>, Theo Kurtén<sup>2</sup>, Hanna Vehkamäki<sup>1</sup>, and Bernhard Reischl<sup>1</sup>**

<sup>1</sup>Institute for Atmospheric and Earth System Research/Physics, Faculty of Science, University of Helsinki, P.O. Box 64, 00014, Helsinki, Finland

<sup>2</sup>Institute for Atmospheric and Earth System Research/Chemistry, Faculty of Science, University of Helsinki, P.O. Box 55, 00014, Helsinki, Finland

**Correspondence:** Roope Halonen (roope.halonen@helsinki.fi)

Published: 4 November 2019

After publication an error was noted in the abstract.

The original sentence published was

“We found that the effective collision cross section of the H<sub>2</sub>SO<sub>4</sub> molecule, as described by an optimized potentials for liquid simulation (OPLS). OPLS all-atom force field, is significantly larger than the hard-sphere diameter assigned to the molecule based on the liquid density of sulfuric acid.”.

The correct sentence is as follows:

“We found that the effective collision cross section of the H<sub>2</sub>SO<sub>4</sub> molecule, as described by an optimized potentials for liquid simulation (OPLS) all-atom force field, is significantly larger than the hard-sphere diameter assigned to the molecule based on the liquid density of sulfuric acid.”.