

Supplement of Atmos. Chem. Phys., 19, 7627–7647, 2019  
<https://doi.org/10.5194/acp-19-7627-2019-supplement>  
© Author(s) 2019. This work is distributed under  
the Creative Commons Attribution 4.0 License.



Atmospheric  
Chemistry  
and Physics  
Open Access  



*Supplement of*

## **Extratropical age of air trends and causative factors in climate projection simulations**


**Petr Šácha et al.**

*Correspondence to:* Petr Šácha (sacha@uvigo.es)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.



**Figure S1.** Time evolution of CMAM zonal mean ozone volume mixing ratio [ $\text{mole}(O_3)/\text{mole}(\text{air})$ ] averaged between 12 and 22gpkm and  $60^\circ$ -  $90^\circ\text{N}$  (blue solid) and  $60^\circ$ -  $90^\circ\text{S}$  (red dashed) smoothed by 12-month running average.



**Figure S2.** CMAM trend of dry density [ $\text{kgm}^{-3}/\text{decade}$  ].Top: Density trends in geopotential height vertical coordinate. Bottom: Density trends in vertical coordinate corrected to the vertical shift of pressure levels. The mean tropopause and turnaround latitude positions in the given period are marked

with black lines. Only the trends in the regions where they exceed the statistical significance of 95% confidence level are plotted.

**Table S1.** Annual trends of the residual circulation strength (RC) in  $10^{-3} \text{kgm}^{-2} \text{s}^{-1} / \text{decade}$ , aging by mixing (AbM), AoA and RCTT ( $\text{days/decade}$ ) averaged over the Ex regions in the Ref, NF and F period. AbM and RCTT trends are computed in a reduced Ref period (1970-2000). Presented are trend values significant at least at the 90% confidence level and also at least at the 80% confidence level (denoted by superscript  $^+$ ), otherwise the cell is left blank.

Period	ExNH							
	AoA	AbM	RCTT	RC	AoA <sup>c</sup>	AbM <sup>c</sup>	RCTT <sup>c</sup>	RC <sup>c</sup>
Ref	-32	-24	-6 <sup>+</sup>	0.19	-30	-23		0.17 <sup>+</sup>
NF	-17	-6	-11	0.17	-11	-3	-9	
F	-31	-21	-10	0.22	-23	-16	-7	
ExSH								
Ref	-41	-50			-38	-49		
NF	-15	-3	-11	0.14	-11		-9	0.01 <sup>+</sup>
F	-33	-25	-9	0.15	-28	-22	-7	0.01 <sup>+</sup>

**Table S2.** The net tropical upwelling proxy (UP) annual trend in  $10^7 \text{kgs}^{-1} / \text{decade}$  in the Ref, NF and F period. Presented are trend values significant at least at the 90% confidence level and also at least at the 80% confidence level (denoted by superscript  $^+$ ), otherwise the cell is left blank. The superscript  $c$  denotes a trend computed in the geopotential height corrected to the vertical shift of pressure levels, the superscript  $^{rel}$  denotes a trend computed in the geopotential height corrected to the tropopause shift.

Period	UP	UP <sup>c</sup>	UP <sup>rel</sup>
Ref	8.6 <sup>+</sup>		
NF	9.7	7.6	8.4
F	12.2	9.1	7.6

**Table S3.** The net tropical upwelling (UPd) annual trend (including density trend in contrast to UP) in  $10^7 \text{kgs}^{-1} / \text{decade}$  in the Ref, NF and F period. Presented are trend values significant at least at the 90% confidence level and also at least at the 80% confidence level (denoted by superscript  $^+$ ), otherwise the cell is left blank.

Period	UPd	UPd <sup>c</sup>	UPd <sup>rel</sup>
Ref	11.6		
NF	13.6		9.7
F	17		

**Table S4.** Seasonal values of correlations between spatial averages of total drag (TD, EPFD + GWD) and  $v^*$  ( $C^{Td-Rc}$ ), EPFD and  $v^*$  ( $C^{EP-Rc}$ ), GWD and  $v^*$  ( $C^{Gw-Rc}$ ) and correlations of EPFD and GWD ( $C^{Gw-Rc}$ ) in the Ref, NF and F period. For the description of spatial average areas please refer to the main text. Presented are only correlation values significant at least at the 90% confidence level, otherwise the significance is given.

Season	ExNH					ExSH				
	DJF	MAM	JJA	SON	Ann	DJF	MAM	JJA	SON	Ann
<b>Ref</b>										
<b>C<sup>Td-Rc</sup></b>	-0.95	-0.95	-0.81	-0.89	-0.92	0.95	0.9	0.92	0.94	0.66
<b>C<sup>EP-Rc</sup></b>	-0.84	-0.87	-0.58	-0.72	-0.84	0.77	0.84	0.82	0.86	0.68
<b>C<sup>Gw-Rc</sup></b>	-0.82	-0.72	-0.36	-0.52	-0.9	0.64	0.37			0.31
<b>NF</b>										
<b>C<sup>Td-Rc</sup></b>	-0.97	-0.94	-0.93	-0.92	-0.92	0.93	0.92	0.89	0.93	0.64
<b>C<sup>EP-Rc</sup></b>	-0.87	-0.81	-0.86	-0.82	-0.83	0.78	0.87	0.69	0.85	0.7
<b>C<sup>Gw-Rc</sup></b>	-0.67	-0.61	-0.63	-0.36	-0.88	0.56	0.45		-0.28	0.21
<b>F</b>										
<b>C<sup>Td-Rc</sup></b>	-0.97	-0.95	-0.93	-0.88	-0.91	0.91	0.92	0.93	0.96	0.64
<b>C<sup>EP-Rc</sup></b>	-0.75	-0.79	-0.78	-0.72	-0.83	0.85	0.83	0.85	0.87	0.71
<b>C<sup>Gw-Rc</sup></b>	-0.62	-0.69	-0.56	-0.39	-0.88	0.4	0.24		-0.26	0.19

**Table S5.** Upward shift corrected trend values of total drag (TD, EPFD + GWD) and its components EPFD and GWD multiplied by  $\rho$  (all in  $10^{-5} \text{kgm}^{-2}\text{s}^{-2}/\text{decade}$ ) and RC ( $10^{-3} \text{kgm}^{-2}\text{s}^{-1}/\text{decade}$ ) averaged over the Ex regions in the Ref, NF and F period. Presented are trend values significant at least at the 90% confidence level and also at least at the 80% confidence level (denoted by superscript  $^+$ ), otherwise the cell is left blank.

Period	Ref					NF					F				
	DJF	MAM	JJA	SON	Ann	DJF	MAM	JJA	SON	Ann	DJF	MAM	JJA	SON	Ann
<b>ExNH</b>															
<b>TD</b>	-3.5	-1.4 <sup>+</sup>	-0.6	-1.3		-1		-0.9							
<b>EPFD</b>	-1.4					-0.6 <sup>+</sup>		-0.7	-0.3 <sup>+</sup>						
<b>GWD</b>	-3	-1.5				-1.1 <sup>+</sup>		-0.3						0.3 <sup>+</sup>	
<b>RC</b>	4	1.3 <sup>+</sup>	1	2.2	0.2	0.1 <sup>+</sup>		0.7	0.7 <sup>+</sup>	0.2			1.4	2	0.3
<b>ExSH</b>															
<b>TD</b>	-2.3	-1.1	1.5	2.2		-1.7			-0.9 <sup>+</sup>	-0.7	-1				-0.6
<b>EPFD</b>		-0.7	1			-0.9			-0.4 <sup>+</sup>	-0.5 <sup>+</sup>	-0.5 <sup>+</sup>				-0.4 <sup>+</sup>
<b>GWD</b>	-0.5	-0.4 <sup>+</sup>	-1.2		-0.6	-0.5			-1.7	-0.6	-0.3 <sup>+</sup>			-0.8	-0.3 <sup>+</sup>
<b>RC</b>	3.2	1.6		-4		1 <sup>+</sup>	1.3	1		0.2	0.8 <sup>+</sup>	0.9 <sup>+</sup>	1.8		0.2

**Table S6.** UPd<sup>c</sup> and UPd<sup>rel</sup> trends in the Ref, NF and F period computed directly by integrating  $\rho w^*$  between the turn-around latitudes (UPd<sup>c</sup><sub>wstar</sub>), indirectly by evaluating the residual mean stream function at the turn-around latitudes using  $\int_{20gpkm}^{45gpkm} \rho v^* dz$  (UPd<sup>c</sup><sub>vstar</sub>) and using  $-\int_{20gpkm}^{45gpkm} \rho \frac{TD}{f} dz$  (UPd<sup>c</sup><sub>DC</sub>). The trends are given in  $10^7 kgs^{-1} / decade$ . Presented are trend values significant at least at the 90% confidence level and also at least at the 80% confidence level (denoted by superscript <sup>+</sup>), otherwise the cell is left blank.

Period	Upwelling														
	Ref					NF					F				
Season	DJF	MAM	JJA	SON	Ann	DJF	MAM	JJA	SON	Ann	DJF	MAM	JJA	SON	Ann
UPd <sup>c</sup> <sub>DC</sub>						0.4 <sup>+</sup>	0.4				0.5	0.6			
UPd <sup>c</sup> <sub>vstar</sub>	24.4	7.6				10.6	4.6	3.6 <sup>+</sup>			6.9	8.2	5.3 <sup>+</sup>	6.9 <sup>+</sup>	5.3 <sup>+</sup>
UPd <sup>c</sup> <sub>wstar</sub>	26.5	7.4				12.8					7.5	8.7	5 <sup>+</sup>		
UPd <sup>rel</sup> <sub>wstar</sub>	8.1 <sup>+</sup>	-37	-37.6	-70.1		16.8	-4.2 <sup>+</sup>	-6.2	-12.8	9.1	-93.3	-44.7	-49.1 <sup>+</sup>		