



Supplement of

The day-to-day co-variability between mineral dust and cloud glaciation: a proxy for heterogeneous freezing

Diego Villanueva et al.

Correspondence to: Diego Villanueva (villanueva@tropos.de)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

For figures S.1 to S.14:

a) Time average at -15°C (averaged in a 12 K range), 2007-2010.

b) Time-zonal average, 2007-2010.

c) Zonal average at -15°C (averaged in a 12 K range), 2007-2010.

S 1-S 3: Frequency phase ratio (FPR) variables.

S 4-S.7: Dust aerosol data from MACC reanalysis.

S.8-S.9: Cloud volume fraction of all and stratiform clouds (CALIPSO-GOCCP).

S.10-S.13: Vertical velocity (MACC), relative humidity (ERA-Interim), isotherm height and temperature (ECMWF-AUX).

S.14-S.16: Sample size distribution.

S.17-S.20: Day-to-day dust-ice statistics without meteorological constrains.



S1. FPR_{DARDAR} [%]



S2. FPR_ALT_{DARDAR} [%]



S3. FPR_{GOCCP} [%]



S4. MACC coarse dust [kg/kg].



S5. Standard deviation of coarse log (dust) MACC [log(kg/kg)].



S6. MACC fine dust [kg/kg].



S7. Standard deviation of fine log (dust) MACC [log(kg/kg)].



S8. Cloud volume fraction [%].



S9. Stratiform cloud volume fraction [%].





S10. MACC vertical velocity [Pa/s] for stratiform clouds.



S11. MACC relative humidity [%] for stratiform clouds.



S12. ECMWF-AUX isotherm height [m] for stratiform clouds.



S13. ECMWF-AUX temperature [K] for stratiform clouds.



S14. Sample size of stratiform clouds [#gridboxes(month, dust decile, temperature, latitude, longitude)].





S15. Sample size of stratiform clouds [#gridboxes(month, dust decile, temperature, latitude, longitude)] for different seasons.















S17. Day-to-day covariance of fine dust mixing-ratio (MACC) and $FPR_{GOCCP} 2007-2010$. The covariance is normalized by the fine dust standard deviation. a) Time mean covariance between 12 monthly covariance values at -15°C (averaged in a 12 K range). b) Time-zonal mean covariance. c) Monthly-zonal mean covariance at -15°C. Red colours denote a high correlation between day-to-day fine dust and ice occurrence. The colour and pattern of the boxes corresponds to the regression lines in S 18 and S 19.



S18. Average cloud phase for the mid-latitude and high-latitude bands for -15 °C (range -21 °C to -9 °C) in the period 2007-2010. The horizontal axis corresponds to the different time deciles (day-to-day variability) of fine (a) and coarse (b) dust mixing-ratio (MACC), calculated for each each 3°C temperature bin and gridbox (1.875°x1.875°) and averaged along each 12 °C temperature range and latitude band (30°x360°). The vertical bars are positioned at each dust decile and show the mean zonal standard deviation within each latitude band. The lines represent the linear regressions (*FPR=A·m^{2/3}+FPR*₀). Only stratiform and non-precipitating clouds were included.



S 19. Same as Fig 7.a but for a) -30 °C (range -36 °C to -24 °C) and b) winter (blue) and summer (red) seasons for the 40-80°N/S latitude bands.



S 20. Average sea/land cloud phase (CALIPSO-GOCCP) a) at -15 °C (range -21 °C to -9 °C) against the MACC deciles of fine dust mixing ratio and b) against temperature for 2007-2010. The vertical lines correspond to the standard deviation between a) Longitudes and b) Months of the year.