

Supplement of Atmos. Chem. Phys., 20, 2781–2794, 2020  
<https://doi.org/10.5194/acp-20-2781-2020-supplement>  
© Author(s) 2020. This work is distributed under  
the Creative Commons Attribution 4.0 License.



*Supplement of*

## **Morphology and size of the particles emitted from a gasoline-direct-injection-engine vehicle and their ageing in an environmental chamber**

**Jiaoping Xing et al.**

*Correspondence to:* Longyi Shao ([shaol@cumtb.edu.cn](mailto:shaol@cumtb.edu.cn)) and Daizhou Zhang ([dzzhang@pu-kumamoto.ac.jp](mailto:dzzhang@pu-kumamoto.ac.jp))

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.


Table S1 Information of various parameters for the vehicles in the experiment

Vehicle ID	GDI-1.4-T
Engine type	GDI
Induction system	turbo
Number of cylinders	4
Displacement/L	1.395
Compression ratio	10.5
After-treatment	TWC
Bore /mm	74.5
Stroke /mm	80.0
Maximum power at engine speed /(kW)/(r/min)	110/(5000~6000)
Maximum torque at engine speed /((N·m)/(r/min))	250/(1750~3000)

Table S2 Properties of fuels used in this experiment.

Fuel	F2
RON	93
MON	84
Density (g/cm <sup>3</sup> )	0.744
Aromatics (%v/v)	36.7
Olefins (%v/v)	15.4
EtOH (%v/v)	0.01
Oxygen (%m/m)	0.02
MMT (mg/L)	<1
Manganese (mg/kg)	<0.1
Sulphur (mg/kg)	6
T10 (°C)	55.4
T50 (°C)	109.9
T90 (°C)	164.3
FBP (°C)	194.4

(a)



(b)




Figure S1. (a) Speed in one Beijing driving cycle under the running test. It is a cycle with a 300-s “cold start” phase followed by a 767-s “hot stabilized running” phase. (b) Sketch of the test system. The number concentration of particles during the tests was monitored by a Combustion Fast Particle Size Spectrometer DMS 500. A single-stage cascade impactor was mounted at the exit of the tailpipe to collect particles on TEM grids with a flow rate 1.0 L/min. A dilution unit was applied to dilute the tailpipe gas into 1/10 concentration using synthetic air.




Figure S2. Size distribution of particles emitted from the GDI-engine gasoline vehicles by the DMS500 instrument.




Figure S3. The numbers of accumulation mode particles emitted from the GDI vehicle under different running states.




Figure S4. Size distribution of particles emitted from the GDI-engine gasoline vehicles by the DMS500 instrument during the Beijing driving cycle.




Figure S5. Number fractions of different particle types in the emission of the GDI vehicle under different running states. N is the number of particles analyzed for the different running states. Data presented as mean  $\pm$  standard deviation,  $n = 3$ .