



Fossil fuel combustion, biomass burning and biogenic sources of fine carbonaceous aerosol in the Carpathian Basin

Imre Salma¹, Anikó Vasanits-Zsigrai¹, Attila Machon², Tamás Varga³, István Major³, Virág Gergely³, and Mihály Molnár³

¹Institute of Chemistry, Eötvös University, Budapest, Hungary

²Air Quality Reference Center, Hungarian Meteorological Service, Budapest, Hungary

³Isotope Climatology and Environmental Research Centre, Institute for Nuclear Research, Debrecen, Hungary

Correspondence: Imre Salma (salma@chem.elte.hu)

Received: 3 September 2019 – Discussion started: 20 November 2019

Revised: 1 March 2020 – Accepted: 18 March 2020 – Published: 9 April 2020

Abstract. Fine-fraction aerosol samples were collected, and air pollutants and meteorological properties were measured in situ in the regional background environment of the Carpathian Basin, a suburban area and central part of its largest city, Budapest, in each season for a 1-year-long time interval. The samples were analysed for PM_{2.5} mass, organic carbon (OC), elemental carbon (EC), water-soluble OC (WSOC), radiocarbon, levoglucosan (LVG) and its stereoisomers, and some chemical elements. Carbonaceous aerosol species made up 36 % of the PM_{2.5} mass, with a modest seasonal variation and with a slightly increasing tendency from the regional background to the city centre (from 32 % to 39 %). A coupled radiocarbon-LVG marker method was applied to apportion the total carbon (TC = OC + EC) into contributions of EC and OC from fossil fuel (FF) combustion (EC_{FF} and OC_{FF}, respectively), EC and OC from biomass burning (BB) (EC_{BB} and OC_{BB}, respectively), and OC from biogenic sources (OC_{BIO}). Fossil fuel combustion showed rather constant daily or monthly mean contributions (of 35 %) to the TC in the whole year in all atmospheric environments, while the daily contributions of BB and biogenic sources changed radically (from < 2 % up to 70 %–85 %) at all locations and over the years. In October, the three major sources contributed equally to the TC in all environments. In January, it was the BB that was the major source, with a share of 70 % at all sites. The contributions from biogenic sources in January were the smallest. In April, FF combustion and biogenic sources were the largest two contributors at all locations with typical shares of 45 %–50 % each. In July, biogenic sources became the major source type with a

monotonically increasing tendency (from 56 % to 72 %) from the city centre to the regional background. The share of BB was hardly quantifiable in July. The EC_{FF} made up more than 90 % of EC in April and July, while in October and January, the contributions of EC_{BB} were considerable. Biomass burning in winter and autumn offers the largest and most considerable potential for improving the air quality in cities as well as in rural areas of the Carpathian Basin.

1 Introduction and objectives

Carbonaceous aerosol constituents make up a major part (e.g. 20 %–60 % in the continental mid-latitudes and up to 90 % in tropical forests) of the PM_{2.5} mass (Kanakidou et al., 2005; Fuzzi et al., 2015). Their largest emission or production source types are fossil fuel (FF) combustion, biomass burning (BB) and biogenic sources (Le Queré et al., 2018). These processes also represent the highest source of certain important aerosol species such as soot (species mainly containing C with imperfect or fragmented graphitic structure; Andreae and Gelencsér, 2006) and of some pollutant or greenhouse gases such as CO, NO_x, CO₂ and volatile organic compounds (VOCs) on a global spatial scale (Wiedinmyer et al., 2011; Von Schneidemesser et al., 2015; Tian et al., 2016). The sources produce both primary and secondary particles, and they are linked to a variety of anthropogenic activities directly or indirectly and in many ways (Hallquist et al., 2009). The perturbations in atmospheric concentrations and chemical, physical and meteorological properties

caused by these sources have important consequences for the Earth system. The related processes include the radiation balance (Lohmann et al., 2000), cloud formation and properties, water cycling, and other biogeochemical cycles (Andreae and Rosenfeld, 2008; Cecchini et al., 2017), atmospheric chemistry and nucleation (Fuzzi et al., 2015; Nozière et al., 2015; Kirkby et al., 2016), atmospheric transport and mixing (Rosenfeld et al., 2019), forest growth and agriculture production (Artaxo et al., 2009; Rap et al., 2015), ecosystems (Cirino et al., 2014), built environment and cultural heritage (Bonazza et al., 2005), and human health and well-being (Lelieveld and Pöschl, 2017; Burnett et al., 2018). Some of the sources, e.g. fuel wood or agricultural residue burnings, are expected to be increased in both domestic and industrial sectors due to their role in decentralised and substitute energy production (Vicente and Alves, 2018). At the same time, the potential disadvantages and risk of these major source types – including BB in particular – have been less recognised (Hays et al., 2003; Chen et al., 2017). It is, therefore, highly relevant to estimate the relative contribution of FF combustion, BB and biogenic sources to major carbonaceous aerosol species, namely to organic carbon (OC) and elemental carbon (EC).

Large quantities, composite character, and spatial and temporal variability of the sources together with atmospheric transformation of their products make the quantification of the source types or their inventory-based source assessment challenging (Nozière et al., 2015). There are several methods to apportion the particulate matter (PM) mass or carbonaceous species among some or all of their major sources. They include (1) source-specific marker methods (Fraser and Lakshmanan, 2000; Szidat et al., 2006, 2009; Minguillón et al., 2011; Zhang et al., 2012; Bernardoni et al., 2013), (2) multi-wavelength optical methods (Sandradewi et al., 2008a, b; Zotter et al., 2017) and (3) various multivariate statistical methods based on online or offline data (Hopke, 2016; Maenhaut et al., 2016). Recently, the third approach also takes advantage of dedicated molecular tracers or fragments from mass spectrometry or advanced optical techniques (Forello et al., 2019; Stefenelli et al., 2019).

The marker methods are beneficial from the point of view that they do not demand many samples or very extensive data sets and that the required analytical data are ordinarily available in related studies. Among the most frequently and successfully adopted markers are radiocarbon (^{14}C , $T_{1/2} = 5730$ years), which is used for quantifying FF combustion, and levoglucosan (LVG, monosaccharide anhydride $\text{C}_6\text{H}_{10}\text{O}_5$), which is utilised for assessing BB. The monosaccharide anhydride analysis often involves the stereoisomers of LVG, namely mannosan (MAN) and galactosan (GAN) in addition to LVG since their concentration ratios can be connected to biomass and wood types (e.g. hardwood or softwood; Fine et al., 2004; Schmidl et al., 2008). The formation, modelling utilisation, atmospheric processes and analytical determinations of these two markers together with their ad-

vantages and limitations were described, evaluated and discussed in detail previously (e.g. Simoneit et al., 1999, 2004; Fraser and Lakshmanan, 2000; Nolte et al., 2001; Pashynska et al., 2002; Zdráhal et al., 2002; Puxbaum et al., 2007; Saarikoski et al., 2008; Caseiro et al., 2009; Fabbri et al., 2009; Szidat et al., 2006, 2009; Favez et al., 2010; Hennigan et al., 2010; Hoffmann et al., 2010; Kourtchev et al., 2011; Piazzalunga et al., 2011; Maenhaut et al., 2012, 2016; Yttri et al., 2015). The coupled radiocarbon-LVG marker method, introduced recently (Salma et al., 2017), is a combination of the two marker methods. It allows the TC ($\text{TC} = \text{OC} + \text{EC}$) to be apportioned among all major source types, and thus among the contributions of EC and OC from FF combustion (EC_{FF} and OC_{FF} , respectively), EC and OC from BB (EC_{BB} and OC_{BB} , respectively), and OC from biogenic sources (OC_{BIO}).

Water-soluble OC (WSOC) is also an important carbonaceous aerosol species because it is considered as an indicator of secondary organic aerosol (SOA) or carbonaceous particles after atmospheric chemical ageing (Claeys et al., 2010). It is related to more oxygenated chemical species than freshly emitted or formed organic constituents, and this class of molecules is expected to contribute substantially to cloud condensation nuclei (CCN) activity of particles and represents potentially larger negative health effects of particulate mass deposited in the human respiratory system due to its solubility (Hallquist et al., 2009; Fuzzi et al., 2015; Nozière et al., 2015).

Despite their overall role together with the health, climate and environmental effects, there are serious gaps in our knowledge on FF combustion, BB and biogenic sources – particularly on biogenic sources in more polluted or urban areas. Information on the properties of the major apportioned or secondary carbonaceous aerosol species and on their relationships with other atmospheric quantities have been missing internationally on extended spatial scales as well as on larger cities. The Carpathian Basin (also known as the Pannonian Basin) is the largest, topographically well separated, orogenic basin in Europe (Salma et al., 2016b). Its land is mostly used for intensive agriculture and farming, while larger forested areas with deciduous, coniferous or mixed wood occur in the inner and bounding mountains. Weather situations within the basin are generally uniform, which makes it advantageous for studying atmospheric phenomena and processes. Budapest with 2.3 million inhabitants in the metropolitan area and with its central geographical location is the largest city in the basin. The mean green space intensity – which indicates the healthy green coverage – for Budapest in 2015 was estimated from Landsat satellite images to be approximately 50 %, with spatial variations from 19 % in the city centre to 55 % in the suburban zone (Tatai et al., 2017).

As part of a research project, we collected aerosol samples in the regional background atmospheric environment of the Carpathian Basin, suburban area and city centre of Budapest

in each season for a 1-year-long time interval and analysed them for various aerosol constituents, which are required for source apportionment. The analytical results were complemented by supporting air pollutant and meteorological data as well. The major objectives of the present paper are to report the main findings of this research; to discuss the properties and contributions of FF combustion, BB and biogenic sources and related atmospheric processes; to interpret the relationships among various variables for different months and environmental types; and to formulate some general conclusions on the air quality of the region and the city.

2 Methods

2.1 Collection of aerosol samples and in situ measurements

The aerosol samples were collected at three sites in Hungary, in a rural background area and at two urban sites in Budapest (Fig. 1). The samplings at the rural location were realised at K-pusztas station ($46^{\circ}57'56''$ N, $19^{\circ}32'42''$ E, 125 m above mean sea level, a.m.s.l.), which is situated on the Great Hungarian Plain in a clearing within a mixed forest of coniferous (60 %) and deciduous trees (30 %) and some grassland (Salma et al., 2016b). The nearest city of Kecskemét (with 110 000 inhabitants) is situated ca. 15 km to the SE of K-pusztas. The station is part of the European monitoring and evaluation programme for the long-range transmission of air pollutants (EMEP network) and represents the largest plain (regional background) of the Carpathian Basin. One of the urban sites was at an open suburban area of residential Budapest at the Marczell György Main Observatory ($47^{\circ}25'46''$ N, $19^{\circ}10'54''$ E, 138 m a.m.s.l.) of the Hungarian Meteorological Service. The collections at the other urban locations were performed at the Budapest platform for Aerosol Research and Training (BpART) Laboratory ($47^{\circ}28'30''$ N, $19^{\circ}03'45''$ E, 115 m a.m.s.l.) of the Eötvös University. The latter site is situated on the bank of the Danube and represents a well-mixed average atmosphere of the city centre (Salma et al., 2016a). Some further details of the sampling campaign are summarised in Table 1.

The aerosol sampling was realised by three identical high-volume devices equipped with $\text{PM}_{2.5}$ inlets (DHA-80, Digitel, Switzerland). The collection substrates were quartz fibre filters with a diameter of 150 mm (QR-100, Advantec, Japan). Daily aerosol samples were taken starting at 00:00 LT (LT = UTC + 1 or daylight-saving time UTC + 2). The sampled air volumes were ca. 720 m³. One field blank sample was also taken at each site and month. All filters were pre-heated at 500 °C for 24 h before the exposure and were stored in a freezer after the collections.

Concentrations of criteria air pollutants, i.e. SO_2 , NO/NO_x , CO , O_3 and PM_{10} mass were obtained from regular stations of the National Air Quality Network. For the



Figure 1. The Carpathian Basin with location of the sampling sites in Budapest (city centre and suburban area) and at K-pusztas station (regional background). This image was created by © Mátyás Gede and used with his permission.

regional background and suburban area, they were measured directly at the sampling sites, while for the city centre, the pollutants were recorded in a distance of 4.5 km in the upwind prevailing direction from the sampling site. The concentrations are measured by UV fluorescence (Ysselbach 43C), chemiluminescence (Thermo 42C), IR absorption (Thermo 48i), UV absorption (Ysselbach 49C) and beta-ray attenuation methods (Thermo FH62-I-R), respectively with a time resolution of 1 h. Local meteorological data including air temperature (T) and relative humidity (RH), wind speed (WS) and global solar electromagnetic radiation (GRad) were acquired using standardised meteorological methods (Vaisala HMP45D humidity and temperature probe, Vaisala WAV15A anemometer, both Finland; CMP3 pyranometer, Kipp & Zonen, the Netherlands) near the sampling sites with a time resolution of 10 min. According to our knowledge, there were no extensive agricultural burns or wildfires in the basin during the actual sampling time intervals, and the BB in the basin is expected to be dominated by biofuel utilisation.

2.2 Analysis of aerosol samples

The PM mass was determined by weighing each filter before and after the sampling on a microbalance with a sensitivity of 10 µg (Cubis MSA225S-000-DA, Sartorius, Germany). The exposed and blank filters were pre-equilibrated before weighing at a T of 19–21 °C and RH of 45 %–50 % for at least 48 h. The measured mass data for the exposed filters were corrected for the field blank values considering the uncertainties in weighing, sampled air volume, RH and some other environmental conditions during the weighing as well. The PM mass data were above the limit of quantitation (LOQ), which was approximately 1 µg m^{−3}.

Table 1. Start and end dates of the sampling periods and number of aerosol samples collected in regional background of the Carpathian Basin, suburban area and city centre of Budapest.

Site type	Time interval: month year	Autumn month: October 2017	Winter month: January 2018	Spring month: April 2018	Summer month: July 2018
Region	Days	18–31	9–22	17–30	17–30
	Samples	14	14	14	14
Suburb	Days	18–31	6–22	17–30	17–1*
	Samples	14	17	14	14
Centre	Days	18–27	10–16	17–23	17–23
	Samples	7	7	7	7

* 1 August 2018.

From each of the filters one or two punches with an area of 1.5 cm^2 were directly analysed using the thermal–optical transmission (TOT) method (Birch and Cary, 1996) using a laboratory OC/EC analyser (Sunset Laboratory, USA) adopting the EUSAAR2 thermal protocol. The measured OC data for the exposed filters were corrected for the field blank values, while the EC on the blanks was negligible. All measured OC and EC data were above the LOQ, which was 0.38 and $0.04\text{ }\mu\text{g m}^{-3}$, respectively.

From each of the filters one or two sections with an area of 2.5 cm^2 were extracted in water, the extracts were filtered, and the filtrates were analysed for WSOC by a Vario TOC cube analyser (Elementar, Germany) in three repetitions with an injected volume of 1 mL each. The measured WSOC data for the exposed filters were corrected for the field blank values. All measured WSOC data were above the LOQ, which was ca. $0.08\text{ }\mu\text{g m}^{-3}$.

From each filter a section with an area of 2 cm^2 was analysed for LVG, MAN and GAN by gas chromatography coupled with mass spectrometry (GC/MS) after trimethylsilylation (Blumberger et al., 2019). The filter sections were extracted repeatedly by dichloromethane-methanol in an ultrasonic bath. The extracts were filtered and spiked with an internal standard (IS) of methyl β -L-arabinopyranoside. The trimethylsilylation was realised by hexamethyldisilazane as silylating agent, pyridine as solvent and trifluoroacetic acid as catalyst at $70\text{ }^\circ\text{C}$. The prepared samples were analysed by a Varian 4000 GC-MS/MS system (USA) with a GC/MS column of a SGE forte BPX5 capillary (length \times inner diameter $15\text{ m} \times 0.25\text{ mm}$; film thickness $0.25\text{ }\mu\text{m}$, SGE, Australia). The quantification was carried out in the selected ion monitoring mode by quantifier ions with mass-to-charge ratios of $m/z = 204$ for LVG and of 217 for MAN, GAN and IS. The LVG data for the exposed filters were corrected for blank values, while MAN and GAN were not detected in the blanks. The LVG amount in the blank filters can be related to the sampling itself, to chemicals used, to various chemical and sample preparation procedures performed, and to the variation of the baseline of the measurement (Maenhaut et

al., 2012). The blanks were the largest with respect to the corrected values in the summer samples, in which the measured LVG amounts were approximately 10 times larger than in the blank filters. In all the other samples, the relative contributions of the blanks were even smaller than this. The LOQ for LVG and MAN was approximately 1.2 ng m^{-3} , while it was approximately 0.5 ng m^{-3} for GAN. All LVG data were above the LOQ, while the MAN and GAN could not be quantified in the summer samples.

Filters collected in parallel on seven overlapping days, i.e. on 18–21, 25 and 26 October; on 10–16 January; on 17–23 April; and on 17–23 July were subjected to C isotope analysis of the TC content by accelerator mass spectrometry (AMS) with an off-line combustion system (Molnár et al., 2013; Janovics et al., 2018). Carbonaceous aerosol species in the eighth section of each filter were oxidised quantitatively to CO_2 gas (Major et al., 2018). This was later introduced into an IonPlus Enviro Mini Carbon Dating System spectrometer (Switzerland) via its dedicated gas ion source interface. The measured results for the exposed filters were corrected for the blank values. The $^{14}\text{C}/^{12}\text{C}$ ratios were also corrected for isotopic fractionation by using the $^{13}\text{C}/^{12}\text{C}$ ratios (Wacker et al., 2010) that were obtained simultaneously in the actual AMS measurements. The $^{14}\text{C}/^{12}\text{C}$ isotope ratios derived were also normalised to that of the oxalic acid II 4990C standard reference material (NIST, USA), and the measurement results were expressed as fraction of modern carbon (f_m), which denotes the $^{14}\text{C}/^{12}\text{C}$ ratio of the samples relative to that of the unperturbed atmosphere in the reference year of 1950 (Burr and Jull, 2009). Since the majority of currently combusted fuel wood was growing after the atmospheric nuclear fusion bomb tests in the late 1950s and early 1960s, the samples were also corrected by a mean factor of 1.08 derived for the Northern Hemisphere (Szidat et al., 2009; Heal et al., 2011). Thus, the fraction of contemporary carbon (f_c) was calculated as $f_c = f_m/1.08$. The same correction factor was also adopted for the TC from biogenic sources, although it is expected to show a somewhat smaller ^{14}C abundance. The differences in the f_c caused by the re-

finer correction factor are ordinarily small when compared to the method uncertainties (Minguillón et al., 2011) and, therefore, this effect was neglected.

A quarter section of each filter was utilised to determine the K (as a possible inorganic tracer for BB), Ni (as a possible tracer for residual oil combustion) and Pb (as a former tracer for vehicles with gasoline engine) content of the aerosol samples by inductively coupled plasma optical emission spectrometry using an iCAP7400 DUO instrument (Thermo Fischer Scientific, Germany). The filter sections were extracted by microwave-assisted HNO_3 – H_2O_2 digestion. The analytical results for the exposed filters were corrected for the blank values. The LOQ values of the elements listed were approximately $0.02\text{ }\mu\text{g m}^{-3}$, 0.4 ng m^{-3} and 0.5 ng m^{-3} , respectively, and most atmospheric concentrations were above them.

2.3 Data evaluation and modelling

Concentrations of organic matter (OM) were derived from the OC data by multiplying them with an organic-aerosol-to-organic-carbon mass conversion factor. This factor is an estimate of the average molecular mass per C atom for OM in general. It is site-dependent and can have seasonal and diurnal variations as well. It is usually derived by indirect considerations (Russell, 2003). Mass conversion factors between 1.2 and 1.4 were estimated for fine atmospheric aerosol in mildly oxidising atmospheric environments (Turpin et al., 2000). Some further studies suggest that a factor of 1.6 ± 0.2 better describes the oxidising urban environments or chemically aged (long-range transported) aerosol (Turpin and Lim, 2001). It should also be noted that the conversion factor is one of the most substantial sources of uncertainty in aerosol chemical mass closure calculations. It was estimated that the relative uncertainty associated with the conversion is approximately 30 % (Maenhaut et al., 2012). In the present study, a factor of 1.4 was adopted for the regional and suburban environments (considering that local or regional sources are mostly substantial or dominating in the Carpathian Basin), and a factor of 1.6 was utilised for the city centre.

The comparisons of atmospheric concentrations, other variables, or their ratios with respect to the months or sites were accomplished by calculating first the ratios on a sample-by-sample or day-by-day basis and then by averaging these individual ratios for the subset under consideration.

The coupled radiocarbon-LVG marker method was utilised to apportion the TC among the EC_{FF} , OC_{FF} , EC_{BB} , OC_{BB} and OC_{BIO} (Salma et al., 2017). The method consists of pragmatic attribution steps, which are realised by multiplications with apportionment factors. The factors are calculated for each sample from measured TC, f_c , EC, OC and LVG concentrations as primary input data and from a general, a priori known EC/OC ratio for BB $[(\text{EC}/\text{OC})_{\text{BB}}]$ and OC/LVG ratio for BB $[(\text{OC}/\text{LVG})_{\text{BB}}]$. Their combined adaptation is related to subsequent and step-wise subtraction

of contemporary TC, EC_{BB} and OC_{BB} from TC on the one hand, and of EC_{FF} from fossil TC on the other hand. The apportionment factors are expressed as follows: $f_1 = f_c$, $f_2 = (\text{OC}/\text{LVG})_{\text{BB}} \times \text{LVG} \times (\text{EC}/\text{OC})_{\text{BB}} / f_1 / \text{TC}$, $f_3 = (\text{OC}/\text{LVG})_{\text{BB}} \times \text{LVG} / f_1 / (1 - f_2) / \text{TC}$ and $f_4 = (\text{EC}/\text{TC} - f_1 \times f_2) / (1 - f_1)$ (Salma et al., 2017). For the $(\text{EC}/\text{OC})_{\text{BB}}$ ratio, we implemented a mean of 17 % derived from a critically evaluated ratio and standard deviation (SD) of $(16 \pm 5)\%$ (Szidat et al., 2006) and from a ratio and SD of $(18 \pm 4)\%$ (Bernardoni et al., 2011, 2013) obtained specifically for wood burning. As far as the $(\text{OC}/\text{LVG})_{\text{BB}}$ ratio is concerned, its actual value depends predominantly on the wood type and burning conditions (Puxbaum et al., 2007). We adopted an $(\text{OC}/\text{LVG})_{\text{BB}}$ ratio of 5.59 (Schmidl et al., 2008). The mean apportionment factors for the different months and site types are summarised in Table S1 in the Supplement. It is the OC_{BIO} and OC_{BB} which are the most sensitively influenced by the input uncertainties. Their relative uncertainty for some individual low concentrations could be up to 40 %–50 %, while it is expected to be approximately 30 % or smaller for the other carbonaceous species.

3 Results and discussion

The results of the study are interpreted in a conservative manner with regard to the months, but they are most likely representative with respect to the seasons as well. This expectation is based on the favourable meteorological conditions during the sample collections and the basin character of the region. General relationships that can exist among the months and atmospheric environments including coupled meteorological and chemical processes need to be surveyed before evaluating the temporal and spatial variability and tendencies in aerosol properties.

3.1 Differences and similarities among months and atmospheric environments

The median concentrations of SO_2 , NO, NO_2 and PM_{10} mass over the sampling time intervals were larger in the city centre than in the suburban area in all months (Table S2). This can mainly be explained by their anthropogenic sources in the city centre, mostly due to the increased intensity and density of road traffic. In contrast, the O_3 level was substantially higher in the suburban area than in the city centre and considerably larger in the regional background than in the suburban area. It tended to show a maximum in July. Such behaviour is typical for a large-scale O_3 formation mechanism. This all suggests that there could be substantial differences in photochemical activity in general, as indicated by O_3 between the regional background and the urban sites except for July. To access the extent of atmospheric dynamics during the sampling campaigns, the mean values and SDs of the same variables are shown in Table S3 as auxiliary data. The variability

ity of the concentrations partially supports the conclusions on the importance of regional meteorology within the basin (Figs. 2–4).

The meteorological data over the sampling time intervals are in accordance with ordinary monthly mean values and denote weather situations without extremes (Table S4). The T data indicate an urban heat island in central Budapest, particularly in winter and autumn. At the regional site, there was snow cover with a thickness from 2 to 4 cm during the sample collections in January for approximately 4 d, while in the Budapest area, there was snow in spots with a thickness of 1–2 cm for 2–3 d. The data suggest that there was somewhat milder weather over the sample collections in January than usually present.

Time series of $\text{PM}_{2.5}$ mass, EC and WSOC over the sampling time intervals in the different months and environments are shown in Figs. 2, 3 and 4, respectively. The $\text{PM}_{2.5}$ mass represents the bulk fine PM; EC is a typical primary aerosol constituent, while WSOC expresses mainly SOA and partially BB products (which also exhibit substantial hygroscopicity and thus water-solubility; Swietlicki et al., 2008). These species are rather different as far as their sources are concerned. Nevertheless, their atmospheric concentrations often changed coherently at the locations with their strongest link in winter. It can likely be explained by the common effects of regional meteorology on atmospheric concentrations especially under anticyclonic weather situations – particularly of boundary layer mixing height – over the Carpathian Basin. It seems that the daily evolution of regional meteorology often has a greater influence on the changes in concentrations than the source intensities if the sources are distributed across a large area (Salma et al., 2004). The strongest connection is related to cold air masses above the Carpathian Basin which generate a lasting T inversion layer (the so-called cold pillow) and which restricts the vertical mixing and results in poor air quality over extended areas of the basin in larger and smaller cities as well as in rural areas.

3.2 Tendencies in aerosol concentrations

Median atmospheric concentrations of the measured aerosol constituents separately for the different months and environmental types are presented in Table 2, while their means and SDs are shown in Table S5. The concentrations are in line with or somewhat smaller than the corresponding results obtained in earlier studies at the same or similar locations, usually for shorter time intervals (Salma and Maenhaut, 2006; Kiss et al., 2002; Salma et al., 2004, 2007, 2013, 2017; Ion et al., 2005; Maenhaut et al., 2005, 2008; Puxbaum et al., 2007; Blumberger et al., 2019). The $\text{PM}_{2.5}$ mass and OC concentrations in the city centre were larger by a mean factor of 1.6–1.7 than in the regional background, while they were similar to the suburban data. Their values in January were usually the largest, and they reached the minimum in July or April.




Figure 2. Time variation of $\text{PM}_{2.5}$ mass for regional background in the Carpathian Basin, suburban area and city centre of Budapest during the aerosol sampling time intervals for different months (a–d).




Figure 3. Time variation of EC for regional background in the Carpathian Basin, suburban area and city centre of Budapest during the aerosol sampling time intervals for different months (a–d).




Figure 4. Time variation of WSOC for regional background in the Carpathian Basin, suburban area and city centre of Budapest during the aerosol sampling time intervals for different months (a–d).

The concentrations of EC increased monotonically according to the environments: regional background, suburban area and city centre location by typical factors of 2 and 3, respectively. In the regional background, the EC data for October and January were similar to each other and they were the largest. In the suburban area, the EC data showed a maximum in January and a minimum in July. In the city centre, the EC levels in October, January and April were similar to each other, and they all showed a minimum in July. These can be explained by the larger intensity of soot emissions from incomplete burning (road vehicles, residential heating and cooking by solid fuel), which is typically of anthropogenic origin. These sources can have either seasonal variability (e.g. residential heating) or constant intensity (traffic or cooking) over a yearly time span.

The WSOC showed maximum medians in January at all sites. In October and July, the urban locations had similar concentrations to each other, while WSOC was somewhat smaller in the regional background. In January, the suburban site exhibited the maximum median concentration. This is explained by larger influence of BB in this environment and month and by the higher water solubility of its products (see Sect. 3.3 and 3.5). In April, the medians had a monotonically increasing tendency from the regional background to the city centre.

The mean atmospheric concentrations of the monosaccharide anhydrides were decreasing in the order of LVG, MAN and GAN. The concentrations of LVG were larger by ca. 1 order of magnitude than for the joint concentrations of MAN and GAN. Their mean ratio was the largest in January and the smallest in October. This could be affected by the share of hardwood burnt in different months (Fine et al., 2004; Schmidl et al., 2008; Maenhaut et al., 2012). The LVG concentration did not vary monotonically with respect to the sites; it was larger in the city centre by a factor of 1.7 than in the regional background and was smaller by approximately 20 % than in the suburban area. This could be related to the spatial distribution of biofuel utilisation (mainly for residential heating) and to atmospheric dispersion of their emission products in the different environments.

As far as the contemporary C is concerned, there were three individual consecutive samples collected in the city centre in October with significantly larger values than any other data in the set. There are several applications of nuclide ^{14}C , mostly in the pharmaceutical/medical and biological academy field in Budapest. They could release radiocarbon of anthropogenic origin into the ambient air (in particular from labelled inorganic compounds such as NaHCO_3). These three data were regarded to be outliers and were excluded from the further evaluation. The centre/suburb f_c ratio in October, however, still remained somewhat higher (1.15) with respect to the other months (for which the ratios were uniformly 1.00). This indicates that the anthropogenic ^{14}C contamination could slightly affect the remaining analytical results as well. Its consequences on the source apportionment are discussed in Sect. 3.5. In October and January, the mean centre/suburb, centre/region and suburb/region ratios were similar to each other (with an overall mean and SD of 1.02 ± 0.10) at all sites, while in April and July they decreased in the order of the ratios above, with means and SDs of 1.04 ± 0.20 , 0.82 ± 0.13 and 0.80 ± 0.15 , respectively. These tendencies are governed by carbonaceous matter of a different origin.

Table 2. Median atmospheric concentrations of $\text{PM}_{2.5}$ mass, elemental carbon (EC), organic carbon (OC), water-soluble organic carbon (WSOC), levoglucosan (LVG), mannosan (MAN), galactosan (GAN), fraction of contemporary total carbon (f_c), and median concentrations of K, Ni and Pb for regional background in the Carpathian Basin, suburban area and city centre of Budapest for different months representing different seasons.

Constituent	Site type	October	January	April	July
$\text{PM}_{2.5}$ mass ($\mu\text{g m}^{-3}$)	Region	12.5	16.5	8.6	10.7
	Suburb	25	26	9.7	11.7
	Centre	28	24	13.3	8.2
EC ($\mu\text{g m}^{-3}$)	Region	0.37	0.36	0.20	0.122
	Suburb	0.45	0.68	0.51	0.35
	Centre	0.99	0.77	0.79	0.37
OC ($\mu\text{g m}^{-3}$)	Region	2.3	3.2	2.0	2.2
	Suburb	4.5	5.4	2.4	2.7
	Centre	6.6	4.6	2.8	2.6
WSOC ($\mu\text{g m}^{-3}$)	Region	1.63	2.0	1.08	1.66
	Suburb	2.4	3.8	1.27	2.0
	Centre	2.7	2.8	1.45	2.0
LVG ($\mu\text{g m}^{-3}$)	Region	0.172	0.40	0.0180	0.0081
	Suburb	0.44	0.71	0.040	0.0124
	Centre	0.38	0.48	0.036	0.0103
MAN (ng m^{-3})	Region	19.2	18.5	2.6	< 1.2
	Suburb	37	39	2.9	< 1.2
	Centre	26	21	4.1	< 1.2
GAN (ng m^{-3})	Region	NA	10.6	1.20	< 0.5
	Suburb	16.4	20	0.61	< 0.5
	Centre	11.7	14.1	1.21	< 0.5
f_c (%)	Region	69	75	61	74
	Suburb	66	74	48	60
	Centre	76	74	48	60
K ($\mu\text{g m}^{-3}$)	Region	0.182	0.23	0.088	0.081
	Suburb	0.22	0.25	0.097	0.075
	Centre	0.26	0.27	0.106	0.057
Ni (ng m^{-3})	Region	0.75	0.68	1.21	1.12
	Suburb	0.88	0.78	1.24	1.09
	Centre	1.10	0.63	1.51	1.08
Pb (ng m^{-3})	Region	3.8	3.0	3.2	2.8
	Suburb	5.8	6.8	4.1	3.5
	Centre	7.5	5.2	4.4	2.3

NA – not available.

The concentrations of K in October, January and April increased monotonically for the regional background, suburb area and city centre. They showed a maximum in January. Its concentrations were the smallest in July and exhibited an opposite tendency as far as the location types are concerned; thus they decreased monotonically for the sites listed above. The concentrations of Ni were similar to each other without any evident tendency, except for its concentrations in April, which seemed to be the largest. The concentration of Pb showed an increasing tendency from the regional background to the urban sites. The present data are smaller than the median levels of 16 ng m^{-3} in the city centre and of 9 ng m^{-3} in the near-city background measured in spring 2002, after the phase-out of leaded gasoline in Hungary in April 1999, and are in line with its overall decreasing trend (Salma and Maenhaut, 2006; Salma et al., 2000).

3.3 Tendencies in concentration ratios

Mean values and SDs of some important concentration ratios separately for the months and different environments are shown in Table 3. The $\text{PM}_{2.5}/\text{PM}_{10}$ mass ratio exhibited strong time dependency. In April and July, the $\text{PM}_{10}-\text{PM}_{2.5}$ fraction particles (coarse mode) made up approximately two-thirds of the particulate mass, while in October and perhaps also in January, the $\text{PM}_{2.5}$ mass prevailed with a similar ratio. These imply and confirm that in spring and summer, the suspension or resuspension of soil, crustal rock, mineral and roadside dust is substantial in Budapest, while in autumn and winter, the aerosol mass levels are more influenced by residential heating, cooking and road traffic (Salma and Maenhaut, 2006).

The contribution of the OM to the $\text{PM}_{2.5}$ mass for the regional background, suburban area and city centre showed little time variation, with annual means and SDs of $(31 \pm 5) \%$, $(32 \pm 6) \%$ and $(35 \pm 7) \%$, respectively, in particular if we consider the uncertainty related to the OM/OC conversion (Sect. 2.3). These balanced contributions are in line with other European results (Puxbaum et al., 2007; Putaud et al., 2010). The mean contributions of EC to the $\text{PM}_{2.5}$ mass were between 1 % and 6 %, with a minimum in the regional background in July. The contributions can change substantially in different microenvironments within a city (e.g. 14 % for a street canyon in central Budapest in spring; Salma et al., 2004; Maenhaut et al., 2005). The carbonaceous particles (OM + EC) in the regional background, suburban area and city centre made up $(32 \pm 5) \%$, $(36 \pm 7) \%$ and $(39 \pm 7) \%$, respectively of the $\text{PM}_{2.5}$ mass as annual means and SDs. Their means revealed limited variability (except for the city centre, where it changed from 33% in January to 48 % in July). The $\text{TC}/\text{PM}_{2.5}$ mass ratios are given as auxiliary information to allow the recalculation of the contributions to the TC shown in Fig. 5 to that to the $\text{PM}_{2.5}$ mass.

The mean WSOC/OC ratios in October showed a monotonically increasing tendency from the city centre to the re-

Table 3. Mean values and SDs for the $\text{PM}_{2.5}/\text{PM}_{10}$ mass, OM/ $\text{PM}_{2.5}$ mass, EC/ $\text{PM}_{2.5}$ mass, TC/ $\text{PM}_{2.5}$ mass, WSOC/OC and OC/EC ratios for regional background in the Carpathian Basin, suburban area and city centre of Budapest for different months representing different seasons.

Ratio	Site type	October	January	April	July
$\text{PM}_{2.5}/\text{PM}_{10}$ mass (%)	Region	64 ± 10	NA	NA	NA
	Suburb	64 ± 4	67 ± 11	30 ± 8	48 ± 8
	Centre	67 ± 9	56 ± 14	32 ± 4	33 ± 7
OM/ $\text{PM}_{2.5}$ mass (%)	Region	33 ± 6	29 ± 5	32 ± 5	33 ± 4
	Suburb	32 ± 9	31 ± 5	32 ± 6	32 ± 5
	Centre	36 ± 5	30 ± 3	30 ± 4	36 ± 5
EC/ $\text{PM}_{2.5}$ mass (%)	Region	3.1 ± 1.4	2.3 ± 0.5	2.3 ± 0.7	1.2 ± 0.3
	Suburb	3.2 ± 1.0	3.1 ± 0.9	4.9 ± 2.2	3.5 ± 1.1
	Centre	4.3 ± 2.4	3.3 ± 0.6	6.4 ± 1.7	4.6 ± 0.9
TC/ $\text{PM}_{2.5}$ mass (%)	Region	28 ± 5	22 ± 4	25 ± 4	23 ± 3
	Suburb	24 ± 7	26 ± 4	27 ± 5	26 ± 4
	Centre	27 ± 5	22 ± 3	28 ± 5	29 ± 10
WSOC/OC (%)	Region	64 ± 11	58 ± 7	54 ± 9	72 ± 5
	Suburb	55 ± 16	64 ± 6	53 ± 7	76 ± 6
	Centre	42 ± 16	59 ± 4	56 ± 9	76 ± 11
OC/EC	Region	8.8 ± 3.2	9.0 ± 1.4	11 ± 3	18 ± 4
	Suburb	8.4 ± 3.5	7.3 ± 1.5	5.4 ± 2.0	7.3 ± 1.6
	Centre	6.3 ± 2.2	5.9 ± 0.7	3.5 ± 0.7	6.7 ± 1.5

gional background. This is just opposite to the atmospheric WSOC concentration (which decreased monotonically). In January, the suburban area exhibited the maximum concentration. This can be explained by intensive BB in the area with respect to the other environments (Sect. 3.5) and with the fact that BB particles possess relatively high hygroscopicity (Swietlicki et al., 2008) and water solubility. In the remaining two months, the shares of the WSOC were similar to each other and varied without an obvious tendency. This can be linked to comparable and large photochemical activity in all environments in April and July (Sect. 3.1). The present ratios are in line with the values reported earlier for the corresponding locations (Kiss et al., 2002; Ion et al., 2005; Maenhaut et al., 2005, 2008; Viana et al., 2006; Puxbaum et al., 2007; Salma et al., 2007). It is noted that the determined OC (and WSOC) concentrations are somewhat method dependent; their ratios can change sensitively, e.g. with the thermal protocol used in the OC/EC TOT analyser for samples containing large amounts of refractory C (Kuhlbusch et al., 2009; Panteliadis et al., 2015).

The highest OC/EC ratios are often linked to atmospheric conditions under which the SOA formation is large. The ratio had a maximum in the regional background in July, which can be associated with large photochemical activity and strong GRad. The ratios for the urban locations did not indicate obvious time tendencies. The formation, composition and properties of SOA and atmospheric humic-like substances (HULIS) together with modelling the air mass trans-

port within the Carpathian Basin are to be dealt with in a separate paper after additional investigations are completed.

Finally, it is noted for completeness that the annual mean LVG/MAN ratios and SDs for the regional background, suburban area and city centre were 13.9 ± 5.9 , 14.3 ± 6.2 and 14.7 ± 5.8 , respectively, and that ca. 40 % of all available individual ratios were larger than the limit of 14.8 derived by Schmidl et al. (2008). The latter value was obtained for the combustion of common hardwood (beech and oak) and softwood species (spruce and larch) in domestic wood stoves in Austria. This means for our samples and conditions, the relationship between the burnt softwood and hardwood mentioned is not applicable for several reasons, e.g. the likely differences in fireplaces and fuel wood in Hungary and mid-European Alpine regions.

3.4 Apportioned carbonaceous species

Median atmospheric concentrations of the apportioned EC_{FF} , EC_{BB} , OC_{FF} , OC_{BB} and OC_{BIO} aerosol constituents derived by the coupled radiocarbon-LVG model separately for the different months and environments are summarised in Table 4. The present values are coherent with the earlier median concentration from late winter to early spring of 2014 at the BpART Laboratory (Salma et al., 2017) and comparable to results for the regional background (Gelencsér et al., 2007; Puxbaum et al., 2007). A more sensible evaluation is to compare the contributions of the apportioned species to TC with other similar studies, which is completed in Sect. 3.5. The uncertainty of the individual apportioned data is larger than for the experimental results (e.g. TC) and, therefore, the substantial differences among their means and their obvious tendencies are only interpreted.

The median concentrations of EC_{FF} were similar to each other in October, April and perhaps in July as well and exhibited a minimum in January. In all months, its concentrations in the urban environments tended to be larger by a factor of 2–3 than in the regional background. The OC_{FF} concentrations at the urban locations were similar to each other in all months, while they tended to be larger than the regional values by a factor of 2–3 in October and July. The EC_{BB} and OC_{BB} concentrations showed a maximum in January and a minimum in July. The concentrations of OC_{BB} in the city centre seemed to be somewhat smaller than in the suburban area, while the latter was larger by a factor of 2–3 than in the regional background in October and April. The concentrations of OC_{BIO} showed a monotonically decreasing tendency from October to July, April and January in all environments. The fluxes of biogenic VOCs (BVOCs) from plants strongly depend on environmental conditions, age of leaves and vegetation, and water and nutrient availability, and it is also affected by the presence of some anthropogenic emissions. Photochemical oxidation reactions of BVOCs, interactions among biogenic and anthropogenic precursors and products, and aerosol formation yield considerations play a

Table 4. Median atmospheric concentration of apportioned elemental carbon from fossil fuel combustion (EC_{FF}) and from biomass burning (EC_{BB}), of apportioned organic carbon from fossil fuel combustion (OC_{FF}), from biomass burning (OC_{BB}) and from biogenic sources (OC_{BIO}) in $\mu\text{g m}^{-3}$ for regional background in the Carpathian Basin, suburban area and city centre of Budapest for different months representing different seasons.

Constituent	Site type	October	January	April	July
EC_{FF}	Region	0.35	0.057	0.23	0.12
	Suburb	0.35	0.10	0.57	0.32
	Centre	0.60	0.24	0.74	0.36
EC_{BB}	Region	0.19	0.34	0.020	0.0076
	Suburb	0.40	0.62	0.050	0.0083
	Centre	0.36	0.46	0.047	0.0095
OC_{FF}	Region	0.85	1.0	0.71	0.53
	Suburb	2.1	1.1	1.0	0.83
	Centre	1.5	1.2	1.0	0.81
OC_{BB}	Region	1.1	2.0	0.12	0.045
	Suburb	2.4	3.6	0.29	0.049
	Centre	2.1	2.7	0.27	0.056
OC_{BIO}	Region	2.0	0.22	1.3	1.8
	Suburb	2.3	0.36	1.2	1.8
	Centre	3.1	0.31	1.3	1.6

rather important role in the process (McFiggans et al., 2019). The tendencies are further discussed after deriving the contributions of the apportioned species to various quantities in Sect. 3.5.

Pearson's coefficients of correlation between the variables were calculated to examine their possible paired relationships. The results should be interpreted with caution since many data sets are not (fully) independent from each other and can be biased by meteorological processes (Sect. 3.1), can be coupled by their potential common sources, or can be influenced jointly by further factors or causes for them. Moreover, interactions among biogenic and anthropogenic VOCs or among organic precursors with rather different SOA yields can significantly enhance or suppress, respectively, the SOA production (Hoyle et al., 2011; McFiggans et al., 2019). Selected coefficients are shown in Table S6. Potassium correlated with both carbonaceous species of BB origin at all locations, while its coefficients with the other variables seemed insignificant (at a level of $p > 0.8$). There was a linear relationship between NO (60 %–70 % of which is emitted by road vehicles in Budapest) and OC_{FF} only in the suburban area. The relationships between T and the apportioned constituents indicated that BB was more intensive under cold weather conditions, while the utilisation of FFs was more constant over the year (campaign). No obvious consistent pattern was observed for FF carbonaceous species (and their contribution to the TC; see Sect. 3.5), which can suggest that




Figure 5. Mean contributions with SDs of elemental carbon from fossil fuel combustion (EC_{FF}) and from biomass burning (EC_{BB}), of organic carbon from fossil fuel combustion (OC_{FF}), from biomass burning (OC_{BB}) and from biogenic sources (OC_{BIO}) to PM_{2.5}-fraction total carbon (TC) in percentage for regional background in the Carpathian Basin, suburban area and city centre of Budapest and for different months representing different seasons. The median atmospheric concentrations of TC are indicated under individual circle charts, while the corresponding mean TC/PM_{2.5} mass ratios are shown in Table 3.

domestic heating is a minor source of OC_{FF} compared for instance to vehicular road traffic.

3.5 Contributions of source types

Fossil fuel combustion showed the most balanced and constant daily or monthly mean contributions to the TC at all sites and over the whole year. Its annual means and SDs for the regional background, suburban area and city centre were (31 ± 7) %, (36 ± 12) % and (36 ± 13) %, respectively. In contrast, the daily mean contributions of BB and biogenic sources changed radically over the year at all locations. For BB, the individual contributions for the atmospheric environments listed above ranged from < 2 % to 73 % (with a median of 10 %), from < 2 % to 73 % (24 %) and from < 2 % to 72 % (19 %), respectively. The analogous daily data for biogenic sources spanned from < 2 % up to 88 % (52 %), from < 2 % to 70 % (35 %) and from < 2 % to 67 % (39 %), respectively.

The monthly mean contributions of various EC and OC species to the TC separately for the different environmental types are shown in Fig. 5 as circle chart diagrams. In October, the three major source types contributed equally to the TC. In January, it was the BB which was the major source,

with a relative share of approximately 60 % at all sites, and its contribution was the largest in this month. The contributions of FF combustion in January were similar to each other for all sites, with a typical share around 25 %. The contributions of biogenic sources were the smallest in this month, although their firm interpretation is limited by the relatively large uncertainties. Their share might be linked to larger temperatures (urban heat island) and less snow coverage in the city centre (Sect. 3.1) than in its surroundings. In April, FF combustion and biogenic sources were the largest two contributors at all locations with typical shares of 45 %–50 % each. The EC_{FF} showed the largest contributions in April, which were increased monotonically according to the location type: region, suburb and centre. In July, biogenic sources became the major contributor, with a monotonically increasing share from the centre to the region.

The overall relative contributions are in good agreement with other similar or accompanying atmospheric studies (Szidat et al., 2006, 2009; Gelencsér et al., 2007; Minguillón et al., 2011; Bernardoni et al., 2013; Bonvalot et al., 2016).

Further conclusions can be derived by focusing on specific contributions of EC_{FF} and EC_{BB} to EC, and of OC_{FF}, OC_{BB} and OC_{BIO} to OC (Figs. 6 and 7, respectively). Elemental carbon is sometimes applied as a marker of automotive emis-




Figure 6. Distribution of mean contributions from FF combustion and BB with SDs to the PM_{2.5}-fraction EC for regional background in the Carpathian Basin, suburban area and city centre of Budapest for different months representing different seasons. The corresponding median atmospheric concentrations of the EC are given in Table 2.

sions mainly from diesel engines in cities of the continental mid-latitude Northern Hemisphere. The present research indicates that in urban ambient air in central Europe, this assumption is, however, valid only in April (spring) and July (summer; when the share of the EC_{FF} was indeed larger than 90 %). In October, the contributions of EC_{BB} can be considerable (up to 40 %–60 %) at urban sites, so they can be by no means negligible. Furthermore, in January, the relative mass of soot particles from BB can be even larger.

3.6 Potentials for air quality

To examine the potentials of the apportioned carbonaceous species for regulatory and legislation purposes, the contributions of the main source types to the PM_{2.5} mass were roughly estimated. It was assumed that the OM/OC conversion factors for the aerosol particles originating from FF




Figure 7. Distribution of mean contributions from FF combustion, BB and biogenic sources (BIO) with SDs to the PM_{2.5}-fraction OC for regional background in the Carpathian Basin, suburban area and city centre of Budapest for different months representing different seasons. The corresponding median atmospheric concentrations of OC are given in Table 2.

combustion, BB and biogenic sources were equal to the conversion factor for the bulk fine-fraction particles, thus 1.4 for the regional background and suburban area, and 1.6 for the city centre (see Sect. 2.3). We are aware that high emissions of some pyrogenic inorganic species such as K, nitrate or sulfate are neglected in the present apportionment model and that the OM/OC conversion factor can also change for organic species from different source types. The present estimates should, therefore, be considered as the first approximation only, and the contribution of BB to the PM_{2.5} mass is likely underestimated. The results obtained are summarised in Table S7. The separate contributions typically represent up to one-fifth or one-quarter of the PM mass as lower estimates and are discussed in Sect. 4. The contributions to the PM mass can be especially valuable when inspecting their tendencies.

The contributions were evaluated as a function of the $\text{PM}_{2.5}$ mass concentration, which is one of the key metrics for air quality considerations. The plots for the monthly mean contributions of OC_{FF} , OC_{BB} and OC_{BIO} to the TC are shown in Fig. 8a–c, respectively. The contributions of FF combustion (Fig. 8a) did not seem to depend substantially on the $\text{PM}_{2.5}$ mass level at any of the locations, so FF exhibits a constant and steady-state importance over various air pollution periods. The share of BB showed an increasing tendency with poor air quality (Fig. 8b). The change rate (the slope of the fitted line, b) was larger for the regional background ($b = 6.7$) and smaller but similar to each other for the two urban sites ($b \approx 2.1$). The trends for the biogenic sources were just the opposite (Fig. 8c); their relative importance decreased by poorer air quality. The tendency was similar again for the two urban sites ($b \approx -1.5$) and substantially larger for the regional background ($b = -6.9$). The tendencies of the EC_{FF} and EC_{BB} were analogous. These results together indicate that BB influences the air quality in the regional background very extensively and it also has a substantial effect on the air quality in the Budapest area, mainly in winter and possibly in the autumn months as well. The conclusions have importance and consequences for the potentials for improving the air quality, further interpreted in Sect. 4.

4 Conclusions

In the present study, the major carbonaceous aerosol species of OC and EC were apportioned among FF combustion, BB and biogenic sources in various types of atmospheric environments of interest in the Carpathian Basin in different months representing the different seasons. The research work is the first extensive adaptation of the coupled ^{14}C -LVG marker method developed recently. The experimental and derived data together with the results achieved were obtained from the first systematic complex research project in this region as far as the spatial scale within the basin and time span (of 1 full year) are concerned. The conclusions represent novel and valuable research contributions for a large area in Central Europe.

The carbonaceous particles made up from 30 % to 48 % of the $\text{PM}_{2.5}$ mass (as monthly mean) depending on the environment and months (representing different seasons). It is the BB in winter that represents the largest potential (with a mass share of > 20 %; Table S7) for improving the air quality both in cities and in rural areas of the basin. It is worth mentioning that all air pollution (smog) alert episodes in Hungary were announced so far exclusively because of the PM_{10} mass limit exceedances and they all happened in winter. Possibilities in controlling various forms of BB for air quality improvements seem to be, therefore, rather relevant due to this coincidence. In the present case for instance, there were 3, 8 and 8 d, respectively (19 d in total) in the subset of 4 × 7 d in the regional background, suburban area and city centre on which




Figure 8. Monthly mean contribution of apportioned OC_{FF} (a), OC_{BB} (b) and OC_{BIO} (c) to TC as a function of the monthly median $\text{PM}_{2.5}$ mass concentration (surrogate or proximity value for the air quality) for regional background in the Carpathian Basin, suburban area and city centre of Budapest. The months are marked by their starting letters. The fitted linear lines are just to guide the eye.

daily mean values exceeded the EU annual $\text{PM}_{2.5}$ limit value of $25 \mu\text{g m}^{-3}$. They all occurred in winter and autumn. If the BB sources (i.e. OC_{BB} and EC_{BB}) had decreased by half of their actual concentrations then the number of exceedance days would reduce to 2, 6 and 5, respectively (13 d in total, thus by 32 %), while a perfect fuel gas aftertreatment of the BB as a source would result in the number of exceedances

of 1, 4 and 5, respectively (10 d in total, thus by 47 %). In addition to carbonaceous particles, some adjunct inorganic constituents are also generated and, more importantly, soil or mineral dust and fly ash particles are also mobilised or blown up into the air due to the combustion or burning process itself. These, on the one hand, can further and substantially enhance the overall mass contributions and potentials of the high-temperature sources (including BB), and, on the other hand, may change somewhat their relative contributions.

The present findings and overall outcome of the research can be utilised as a background in modifying the municipal air quality regulations and action plans in Budapest, which are currently under preparation, and for inspiring the users of various household heat appliances for the societal implications of atmospheric aerosol. The apportioned contributions can also be used as a starting point for climate-related projects as far as both the regional issues in the Carpathian Basin, including the large-scale substitution of coal by forest biomass and the urban climate in Budapest, are concerned.

Fossil fuel combustion is an abundant source of PM mass (with a share of > 20 %; Table S7) only at urban sites and only in April and July. Resuspension or suspension of road and surface dust by moving vehicles can again represent a substantial auxiliary increment for FF contribution. Biogenic sources are normally considered as natural processes or as dominated by natural processes, and, therefore and strictly speaking, they are not associated with the issue of air pollution. It is expected that the unaccounted PM_{2.5} mass contains secondary inorganic aerosol particles, mostly sulfates, nitrates and elements, and soil or mineral and crustal rock dust particles as well. These constituents should definitely be revisited and taken into account in further source apportionment research.

Another challenge in health-related or air-quality-type assessment studies is to refine the apportionment within the major source types with burning of plastics, domestic or agricultural waste (garbage), coal, and stained wood in households through identification of their appropriate tracers and via quantification of various emission factors of their specific sources, e.g. using advanced coupled mass spectrometry or optical methods combined with powerful statistical data treatment. These additional combustion categories deserve more investigation since many of them seem to be prevalent and of increasing volume in the studied geographical area, and they produce some specific air pollutants or air toxics which can present serious risk to human health, well-being and the environment.

Data availability. Raw data are available from the corresponding author on reasonable request.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-20-4295-2020-supplement>.

Author contributions. IS designed the research study. AM performed most sample collections; AM and AVZ performed most analytical measurements and data treatment. TV, IM, VG and MM accomplished the isotope measurements and data treatment. IS and MM interpreted the results. IS prepared the paper with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The authors are grateful to Gergő Farkas, Veronika Varga, Debóra Varga and Péter Varga of the Eötvös University for their help in collecting the aerosol samples and their chemical analyses. The map in Fig. 1 was created by Mátyás Gede of the Eötvös University, Department of Cartography and Geoinformatics. The project was realised within the frame of the COST COLOSSAL Action (CA16109).

Financial support. This research has been supported by the National Research, Development and Innovation Office, Hungary (grant nos. K116788 and K132254), and the European Regional Development Fund and the Hungarian Government (grant nos. GINOP-2.3.2-15-2016-00028 and GINOP-2.3.2-15-2016-00009).

Review statement. This paper was edited by Nikolaos Mihalopoulos and reviewed by four anonymous referees.

References

- Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131–3148, <https://doi.org/10.5194/acp-6-3131-2006>, 2006.
- Andreae, M. O. and Rosenfeld, D.: Aerosol-cloud-precipitation interactions. Part 1. The nature and sources of cloud-active aerosols, *Earth-Sci. Rev.*, 89, 13–41, 2008.
- Artaxo, P., Rizzo, L. V., Paixao, M., de Lucca, S., Oliveira, P. H., Lara, L. L., Wiedemann, K. T., Andreae, M. O., Holben, B., Schafer, J., Correia, A. L., and Pauliquevis, T. M.: Aerosol particles in Amazonia: Their composition, role in the radiation balance, cloud formation, and nutrient cycles, in: *Amazonia and Global Change*, edited by: Keller, M., Bustamante, M., Gash, J., and Dias, P. S., 233–250, Geophysical Monograph Series, AGU, Washington, DC, 2009.
- Bernardoni, V., Vecchi, R., Valli, G., Piazzalunga, A., and Fermo, P.: PM₁₀ source apportionment in Milan (Italy) using time-resolved data, *Sci. Total Environ.*, 409, 4788–4795, 2011.
- Bernardoni, V., Calzolari, G., Chiari, M., Fedi, M., Lucarelli, F., Nava, S., Piazzalunga, A., Riccobono, F., Taccetti, F., Valli, G., and Vecchi, R.: Radiocarbon analysis on organic and elemental carbon in aerosol samples and source apportionment at an urban site in Northern Italy, *J. Aerosol Sci.*, 56, 88–99, 2013.

- Birch, M. E. and Cary, R. A.: Elemental carbon-based method for monitoring occupational exposures to particulate diesel exhaust, *Aerosol Sci. Tech.*, 25, 221–241, 1996.
- Blumberger, Z. I., Vasanits-Zsigrai, A., Farkas, G., and Salma, I.: Mass size distribution of major monosaccharide anhydrides and mass contribution of biomass burning, *Atmos. Res.*, 220, 1–9, 2019.
- Bonazza, A., Sabbioni, C., and Ghedini, N.: Quantitative data on carbon fractions in interpretation of black crusts and soiling on European built heritage, *Atmos. Environ.*, 39, 2607–2618, 2005.
- Bonvalot, L., Tuna, T., Fagault, Y., Jaffrezo, J.-L., Jacob, V., Chevrier, F., and Bard, E.: Estimating contributions from biomass burning, fossil fuel combustion, and biogenic carbon to carbonaceous aerosols in the Valley of Chamonix: a dual approach based on radiocarbon and levoglucosan, *Atmos. Chem. Phys.*, 16, 13753–13772, <https://doi.org/10.5194/acp-16-13753-2016>, 2016.
- Burnett, R., Chen, H., Szyszkowicz, M., Fann, N., Hubbell, B., Pope, C. A., Apte, J. S., Brauer, M., Cohen, A., Weichenthal, S., Coggin, J., Di, Q., Brunekreef, B., Frostad, J., Lim, S. S., Kan, H., Walker, K. D., Thurston, G. D., Hayes, R. B., Lim, C. C., Turner, M. C., Jerrett, M., Krewski, D., Gapstur, S. M., Diver, W. R., Ostro, B., Goldberg, D., Crouse, D. L., Martin, R. V., Peters, P., Pinault, L., Tjepkema, M., van Donkelaar, A., Villeneuve, P. J., Miller, A. B., Yin, P., Zhou, M., Wang, L., Janssen, N. A. H., Marra, M., Atkinson, R. W., Tsang, H., Quoc Thach, T., Cannon, J. B., Allen, R. T., Hart, J. E., Laden, F., Cesaroni, G., Forastiere, F., Weinmayr, G., Jaensch, A., Nagel, G., Concin, H., and Spadaro, J. V.: Global estimates of mortality associated with long-term exposure to outdoor fine particulate matter, *P. Natl. Acad. Sci. USA*, 115, 9592–9597, 2018.
- Burr, G. S. and Jull, A. J. T.: Accelerator mass spectrometry for radiocarbon research, in: *Encyclopedia of mass spectrometry*, edited by: Gross, M. L. and Caprioli, R., Elsevier, Amsterdam, 2009.
- Caseiro, A., Bauer, H., Schmidl, C., Pio, C. A., and Puxbaum, H.: Wood burning impact on PM₁₀ in three Austrian regions, *Atmos. Environ.*, 43, 2186–2195, 2009.
- Cecchini, M. A., Machado, L. A. T., Andreae, M. O., Martin, S. T., Albrecht, R. I., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Fütterer, D., Jurkat, T., Mahnke, C., Minikin, A., Molleker, S., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Voigt, C., Weinzierl, B., and Wendisch, M.: Sensitivities of Amazonian clouds to aerosols and updraft speed, *Atmos. Chem. Phys.*, 17, 10037–10050, <https://doi.org/10.5194/acp-17-10037-2017>, 2017.
- Chen, J., Li, Ch., Ristovski, Z., Milic, A., Gu, Y., Islam, M. S., Wang, S., Hao, J., Zhang, H., He, C., Guo, H., Fu, H., Miljevic, B., Morawska, L., Thai, P., Lam, Y. F., Pereira, G., Ding, A., Huang, X., and Dumka, U. C.: A review of biomass burning: Emissions and impacts on air quality, health and climate in China, *Sci. Total Environ.*, 579, 1000–1034, 2017.
- Cirino, G. G., Souza, R. A. F., Adams, D. K., and Artaxo, P.: The effect of atmospheric aerosol particles and clouds on net ecosystem exchange in the Amazon, *Atmos. Chem. Phys.*, 14, 6523–6543, <https://doi.org/10.5194/acp-14-6523-2014>, 2014.
- Claeys, M., Kourtchev, I., Pashynska, V., Vas, G., Vermeylen, R., Wang, W., Cafmeyer, J., Chi, X., Artaxo, P., Andreae, M. O., and Maenhaut, W.: Polar organic marker compounds in atmospheric aerosols during the LBA-SMOCC 2002 biomass burning experiment in Rondônia, Brazil: sources and source processes, time series, diel variations and size distributions, *Atmos. Chem. Phys.*, 10, 9319–9331, <https://doi.org/10.5194/acp-10-9319-2010>, 2010.
- Fabbri, D., Torri, C., Simoneit, B. R. T., Marynowski, L., Rushdi, A. I., and Fabiańska, M. J.: Levoglucosan and other cellulose and lignin markers in emissions from burning of Miocene lignites, *Atmos. Environ.*, 43, 2286–2295, 2009.
- Favez, O., El Haddad, I., Piot, C., Boréave, A., Abidi, E., Marchand, N., Jaffrezo, J.-L., Besombes, J.-L., Personnaz, M.-B., Sciare, J., Wortham, H., George, C., and D’Anna, B.: Inter-comparison of source apportionment models for the estimation of wood burning aerosols during wintertime in an Alpine city (Grenoble, France), *Atmos. Chem. Phys.*, 10, 5295–5314, <https://doi.org/10.5194/acp-10-5295-2010>, 2010.
- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the fireplace combustion of wood types grown in the Midwestern and Western United States, *Environ. Eng. Sci.*, 21, 387–409, 2004.
- Forello, A. C., Bernardoni, V., Calzolari, G., Lucarelli, F., Massabò, D., Nava, S., Pileci, R. E., Prati, P., Valentini, S., Valli, G., and Vecchi, R.: Exploiting multi-wavelength aerosol absorption coefficients in a multi-time resolution source apportionment study to retrieve source-dependent absorption parameters, *Atmos. Chem. Phys.*, 19, 11235–11252, <https://doi.org/10.5194/acp-19-11235-2019>, 2019.
- Fraser, M. P. and Lakshmanan, K.: Using levoglucosan as a molecular marker for the long range transport of biomass combustion aerosols, *Environ. Sci. Technol.*, 34, 4560–4564, 2000.
- Fuzzi, S., Baltensperger, U., Carslaw, K., Decesari, S., Denier van der Gon, H., Facchini, M. C., Fowler, D., Koren, I., Langford, B., Lohmann, U., Nemitz, E., Pandis, S., Riipinen, I., Rudich, Y., Schaap, M., Slowik, J. G., Spracklen, D. V., Vignati, E., Wild, M., Williams, M., and Gilardoni, S.: Particulate matter, air quality and climate: lessons learned and future needs, *Atmos. Chem. Phys.*, 15, 8217–8299, <https://doi.org/10.5194/acp-15-8217-2015>, 2015.
- Gelencsér, A., May, B., Simpson, D., Sánchez-Ochoa, A., Kasper-Giebl, A., Puxbaum, H., Caseiro, A., Pio, C., and Legrand, M.: Source apportionment of PM_{2.5} organic aerosol over Europe: primary/secondary, natural/anthropogenic, and fossil/biogenic origin, *J. Geophys. Res.*, 112, D23S04, <https://doi.org/10.1029/2006JD008094>, 2007.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, <https://doi.org/10.5194/acp-9-5155-2009>, 2009.
- Hays, M. D., Smith, N. D., Kinsey, J., Dong, Y., and Kariher, P.: Polycyclic aromatic hydrocarbon size distributions in aerosols from appliances of residential wood combustion as determined by direct thermal desorption-GC/MS, *J. Aerosol Sci.*, 34, 1061–1084, 2003.

- Heal, M. R., Naysmith, Ph., Cook, G. T. Xu, S., Duran, T. R., and Harrison, R. M.: Application of ^{14}C analyses to source apportionment of carbonaceous $\text{PM}_{2.5}$ in the UK, *Atmos. Environ.*, 45, 2341–2348, 2011.
- Hennigan, C. J., Sullivan, A. P., Collett Jr., J. L., and Robinson, A. L.: Levoglucosan stability in biomass burning particles exposed to hydroxyl radicals, *Geophys. Res. Lett.*, 37, L09806, <https://doi.org/10.1029/2010GL043088>, 2010.
- Hoffmann, D., Tilgner, A., Iinuma, Y., and Herrmann, H.: Atmospheric stability of levoglucosan: A detailed laboratory and modelling study, *Environ. Sci. Technol.*, 44, 694–699, 2010.
- Hopke, Ph. K.: Review of receptor modeling methods for source apportionment, *J. Air Waste Manage.*, 66, 237–259, 2016.
- Hoyle, C. R., Boy, M., Donahue, N. M., Fry, J. L., Glasius, M., Guenther, A., Hallar, A. G., Huff Hartz, K., Petters, M. D., Petäjä, T., Rosenoern, T., and Sullivan, A. P.: A review of the anthropogenic influence on biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 11, 321–343, <https://doi.org/10.5194/acp-11-321-2011>, 2011.
- Ion, A. C., Vermeylen, R., Kourtchev, I., Cafmeyer, J., Chi, X., Gelencsér, A., Maenhaut, W., and Claeys, M.: Polar organic compounds in rural $\text{PM}_{2.5}$ aerosols from K-puszt, Hungary, during a 2003 summer field campaign: Sources and diel variations, *Atmos. Chem. Phys.*, 5, 1805–1814, <https://doi.org/10.5194/acp-5-1805-2005>, 2005.
- Janovics, R., Futó, I., and Molnár, M.: Sealed tube combustion method with MnO_2 for AMS ^{14}C measurement, *Radiocarbon*, 60, 1347–1355, 2018.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, <https://doi.org/10.5194/acp-5-1053-2005>, 2005.
- Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J., Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A. D., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipilä, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533, 521–526, 2016.
- Kiss, G., Varga, B., Galambos, I., and Ganszky, I.: Characterization of water-soluble organic matter isolated from atmospheric fine aerosol, *J. Geophys. Res.*, 107, 8339, <https://doi.org/10.1029/2001JD000603>, 2002.
- Kourtchev, I., Hellebust, S., Bell, J. M., O'Connor, I. P., Healy, R. M., Allanic, A., Healy, D., Wenger, J. C., and Sodeau, J. R.: The use of polar organic compounds to estimate the contribution of domestic solid fuel combustion and biogenic sources to ambient levels of organic carbon and $\text{PM}_{2.5}$ in Cork Harbour, Ireland, *Sci. Total Environ.*, 409, 2143–2155, 2011.
- Kuhlbusch, Th. A. J., Borowiak, A., Gelencsér, A., Genberg, J., Gladtko, D., Maenhaut, W., Pio, C., Popovicheva, O., Putaud, J. P., Quincey, P., Sciare, J., ten Brink, H., Viana, M., and Yttri, K. E.: Measurement of elemental and organic carbon in Europe, JRC Scientific and Technical Reports, European Commission, Joint Research Centre, Ispra, 2009.
- Lelieveld, J. and Pöschl, U.: Chemists can help to solve the air-pollution health crisis, *Nature*, 551, 291–293, 2017.
- Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Arneeth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Doney, S. C., Gkritzalis, T., Goll, D. S., Harris, I., Haverd, V., Hoffman, F. M., Hoppema, M., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Johannessen, T., Jones, C. D., Kato, E., Keeling, R. F., Goldewijk, K. K., Landschützer, P., Lefèvre, N., Lienert, S., Liu, Z., Lombardozzi, D., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., Neill, C., Olsen, A., Ono, T., Patra, P., Peregon, A., Peters, W., Peylin, P., Pfeil, B., Pierrot, D., Poulter, B., Rehdner, G., Resplandy, L., Robertson, E., Rocher, M., Rödenbeck, C., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Steinhoff, T., Sutton, A., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., Wright, R., Zaehle, S., and Zheng, B.: Global Carbon Budget 2018, *Earth Syst. Sci. Data*, 10, 2141–2194, <https://doi.org/10.5194/essd-10-2141-2018>, 2018.
- Lohmann, U., Feichter, J., Penner, J., and Leaitch, R.: Indirect effect of sulfate and carbonaceous aerosols: A mechanistic treatment, *J. Geophys. Res.*, 105, 12193–12206, 2000.
- Maenhaut, W., Raes, N., Chi, X., Cafmeyer, J., Wang, W., and Salma, I.: Chemical composition and mass closure for fine and coarse aerosols at a kerbside in Budapest, Hungary, in spring 2002, *X-ray Spectrom.*, 34, 290–296, 2005.
- Maenhaut, W., Raes, N., Chi, X., Cafmeyer, J., and Wang, W.: Chemical composition and mass closure for $\text{PM}_{2.5}$ and PM_{10} aerosols at K-puszt, Hungary, in summer 2006, *X-ray Spectrom.*, 37, 193–197, 2008.
- Maenhaut, W., Vermeylen, R., Claeys, M., Vercauteren, J., Mathieu, C., and Roekens, E.: Assessment of the contribution from wood burning to the PM_{10} aerosol in Flanders, Belgium, *Sci. Total Environ.*, 437, 226–236, 2012.
- Maenhaut, W., Vermeylen, R., Claeys, M., Vercauteren, J., and Roekens, E.: Sources of the PM_{10} aerosol in Flanders, Belgium, and re-assessment of the contribution from wood burning, *Sci. Total Environ.*, 562, 550–560, 2016.
- Major, I., Gyökös, B., Túri, M., Futó, I., Filep, Á., Hoffer, A., Furu, E., Jull, A. J. T., and Molnár, M.: Evaluation of an automated EA-IRMS method for total carbon analysis of atmospheric aerosol at HEKAL, *J. Atmos. Chem.*, 75, 85–96, 2018.
- McFiggans, G., Mentel, T. F., Wildt, J., Pullinen, I., Kang, S., Kleist, E., Schmitt, S., Springer, M., Tillmann, R., Wu, C., Zhao, D., Hallquist, M., Faxon, C., Le Breton, M., Hallquist, A. M., Simpson, D., Bergstroem, R., Jenkin, M. E., Ehn, M., Thornton, J.

- A., Alfara, M. R., Bannan, T. J., Percival, C. J., Priestley, M., Topping, D., and Kiendler-Scharr, A.: Secondary organic aerosol reduced by mixture of atmospheric vapours, *Nature*, 565, 587–593, 2019.
- Minguillón, M. C., Perron, N., Querol, X., Szidat, S., Fahrni, S. M., Alastuey, A., Jimenez, J. L., Mohr, C., Ortega, A. M., Day, D. A., Lanz, V. A., Wacker, L., Reche, C., Cusack, M., Amato, F., Kiss, G., Hoffer, A., Decesari, S., Moretti, F., Hillamo, R., Teinilä, K., Seco, R., Peñuelas, J., Metzger, A., Schallhart, S., Müller, M., Hansel, A., Burkhardt, J. F., Baltensperger, U., and Prévôt, A. S. H.: Fossil versus contemporary sources of fine elemental and organic carbonaceous particulate matter during the DAURE campaign in Northeast Spain, *Atmos. Chem. Phys.*, 11, 12067–12084, <https://doi.org/10.5194/acp-11-12067-2011>, 2011.
- Molnár, M., Rinyu, L., Veres, M., Seiler, M., Wacker, L., and Synal, H.-A.: EnvironMICADAS: a mini ^{14}C AMS with enhanced gas ion source interface in the Hertelendi Laboratory of Environmental Studies, Hungary, *Radiocarbon*, 55, 338–344, 2013.
- Nolte, C. G., Schauer, J. J., Cass, G. R., and Simoneit, B. R. T.: Highly polar organic compounds present in wood smoke and in the ambient atmosphere, *Environ. Sci. Technol.*, 35, 1912–1919, 2001.
- Nozière, B., Kalberer, M., Claeys, M., Allan, J., D’Anna, B., Decesari, S., Finessi, E., Glasius, M., Grgić, I., Hamilton, J. F., Hoffmann, T., Iinuma, Y., Jaoui, M., Kahnt, A., Kampf, C. J., Kourtchev, I., Maenhaut, W., Marsden, N., Saarikoski, S., Schnelle-Kreis, J., Surratt, J. D., Szidat, S., Szmigielski, R., and Wisthaler, A.: The molecular identification of organic compounds in the atmosphere: state of the art and challenges, *Chem. Rev.*, 115, 3919–3983, 2015.
- Panteliadis, P., Hafkenscheid, T., Cary, B., Diapouli, E., Fischer, A., Favez, O., Quincey, P., Viana, M., Hitzenberger, R., Vecchi, R., Saraga, D., Sciare, J., Jaffrezzo, J. L., John, A., Schwarz, J., Giannoni, M., Novak, J., Karanasiou, A., Fermo, P., and Maenhaut, W.: ECOC comparison exercise with identical thermal protocols after temperature offset correction – instrument diagnostics by in-depth evaluation of operational parameters, *Atmos. Meas. Tech.*, 8, 779–792, <https://doi.org/10.5194/amt-8-779-2015>, 2015.
- Pashynska, V., Vermeylen, R., Vas, G., Maenhaut, W., and Claeys, M.: Development of a gas chromatography/ion trap mass spectrometry method for the determination of levoglucosan and saccharidic compounds in atmospheric aerosols, Application to urban aerosols, *J. Mass Spectrom.*, 37, 1249–125, 2002.
- Piazzalunga, A., Belis, C., Bernardoni, V., Cazzuli, O., Fermo, P., Valli, G., and Vecchi, R.: Estimates of wood burning contribution to PM by the macro-tracer method using tailored emission factors, *Atmos. Environ.*, 45, 6642–6649, 2011.
- Putaud, J.-P., Van Dingenen, R., Alastuey, A., Bauer, H., Birmili, W., Cyrys, J., Flentje, H., Fuzzi, S., Gehrig, R., Hansson, H. C., Harrison, R. M., Herrmann, H., Hitzenberger, R., Hügl, C., Jones, A. M., Kasper-Giebl, A., Kiss, G., Kousa, A., Kuhlbusch, T. A. J., Löschau, G., Maenhaut, W., Molnár, A., Moreno, T., Pekkanen, J., Perrino, C., Pitz, M., Puxbaum, H., Querol, X., Rodriguez, S., Salma, I., Schwarz, J., Smolik, J., Schneider, J., Spindler, G., ten Brink, H., Tursic, J., Viana, M., Wiedensohler, A., and Raes, F.: A European aerosol phenomenology – 3: physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe, *Atmos. Environ.*, 44, 1308–1320, 2010.
- Puxbaum, H., Caseiro, A., Sanchez-Ochoa, A., Kasper-Giebl, A., Claeys, M., Gelencsér, A., Legrand, M., Preunkert, S., and Pio, C.: Levoglucosan levels at background sites in Europe for assessing the impact of biomass combustion on the European aerosol background, *J. Geophys. Res.-Atmos.*, 112, D23S05, <https://doi.org/10.1029/2006JD008114>, 2007.
- Rap, A., Spracklen, D. V., Mercado, L., Reddington, C. L., Haywood, J. M., Ellis, R. J., Phillips, O. L., Artaxo, P., Bonal, D., Coupe, N. R., and Butt, N.: Fires increase Amazon forest productivity through increases in diffuse radiation, *Geophys. Res. Lett.*, 42, 4654–4662, 2015.
- Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T., and Yu, S.: Aerosol-driven droplet concentrations dominate coverage and water of oceanic low-level clouds, *Science*, 363, eaav0566, <https://doi.org/10.1126/science.aav0566>, 2019.
- Russell, L. M.: Aerosol organic-mass-to-organic-carbon ratio measurements, *Environ. Sci. Technol.*, 37, 2982–2987, 2003.
- Saarikoski, S., Timonen, H., Saarnio, K., Aurela, M., Järvi, L., Keronen, P., Kerminen, V.-M., and Hillamo, R.: Sources of organic carbon in fine particulate matter in northern European urban air, *Atmos. Chem. Phys.*, 8, 6281–6295, <https://doi.org/10.5194/acp-8-6281-2008>, 2008.
- Salma, I. and Maenhaut, W.: Changes in chemical composition and mass of atmospheric aerosol pollution between 1996 and 2002 in a Central European city, *Environ. Pollut.*, 143, 479–488, 2006.
- Salma, I., Maenhaut, W., Dubtsov, S., Zemplén-Papp, É., and Zárny, Gy.: Impact of phase out of leaded gasoline on the air quality in Budapest, *Microchem. J.*, 67, 127–133, 2000.
- Salma, I., Chi, X., and Maenhaut, W.: Elemental and organic carbon in urban canyon and background environments in Budapest, Hungary, *Atmos. Environ.*, 38, 27–36, 2004.
- Salma, I., Ocskay, R., Chi, X., and Maenhaut, W.: Sampling artefacts, concentrations and chemical composition of fine water-soluble organic carbon and humic-like substances in a continental urban atmospheric environment, *Atmos. Environ.*, 41, 4106–4118, 2007.
- Salma, I., Mészáros, T., and Maenhaut, W.: Mass size distribution of carbon in atmospheric humic-like substances and water-soluble organic carbon for an urban environment, *J. Aerosol Sci.*, 56, 53–60, 2013.
- Salma, I., Németh, Z., Weidinger, T., Kovács, B., and Kristóf, G.: Measurement, growth types and shrinkage of newly formed aerosol particles at an urban research platform, *Atmos. Chem. Phys.*, 16, 7837–7851, <https://doi.org/10.5194/acp-16-7837-2016>, 2016a.
- Salma, I., Németh, Z., Kerminen, V. M., Aalto, P., Nieminen, T., Weidinger, T., Molnár, Á., Imre, K., and Kulmala, M.: Regional effect on urban atmospheric nucleation, *Atmos. Chem. Phys.*, 16, 8715–8728, 2016b.
- Salma, I., Németh, Z., Weidinger, T., Maenhaut, W., Claeys, M., Molnár, M., Major, I., Ajtai, T., Utry, N., and Bozóki, Z.: Source apportionment of carbonaceous chemical species to fossil fuel combustion, biomass burning and biogenic emissions by a coupled radiocarbon–levoglucosan marker method, *Atmos. Chem. Phys.*, 17, 13767–13781, <https://doi.org/10.5194/acp-17-13767-2017>, 2017.

- Sandradewi, J., Prévôt, A. S. H., Szidat, S., Perron, N., Alfara, R. M., Lanz, V. A., Weingartner, E., and Baltensperger, U.: Using aerosol light absorption measurements for the quantitative determination of wood burning and traffic emission contributions to particulate matter, *Environ. Sci. Technol.*, 42, 3316–3323, 2008a.
- Sandradewi, J., Prévôt, A. S. H., Weingartner, E., Schmidhauser, R., Gysel, M., and Baltensperger, U.: A study of wood burning and traffic aerosols in an Alpine valley using a multi-wavelength Aethalometer, *Atmos. Environ.*, 42, 101–112, 2008b.
- Schmidl, C., Marr, L. L., Caseiro, A., Kotianova, P., Berner, A., Bauer, H., Kasper-Giebl, A., and Puxbaum, H.: Chemical characterisation of fine particle emissions from wood stove combustion of common woods growing in mid-European Alpine regions, *Atmos. Environ.*, 42, 126–141, 2008.
- Simoneit, B. R. T., Schauer, J. J., Nolte, C. G., Oros, D. R., Elias, V. O., Fraser, M. P., Rogge, W. F., and Cass, G. R.: Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles, *Atmos. Environ.*, 33, 173–182, 1999.
- Simoneit, B. R. T., Elias, V. O., Kobayashi, M., Kawamura, K., Rushdi, A. I., Medeiros, P. M., Rogge, W. F., and Didyk, B. M.: Sugars-dominant water-soluble organic compounds in soils and characterization as tracers in atmospheric particulate matter, *Environ. Sci. Technol.*, 38, 5939–5949, 2004.
- Stefenelli, G., Pospisilova, V., Lopez-Hilfiker, F. D., Daellenbach, K. R., Hüglin, C., Tong, Y., Baltensperger, U., Prévôt, A. S. H., and Slowik, J. G.: Organic aerosol source apportionment in Zurich using an extractive electrospray ionization time-of-flight mass spectrometer (EESI-TOF-MS) – Part I: Biogenic influences and day–night chemistry in summer, *Atmos. Chem. Phys.*, 19, 14825–14848, <https://doi.org/10.5194/acp-19-14825-2019>, 2019.
- Swietlicki, E., Hansson, H. C., Hämeri, K., Svenningsson, B., Massling, A., McFiggans, G., McMurry, P. H., Petäjä, T., Tunved, P., Gysel, M., Topping, D., Weingartner, E., Baltensperger, U., Rissler, J., Wiedensohler, A., and Kulmala, M.: Hygroscopic properties of submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various environments – a review, *Tellus B*, 60, 432–469, 2008.
- Szidat, S., Jenk, T. M., Synal, H. A., Kalberer, M., Wacker, L., Hajdas, I., Kasper-Giebl, A., and Baltensperger, U.: Contributions of fossil fuel, biomass-burning, and biogenic emissions to carbonaceous aerosols in Zurich as traced by ^{14}C , *J. Geophys. Res.*, 111, D07206, <https://doi.org/10.1029/2005JD006590>, 2006.
- Szidat, S., Ruff, M., Perron, N., Wacker, L., Synal, H.-A., Hallquist, M., Shannigrahi, A. S., Yttri, K. E., Dye, C., and Simpson, D.: Fossil and non-fossil sources of organic carbon (OC) and elemental carbon (EC) in Göteborg, Sweden, *Atmos. Chem. Phys.*, 9, 1521–1535, <https://doi.org/10.5194/acp-9-1521-2009>, 2009.
- Tatai, Zs., Szőke, B., and Körmendi, K.: Budapest zöldinfrastruktúra koncepciója (Concept on green infrastructure of Budapest), Municipality of Budapest, BFTV Kft., Budapest, 2017 (in Hungarian).
- Tian, H., Lu, C., Ciais, P., Michalak, A. M., Canadell, J. G., Saikawa, E., Huntzinger, D. N., Gurney, K. R., Sitch, S., Zhang, B., Yang, J., Bousquet, P., Bruhwiler, L., Chen, G., Dlugokencky, E., Friedlingstein, P., Melillo, J., Pan, S., Poulter, B., Prinn, R., Saunio, M., Schwalm, C. R., and Wofsy, S. C.: The terrestrial biosphere as a net source of greenhouse gases to the atmosphere, *Nature*, 531, 225–228, 2016.
- Turpin, B. J. and Lim, H.-J.: Species contributions to $\text{PM}_{2.5}$ mass concentrations: revisiting common assumptions for estimating organic mass, *Aerosol Sci. Tech.*, 35, 602–610, 2001.
- Turpin, B. J., Saxena, P., and Andrews, E.: Measuring and simulating particulate organics in the atmosphere: problems and prospects, *Atmos. Environ.*, 34, 2983–3013, 2000.
- Viana, M., Chi, X., Maenhaut, W., Cafmeyer, J., Querol, X., Alastuey, A., Mikuška, P., and Večera, Z.: Influence of sampling artefacts on measured PM, OC, and EC levels in carbonaceous aerosols in an urban area, *Aerosol Sci. Tech.*, 40, 107–117, 2006.
- Vicente, E. D. and Alves, C. A.: An overview of particulate emissions from residential biomass combustion, *Atmos. Res.*, 199, 159–185, 2018.
- Von Schneidmesser, E., Monks, P. S., Allan, J. D., Bruhwiler, L., Forster, P., Fowler, D., Lauer, A., Morgan, W. T., Paasonen, P., Righi, M., Sindelarova, K., and Sutton, M. A.: Chemistry and the linkages between air quality and climate change, *Chem. Rev.*, 115, 3856–3897, 2015.
- Wacker, L., Bonani, G., Friedrich, M., Hajdas, I., Kromer, B., Némec, M., Ruff, M., Suter, M., Synal, H.-A., and Vockenhuber, C.: MICADAS: Routine and high-precision radiocarbon dating, *Radiocarbon*, 52, 252–262, 2010.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625–641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.
- Yttri, K. E., Schnelle-Kreis, J., Maenhaut, W., Abbaszade, G., Alves, C., Bjerke, A., Bonnier, N., Bossi, R., Claeys, M., Dye, C., Evtyugina, M., García-Gacio, D., Hillamo, R., Hoffer, A., Hyder, M., Iinuma, Y., Jaffrezo, J.-L., Kasper-Giebl, A., Kiss, G., López-Mahia, P. L., Pio, C., Piot, C., Ramirez-Santa-Cruz, C., Sciare, J., Teinilä, K., Vermeylen, R., Vicente, A., and Zimmermann, R.: An intercomparison study of analytical methods used for quantification of levoglucosan in ambient aerosol filter samples, *Atmos. Meas. Tech.*, 8, 125–147, <https://doi.org/10.5194/amt-8-125-2015>, 2015.
- Zdráhal, Z., Oliveira, J., Vermeylen, R., Claeys, M., and Maenhaut, W.: Improved method for quantifying levoglucosan and related monosaccharide anhydrides in atmospheric aerosols and application to samples from urban and tropical locations, *Environ. Sci. Technol.*, 36, 747–753, 2002.
- Zhang, Y. L., Perron, N., Ciobanu, V. G., Zotter, P., Minguilón, M. C., Wacker, L., Prévôt, A. S. H., Baltensperger, U., and Szidat, S.: On the isolation of OC and EC and the optimal strategy of radiocarbon-based source apportionment of carbonaceous aerosols, *Atmos. Chem. Phys.*, 12, 10841–10856, <https://doi.org/10.5194/acp-12-10841-2012>, 2012.
- Zotter, P., Herich, H., Gysel, M., El-Haddad, I., Zhang, Y., Močnik, G., Hüglin, C., Baltensperger, U., Szidat, S., and Prévôt, A. S. H.: Evaluation of the absorption Ångström exponents for traffic and wood burning in the Aethalometer-based source apportionment using radiocarbon measurements of ambient aerosol, *Atmos. Chem. Phys.*, 17, 4229–4249, <https://doi.org/10.5194/acp-17-4229-2017>, 2017.