

Supplement of Atmos. Chem. Phys., 20, 8939–8951, 2020
<https://doi.org/10.5194/acp-20-8939-2020-supplement>
© Author(s) 2020. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Atmospheric fate of two relevant unsaturated ketoethers: kinetics, products and mechanisms for the reaction of hydroxyl radicals with (*E*)-4-methoxy-3-buten-2-one and (1*E*)-1-methoxy-2-methyl-1-penten-3-one

Rodrigo Gastón Gibilisco et al.

Correspondence to: Rodrigo Gastón Gibilisco (gibilisco@uni-wuppertal.de) and Iustinian Gabriel Bejan (iustinian.bejan@uaic.ro)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Supplementary Information

Content Summary

Fig. S1: Residual spectrum obtained on the TMBO + OH radical reaction (Fig. 3 trace H) focused in the frequency region between 3400 cm^{-1} and 3800 cm^{-1} .

Fig. S2: Concentrations of carbonyls formed against reacted TMBO in the reaction with OH radicals.

Fig. S3: Residual spectrum obtained in the MMPO + OH reaction (Fig. 4 trace H) in the frequency region between 3400 cm^{-1} and 3800 cm^{-1} .

Fig. S4: Concentrations of carbonyls formed against reacted MMPO in the reaction with OH radicals.




Figure S1. Residual spectrum obtained in the TMBO + OH reaction (Fig. 3 trace H) in the frequency region between 3400 cm⁻¹ and 3800 cm⁻¹.




Figure S2. Example of the concentrations of products formed against reacted TMBO in the reaction with OH radicals.




Figure S3. Residual spectrum obtained in the MMPO + OH reaction (Fig. 4 trace H) in the frequency region between 3400 cm^{-1} and 3800 cm^{-1} .




Figure S4. Example of the concentrations of products formed against reacted MMPO in the reaction with OH radicals.