



Supplement of

Measurement report: Observation-based formaldehyde production rates and their relation to OH reactivity around the Arabian Peninsula

Dirk Dienhart et al.

Correspondence to: Dirk Dienhart (d.dienhart@mpic.de) and Horst Fischer (horst.fischer@mpic.de)

The copyright of individual parts of the supplement might differ from the article licence.




Figure S1: Timelines of the contribution of inorganic compounds (NO , NO_2 , SO_2 , O_3) and further reactants (CO , HCHO) to the OH reactivity $\text{R}(\text{OH})$. Removal of the non- HCHO yielding reactions resulted in the so-called effective OH reactivity ($\text{R}(\text{OH})_{\text{eff}}$).

5 **Table S1:** Trace gases used for removal of the non- HCHO yielding reactions from $\text{R}(\text{OH})$ to calculate the effective OH reactivity ($\text{R}(\text{OH})_{\text{eff}}$) with IUPAC preferred values for their reaction constants.

Compound	$k_{M+\text{OH}}$ [cm 3 molecules $^{-1}\text{s}^{-1}$]	Max. OH reactivity [s $^{-1}$] (Pfannerstill et al. 2019)
NO	9.70E-12	29.3
NO_2	9.80E-12	6.94
O_3	7.30E-14	0.26
SO_2	9.30E-13	0.33
CO	1.44E-13	1.16
HCHO	8.50E-12	2.33




Figure S2: Scatter plots with bivariate fits (York et al., 2004) of the product $[\text{OH}] \times R(\text{OH})$ versus the HCHO production rate, subdivided into the different regions probed during the AQABA cruise.

Table S2: Chemical compounds used for calculating the speciated OH reactivities ($R(OH)_{\text{alkanes}}$, $R(OH)_{\text{alkenes}}$, $R(OH)_{\text{OVOCs}}$, $R(OH)_{\text{aromatics}}$) based on Pfannerstill et al. (2019):

Compound/ protonated mass (grey shaded compounds are gas standard calibrated with low uncertainty)	Chem- ical formula	k_{M+OH} [cm ³ molecules ⁻¹ s ⁻¹]	Instrument	Compounds attributed to protonated mass	If no unambiguous compound identification: reaction rate coefficient derived from...	Literature reference for reaction rate coefficient	Max. OH reactivity during AQABA [s ⁻¹]	Comment
Alkanes								
Methane	CH ₄	6.45E-15	Picarro (Cavity ring- down spectroscopy)			IUPAC preferred value	0.35	
Ethane	C ₂ H ₆	2.40E-13	GC-FID			IUPAC preferred value	0.26	
Propane	C ₃ H ₈	1.10E-12	GC-FID			IUPAC preferred value	1.33	
2-Methylpropane	C ₄ H ₁₀	2.12E-12	GC-FID			Atkinson and Arey, 2003	0.68	
n-Butane	C ₄ H ₁₀	2.35E-12	GC-FID			IUPAC preferred value	1.42	
2-Methylbutane	C ₅ H ₁₂	3.60E-12	GC-FID			Atkinson and Arey, 2003	0.92	
n-pentane	C ₅ H ₁₂	3.80E-12	GC-FID			Atkinson and Arey, 2003	1.03	
2-Methylpentane	C ₆ H ₁₄	5.20E-12	GC-FID			Atkinson and Arey, 2003	0.11	
n-hexane	C ₆ H ₁₄	5.20E-12	GC-FID			Atkinson and Arey, 2003	0.16	
2,2,4- Trimethylpentane	C ₈ H ₁₈	3.34E-12	GC-FID			Atkinson and Arey, 2003	0.004	
n-heptane	C ₇ H ₁₆	6.76E-12	GC-FID			Atkinson and Arey, 2003	0.06	
Octane	C ₈ H ₁₈	8.11E-12	GC-FID			Atkinson and Arey, 2003	0.04	
Alkenes and Alkynes								
Ethene	C ₂ H ₄	7.50E-12	GC-FID			IUPAC preferred value	2.26	
Propene	C ₃ H ₆	2.90E-11	GC-FID			IUPAC preferred value	1.11	
trans-2-butene	C ₄ H ₈	6.40E-11	GC-FID			IUPAC preferred value	0.11	
1-Butene	C ₄ H ₈	3.14E-11	GC-FID			IUPAC preferred value	0.26	
1-Pentene	C ₅ H ₁₀	3.14E-11	GC-FID			Atkinson and Arey, 2003	0.10	
Isoprene	C ₅ H ₈	1.00E-10	GC-FID			IUPAC preferred value	0.76	
27.023	C ₂ H ₂	7.50E-13	PTR-ToF-MS	Ethyne		IUPAC preferred value	0.001	
41.0386	C ₃ H ₄	6.95E-13	PTR-ToF-MS	Propyne; 1,2-Propadiene	average	Warnatz, 1984; Atkinson and Arey, 2003	0.27	
67.0543	C ₅ H ₆	5.16E-11	PTR-ToF-MS	1,3-Cyclopentadiene; 3-penten-2-yne	average	Grosjean and Williams, 1992; Boodaghians et al., 1987	0.35	

69.0699	C ₅ H ₈	5.10E-11	PTR-ToF-MS	1-Pentyne; 1,2-Pentadiene; 1,4-Pentadiene; 1,3-Pentadiene; Cyclopentene	average	Boodaghians et al., 1987; Rogers, 1989; Ohta, 1983; Atkinson and Arey, 2003	1.92	C ₅ dienes. Isoprene measured by GC-FID was subtracted
83.0856	C ₆ H ₁₀	8.69E-11	PTR-ToF-MS	Cyclohexene, 1-methyl-cyclopentene; 2,3-dimethylbutadiene; 2-methylpentadiene; hexyne; 2-methyl-1,4-pentadiene; 3-methyl-1,3-pentadiene; 4-methyl-1,3-pentadiene; 1,2-hexadiene; 2-4-hexadiene; 1,4-hexadiene; 1,3-hexadiene; 1,5-Hexadiene	average	Atkinson and Arey, 2003; Boodaghians et al., 1987	3.13	C ₆ dienes
85.1012	C ₆ H ₁₂	6.63E-11	PTR-ToF-MS	2-Methyl-2-pentene; 2-Methyl-1-pentene; 3-Methyl-2-pentene; 2-Ethyl-1-butene; 2,3-Dimethyl-2-butene; 1-Hexene; 2-Hexene; 3-Hexene; 4-Methyl-2-pentene	average	Atkinson and Arey, 2003 Ohta, 1983; Grosjean and Williams, 1992	0.57	C ₆ alkenes
95.0856	C ₇ H ₁₀	4.68E-11	PTR-ToF-MS	2-Norbornene		Atkinson et al., 1983	0.76	
97.1012	C ₇ H ₁₂	7.98E-11	PTR-ToF-MS	1-methyl-cyclohexene; cycloheptene; 2-methyl-1,5-hexadiene; 2,4-dimethyl-1,3-pentadiene	average	Aschmann et al., 2012; Atkinson and Arey, 2003; Grosjean and Williams, 1992	2.27	C ₇ dienes
109.1012	C ₈ H ₁₂	1.07E-10	PTR-ToF-MS	Bicyclo[2.2.2]oct-2-ene; 1,2,3-Trimethylcyclopentadiene	average	Atkinson and Arey, 2003; Grosjean and Williams, 1992	3.82	
111.1169	C ₈ H ₁₄	1.20E-10	PTR-ToF-MS	Z-Cyclooctene; 2,5-Dimethyl-2,4-hexadiene; 2,5-Dimethyl-1,5-hexadiene		Aschmann et al., 2012; Atkinson and Arey, 2003	1.89	C ₈ dienes
137.1325	C ₁₀ H ₁₆	3.73E-10	PTR-ToF-MS	1,3,7-Octatriene; α -Pinene	average	IUPAC preferred value; Peeters et al., 2007	3.07	
205.1951	C ₁₅ H ₂₄	1.01E-10	PTR-ToF-MS	β -caryophyllene; α -copaene; α -cedrene; α -humulene; E- β -farnesene; Longifolene		Atkinson and Arey, 2003; IUPAC preferred values	1.10	Sesquiterpenes
Aromatic Compounds								
m- and p-Xylene	C ₈ H ₁₀	1.87E-11	GC-FID		average	Atkinson and Arey, 2003	0.26	
69.0335	C ₄ H ₄ O	4.04E-11	PTR-ToF-MS	Furan		Atkinson, 1986	0.28	
Benzene	C ₆ H ₆	1.20E-12	GC-FID			IUPAC preferred value	0.03	
80.0495	C ₅ H ₅ N	5.31E-13	PTR-ToF-MS	Pyridine		Yeung and Elrod, 2003	0.002	

83.0492	C ₅ H ₆ O	9.00E-11	PTR-ToF-MS	3-methyl-furan		Atkinson et al., 1989	0.79	
Toluene	C ₇ H ₈	5.60E-12	GC-FID			IUPAC preferred value	0.23	
94.0652	C ₆ H ₇ N	5.68E-11	PTR-ToF-MS	Aniline; 4-methyl-pyridine; 3-methyl-pyridine; 2-methyl-pyridine	average of methylpyridines and aniline rate coefficients	Witte et al., 1986; Yeung and Elrod, 2003	0.32	
95.0492	C ₆ H ₆ O	2.80E-11	PTR-ToF-MS	Phenol		IUPAC preferred value	0.87	
97.0648	C ₆ H ₈ O	1.25E-10	PTR-ToF-MS	2,5-dimethylfuran		Aschmann et al., 2011	0.86	
105.0731	C ₈ H ₈	5.80E-11	PTR-ToF-MS	Styrene		Atkinson and Arey, 2003	0.66	
107.0492	C ₇ H ₆ O	1.26E-11	PTR-ToF-MS	Benzaldehyde		IUPAC preferred value	0.23	
107.0855	C ₈ H ₁₀	1.36E-11	PTR-ToF-MS	o-Xylene		Atkinson and Arey, 2003	0.35	m- and p-Xylene mixing ratios measured by GC-FID were subtracted
108.0808	C ₇ H ₉ N	9.60E-12	PTR-ToF-MS	2,6-dimethylpyridine; 2,4-dimethylpyridine; 2,3-dimethylpyridine; 2,5-dimethylpyridine	average	Yeung and Elrod, 2003	0.20	
119.0856	C ₉ H ₁₀	5.95E-11	PTR-ToF-MS	α -methylstyrene; β -methylstyrene	average	Bignozzi et al., 1981	0.72	
121.0648	C ₈ H ₈ O	1.61E-11	PTR-ToF-MS	4-methylbenzaldehyde; 3-methylbenzaldehyde; 2-methylbenzaldehyde; acetophenone	average	Clifford et al., 2005; Aschmann et al., 2010	0.20	
121.1012	C ₉ H ₁₂	5.70E-11	PTR-ToF-MS	1,3,5-trimethylbenzene		Atkinson and Arey, 2003	0.91	
123.0441	C ₇ H ₆ O ₂	1.20E-12	PTR-ToF-MS	benzoic acid	rate coefficient of benzene	IUPAC preferred value	0.005	
125.0961	C ₈ H ₁₂ O	4.04E-11	PTR-ToF-MS	2,3,4,5-tetramethylfuran	rate coefficient of furan	Atkinson, 1986	0.14	
129.0699	C ₁₀ H ₈	2.30E-11	PTR-ToF-MS	naphthalene		Atkinson and Arey, 2003	0.07	
131.0856	C ₁₀ H ₁₀	2.30E-11	PTR-ToF-MS	dihydronaphthalene	rate coefficient of naphthalene	Atkinson and Arey, 2003	0.07	
133.1012	C ₁₀ H ₁₂	3.37E-11	PTR-ToF-MS	Tetralin; 2-methyl-1-propenylbenzene	average	Atkinson and Arey, 2003; Chiorboli et al., 1983	0.19	
135.0805	C ₉ H ₁₀ O	3.0E-11	PTR-ToF-MS	Dimethylbenzaldehyde	average	Tse et al., 1997; Aschmann et al., 2010	0.18	

135.1169	C ₁₀ H ₁₄	3.70E-11	PTR-ToF-MS	tert-butylbenzene; p-cymene; 1,2,3,5-tetramethylbenzene; 1,2,3,4-tetramethylbenzene	average	Alarcón et al., 2015; Atkinson and Arey, 2003	1.02	
143.0856	C ₁₁ H ₁₀	5.08E-11	PTR-ToF-MS	2-methyl-Naphthalene; 1-methyl-naphthalene	average	Atkinson and Aschmann, 1986; Phousongphouang and Arey, 2002	0.94	
147.1169	C ₁₁ H ₁₄	5.80E-11	PTR-ToF-MS	4-isopropyl styrene; 3-pentenyl benzene	rate coefficient of styrene	Atkinson and Arey, 2003	1.15	
149.1325	C ₁₁ H ₁₆	1.10E-10	PTR-ToF-MS	Pentamethylbenzene		Alarcón et al., 2015	2.23	
161.1325	C ₁₂ H ₁₆	5.80E-11	PTR-ToF-MS	4-tert-butylstyrene	rate coefficient of styrene	Atkinson and Arey, 2003	1.31	
163.1482	C ₁₂ H ₁₈	1.13E-10	PTR-ToF-MS	Hexamethylbenzene		Atkinson and Arey, 2003	1.30	
175.1482	C ₁₃ H ₁₈	3.85E-11	PTR-ToF-MS	1-2,2-dimethyl-1-methylenepropyl-4-methylbenzene; 2-tert-butyl-indan	average of indan and styrene rate coefficients	Atkinson and Arey, 2003	1.02	
189.1638	C ₁₄ H ₂₀	4.60E-11	PTR-ToF-MS	3,5-dimethyl-t-butylstyrene; 1-t-butyltetralin	average of styrene and tetralin rate coefficients	Atkinson and Arey, 2003	0.591	
197.1325	C ₁₅ H ₁₆	4.15E-12	PTR-ToF-MS	1,1'-1,3-propanediylbis-Benzene; 4-1-methylethyl-1,1'-Biphenyl	average of benzene and biphenyl	Atkinson and Arey, 2003; IUPAC preferred value	0.003	
199.1482	C ₁₅ H ₁₈	2.30E-11	PTR-ToF-MS	substituted naphthalenes	rate coefficient of naphthalene	Atkinson and Arey, 2003	0.05	
OVOCs								
33.0335	CH ₃ OH	9.00E-13	PTR-ToF-MS	methanol		IUPAC preferred value	0.24	
45.0335	C ₂ H ₄ O	1.50E-11	PTR-ToF-MS	acetaldehyde		IUPAC preferred value	2.21	
47.0128	HCOOH	4.50E-13	PTR-ToF-MS	formic acid		IUPAC preferred value	0.07	
57.0335	C ₃ H ₄ O	1.13E-11	PTR-ToF-MS	Methyl tert-butyl ether (MTBE); acrolein	average	Atkinson and Arey, 2003	0.87	
59.0491	C ₃ H ₆ O	1.80E-13	PTR-ToF-MS	acetone		IUPAC preferred value	0.16	
61.0285	C ₂ H ₄ O ₂	6.90E-13	PTR-ToF-MS	acetic acid		IUPAC preferred value	0.13	
63.0441	C ₂ H ₆ O ₂	1.45E-11	PTR-ToF-MS	1,2-ethanediol		IUPAC preferred value	0.18	
65.0233	CH ₄ O ₃	7.10E-12	PTR-ToF-MS	Hydroxymethyl hydroperoxide		Allen et al., 2018	0.14	
71.0491	C ₄ H ₆ O	2.45E-11	PTR-ToF-MS	Methyl vinyl ketone; methacroleine; isoprene hydroxy hydroperoxide	average	IUPAC preferred values; St Clair et al., 2016	0.42	

73.0284	C ₃ H ₄ O ₂	1.30E-11	PTR-ToF-MS	Methyl glyoxal		IUPAC preferred value	0.21	
73.0648	C ₄ H ₈ O	4.87E-11	PTR-ToF-MS	butanal; 2-methyl-propanal; tetrahydrofuran; 2-butanone MEK; 2-butene-1-ol; ethoxy ethene; 3-butene-1-ol; 2-methyl-2-propene-1-ol	average	IUPAC preferred value; Cometto et al., 2008; Moriarty et al., 2003; Audley et al., 1981; Zhou et al., 2006	3.11	C ₄ carbonyls
75.0441	C ₃ H ₆ O ₂	2.85E-12	PTR-ToF-MS	hydroxyacetone; propanoic acid	average	IUPAC preferred values	0.10	
77.0598	C ₃ H ₈ O ₂	1.62E-11	PTR-ToF-MS	2-methoxyethanol; 1,2-propanediol		IUPAC preferred value, Porter et al., 1997	0.07	
85.0648	C ₅ H ₈ O	4.43E-11	PTR-ToF-MS	Cyclopentanone; E-2-pentenal; 3-methyl-3-butene-2-one; E-2-methyl-2-butenal; 3-methyl-2-butenal; 3-penten-2-one; 1-penten-3-one; 3,6-dihydro-2H-pyran	average	Atkinson and Arey, 2003; Grosjean and Williams, 1992; Jiménez et al., 2009; Davis et al., 2007	0.52	
87.0441	C ₄ H ₆ O ₂	2.40E-11	PTR-ToF-MS	2,3-Butanedione; acetic acid ethenyl ester; 2-propenoic acid methyl ester; 4-hydroxy-2-butenal	average	Baker et al., 2005a; Teruel et al., 2006; Blanco et al., 2009; Dagaut et al., 1988	0.61	C ₄ dioxygenates
87.0805	C ₅ H ₁₀ O	4.63E-11	PTR-ToF-MS	Pentanal; 3-methyl-2-butanone; 3-pantanone; 2-pantanone; 1-vinyloxypropane; 2-Methyl-3-buten-2-ol MBO; 3-methyl-3-buten-1-ol; 2-methylbutanal; 3-methylbutanal; 1-penten-3-ol; Z-2-penten-1-ol	average	Zhou et al., 2006; IUPAC; Cometto et al., 2008; Atkinson and Arey, 2003; Orlando et al., 2001	2.48	
89.0234	C ₃ H ₄ O ₃	1.20E-13	PTR-ToF-MS	pyruvic acid		IUPAC preferred value	0.001	
89.0598	C ₄ H ₈ O ₂	6.47E-12	PTR-ToF-MS	1,4-dioxane; formic acid 1-methylethyl ester; ethyl acetate; butyric acid; acetoin; 1-hydroxy-3-butanone	average	Picquet et al., 1998; Aschmann et al., 2000; IUPAC; Moriarty et al., 2003; Pimentel et al., 2010	0.10	
91.039	C ₃ H ₆ O ₃	9.85E-13	PTR-ToF-MS	carbonic acid dimethyl ester; methyoxyethyl formate	average	Katrib et al., 2002; O'Donnell et al., 2004	0.003	
99.0805	C ₆ H ₁₀ O	4.76E-11	PTR-ToF-MS	cyclohexanone; 5-hexen-2-one; z-3-methyl-3-penten-2-one; 4-methyl-3-penten-2-one; E-2-hexenal; Z-3-hexenal; 2-ethyl-2-butenal; E-4-hexene-3-one	average	Atkinson and Arey, 2003; Grosjean and Williams, 1992; Wang et al., 2010; Gao et al., 2009; Xing et al., 2012	0.59	

101.0598	C ₅ H ₈ O ₂	3.17E-11	PTR-ToF-MS	Methyl methacrylate; E-2-butenoic acid methyl ester; acetylacetone; 2-propenoic acid ethyl ester; glutaraldehyde; vinyl propionate; isopropenyl acetate; n-propenyl acetate	average	Teruel et al., 2006; Teruel et al., 2012; Rogers, 1989; Ferrari et al., 1996; Dagaut et al., 1988	1.52	C ₅ dioxygenates
101.0961	C ₆ H ₁₂ O	4.11E-11	PTR-ToF-MS	Oxepane; 2-hexanone; hexanal; Z-3-hexen-1-ol; t-butyl vinyl ether; 2-methylpentanal; 3-methylpentanal; 4-methylpentanal; 3,3-dimethylbutanal; 2-ethylbutanal	average	Moriarty et al., 2003; Atkinson and Arey, 2003; Zhou et al., 2012; Le Calvé et al., 1998	1.15	C ₆ oxygenates/ carbonyls
103.0754	C ₅ H ₁₀ O ₂	8.71E-12	PTR-ToF-MS	butanoic acid methyl ester; n-propyl acetate; 1-hydroxy-2-methyl-3-butanone; 1-hydroxy-3-methyl-2-butanone; 1,2-dioxepane; 5-hydroxy-2-pentanone	average	Schütze et al., 2010; Ferrari et al., 1996; Moriarty et al., 2003; Aschmann et al., 2000; Baker et al., 2005b	0.14	
113.0598	C ₆ H ₈ O ₂	5.80E-11	PTR-ToF-MS	E-hexendione; Z-hexendione	average	Tuazon et al., 1985	0.23	
113.0961	C ₇ H ₁₂ O	2.52E-11	PTR-ToF-MS	E-2-Hepten-1-al; cycloheptanone	average	Davis et al., 2007; Atkinson and Arey, 2003	0.16	
115.0754	C ₆ H ₁₀ O ₂	3.40E-11	PTR-ToF-MS	2,5-hexanedione; ethyl-2-butenoate; methacrylic acid ethyl ester	average	Dagaut et al., 1988; Teruel et al., 2012; Blanco et al., 2006	0.97	
117.0911	C ₆ H ₁₂ O ₂	7.42E-12	PTR-ToF-MS	4-hydroxy-3-hexanone; acetic acid butyl ester; butanoic acid ethyl ester; methyl valerate	average	Schütze et al., 2010; Aschmann et al., 2000; Veillerot et al., 1996; Ferrari et al., 1996	0.13	
127.1118	C ₈ H ₁₄ O	9.88E-11	PTR-ToF-MS	E-2-octenal; cyclooctanone; 6-methyl-5-hepten-2-one	average	Gao et al., 2009; Smith et al., 1996	1.87	
129.1274	C ₈ H ₁₆ O	1.10E-11	PTR-ToF-MS	2-octanone		Atkinson and Arey, 2003	0.03	
141.1274	C ₉ H ₁₆ O	4.35E-11	PTR-ToF-MS	nonenal		Gao et al., 2009	0.07	
143.1067	C ₈ H ₁₄ O ₂	7.08E-11	PTR-ToF-MS	butyl methacrylate		Blanco et al., 2009	1.14	
143.1431	C ₉ H ₁₈ O	2.41E-11	PTR-ToF-MS	nonanal, 1-nonanone	average	Bowman et al., 2003; Wallington and Kurylo, 1987	0.11	
201.1485	C ₁₁ H ₂₀ O ₃	7.35E-12	PTR-ToF-MS	C ₁₁ oxo esters e.g. 9-oxo-nonanoic acid ethyl ester	rate coefficient of acetic acid n-pentyl ester	Williams et al., 1993	0.06	

References for reaction rate coefficients

- Alarcón, P., Bohn, B., and Zetzsch, C.: Kinetic and mechanistic study of the reaction of OH radicals with methylated benzenes: 1,4-dimethyl-, 1,3,5-trimethyl-, 1,2,4,5-, 1,2,3,5- and 1,2,3,4-tetramethyl-, pentamethyl-, and hexamethylbenzene, *Physical chemistry chemical physics PCCP*, 17, 13053–13065, doi:10.1039/c5cp00253b, 2015.
- Allen, H. M., Crounse, J. D., Bates, K. H., Teng, A. P., Krawiec-Thayer, M. P., Rivera-Rios, J. C., Keutsch, F. N., St Clair, J. M., Hanisco, T. F., Møller, K. H., Kjaergaard, H. G., and Wennberg, P. O.: Kinetics and Product Yields of the OH Initiated Oxidation of Hydroxymethyl Hydroperoxide, *The journal of physical chemistry. A*, 122, 6292–6302, doi:10.1021/acs.jpca.8b04577, 2018.
- Aschmann, S. M., Arey, J., and Atkinson, R.: Atmospheric Chemistry of Selected Hydroxycarbonyls, *J. Phys. Chem. A*, 104, 3998–4003, doi:10.1021/jp9939874, 2000.
- Aschmann, S. M., Arey, J., and Atkinson, R.: Extent of H-atom abstraction from OH+p-cymene and upper limits to the formation of cresols from OH+m-xylene and OH+p-cymene, *ATMOSPHERIC ENVIRONMENT*, 44, 3970–3975, doi:10.1016/j.atmosenv.2010.06.059, 2010.
- Aschmann, S. M., Arey, J., and Atkinson, R.: Kinetics and products of the reactions of OH radicals with cyclohexene, 1-methyl-1-cyclohexene, cis-cyclooctene, and cis-cyclodecene, *The journal of physical chemistry. A*, 116, 9507–9515, doi:10.1021/jp307217m, 2012.
- Aschmann, S. M., Nishino, N., Arey, J., and Atkinson, R.: Kinetics of the reactions of OH radicals with 2- and 3-methylfuran, 2,3- and 2,5-dimethylfuran, and E- and Z-3-hexene-2,5-dione, and products of OH + 2,5-dimethylfuran, *Environmental science & technology*, 45, 1859–1865, doi:10.1021/es103207k, 2011.
- Atkinson, R.: Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions, *Chemical reviews*, 86, 69–201, doi:10.1021/cr00071a004, 1986.
- Atkinson, R. and Arey, J.: Atmospheric degradation of volatile organic compounds, *Chemical reviews*, 103, 4605–4638, doi:10.1021/cr0206420, 2003.
- Atkinson, R. and Aschmann, S. M.: Kinetics of the reactions of naphthalene, 2-methylnaphthalene, and 2,3-dimethylnaphthalene with OH radicals and with O₃ at 295 ± 1 K, *Int. J. Chem. Kinet.*, 18, 569–573, doi:10.1002/kin.550180507, 1986.
- Atkinson, R., Aschmann, S. M., and Carter, W. P. L.: Effects of ring strain on gas-phase rate constants. 2. OH radical reactions with cycloalkenes, *Int. J. Chem. Kinet.*, 15, 1161–1177, doi:10.1002/kin.550151105, 1983.
- Atkinson, R., Aschmann, S. M., Tuazon, E. C., Arey, J., and Zielinska, B.: Formation of 3-Methylfuran from the gas-phase reaction of OH radicals with isoprene and the rate constant for its reaction with the OH radical, *Int. J. Chem. Kinet.*, 21, 593–604, doi:10.1002/kin.550210709, 1989.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O_x, HO_x, NO_x and SO_x species, *Atmos. Chem. Phys.*, 4, 1461–1738, doi:10.5194/acp-4-1461-2004, 2004.
- Atkinson, R., Tuazon, E. C., Wallington, T. J., Aschmann, S. M., Arey, J., Winer, A. M., and Pitts, J. N.: Atmospheric chemistry of aniline, N,N-dimethylaniline, pyridine, 1,3,5-triazine, and nitrobenzene, *Environ. Sci. Technol.*, 21, 64–72, doi:10.1021/es00155a007, 1987.
- Audley, G. J., Baulch, D. L., and Campbell, I. M.: Gas-phase reactions of hydroxyl radicals with aldehydes in flowing H₂O₂+NO₂+CO mixtures, *J. Chem. Soc., Faraday Trans. 1*, 77, 2541, doi:10.1039/f19817702541, 1981.
- Baker, J., Arey, J., and Atkinson, R.: Formation and Reaction of Hydroxycarbonyls from the Reaction of OH Radicals with 1,3-Butadiene and Isoprene, *Environ. Sci. Technol.*, 39, 4091–4099, doi:10.1021/es047930t, 2005a.
- Baker, J., Arey, J., and Atkinson, R.: Rate constants for the reactions of OH radicals with a series of 1,4-hydroxyketones, *Journal of Photochemistry and Photobiology A: Chemistry*, 176, 143–148, doi:10.1016/j.jphotochem.2005.07.022, 2005b.
- Bignozzi, C. A., Maldotti, A., Chiorboli, C., Bartocci, C., and Carassiti, V.: Kinetics and mechanism of reactions between aromatic olefins and hydroxyl radicals, *Int. J. Chem. Kinet.*, 13, 1235–1242, doi:10.1002/kin.550131204, 1981.
- Blanco, M. B., Bejan, I., Barnes, I., Wiesen, P., and Teruel, M. A.: OH-initiated degradation of unsaturated esters in the atmosphere: Kinetics in the temperature range of 287–313 K, *The journal of physical chemistry. A*, 113, 5958–5965, doi:10.1021/jp901755x, 2009.
- Blanco, M. B., Taccone, R. A., Lane, S. I., and Teruel, M. A.: On the OH-initiated degradation of methacrylates in the troposphere: Gas-phase kinetics and formation of pyruvates, *Chemical Physics Letters*, 429, 389–394, doi:10.1016/j.cplett.2006.08.088, 2006.
- Boodaghians, R. B., Hall, I. W., Toby, F. S., and Wayne, R. P.: Absolute determinations of the kinetics and temperature dependences of the reactions of OH with a series of alkynes, *J. Chem. Soc., Faraday Trans. 2*, 83, 2073, doi:10.1039/f29878302073, 1987.
- Bowman, J. H., Barket, D. J., and Shepson, P. B.: Atmospheric Chemistry of Nonanal, *Environ. Sci. Technol.*, 37, 2218–2225, doi:10.1021/es026220p, 2003.
- Chiorboli, C., Bignozzi, C. A., Maldotti, A., Giardini, P. F., Rossi, A., and Carassiti, V.: Rate constants for the gas-phase reactions of OH radicals with dimethylstyrene and acetone. Mechanism of dimethylstyrene NO_x - air photooxidation, *Int. J. Chem. Kinet.*, 15, 579–586, doi:10.1002/kin.550150608, 1983.

- Clifford, G. M., Thüner, L. P., Wenger, J. C., and Shallcross, D. E.: Kinetics of the gas-phase reactions of OH and NO₃ radicals with aromatic aldehydes, *Journal of Photochemistry and Photobiology A: Chemistry*, 176, 172–182, doi:10.1016/j.jphotochem.2005.09.022, 2005.
- Cometto, P. M., Dalmasso, P. R., Taccone, R. A., Lane, S. I., Oussar, F., Daëlle, V., Mellouki, A., and Le Bras, G.: Rate coefficients for the reaction of OH with a series of unsaturated alcohols between 263 and 371 K, *The journal of physical chemistry. A*, 112, 4444–4450, doi:10.1021/jp7111186, 2008.
- Dagaut, P., Wallington, T. J., Liu, R., and Kurylo, M. J.: A kinetic investigation of the gas-phase reactions of hydroxyl radicals with cyclic ketones and diones: Mechanistic insights, *J. Phys. Chem.*, 92, 4375–4377, doi:10.1021/j100326a026, 1988.
- Davis, M. E., Gilles, M. K., Ravishankara, A. R., and Burkholder, J. B.: Rate coefficients for the reaction of OH with E-2-pentenal, E-2-hexenal, and E-2-heptenal, *Physical chemistry chemical physics CCP*, 9, 2240–2248, doi:10.1039/b700235a, 2007.
- Ferrari, C., Roche, A., Jacob, V., Foster, P., and Baussand, P.: Kinetics of the reaction of OH radicals with a series of esters under simulated conditions at 295 K, *Int. J. Chem. Kinet.*, 28, 609–614, doi:10.1002/SICI1097-4601199628:8<609::AID-KIN6>3.0.CO;2-Z, 1996.
- Gao, T., Andino, J. M., Rivera, C. C., and Márquez, M. F.: Rate constants of the gas-phase reactions of OH radicals with trans -2-hexenal, trans -2-octenal, and trans -2-nonenal, *Int. J. Chem. Kinet.*, 41, 483–489, doi:10.1002/kin.20424, 2009.
- Grosjean, D. and Williams, E. L.: Environmental persistence of organic compounds estimated from structure-reactivity and linear free-energy relationships. Unsaturated aliphatics, *Atmospheric Environment. Part A. General Topics*, 26, 1395–1405, doi:10.1016/0960-16869290124-4, 1992.
- IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation: Datasheets - gas phase: <http://iupac.pole-ether.fr/>, last access: 10 December 2018.
- Jiménez, E., Lanza, B., Antiñolo, M., and Albaladejo, J.: Influence of temperature on the chemical removal of 3-methylbutanal, trans-2-methyl-2-butenal, and 3-methyl-2-butenal by OH radicals in the troposphere, *ATMOSPHERIC ENVIRONMENT*, 43, 4043–4049, doi:10.1016/j.atmosenv.2009.05.005, 2009.
- Katrib, Y., Deiber, G., Mirabel, P., Le Calvé, S., George, C., Mellouki, A., and Le Bras, G.: Atmospheric Loss Processes of Dimethyl and Diethyl Carbonate, *Journal of Atmospheric Chemistry*, 43, 151–174, doi:10.1023/A:1020605807298, 2002.
- Le Calvé, S., Hitier, D., Le Bras, G., and Mellouki, A.: Kinetic Studies of OH Reactions with a Series of Ketones, *J. Phys. Chem. A*, 102, 4579–4584, doi:10.1021/jp980848y, 1998.
- Moriarty, J., Sidebottom, H., Wenger, J., Mellouki, A., and Le Bras, G.: Kinetic Studies on the Reactions of Hydroxyl Radicals with Cyclic Ethers and Aliphatic Diethers, *J. Phys. Chem. A*, 107, 1499–1505, doi:10.1021/jp021267i, 2003.
- O'Donnell, S. M., Sidebottom, H. W., Wenger, J. C., Mellouki, A., and Le Bras, G.: Kinetic Studies on the Reactions of Hydroxyl Radicals with a Series of Alkoxy Esters, *J. Phys. Chem. A*, 108, 7386–7392, doi:10.1021/jp048782w, 2004.
- Ohta, T.: Rate constants for reactions of diolefins with hydroxyl radicals in the gas phase. Estimate of the rate constants from those for monoolefins, *J. Phys. Chem.*, 87, 1209–1213, doi:10.1021/j100230a023, 1983.
- Orlando, J. J., Tyndall, G. S., and Ceazan, N.: Rate Coefficients and Product Yields from Reaction of OH with 1-Penten-3-ol, Z -2-Penten-1-ol, and Allyl Alcohol 2-Propen-1-ol, *J. Phys. Chem. A*, 105, 3564–3569, doi:10.1021/jp0041712, 2001.
- Peeters, J., Boullart, W., Pultau, V., Vandenbergk, S., and Vereecken, L.: Structure-activity relationship for the addition of OH to polyalkenes: Site-specific and total rate constants, *J. Phys. Chem. A*, 111, 1618–1631, doi:10.1021/jp066973o, 2007.
- Phousongphouang, P. T. and Arey, J.: Rate Constants for the Gas-Phase Reactions of a Series of Alkylnaphthalenes with the OH Radical, *Environ. Sci. Technol.*, 36, 1947–1952, doi:10.1021/es011434c, 2002.
- Picquet, B., Heroux, S., Chebbi, A., Doussin, J.-F., Durand-Jolibois, R., Monod, A., Loirat, H., and Carlier, P.: Kinetics of the reactions of OH radicals with some oxygenated volatile organic compounds under simulated atmospheric conditions, *Int. J. Chem. Kinet.*, 30, 839–847, doi:10.1002/SICI1097-4601199830:11<839::AID-KIN6>3.0.CO;2-W, 1998.
- Pimentel, A. S., Tyndall, G. S., Orlando, J. J., Hurley, M. D., Wallington, T. J., Sulbaek Andersen, M. P., Marshall, P., and Dibble, T. S.: Atmospheric chemistry of isopropyl formate and tert-butyl formate, *Int. J. Chem. Kinet.*, 42, 479–498, doi:10.1002/kin.20498, 2010.
- Porter, E., Wenger, J., Treacy, J., Sidebottom, H., Mellouki, A., Téton, S., and LeBras, G.: Kinetic Studies on the Reactions of Hydroxyl Radicals with Diethers and Hydroxyethers, *J. Phys. Chem. A*, 101, 5770–5775, doi:10.1021/jp971254i, 1997.
- Rogers, J. D.: Rate constant measurements for the reaction of the hydroxyl radical with cyclohexene, cyclopentene, and glutaraldehyde, *Environ. Sci. Technol.*, 23, 177–181, doi:10.1021/es00179a006, 1989.
- Schütze, N., Zhong, X., Kirschbaum, S., Bejan, I., Barnes, I., and Benter, T.: Relative kinetic measurements of rate coefficients for the gas-phase reactions of Cl atoms and OH radicals with a series of methyl alkyl esters, *ATMOSPHERIC ENVIRONMENT*, 44, 5407–5414, doi:10.1016/j.atmosenv.2009.08.011, 2010.
- Smith, A. M., Rigler, E., Kwok, E. S. C., and Atkinson, R.: Kinetics and Products of the Gas-Phase Reactions of 6-Methyl-5-hepten-2-one and trans -Cinnamaldehyde with OH and NO₃ Radicals and O₃ at 296 ± 2 K, *Environ. Sci. Technol.*, 30, 1781–1785, doi:10.1021/es950871m, 1996.

- St Clair, J. M., Rivera-Rios, J. C., Crounse, J. D., Knap, H. C., Bates, K. H., Teng, A. P., Jørgensen, S., Kjaergaard, H. G., Keutsch, F. N., and Wennberg, P. O.: Kinetics and Products of the Reaction of the First-Generation Isoprene Hydroxy Hydroperoxide ISOPOOH with OH, *The journal of physical chemistry. A*, 120, 1441–1451, doi:10.1021/acs.jpca.5b06532, 2016.
- Teruel, M. A., Benitez-Villalba, J., Caballero, N., and Blanco, M. B.: Gas-phase oxidation of methyl crotonate and ethyl crotonate. kinetic study of their reactions toward OH radicals and Cl atoms, *The journal of physical chemistry. A*, 116, 6127–6133, doi:10.1021/jp2113889, 2012.
- Teruel, M. A., Lane, S. I., Mellouki, A., Solignac, G., and Le Bras, G.: OH reaction rate constants and UV absorption cross-sections of unsaturated esters, *ATMOSPHERIC ENVIRONMENT*, 40, 3764–3772, doi:10.1016/j.atmosenv.2006.03.003, 2006.
- Tse, C. W., Flagan, R. C., and Seinfeld, J. H.: Rate constants for the gas-phase reaction of the hydroxyl radical with a series of dimethylbenzaldehydes and trimethylphenols at atmospheric pressure, *Int. J. Chem. Kinet.*, 29, 523–525, doi:10.1002/SICI1097-4601199729:7<523::AID-KIN6>3.0.CO;2-W, 1997.
- Tuazon, E. C., Atkinson, R., and Carter, W. P.: Atmospheric chemistry of cis- and trans-3-hexene-2,5-dione, *Environmental science & technology*, 19, 265–269, doi:10.1021/es00133a009, 1985.
- Veillerot, M., Foster, P., Guillermo, R., and Galloo, J. C.: Gas-phase reaction of n-butyl acetate with the hydroxyl radical under simulated tropospheric conditions: Relative rate constant and product study, *Int. J. Chem. Kinet.*, 28, 235–243, doi:10.1002/SICI1097-4601199628:4<235::AID-KIN1>3.0.CO;2-W, 1996.
- Wallington, T. J. and Kurylo, M. J.: Flash photolysis resonance fluorescence investigation of the gas-phase reactions of hydroxyl radicals with a series of aliphatic ketones over the temperature range 240–440 K, *J. Phys. Chem.*, 91, 5050–5054, doi:10.1021/j100303a033, 1987.
- Wang, K., Ge, M., and Wang, W.: Kinetics of the gas-phase reactions of 5-hexen-2-one with OH and NO₃ radicals and O₃, *Chemical Physics Letters*, 490, 29–33, doi:10.1016/j.cplett.2010.03.023, 2010.
- Warnatz, J.: Rate Coefficients in the C/H/O System, in: *Combustion Chemistry*, Gardiner, W. C. Ed., Springer US, New York, NY, 197–360, 1984.
- Williams, D. C., O'Rji, L. N., and Stone, D. A.: Kinetics of the reactions of OH radicals with selected acetates and other esters under simulated atmospheric conditions, *Int. J. Chem. Kinet.*, 25, 539–548, doi:10.1002/kin.550250704, 1993.
- Witte, F., Urbanik, E., and C, Z.: Temperature dependence of the rate constants for the addition of hydroxyl to benzene and to some monosubstituted aromatics aniline, bromobenzene, and nitrobenzene and the unimolecular decay of the adducts. Part 2. Kinetics into a quasi-equilibrium, *J. Phys. Chem.*, 90, 3251–3259, doi:10.1021/j100405a040, 1986.
- Xing, J.-H., Ono, M., Kuroda, A., Obi, K., Sato, K., and Imamura, T.: Kinetic study of the daytime atmospheric fate of Z-3-hexenal, *The journal of physical chemistry. A*, 116, 8523–8529, doi:10.1021/jp303202h, 2012.
- Yeung, L. Y. and Elrod, M. J.: Experimental and Computational Study of the Kinetics of OH + Pyridine and Its Methyl- and Ethyl-Substituted Derivatives, *J. Phys. Chem. A*, 107, 4470–4477, doi:10.1021/jp027389s, 2003.
- Zhou, S., Barnes, I., Zhu, T., Bejan, I., and Benter, T.: Kinetic study of the gas-phase reactions of OH and NO₃ radicals and O₃ with selected vinyl ethers, *J. Phys. Chem. A*, 110, 7386–7392, doi:10.1021/jp061431s, 2006.
- Zhou, S., Barnes, I., Zhu, T., and Benter, T.: Kinetic study of gas-phase reactions of OH and NO₃ radicals and O₃ with iso-butyl and tert-butyl vinyl ethers, *The journal of physical chemistry. A*, 116, 8885–8892, doi:10.1021/jp305992a, 2012.