



Supplement of

Pre-deliquescent water uptake in deposited nanoparticles observed with in situ ambient pressure X-ray photoelectron spectroscopy

Jack J. Lin et al.

Correspondence to: Nønne L. Prisle (nonne.prisle@oulu.fi) and Samuli Urpelainen (samuli.urpelainen@oulu.fi)

The copyright of individual parts of the supplement might differ from the article licence.

S1 Spectra fitting and uncertainty

In this section, we explain the constraints used during the fitting of the measured XPS spectra. The peaks are represented with Voigt functions, or the convolution of a Lorentzian and a Gaussian shape. The peak shape has a Lorentzian contribution from the X-ray source, which uses the decay of Al $K\alpha$ and Mg $K\alpha$ core-holes to generate the X-rays. Physically, the Lorentzian shape represents the lifetime broadening effect experienced by a photoelectron. The Gaussian shape represents measurement uncertainties². Other effects may induce asymmetric broadening, but an asymmetry parameter was usually not necessary to explain the peak shapes fitted here.

We also present the results of Monte Carlo error analysis of the fit results. The Monte Carlo analysis is done by creating a noiseless version of the fit and repeating the fitting procedure with random noise added to the noiseless spectrum. For the Monte Carlo analysis here, the fits were repeated either 100 or 200 times. We report the error in a given fit parameter as $\pm 2\sigma$, where σ is the standard deviation of the fit parameters from the set of Monte Carlo simulations.

S1.1 NaCl

Measurements were made on NaCl particles at UHV conditions as well as at 0, 2, and 5 mbar water vapor pressure inside the high pressure cell. Another set of measurements was taken at 0 mbar water vapor pressure after dosing with water vapor.

S1.1.1 Na 1s

Fit parameters for Na 1s spectra are shown in Table S1.

Table S1: Fitting parameters with uncertainty estimates from Monte Carlo simulations for Na 1s spectra measured from NaCl particles.

P_{H_2O} (mbar)	Peak number	Binding energy (eV)	FWHM (eV)			Asym metry	Area (a.u.)	Monte Carlo iterations
			Lorentzian	Gaussian	Total			
0	1	1074.64 \pm 0.02	570.99 \pm 175.70	1709.38 \pm 135.07	2034.93 \pm 221.62	0 \pm 0	4875.78 \pm 201.49	200
2	1	1072.42 \pm 0.05	64.35 \pm 145.80	1980.93 \pm 132.29	2015.05 \pm 196.87	0 \pm 0	1022.05 \pm 63.74	200
5	1	1072.70 \pm 0.11	1.28 \pm 0.01	2714.72 \pm 911.72	2714.75 \pm 911.72	0 \pm 0	217.08 \pm 78.77	200
0*	1	1074.43 \pm 0.02	41.70 \pm 33.13	2078.67 \pm 46.20	2099.05 \pm 56.86	0 \pm 0	3800.02 \pm 101.13	200
	2	1072.54 \pm 0.06					268.48 \pm 22.61	

*Measured after dosing.

S1.1.2 Cl 2p

The Cl 2p spectra are split into two peaks from the Cl 2p_{1/2} and Cl 2p_{3/2} spin orbit components. The separation between the two is set to 1.6 eV, and the intensity of the 1/2 component fixed to half that of the 3/2 component. Both peaks are assumed to have the same line shape, meaning the peak area of the 1/2 component is also half that of the 3/2 component. Fit parameters for Cl 2p spectra are shown in Table S2.

Table S2: Fitting parameters with uncertainty estimates from Monte Carlo simulations for Cl 2p spectra measured from NaCl particles.

P _{H₂O} (mbar)	Peak number	Binding energy (eV)	FWHM (eV)			Asym metry	Area (a.u.)	Monte Carlo iterations
			Lorentzian	Gaussian	Total			
0 (UHV)	1 2	201.28 ± 0.01 202.88 ± 0.01	455.45±7.80	1203.56±11.32	1465.30±13.74	0±0	18 580.90 ± 118.06 9290.45 ± 59.03	100
0	1 2	201.41 ± 0.03 203.01 ± 0.03	127.98±127.87	1486.22±111.91	1555.66±169.93	0±0	1266.35 ± 51.72 633.18 ± 25.86	100
2	1 2	199.27 ± 0.01 200.87 ± 0.01	9.61±0.01	1776.11±39.85	1778.17±39.85	0±0	632.06 ± 13.99 316.03 ± 6.99	100
5	1 2	199.00 ± 0.10 200.60 ± 0.10	4.64±24.57	1416.94±141.16	1417.57±143.29	0±0	288.30 ± 30.99 144.15 ± 19.68	100
0*	1 2	201.27 ± 0.00 202.87 ± 0.00	214.68±2.99	1388.00±19.23	1506.21±19.47	0±0	1084.50 ± 13.46 542.25 ± 6.73	100

*Measured after dosing.

S1.2 Sucrose

Sucrose particles were measured at 0, 0.2, 1, and 5 mbar water vapor pressure.

S1.2.1 C 1s

Sucrose C 1s spectra contain three main peaks corresponding to distinct carbon atoms in C-C/C-H, C-O and O-C-O bonds. All peaks were forced to have the same symmetric Voigt line shape, as we expect the line shapes to be characterized by the natural broadening of the Mg K_{α} X-ray emission lines (680 meV) and the instrumental broadening of the electron analyzer (approximately 1000 meV). During fitting of the spectra, the separation between the C-O and O-C-O carbon peaks was fixed to 1.4 eV at all UHV vacuum conditions below 10^{-8} mbar, based on previous XPS studies of disaccharides by Stevens and Schroeder³. The fitted C-O and O-C-O peaks were, however, not linked to the C-C peak, as there are no purely C-C bound carbon in sucrose itself. The intensity ratios of the peaks were left unconstrained during fitting. Results of the fitting are shown in Table S3.

Table S3: Fitting parameters with uncertainty estimates from Monte Carlo simulations for C 1s spectra measured from sucrose particles.

P_{H_2O} (mbar)	Peak number	Binding energy (eV)	FWHM (eV)			Asym metry	Area (a.u.)	Monte Carlo iterations
			Lorentzian	Gaussian	Total			
0	1	285.90 ± 0.00					4207.45 ± 14.71	100
	2	287.36 ± 0.00	1.16±0.00	2199.86±7.70	2199.89±7.70	0±0	728.57 ± 2.56	
	3	288.76 ± 0.00					262.60 ± 0.91	
0.2	1	285.65 ± 0.02					3734.51 ± 112.12	100
	2	287.36 ± 0.06	234.81±102.91	1821.81±87.39	1950.42±135.01	0±0	607.31 ± 24.92	
	3	288.76 ± 0.06					316.91 ± 11.79	
1	1	285.57 ± 0.04					2745.13 ± 139.91	100
	2	287.10 ± 0.07	223.93±198.68	1791.20±154.98	1913.74±251.98	0±0	416.77 ± 19.78	
	3	288.50 ± 0.07					306.95 ± 13.54	
5	1	285.57 ± 0.08					582.33 ± 43.63	100
	2	287.56 ± 0.09	1.96±2.29	1950.26±127.58	1950.35±127.60	0±0	171.54 ± 11.71	
	3	288.96 ± 0.09					78.24 ± 5.06	

S1.2.2 O 1s

The sucrose O 1s spectra have two peaks arising from O-C-O and C-O components in the sucrose molecule. Both peaks were fixed to have the same symmetric line shape separated by 0.7 eV that was fixed for all humidity conditions³. With the introduction of water vapor, a third peak appears and grows in intensity with increasing RH. The separation between the two sucrose C-O and O-C-O oxygen peaks was fixed at 0.7 eV in all humidity conditions. Details of the fit results are shown in Table S4.

Table S4: Fitting parameters with uncertainty estimates from Monte Carlo simulations for O 1s spectra measured from sucrose particles.

P_{H_2O} (mbar)	Peak number	Binding energy (eV)	FWHM (eV)			Asym metry	Area (a.u.)	Monte Carlo iterations
			Lorentzian	Gaussian	Total			
0	1	533.02 ± 0.00					1546.23 ± 10.19	100
	2	533.72 ± 0.00	7.78±0.07	2209.60±14.72	2210.74±14.73	0±0	1320.93 ± 8.65	
0.2	1	533.01 ± 0.05					1819.54 ± 74.83	100
	2	533.71 ± 0.05	0.02±0.63	2405.64±85.68	2405.64±85.68	0±0	796.45 ± 33.70	
	3	536.05 ± 0.03	61.89 ± 57.07	1437.91 ± 53.95	1471.17 ± 78.53		1991.47 ± 84.85	
1	1	532.85 ± 0.05					1491.24 ± 62.35	100
	2	533.55 ± 0.05	0.08±0.40	2562.56±93.22	2562.56±93.22	0±0	919.39 ± 40.41	
	3	536.06 ± 0.02	56.18 ± 125.92	1455.71 ± 84.11	1485.80 ± 151.43		4073.48 ± 193.41	
5	1	532.45 ± 0.12					277.42 ± 16.56	100
	2	533.15 ± 0.12	80.00±0.00	2500.00±0.00	2542.37±0.00	0±0	325.69 ± 10.64	
	3	536.03 ± 0.01	518.36 ± 30.49	1206.97 ± 34.41	1507.75 ± 45.98		8027.81 ± 133.41	

S1.3 Malonic acid

Malonic acid particles were measured at 0, 0.2, and 1 mbar water vapor pressure. Due to time constraints, measurements at 0 mbar water vapor pressure after dosing were not performed.

S1.3.1 C 1s

The malonic acid C 1s has three main peaks corresponding to C-C/C-H bound carbon originating from adventitious carbon, COOH and C-C carbon from malonic acid particles. The adventitious carbon C-C/C-H and the malonic acid COOH peaks were not fixed in energy, whereas the malonic acid C-C peak was fixed at 4.0 eV lower binding energy than the COOH peak. The intensity of the attributed malonic acid C-C peak was fixed to half of that of the COOH peak, reflecting the expected stoichiometry in malonic acid molecules. The decomposition peaks DP1 and DP2 observed by Ferreira Jr. et al.¹ appear in the 0 mbar spectrum and were fixed to be at 1.5 eV (DP1) and 1.2 eV (DP2) higher binding energies than the malonic acid C-C and COOH carbon peaks, respectively. Upon exposure to water vapor, the DP2 feature disappears but a feature near the location of DP1 remains. This feature was let to vary freely in energy and intensity, but the shape was fixed to be the same as for the other lines in the spectra. The fitting results are shown in Table S5.

Table S5: Fitting parameters with uncertainty estimates from Monte Carlo simulations for C 1s spectra measured from malonic acid particles.

P_{H_2O} (mbar)	Peak number	Binding energy (eV)	FWHM (eV)			Asym metry	Area (a.u.)	Monte Carlo iterations
			Lorentzian	Gaussian	Total			
0	1	285.75 ± 0.00					68 477.98 ± 49.32	100
	2	290.02 ± 0.00	674.72±0.49	1525.08±1.16	1917.57±1.27		13 213.58 ± 9.70	
	3	286.02 ± 0.00				0±0	6606.79 ± 4.85	
	4	291.52 ± 0.00	609.88 ± 0.00	1980.96 ± 0.00	2326.98 ± 0.00		2931.03 ± 0.99	
	5	287.22 ± 0.00	609.88 ± 0.00	1500.00 ± 0.00	1852.31 ± 0.00		1061.54 ± 0.32	
0.2	1	285.48 ± 0.00					3513.38 ± 7.58	100
	2	289.64 ± 0.00					443.65 ± 0.96	
	3	285.64 ± 0.00	512.75±1.18	1531.06±3.20	1823.45±3.41	0±0	221.83 ± 0.48	
	4	287.05 ± 0.00					402.31 ± 0.85	
1	1	285.22 ± 0.00					2215.90 ± 8.20	100
	2	289.07 ± 0.00					280.55 ± 1.01	
	3	285.07 ± 0.00	315.40±0.79	1498.03±5.75	1673.49±5.80	0±0	140.27 ± 0.51	
	4	286.50 ± 0.00					382.13 ± 1.36	

S1.3.2 O 1s

Fit parameters for malonic acid O 1s spectra are shown in Table S6.

Table S6: Fitting parameters with uncertainty estimates from Monte Carlo simulations for O 1s spectra measured from malonic acid particles.

P_{H_2O} (mbar)	Peak number	Binding energy (eV)	FWHM (eV)			Asym metry	Area (a.u.)	Monte Carlo iterations
			Lorentzian	Gaussian	Total			
0	1	532.33 ± 0.00					23 867.29 ± 416.63	200
	2	533.28 ± 0.00					17 252.85 ± 162.48	
	3	537.37 ± 0.02	129.45±8.21		1555.64±15.18	0±0	1271.58 ± 5.83	
	4	538.33 ± 0.02		1485.39±12.77			920.38 ± 22.85	
	5	546.48 ± 0.02	3491.24 ± 47.65		4146.51 ± 49.34	0.21 ± 0.01	46 576.75 ± 529.69	
	6	523.10 ± 0.01	1508.82 ± 95.72		2997.23 ± 96.57	0.99 ± 0.06	7618.32 ± 419.29	
0.2	1	532.07 ± 0.01					1752.00 ± 30.48	200
	2	533.27 ± 0.01	129.45±8.21		1451.89±25.03	0±0	1386.19 ± 23.44	
	3	535.78 ± 0.02		1381.57±23.65			645.20 ± 10.51	
	4	546.66 ± 0.03	3491.24 ± 47.65		4079.17 ± 53.20	0.21 ± 0.01	2441.17 ± 28.55	
	5	523.99 ± 0.06	5.68 ± 0.36		1484.24 ± 23.65	-17.64 ± 0.18	229.39 ± 3.43	
1	1	531.84 ± 0.02					1355.54 ± 22.49	200
	2	533.04 ± 0.02	129.45±8.21		1451.89±25.03	0±0	1229.71 ± 20.37	
	3	535.78 ± 0.00		1381.57±23.65			3301.69 ± 52.93	
	4	546.33 ± 0.03	3491.24 ± 47.65		4079.17 ± 53.20	0.21 ± 0.01	1229.71 ± 20.37	
	5	527.13 ± 0.03	1328.21 ± 84.26		2224.08 ± 87.52	0 ± 0	480.43 ± 16.81	

S2 Additional spectra from NaCl

S2.1 Na 1s temperature dependence

Measured Na 1s core level spectra from deposited NaCl nanoparticles at various temperatures are shown in Fig. S1. The spectra are all taken at 0 mbar water vapor pressure, both before and after sample was exposed to water vapor in the ambient pressure cell.




Figure S1: Change in Na 1s spectra of deposited NaCl aerosol particles with temperature. Photon energy was 1486.6 eV from the Al anode.

S2.2 UHV spectra

Spectra for Na 1s and Cl 2p from NaCl particles measured at UHV conditions are shown in Fig. S2 and Fig. S3, respectively.




Figure S2: Na 1s spectrum measured at UHV conditions. Photon energy was 1486.6 eV from the Al anode.




Figure S3: Cl 2p spectrum measured at UHV conditions. Photon energy was 1486.6 eV from the Al anode.

References

- [1] Ferreira Jr., J. M., Trindade, G. F., Tshulu, R., Watts, J. F., and Baker, M. A.: Dicarboxylic acids analysed by x-ray photoelectron spectroscopy, Part I - propanedioic acid anhydrous, *Surface Science Spectra*, 24, 011 101–8, 2017.
- [2] Jain, V., Biesinger, M. C., and Linford, M. R.: The Gaussian-Lorentzian Sum, Product, and Convolution (Voigt) functions in the context of peak fitting X-ray photoelectron spectroscopy (XPS) narrow scans, *Applied Surface Science*, 447, 548–553, 2018.
- [3] Stevens, J. S. and Schroeder, S. L. M.: Quantitative analysis of saccharides by X-ray photoelectron spectroscopy, *Surface and Interface Analysis*, 41, 453–462, 2009.