



Supplement of

Trends, composition, and sources of carbonaceous aerosol at the Birkenes Observatory, northern Europe, 2001–2018

Karl Espen Yttri et al.

Correspondence to: Karl Espen Yttri (key@nilu.no)

The copyright of individual parts of the supplement might differ from the article licence.

S1. Quality assurance

The OC/EC data are not field blank corrected, in accordance with the standard operating procedure provided by EMEP (Yttri et al., 2007a; EMEP, 2014). The positive sampling artefact of OC for weekly samples collected at Birkenes has been quantified on a campaign basis using the QBQ (Quartz fibre filter Behind Quartz fibre filter) approach (McDow and Huntzicker, 1990; Turpin et al., 1994) in summer ($18\pm4\%$; Yttri et al., 2011b), fall ($19\pm7\%$; Yttri et al., 2019), and winter/spring ($24\pm13\%$; Yttri et al., 2019) but only for PM_{10} . For OC in $PM_{2.5}$, which at Birkenes is obtained from an identical and co-located sampler, operating at the same filter face velocity as the PM_{10} sampler, the positive sampling artefact is considered equally large, whereas its relative importance is slightly higher. The negative sampling artefact has not been addressed.

OC/EC analysis was performed within 2 months after the filter samples were collected and according to the Quartz (2001–2008) and the EUSAAR-2 (from 2008) temperature programs. EUSAAR-2 is designed to reduce the inherited uncertainties associated with splitting of OC and EC, e.g. by preventing premature burn-off of EC (Cavalli et al., 2010). The uncertainty associated with repeated OC/EC analyzes of a filter sample is typically $<10\%$, which includes both analytical uncertainty and heterogenic distribution of the deposited aerosol particles on the filter sample.

The laser's ability to detect changes in the transmittance of a filter sample high in initial EC is crucial to obtain a correct value for EC (and OC). $15\text{ }\mu\text{g EC cm}^{-2}$ has been suggested as an upper limit (Subramanian et al., 2004; Wallén et al., 2010) but this value is likely to vary. The nine filter samples (out of nearly 1800) with an EC content exceeding $15\text{ }\mu\text{g C cm}^{-2}$ in the current dataset were considered valid. Further, a non-biased separation between OC and EC requires that either pyrolytic carbon (PC) evolves before EC during analysis or that PC and EC have the same light absorption coefficient. It is well known that this is not always the case (Yang and Yu, 2002) and there is a lack of information on the magnitude of this imperfection.

Deviation from the protocol-defined temperature steps will affect the analysis results of the TOA instrument (Chow et al., 2005; Panteliadis et al., 2015) and temperature offsets ranging from $-93\text{ }^{\circ}\text{C}$ to $+100\text{ }^{\circ}\text{C}$ per temperature step have been reported (Panteliadis et al., 2015). Thus, calibration by the temperature calibration kit available from the instrument manufacturer (Sunset laboratory Inc) since 2012 is strongly recommended. Temperature calibration was implemented as part of the regular QA/QC procedures for thermal-optical analysis in 2013.

A comparison of the two temperature programmes used for the Birkenes time series was performed for $PM_{2.5}$ filter samples collected at Birkenes in 2014, using temperature calibrated versions of both Quartz and EUSAAR-2. There was a good agreement between the two temperature programs for TC and OC, i.e. close to the expected uncertainty associated with analysis and sampling, whereas for EC the difference was pronounced (Table S 17), although in close correspondence with that previously reported by Panteliadis et al. (2015). Note that OC and EC data for the period 2001–2007 discussed in the main are text not corrected according to Eq. (S 18–20) (Table S 17), except for the

purpose of trend calculations.

Field blanks did not contain monosaccharide anhydrides, sugars, sugar-alcohols or 2-methyltetrols in noticeable amounts. Filter samples for which the content was below the limit of detection (LOD) but > 0 , were considered valid and included when calculating the annual and seasonal means. Organic tracers were analyzed within 1 year after collection of the aerosol filter samples. The uncertainty (analytical and sampling uncertainty) associated with measurements of monosaccharide anhydrides is within 10 – 15 % (Yttri et al., 2015). A similar range of uncertainty is expected for the other organic tracers.

Mass concentrations of PM_{10} and $PM_{2.5}$ were field blank corrected. The overall uncertainty associated with determination of the PM_{10} and $PM_{2.5}$ mass concentration is $< 5\%$. The monitoring of major ions and trace elements follows the guidelines by EMEP (EMEP, 2014) and are within the data quality objective of the network: 15–25% uncertainty for the combined sampling and analysis of major ions and 30% for heavy metals.

S2. Calculation of trends - Statistical approach

The Mann-Kendall test (Mann, 1945; Kendall, 1975; Gilbert, 1987) was used for calculating the significance of the trend and if a significant trend was found, the Theil-Sen slope (Theil, 1950; Sen, 1968; Gilbert, 1987) was calculated. This procedure has been widely used in atmospheric science, like in the recent TOAR project analysing global surface ozone trends (e.g. Fleming et al., 2018; Lefohn et al., 2018), in the review of the EMEP observations (Tørseth et al., 2012) and in numerous other observation based papers (Aas et al. 2019; Ciarelli et al., 2019; Theobald et al., 2019; Masiol et al., 2019; Collaud Coen et al., 2020).

The Mann-Kendall test is a non-parametric test that does not rely on any assumptions of distribution and is therefore well suited for atmospheric data that often deviate from normality and contain outliers that would hamper a standard linear regression. The basics of the Mann-Kendall test is to count the signs of all forward concentration differences in time, and if there is a sufficient overweight of positive or negative differences, the 0-hypothesis (H_0) of no trend could be rejected. The S statistic given below contains the sum of all the signs based on the observed values y_i at time i :

$$S = \sum_{j=1}^{n-1} \sum_{i=j+1}^n \text{sign}(y_j - y_i) \quad \text{Eq. S1}$$

This statistic together with the number of samples and the number of ties in the data were used to calculate the p value as given by Gilbert (1987). In our work, we assumed significant trends when $p < 0.05$.

With $p < 0.05$ H_0 was rejected and the value of the trend was estimated by the Theil-Sen slope estimator:

$$\beta = \text{median} \left(\frac{y_j - y_i}{t_j - t_i} \right), \quad j > i \quad \text{Eq. S2}$$

where t_i denotes the time i of the observed value y_i .

The Theil-Sen slope is simply the median of all the forward concentration gradients. In addition to the slope, the 2σ confidence intervals were calculated according to Gilbert (1987), providing the 95 % confidence range of the slopes.

The Mann-Kendall test and Theil-Sen slope estimation were applied to all species and ratios discussed in this work. These calculations were based on the seasonal and annual mean values, separately, as presented below. For the ratios, $r = x/y$ (e.g. the fraction of NO_3^- in PM_{10}), we based the calculations on the ratios of the seasonal means and not on the seasonal means of the ratios, i.e.:

$$r = \frac{x}{y}, \text{ where } x = \frac{1}{n} \sum (x_i) \text{ and } y = \frac{1}{n} \sum (y_i) \quad \text{Eq. S3}$$

For all cases where the 0-hypothesis (H_0) could be rejected, the Theil-Sen slopes were calculated, and this slope was further transferred into the relative trend by dividing the trend (β) by the mean of the observed values:

$$\beta_{\text{rel}} = \frac{\beta}{\left[\frac{1}{n} \sum (y_i) \right]}, \text{ where } y_i = \text{observed concentration or ratio at time } i \quad \text{Eq. S4}$$

S3. Absorption coefficient measurements and source apportionment

The absorption coefficient (B_{Abs}) was measured using the multi wavelength ($\lambda=370; 470; 520; 590; 660; 880; 950$ nm) aethalometer (AE33, Magee Scientific), operating behind a PM_{10} inlet. We calculate absorption coefficients (B_{Abs}) according to Drinovec et al. (2015):

$$B_{\text{Abs}}(\lambda) = \frac{A \cdot \left(\frac{ATN_{t2}(\lambda) - ATN_{t1}(\lambda)}{100} \right)}{Q \cdot C \cdot (1 - \zeta) \cdot \left(1 - k(\lambda) \cdot (ATN_{t2}(\lambda) - ATN_{\text{ref}}(\lambda)) \right) \cdot (t_2 - t_1)} \quad \text{Eq. S5}$$

where ATN = attenuation at time $t = 1$ and $t = 2$, and of the reference spot ref , Q is the instrument flow rate on spot 1, A is the filter spot area, k is the loading compensation parameter from the 2 spot compensation algorithm. Here we neglect lateral air flow losses (ζ) and the scattering compensation C since these are not wavelength dependent in Eq. (S5) and hence do not affect source apportionment based on wavelength dependence, while conversion to eBC via co-located filter measurements of EC also results in compensation of these parameters using:

$$eBC(\lambda) = B_{Abs}(\lambda) / \alpha_{effective}(\lambda) \quad Eq. S6$$

where $\alpha_{effective}$ is an effective mass absorption cross section (α) incorporating scattering and lateral flow losses:

$$\alpha_{effective}(\lambda) = \alpha(\lambda) \times c \times (1 - \zeta) \quad Eq. S7$$

Hence $\alpha_{effective}$ is a conversion factor between B_{Abs} and eBC and has no physical meaning beyond this.

The AE33 of this study automatically generates $B_{Abs}(\lambda)$ at 1-minute resolution. However, as discussed by Springston et al. (2007) and Backman et al. (2017), the time interval ($t_2 - t_1$) Eq.(S5) can be adjusted to any integer multiple of the base resolution in post-processing. Here we adopt the approach of Backman et al. (2017), fixing the time interval to 1 hour and calculating $B_{Abs}(\lambda)$ according to Eq. (S5). In case one or more filter advances occurred within the one-hour interval, data from each individual filter spot falling within the interval were treated separately and a time-weighted average recorded for that hour. The advantage of this technique is enhanced noise reduction, i.e. using the one-hour interval approach the noise reduction is proportional to as much as $1/n$ (where n are the measurement points), rather than $1/\sqrt{n}$, attainable via signal averaging.

Here we performed source apportionment of aethalometer data using the *aethalometer model* (Sandradewi et al., 2008). Assuming two sources contribute to total Babs ($B_{Abs,Tot}$), i.e. fossil fuel combustion ($B_{Abs,ff}$) and biomass burning ($B_{Abs,bb}$):

$$B_{Abs,Tot} = B_{Abs,ff} + B_{Abs,bb} \quad Eq. S8$$

Then, using a wavelength pair, here $\lambda_1=470$ nm and $\lambda_2=880$ nm,

$$B_{Abs,bb}(\lambda_2) = \frac{B_{Abs}(\lambda_1) - B_{Abs}(\lambda_2) \cdot \left(\frac{\lambda_1}{\lambda_2}\right)^{-\alpha_{ff}}}{\left(\frac{\lambda_1}{\lambda_2}\right)^{-\alpha_{bb}} - \left(\frac{\lambda_1}{\lambda_2}\right)^{-\alpha_{ff}}} \quad Eq. S9 \text{ and}$$

$$B_{Abs,ff}(\lambda_2) = \frac{B_{Abs}(\lambda_1) - B_{Abs}(\lambda_2) \cdot \left(\frac{\lambda_1}{\lambda_2}\right)^{-\alpha_{bb}}}{\left(\frac{\lambda_1}{\lambda_2}\right)^{-\alpha_{ff}} - \left(\frac{\lambda_1}{\lambda_2}\right)^{-\alpha_{bb}}} \quad Eq. S10$$

where α_{ff} and α_{bb} are the absorption Ångström exponents (AAE) for fossil fuel and biomass burning, respectively. Note that when using this approach, the AAEs must be assumed *a priori*, while the data are not fitted or error weighted, which can lead to negative values in the resulting time series of the factors due to uncertainty in the AAEs e.g. Grange et al. (2020).

Here we also used positive matrix factorisation (PMF) to distinguish between the two sources in Eq. (S8). The theory of PMF is detailed elsewhere (Paatero and Tapper, 1994) Briefly, a matrix of measurement data X is represented by a bilinear model comprising factor profiles F (rows), factor time series G (columns) and a residual matrix E :

$$X = G \cdot F + E \quad Eq. S11$$

In PMF factors are found using a least-squares fitting routine in which the object function Q , i.e. the square of residuals e weighted to uncertainty σ , is minimised across all cells (rows i - m , columns j - n)

$$Q^m = \sum_{i=1}^m \sum_{j=1}^n \left(\frac{e_{ij}}{\sigma_{ij}} \right)^2 \quad \text{Eq. S12}$$

Here, we use the source finder (SoFi, (Canonaco et al., 2013)) toolkit ref, to call PMF (To model the error matrix σ_{ij} we use the clean air test function of the AE33 to determine the standard deviation of the attenuation of the blank $\delta_{ATN_{air}}$, calculating σ_{ij} , using:

$$\sigma_{ij} = \sqrt{f_A^2 + f_Q^2 + 2 \left(\frac{\delta_{ATN_{air}}(\lambda_j)}{ATN_i(\lambda_j)} \right)^2 + \left(\frac{\delta_{ATN_{air}}(\lambda_j)}{ATN_{i-1}(\lambda_j)} \right)^2 + \left(\frac{\delta_{ATN_{air}}(\lambda_j)}{ATN_{ref}(\lambda_j)} \right)^2} \cdot B_{Abs,i}(\lambda_j) \quad \text{Eq. S13}$$

where f_A and f_Q are the fractional uncertainties in the spot area and the flow rate, respectively (both 0.015 according to Backman et al., 2017). Clean air tests were performed only periodically. Therefore, to generate an error estimate for all time points, we interpolated (bilinear interpolation) between the clean air tests to generate the full error matrix, accounting for drift in $\delta_{ATN_{air}}$. Points before and after the last clean air test were calculated using the first and last values of $\delta_{ATN_{air}}$, respectively.

According to Eq. (S11), X could be represented by any combination of G and F , i.e. the PMF model has *rotational ambiguity*. In practice, many rotations produce negative values and are thus forbidden. Nevertheless, many rotations and local minima in Eq. (S11) are likely to exist. To assess this, we generated multiple ($n=2000$) bootstrap replacement matrices (block size 24 to conserve diurnal variation if present), running PMF on each matrix 5 times for a total of 10000 runs. PMF settings are shown in Table S 2.

We import all 2000 files generated using SoFi for each factor solution. To map the factors, we calculated an effective AAE from the factor profiles α_F , using

$$\alpha_F = - \frac{\log \left(\frac{F_{j=2}}{F_{j=6}} \right)}{\log(470/880)} \quad \text{Eq. S14}$$

sorting factors and time series from each run from low to high with respect to α_F . Binning the effective AAEs from each factor also provides a convenient means to investigate the solution space for rotational ambiguity.

S4. Positive matrix factorisation applied to filter data

We performed PMF for samples collected in 2016-2018 (151 samples), with the following as input data: OC (in PM_{2.5} and PM_{10-2.5}), EC (in PM₁₀), levoglucosan, mannosan, galactosan, arabinol, mannitol, trehalose, glucose, V, Mn, Ti, Fe, Co, Ni, Cu, Zn, As, Cd, and Pb (all in PM₁₀), SO₄²⁻, NO₃⁻, NH₄⁺, Ca²⁺, Mg²⁺, K⁺, Na⁺, Cl⁻ (open filter face).

Table S 3 shows miscellaneous settings of the PMF analysis of these data. The input data and error estimates were prepared using the procedure suggested by Polissar et al. (1998) and Norris et al. (2014), see also Table S 3 for miscellaneous settings including missing data treatment and assessment of the PMF performance.

If the concentration was greater than the LOD, the calculation was based on a user provided fraction of the concentration and LOD:

$$Unc = \sqrt{(Error\ Fraction \times Concentration)^2 + (\frac{1}{2} \times LOD)^2} \quad Eq. S15$$

The analytical uncertainties (20%) as error fraction of OC, EC, organic tracers, ions, and elements were used to determine the corresponding error estimates. Based on given understanding of OC sources, 2–10 factors with random seeds were examined, and 7 factors were determined based on: 1) The decrease in Q/Q_{exp} was larger than the relative change in number of factors up to 7; 2) All factors could be interpreted; 3) All factors were distinct.

To assess the statistical uncertainty in the model we performed repeated analyses on bootstrap-resampled matrices. A base profile was generated from a manually mapped average of 50 runs. From each bootstrap run, we fitted all 7 bootstrap factors vs all 7 factors from the base run profile (representing a 7×7 matrix of r² values). We then mapped the bootstrap factors in order of the r² value: The highest value was assumed to be a match, then then the next highest value excluding both previously mapped factors to any other factor (representing a 6×6 matrix of r² values), and so on. This was to avoid any factors being mapped twice.

The minimal robust and true Q values of the base run were 5507.9 and 5580.8, respectively. All the (error) scaled residuals were within ±5 and > 97.8% within ± 3, normally distributed and centred around zero. The average Q/Q_{exp} was 1.2. We also observe no structure in the residuals, which were evenly distributed between measurements from different instruments (i.e. we did not observe factors representing groups of compounds by instrument type, Figure S 3).

S5. Emission ratios used to calculate OC and EC from biomass burning

Emission ratios derived from ambient data are a good alternative to direct emission measurements, accounting for the aggregate effects of fuel type and combustion conditions, but results will nevertheless vary from region to region (e.g. Zotter et al., 2014). Here, we used ratios from our PMF analysis (Table 1) to calculate carbonaceous aerosol from biomass burning for 2008–2018. The levoglucosan to

mannosan ratio is rather consistent between seasons, with the values for summer (5.1 ± 0.9) and fall (5.2 ± 0.7) being slightly lower than for winter (5.4 ± 0.8) and spring (6.0 ± 0.7). This might indicate that emissions from one source of biomass burning (wood burning for residential heating) dominate for all seasons, supporting the use of one levoglucosan to OC (and EC) ratio for calculations. The lower levoglucosan to mannosan ratio observed in summer and fall might indicate increased influence of wild and agricultural fires, but the magnitude of these sources remains speculative, except during severe episodes, e.g. in August 2002, May and September 2006, and June 2008.

S6. Levels of PBAP and BSOA organic tracers

The annual mean concentration of the PBAP tracers ranged from 2.8–3.4 ng m⁻³ (trehalose) to 4.8–5.8 ng m⁻³ (arabitol) (2016–2018) (Figure 6, Table S 15). Levels were elevated in the vegetative season, particularly in summer and fall. Mannitol and arabitol were highly correlated ($R^2=0.85$), underlining their common origin, and the mannitol to arabitol ratio (0.9 ± 0.2) corresponds well with previously reported results for these fungal spore tracers (e.g. Bauer et al., 2008; Yttri et al., 2007b; Yttri et al. 2011 a, b).

The annual mean concentration of 2-methylerythritol ($0.365\text{--}0.441$ ng m⁻³) (2016–2018) was higher than that of 2-methylthreitol ($0.105\text{--}162$ ng m⁻³), and the two isomers were highly correlated ($R^2=0.915$), which is consistent with other studies (e.g., Ion et al., 2005; Kourtchev et al., 2005; Edney et al., 2005; El Haddad et al., 2011; Alier et al., 2013). 2-methyltetrols were elevated in the period when deciduous trees have leaves (transition May/June to early October).

References Supplementary

Aas, W., Mortier, A., Bowersox, V., Cherian, R., Faluvegi, G., Fagerli, H., Hand, J., Klimont, Z., Galy-Lacaux, C., Lehmann, C. M. B., Myhre, C. L., Myhre, G., Olivie, D., Sato, K., Quaas, J., Rao, P. S. P., Schulz, M., Shindell, D., Skeie, R. B., Stein, A., Takemura, T., Tsyro, S., Vet, R., and Xu, X. B.: Global and regional trends of atmospheric sulfur (vol 9, 953, 2019), *Sci. Rep.*, 10, 10.1038/s41598-020-62441-w, 2020.

Alier, M., van Drooge, B. L., Dall'Osto, M., Querol, X., Grimalt, J. O., and Tauler, R.: Source apportionment of submicron organic aerosol at an urban background and a road site in Barcelona (Spain) during SAPUSS, *Atmos. Chem. Phys.*, 13, 10353-10371, 10.5194/acp-13-10353-2013, 2013.

Backman, J., Schmeisser, L., Virkkula, A., Ogren, J. A., Asmi, E., Starkweather, S., Sharma, S., Eleftheriadis, K., Uttal, T., Jefferson, A., Bergin, M., Makshtas, A., Tunved, P., and Fiebig, M.: On Aethalometer measurement uncertainties and an instrument correction factor for the Arctic, *Atmos. Meas. Techn.*, 10, 5039-5062, 10.5194/amt-10-5039-2017, 2017.

240 Bauer, H., Claeys, M., Vermeylen, R., Schueller, E., Weinke, G., Berger, A., and Puxbaum, H.: Arabitol
 241 and mannitol as tracers for the quantification of airborne fungal spores, *Atmos. Environ.*, 42, 588-593,
 242 10.1016/j.atmosenv.2007.10.013, 2008.

243 Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U., and Prevot, A. S. H.: SoFi, an IGOR-based
 244 interface for the efficient use of the generalized multilinear engine (ME-2) for the source apportionment:
 245 ME-2 application to aerosol mass spectrometer data, *Atmos. Meas. Tech.*, 6, 3649-3661, 10.5194/amt-
 246 6-3649-2013, 2013.

247 Cavalli, F., Viana, M., Yttri, K. E., Genberg, J., and Putaud, J.-P.: Toward a standardised thermal-optical
 248 protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol, *Atmos.*
 249 *Meas. Tech.*, 3, 79-89, 2010.

250 Chow, J. C., Watson, J. G., Chen, L. W. A., Paredes-Miranda, G., Chang, M. C. O., Trimble, D., Fung,
 251 K. K., Zhang, H., and Yu, J. Z.: Refining temperature measures in thermal/optical carbon analysis,
 252 *Atmos. Chem. Phys.*, 5, 2961-2972, 10.5194/acp-5-2961-2005, 2005.

253 Ciarelli, G., Theobald, M. R., Vivanco, M. G., Beekmann, M., Aas, W., Andersson, C., Bergstrom, R.,
 254 Manders-Groot, A., Couvidat, F., Mircea, M., Tsyro, S., Fagerli, H., Mar, K., Raffort, V., Roustan, Y.,
 255 Pay, M. T., Schaap, M., Kranenburg, R., Adani, M., Briganti, G., Cappelletti, A., D'Isidoro, M.,
 256 Cuvelier, C., Cholakian, A., Bessagnet, B., Wind, P., and Colette, A.: Trends of inorganic and organic
 257 aerosols and precursor gases in Europe: insights from the EURODELTA multi-model experiment over
 258 the 1990-2010 period, *Geosci. Model Dev.*, 12, 4923-4954, 10.5194/gmd-12-4923-2019, 2019.

259 Coen, M. C., Andrews, E., Alastuey, A., Arsov, T. P., Backman, J., Brem, B. T., Bukowiecki, N., Couret,
 260 C., Eleftheriadis, K., Flentje, H., Fiebig, M., Gysel-Beer, M., Hand, J. L., Hoffer, A., Hooda, R.,
 261 Hueglin, C., Joubert, W., Keywood, M., Kim, J. E., Kim, S. W., Labuschagne, C., Lin, N. H., Lin, Y.,
 262 Myhre, C. L., Luoma, K., Lyamani, H., Marinoni, A., Mayol-Bracero, O. L., Mihalopoulos, N., Pandolfi,
 263 M., Prats, N., Prenni, A. J., Putaud, J. P., Ries, L., Reisen, F., Sellegri, K., Sharma, S., Sheridan, P.,
 264 Sherman, J. P., Sun, J. Y., Titos, G., Torres, E., Tuch, T., Weller, R., Wiedensohler, A., Zieger, P., and
 265 Laj, P.: Multidecadal trend analysis of in situ aerosol radiative properties around the world, *Atmos.*
 266 *Chem. Phys.*, 20, 8867-8908, 10.5194/acp-20-8867-2020, 2020.

267 Drinovec, L., Mocnik, G., Zotter, P., Prevot, A. S. H., Ruckstuhl, C., Coz, E., Rupakheti, M., Sciare, J.,
 268 Muller, T., Wiedensohler, A., and Hansen, A. D. A.: The "dual-spot" Aethalometer: an improved
 269 measurement of aerosol black carbon with real-time loading compensation, *Atmos. Meas. Tech.*, 8,
 270 1965-1979, 10.5194/amt-8-1965-2015, 2015.

271 Edney, E. O., Kleindienst, T. E., Jaoui, M., Lewandowski, M., Offenberger, J. H., Wang, W., and Claeys,

272 M.: Formation of 2-methyl tetrols and 2-methylglyceric acid in secondary organic aerosol from
 273 laboratory irradiated isoprene/NOX/SO2/air mixtures and their detection in ambient PM2.5 samples
 274 collected in the eastern United States, *Atmos. Environ.*, 39, 5281-5289,
 275 10.1016/j.atmosenv.2005.05.031, 2005.

276 El Haddad, I., Marchand, N., Temime-Roussel, B., Wortham, H., Piot, C., Besombes, J. L., Baduel, C.,
 277 Voisin, D., Armengaud, A., and Jaffrezo, J. L.: Insights into the secondary fraction of the organic aerosol
 278 in a Mediterranean urban area: Marseille, *Atmos. Chem. Phys.*, 11, 2059-2079, 10.5194/acp-11-2059-
 279 2011, 2011.

280 EMEP: Standard Operating Procedures for thermal-optical analysis of atmospheric particulate organic
 281 and elemental carbon, in: EMEP manual for sampling and chemical analysis, EMEP/CCC 01/2014,
 282 NILU, Kjeller, Norway, chapter 4.22, 2014.

283 Fleming, Z. L., Doherty, R. M., von Schneidmesser, E., Malley, C. S., Cooper, O. R., Pinto, J. P.,
 284 Colette, A., Xu, X. B., Simpson, D., Schultz, M. G., Lefohn, A. S., Hamad, S., Moolla, R., Solberg, S.,
 285 and Feng, Z. Z.: Tropospheric Ozone Assessment Report: Present-day ozone distribution and trends
 286 relevant to human health, *Elem. Sci. Anth.*, 6, 10.1525/elementa.273, 2018.

287 Grange, S. K., Lotscher, H., Fischer, A., Emmenegger, L., and Hueglin, C.: Evaluation of equivalent
 288 black carbon source apportionment using observations from Switzerland between 2008 and 2018,
 289 *Atmos. Meas. Tech.*, 13, 1867-1885, 10.5194/amt-13-1867-2020, 2020.

290 Gilbert, R. O.: Statistical Methods for Environmental Pollution Monitoring, Wiley, NY, United States,
 291 pp. 336, 1987.

292 Ion, A. C., Vermeylen, R., Kourtchev, I., Cafmeyer, J., Chi, X., Gelencser, A., Maenhaut, W., and
 293 Claeys, M.: Polar organic compounds in rural PM_{2.5} aerosols from K-puszt, Hungary, during a 2003
 294 summer field campaign: Sources and diel variations, *Atmos. Chem. Phys.*, 5, 1805-1814, 10.5194/acp-
 295 5-1805-2005, 2005.

296 Kendall, M. G.: Rank correlation methods, 4th edition, Charles Griffin, London, 1975.

297 Kourtchev, I., Ruuskanen, T., Maenhaut, W., Kulmala, M., and Claeys, M.: Observation of 2-
 298 methyltetrols and related photo-oxidation products of isoprene in boreal forest aerosols from Hyytiala,
 299 Finland, *Atmos. Chem. Phys.*, 5, 2761-2770, 10.5194/acp-5-2761-2005, 2005.

300 Lefohn, A. S., Malley, C. S., Smith, L., Wells, B., Hazucha, M., Simon, H., Naik, V., Mills, G., Schultz,
 301 M. G., Paoletti, E., De Marco, A., Xu, X. B., Zhang, L., Wang, T., Neufeld, H. S., Musselman, R. C.,

302 Tarasick, D., Brauer, M., Feng, Z. Z., Tang, H. Y., Kobayashi, K., Sicard, P., Solberg, S., and Gerosa,
303 G.: Tropospheric ozone assessment report: Global ozone metrics for climate change, human health, and
304 crop/ecosystem research, *Elem. Sci. Anth.*, 6, 10.1525/elementa.279, 2018.

305 Mann, H. B.: Non-parametric tests against trend, *Econometrica* 13:163-171, 1945.

306 Masiol, M., Squizzato, S., Rich, D. Q., and Hopke, P. K.: Long-term trends (2005-2016) of source
307 apportioned PM_{2.5} across New York State, *Atmos. Environ.*, 201, 110-120,
308 10.1016/j.atmosenv.2018.12.038, 2019.

309 McDow, S. R. and Huntzicker, J. J.: Vapor adsorption artifact in the sampling of organic aerosol: face
310 velocity effects, *Atmos. Environ.*, 24A, 2563–2571, 10.1016/0960-1686(90)90134-9, 1990.

311 Norris, G., Duvall, R., Brown, S., and Bai, S.: EPA Positive Matrix Factorization (PMF) 5.0
312 Fundamentals and User Guide, U.S. Environmental Protection Agency, Washington, DC, 20460 (i-124,
313 EPA/600/R-14/108, April), 2014.

314 Paatero, P. and Tapper, U.: Positive Matrix Factorization – A nonnegative factor model with optimal
315 utilization of error-estimates of data values. *Environmetrics*, 5, 111-126, DOI:
316 10.1002/env.3170050203, 1994.

317 Panteliadis, P., Hafkenscheid, T., Cary, B., Diapouli, E., Fischer, A., Favez, O., Quincey, P., Viana, M.,
318 Hitzenberger, R., Vecchi, R., Saraga, D., Sciare, J., Jaffrezo, J. L., John, A., Schwarz, J., Giannoni, M.,
319 Novak, J., Karanasiou, A., Fermo, P., and Maenhaut, W.: ECOC comparison exercise with identical
320 thermal protocols after temperature offset correction - instrument diagnostics by in-depth evaluation of
321 operational parameters, *Atmos. Meas. Tech.*, 8, 779-792, 10.5194/amt-8-779-2015, 2015.

322 Pio, C. A., Legrand, M., Oliveira, T., Afonso, J., Santos, C., Caseiro, A., Fialho, P., Barata, F., Puxbaum,
323 H., Sanchez-Ochoa, A., Kasper-Giebl, A., Gelencser, A., Preunkert, S., and Schock, M.: Climatology
324 of aerosol composition (organic versus inorganic) at nonurban sites on a west-east transect across
325 Europe, *J. Geophys. Res.-Atmos.*, 112, 10.1029/2006jd008038, 2007.

326 Polissar, A. V., Hopke, P. K., and Paatero, P.: Atmospheric aerosol over Alaska - 2. Elemental
327 composition and sources, *J. Geophys. Res.-Atmos.*, 103, 19045-19057, 10.1029/98jd01212, 1998.

328 Sandradewi, J., Prevot, A. S. H., Szidat, S., Perron, N., Alfarra, M. R., Lanz, V. A., Weingartner, E.,
329 and Baltensperger, U.: Using aerosol light absorption measurements for the quantitative determination
330 of wood burning and traffic emission contributions to particulate matter, *Environ. Sci. Technol.*, 42,
331 3316-3323, 10.1021/es702253m, 2008.

332

333 Sen, P. K.: Estimates of the regression coefficient based on Kendall's Tau. *J. Am., Stat. Assoc.* 63 (324),
 334 1379–1389, <https://doi.org/10.2307/2285891>, 1968.

335 Springston, S. R. and Sedlacek, A. J.: Noise characteristics of an instrumental particle absorbance
 336 technique, *Aerosol Sci. and Technol.*, 41, 1110-1116, 10.1080/02786820701777457, 2007.

337 Subramanian, R., Khlystov, A. Y., Cabada, J. C., and Robinson, A. L.: Positive and negative artifacts in
 338 particulate organic carbon measurements with denuded and undenuded sampler configurations, *Aerosol*
 339 *Sci. Technol.*, 38, 27-48, 10.1080/02786820390229354, 2004.

340 Theil, H.: A rank-invariant method of linear and polynomial regression analysis. *Proc. R. Netherlands*,
 341 *Acad. Sci.* 53, 386–392, https://doi.org/10.1007/978-94-011-2546-8_20, 1950.

342 Theobald, M. R., Vivanco, M. G., Aas, W., Andersson, C., Ciarelli, G., Couvidat, F., Cuvelier, K.,
 343 Manders, A., Mircea, M., Pay, M. T., Tsyro, S., Adani, M., Bergstrom, R., Bessagnet, B., Briganti, G.,
 344 Cappelletti, A., D'Isidoro, M., Fagerli, H., Mar, K., Otero, N., Raffort, V., Roustan, Y., Schaap, M.,
 345 Wind, P., and Colette, A.: An evaluation of European nitrogen and sulfur wet deposition and their trends
 346 estimated by six chemistry transport models for the period 1990-2010, *Atmos. Chem. Phys.*, 19, 379-
 347 405, 10.5194/acp-19-379-2019, 2019.

348 Torseth, K., Aas, W., Breivik, K., Fjaeraa, A. M., Fiebig, M., Hjellbrekke, A. G., Myhre, C. L., Solberg,
 349 S., and Yttri, K. E.: Introduction to the European Monitoring and Evaluation Programme (EMEP) and
 350 observed atmospheric composition change during 1972-2009, *Atmos. Chem. Phys.*, 12, 5447-5481,
 351 10.5194/acp-12-5447-2012, 2012.

352 Turpin, B. J., Huntzicker, J. J., and Hering, S. V.: Investigation of organic aerosol sampling artifacts in
 353 the Los-Angeles basin, *Atmos. Environ.*, 28, 3061-3071, 10.1016/1352-2310(94)00133-6, 1994.

354 Wallen, A., Liden, G., and Hansson, H. C.: Measured Elemental Carbon by Thermo-Optical
 355 Transmittance Analysis in Water-Soluble Extracts from Diesel Exhaust, Woodsmoke, and Ambient
 356 Particulate Samples, *J. Occup. Environ. Hyg.*, 7, 35-45, 10.1080/15459620903368859, 2010.

357 Yang, H. and Yu, J. Z.: Uncertainties in charring correction in the analysis of elemental and organic
 358 carbon in atmospheric particles by thermal/optical methods, *Environ. Sci. Technol.*, 36, 5199-5204,
 359 10.1021/es025672z, 2002.

360 Yttri, K. E., Dye, C., and Kiss, G.: Ambient aerosol concentrations of sugars and sugar-alcohols at four
 361 different sites in Norway, *Atmos. Chem. Phys.*, 7, 4267-4279, 2007b.

362 Yttri, K. E., Aas, W., Bjerke, A., Cape, J. N., Cavalli, F., Ceburnis, D., Dye, C., Emblico, L., Facchini,
 363 M. C., Forster, C., Hanssen, J. E., Hansson, H. C., Jennings, S. G., Maenhaut, W., Putaud, J. P., and
 364 Torseth, K.: Elemental and organic carbon in PM₁₀: a one year measurement campaign within the
 365 European Monitoring and Evaluation Programme EMEP, *Atmos. Chem. Phys.*, 7, 5711-5725, 2007a.

366 Yttri, K. E., Simpson, D., Bergstrom, R., Kiss, G., Szidat, S., Ceburnis, D., Eckhardt, S., Hueglin, C.,
 367 Nojgaard, J. K., Perrino, C., Pisso, I., Prevot, A. S. H., Putaud, J. P., Spindler, G., Vana, M., Zhang, Y.
 368 L., and Aas, W.: The EMEP Intensive Measurement Period campaign, 2008-2009: characterizing
 369 carbonaceous aerosol at nine rural sites in Europe, *Atmos. Chem. Phys.*, 19, 4211-4233, 10.5194/acp-
 370 19-4211-2019, 2019.

371 Yttri, K. E., Simpson, D., Nojgaard, J. K., Kristensen, K., Genberg, J., Stenstrom, K., Swietlicki, E.,
 372 Hillamo, R., Aurela, M., Bauer, H., Offenberg, J. H., Jaoui, M., Dye, C., Eckhardt, S., Burkhardt, J. F.,
 373 Stohl, A., and Glasius, M.: Source apportionment of the summer time carbonaceous aerosol at Nordic
 374 rural background sites, *Atmos. Chem. Phys.*, 11, 13339-13357, 10.5194/acp-11-13339-2011, 2011b.

375 Yttri, K. E., Simpson, D., Stenstrom, K., Puxbaum, H., and Svendby, T.: Source apportionment of the
 376 carbonaceous aerosol in Norway - quantitative estimates based on C-14, thermal-optical and organic
 377 tracer analysis, *Atmos. Chem. Phys.*, 11, 9375-9394, 10.5194/acp-11-9375-2011, 2011a.

378 Yttri, K. E., Schnelle-Kreis, J., Maenhaut, W., Abbaszade, G., Alves, C., Bjerke, A., Bonnier, N., Bossi,
 379 R., Claeys, M., Dye, C., Evtyugina, M., Garcia-Gacio, D., Hillamo, R., Hoffer, A., Hyder, M., Iinuma,
 380 Y., Jaffrezo, J.-L., Kasper-Giebl, A., Kiss, G., Lopez-Mahia, P. L., Pio, C., Piot, C., Ramirez-Santa-
 381 Cruz, C., Sciare, J., Teinila, K., Vermeylen, R., Vicente, A., and Zimmermann, R.: An intercomparison
 382 study of analytical methods used for quantification of levoglucosan in ambient aerosol filter samples,
 383 *Atmos. Meas. Tech.*, 8, 125-147, 10.5194/amt-8-125-2015, 2015.

384 Zotter, P., Ciobanu, V. G., Zhang, Y. L., El-Haddad, I., Macchia, M., Daellenbach, K. R., Salazar, G.
 385 A., Huang, R. J., Wacker, L., Hueglin, C., Piazzalunga, A., Fermo, P., Schwikowski, M., Baltensperger,
 386 U., Szidat, S., and Prevot, A. S. H.: Radiocarbon analysis of elemental and organic carbon in Switzerland
 387 during winter-smog episodes from 2008 to 2012-Part 1: Source apportionment and spatial variability,
 388 *Atmos. Chem. Phys.*, 14, 13551-13570, 10.5194/acp-14-13551-2014, 2014.

Supplementary Figures




Figure S 1: Annual and seasonal ambient mean (point), 10th to 25th percentile (bar), 50th percentile (line), 75th to 90th percentile (whisker) temperature (left panel) and precipitation (right panel) at the Birkenes Observatory, 2001–2018.




Figure S 2: Age spectra for June 2008 for the CO tracer originating from wildfires calculated for the Birkenes Observatory. We ran a 20 days FLEXPART simulation backwards in time, releasing 40 000 particles for the Birkenes Observatory on a 3-hourly basis, using ECMWF meteorology. With daily MODIS information of burned area we constructed a CO emission inventory, which we combined with the model simulation. With this approach we achieved a time series of CO from wildfires with a 3-hourly time resolution. Additionally, we split the modeled CO concentrations by age. The spectrum goes from 1 to 20 days after release according to the bar. This approach is described in more detail in Stohl et al. (2007). For most of June 2008, concentrations of a few ppb were calculated for the Birkenes Observatory, except for 11–12 of June (100 ppb). The age of the airmasses were only 1 day, which means that the CO was released on a location less than 24 hours before it reached the site.




Figure S 3: Average error scaled residuals for each of the variables in the PMF solution presented in this paper

Supplementary Tables

Table S 1: Species quantification ion, molecular weight and formula, and internal/recovery and quantification standards used for identification and quantification.

Species	Quantitation ion M-H(+)	Molecular weight	Molecular formula	Internal/ Recovery standard	Quantification standard
<i>Monosaccharide anhydrides</i>					
Galactosan	161.046	162.141	C ₆ H ₁₀ O ₅	¹³ C ₆ -Galactosan (CIL; 98%; Andover, MA)	Galactosan (Sigma; purity not given; Product of England)
Mannosan	161.046	162.141	C ₆ H ₁₀ O ₅	¹³ C ₆ -Levoglucosan ¹ ¹³ C ₆ -Galactosan ¹	Mannosan (Sigma; Approx 98%; Product of England)
Levoglucosan	161.046	162.141	C ₆ H ₁₀ O ₅	¹³ C ₆ -Levoglucosan (CIL; 98%; city not given)	Levoglucosan (Aldrich; 99%; Product of Switzerland)
<i>Sugar-alcohols</i>					
Mannitol	181.072	182.172	C ₆ H ₁₄ O ₆	¹³ C ₆ -Mannitol (Omnicon Biochemicals Inc; 99.70%; South Bend, Indiana)	Mannitol (ICN Biochemicals; ACS reagent grade; Aurora, Ohio)
Arabitol	151.061	152.15	C ₅ H ₁₂ O ₆	¹³ C ₅ -Arabitol (Omnicon Biochemicals Inc; 99.60%; South Bend, Indiana)	Arabitol (ICN Biochemicals; purity not given; Aurora, Ohio)
<i>2-methyltetrols</i>					
2-Methylerythritol	135.066	136.147	C ₅ H ₁₂ O ₄	¹³ C ₆ -Galactosan (CIL; 98%; Andover, MA)	2-Methylerythritol, Produced at UNC ²
2-Methylthreitol	135.066	136.147	C ₅ H ₁₂ O ₄	¹³ C ₆ -Galactosan (Brand; purity; city)	2-Methylthreitol, Produced at UNC ²
<i>Dimeric sugars</i>					
Trehalose	341.109	342.296	C ₁₂ H ₂₂ O ₁₁	¹³ C ₁₂ -Trehalose (Omnicon Biochemicals Inc; 99.70%; South Bend, Indiana)	Trehalose (Fluka; <99.5%; Packed in Switzerland)
<i>Monomeric sugars</i>					
Glucose	179.0561	180.16	C ₆ H ₁₂ O ₆	¹³ C ₁ -Glucose (CIL; 99%; Andover, MA)	Glucose (Sigma; purity not given; city not given)

1.C-labelled mannosan is not commercially available, hence we used the average of ¹³C₆-Levoglucosan ¹³C₆-Galactosan to calculate the recovery of mannosan.

2.Standard produced by University of North Carolina (UNC)

6 **Table S 2: Settings used for absorption coefficient PMF analysis.**

Parameter	Setting	7
Data matrix dimensions i×j	5240×7	
Missing data treatment	Rows removed	
Number of factors	2	
Factor constraints	None	
Robust mode setting	4	
Seed	Random	
Bootstrap replacement runs	2000	
Block size	24	
Repeat runs (per bootstrap)	5	

8

9

10 **Table S 3: Miscellaneous settings of PMF analysis.**

Parameter	Setting
Data matrix dimensions $i \times j$	151×34
Missing data and below detection limit data	Replaced with geometric mean concentration
Missing data error estimate	Replaced with $4 \times$ geometric mean concentration
Missing data error estimate	$5/6 \times$ limit of detection
Number of factors	7
Factor constraints	None
Robust mode setting	4
Seed	Random
Bootstrap replacement runs	5000
Block size	1 row
Repeat runs (per bootstrap)	5

11

Table S4: Contribution weighted relative profiles for PMF-derived factors (%).

	Mineral Dust (MIN)	Traffic/Industry (TRA/IND)	Biogenic Secondary Organic Aerosol (BSOA _{SRT})	Primary Biological Aerosol Particle (PBAP)	Sea salt aerosol (SS)	Biomass burning (BB)	Ammonium Nitrate (NH ₄ NO ₃)
PM_{2.5} OC	31.3	9.7	9.1	15.6	0.9	16.7	16.6
PM₁₀ EC	12.8	50.0	0.0	2.6	0.0	21.2	13.4
PM_{10-2.5} OC	12.6	3.5	12.5	53.0	0.1	6.3	12.0
Levogluconan	0.2	0.0	1.5	0.5	0.0	97.8	0.0
Mannosan	0.0	0.5	1.5	1.5	0.0	95.6	0.9
Galactosan	0.0	3.5	0.0	0.0	1.1	95.5	0.0
2-methylerythritol	0.5	0.7	95.9	1.8	0.0	0.4	0.7
2-methylthreitol	0.6	1.3	91.5	3.2	0.7	1.5	1.2
Glucose	1.0	0.6	6.5	81.6	2.0	5.2	3.1
Mannitol	0.1	0.3	6.1	91.3	1.7	0.0	0.5
Arabitol	0.0	0.0	8.6	89.5	1.3	0.5	0.0
Trehalose	0.5	0.0	3.3	93.5	1.1	0.7	0.9
Ammonium	2.1	13.9	4.8	0.0	1.5	1.0	76.7
Calcium	39.0	6.0	7.7	3.9	34.9	1.0	7.6
Chloride	0.0	0.0	0.0	0.0	96.2	0.2	3.5
Magnesium	5.8	2.7	3.9	1.8	79.0	0.6	6.2
Nitrate	0.8	3.6	5.7	3.4	17.1	1.7	67.8
Potassium	3.6	12.7	4.6	8.3	28.4	10.4	32.0
Sodium	2.2	5.1	2.7	0.0	86.8	0.3	3.0
Sulfate	3.9	19.6	17.2	3.3	17.4	3.4	35.2
NO₃⁻	7.2	15.0	4.6	4.6	20.0	19.0	29.7
V	14.1	70.1	10.1	0.0	0.0	0.0	5.8
Mn	51.9	38.6	0.6	5.7	2.4	0.8	0.0
Ti	93.4	0.5	0.6	0.0	3.7	1.8	0.0
Fe	74.5	18.4	0.0	3.2	0.7	1.0	2.1
Co	42.6	42.1	1.3	3.3	5.0	2.4	3.4
Ni	17.1	68.6	3.0	4.5	1.7	2.1	2.9

	Mineral Dust (MIN)	Traffic/Industry (TRA/IND)	Biogenic Secondary Organic Aerosol (BSOA _{SRT})	Primary Biological Aerosol Particle (PBAP)	Sea salt aerosol (SS)	Biomass burning (BB)	Ammonium Nitrate (NH ₄ NO ₃)
Cu	19.9	61.9	3.3	3.7	5.7	1.3	4.2
Zn	4.3	81.5	1.4	0.6	0.0	5.8	6.4
As	3.2	78.4	1.7	6.2	0.0	4.8	5.7
Cd	5.4	80.5	0.1	2.3	1.6	6.4	3.8
Pb	4.5	83.9	0.8	0.0	1.5	2.8	6.4

14 **Table S 5: Annual and seasonal mean concentrations of EC and OC in PM₁₀, PM_{2.5} and PM_{10-2.5} at Birkenes for 2001–**
15 **2018 (Unit: $\mu\text{g C m}^{-3}$).**

	PM ₁₀		EC		PM _{10-2.5}		PM _{2.5}		EC	
	OC	Capture		Capture	OC	Capture	OC	Capture		Capture
2001	0.96	56	0.14	56	0.08	45	0.93	56	0.15	56
DJF	0.60	59	0.12	59	NA	34	0.64	59	0.12	59
MAM	0.98	92	0.16	92	0.03	73	1.00	92	0.18	92
JJA	2.34	26	0.16	26	0.24	26	2.1	26	0.20	26
SON	0.58	48	0.10	48	0.09	48	0.49	48	0.11	48
2002	1.01	85	0.14	85	0.190	80	0.89	91	0.12	91
DJF	0.53	95	0.12	95	0.05	88	0.49	95	0.12	95
MAM	1.30	92	0.17	92	0.12	92	1.19	92	0.14	92
JJA	1.77	68	0.20	68	0.48	68	1.4	91	0.14	91
SON	0.59	86	0.10	86	0.16	73	0.49	85	0.09	85
2003	1.01	82	0.10	82	0.23	77	0.81	81	0.11	81
DJF	0.83	85	0.11	85	0.06	76	0.84	85	0.12	85
MAM	1.13	86	0.13	86	0.21	76	1.02	86	0.15	86
JJA	1.26	85	0.08	85	0.44	85	0.82	85	0.09	85
SON	0.75	70	0.09	70	0.22	70	0.53	70	0.09	70
2004	0.82	85	0.10	85	0.27	81	0.57	84	0.09	84
DJF	0.57	86	0.08	86	0.09	86	0.48	86	0.08	86
MAM	1.13	86	0.12	86	0.28	73	0.79	86	0.12	86
JJA	0.99	79	0.10	79	0.38	79	0.61	79	0.08	79
SON	0.71	86	0.08	86	0.32	86	0.39	86	0.08	80
2005	0.85	79	0.14	79	0.29	75	0.60	80	0.12	80
DJF	0.46	64	0.10	64	0.06	51	0.53	71	0.11	71
MAM	0.78	85	0.13	86	0.15	85	0.63	85	0.12	85
JJA	0.92	86	0.09	85	0.33	86	0.59	86	0.08	86
SON	1.16	79	0.25	79	0.53	79	0.62	79	0.16	79
2006	1.07	78	0.13	78	0.33	75	0.77	78	0.13	78
DJF	0.79	86	0.12	86	0.08	79	0.71	86	0.17	86
MAM	0.95	85	0.08	85	0.16	78	0.82	85	0.14	85
JJA	1.43	57	0.12	57	0.48	57	0.96	57	0.10	57
SON	1.24	86	0.19	86	0.60	86	0.64	86	0.11	86
2007	0.82	79	0.15	77	0.21	77	0.61	77	0.13	77
DJF	0.58	58	0.17	58	0.08	58	0.50	58	0.17	58
MAM	0.99	86	0.18	85	0.17	86	0.82	86	0.15	86
JJA	1.03	86	0.13	86	0.39	79	0.66	79	0.10	79
SON	0.60	86	0.13	79	0.18	86	0.41	86	0.10	86
2008	0.75	87	0.09	84	0.24	75	0.53	88	0.08	88
DJF	0.44	90	0.08	83	0.07	82	0.38	90	0.09	84
MAM	0.79	86	0.11	92	0.20	79	0.61	86	0.10	86
JJA	1.27	86	0.08	86	0.46	86	0.81	86	0.06	86
SON	0.51	90	0.09	82	0.23	82	0.32	90	0.08	97
2009	0.79	98	0.10	98	0.23	77	0.58	96	0.09	96
DJF	0.56	100	0.12	100	0.06	69	0.47	92	0.11	92
MAM	0.81	92	0.11	85	0.11	85	0.74	92	0.10	92
JJA	1.1	100	0.09	100	0.40	100	0.71	100	0.07	100
SON	0.68	100	0.09	100	0.28	100	0.40	100	0.07	100
2010	0.90	94	0.11	94	0.24	79	0.67	96	0.10	96
DJF	0.95	92	0.14	92	0.09	40	0.86	92	0.16	92
MAM	0.82	85	0.09	85	0.18	82	0.61	100	0.08	100
JJA	1.02	100	0.09	100	0.34	100	0.68	100	0.07	100
SON	0.79	100	0.11	100	0.24	100	0.51	92	0.09	92
2011	0.92	98	0.11	92	0.26	94	0.68	98	0.11	96
DJF	0.60	100	0.11	100	0.09	92	0.51	92	0.11	100
MAM	0.99	93	0.10	70	0.27	93	0.73	100	0.10	85
JJA	1.07	100	0.07	100	0.37	100	0.69	100	0.07	100
SON	1.04	100	0.17	100	0.30	92	0.77	100	0.16	100
2012	0.56	89	0.08	86	0.10	79	0.50	90	0.08	90

	PM ₁₀				PM _{10-2.5}				PM _{2.5}			
	OC	Capture	EC	Capture	OC	Capture	OC	Capture	EC	Capture	EC	Capture
DJF	0.52	100	0.09	100	0.04	84	0.49	100	0.09	100		
MAM	0.58	85	0.09	85	0.03	70	0.63	70	0.11	70		
JJA	0.78	70	0.07	70	0.18	70	0.59	100	0.07	100		
SON	0.56	100	0.07	92	0.14	92	0.30	92	0.07	92		
2013	0.76	92	0.09	96	0.21	90	0.57	98	0.08	96		
DJF	0.49	92	0.10	92	0.05	68	0.47	91	0.09	91		
MAM	0.79	100	0.10	100	0.15	100	0.63	100	0.09	100		
JJA	1.16	100	0.07	92	0.37	100	0.79	100	0.07	92		
SON	0.58	100	0.07	100	0.22	92	0.39	100	0.07	100		
2014	0.91	98	0.09	100	0.29	94	0.65	96	0.08	96		
DJF	0.61	100	0.10	100	0.08	76	0.59	84	0.11	84		
MAM	0.91	100	0.10	100	0.23	100	0.69	100	0.09	100		
JJA	1.10	100	0.05	100	0.35	100	0.75	100	0.05	100		
SON	1.20	100	0.10	100	0.47	100	0.55	100	0.09	100		
2015	0.72	98	0.09	98	0.19	85	0.52	88	0.08	88		
DJF	0.44	100	0.06	100	0.11	100	0.34	100	0.06	100		
MAM	0.59	92	0.10	92	0.11	79	0.50	92	0.08	92		
JJA	1.01	100	0.09	100	0.35	63	0.66	63	0.08	63		
SON	0.83	100	0.10	100	0.22	99	0.62	99	0.10	99		
2016	0.73	100	0.06	100	0.21	95	0.54	100	0.06	100		
DJF	0.44	100	0.07	100	0.07	86	0.37	100	0.06	100		
MAM	0.83	100	0.07	100	0.21	92	0.64	100	0.07	100		
JJA	0.98	100	0.04	100	0.33	100	0.65	100	0.05	100		
SON	0.68	100	0.07	100	0.20	100	0.48	100	0.07	100		
2017	0.72	94	0.05	94	0.25	79	0.52	94	0.05	94		
DJF	0.57	100	0.07	100	0.11	63	0.53	100	0.07	100		
MAM	0.65	92	0.05	92	0.14	84	0.52	92	0.05	92		
JJA	0.92	86	0.03	86	0.34	86	0.58	86	0.04	86		
SON	0.77	100	0.06	100	0.38	83	0.47	100	0.05	100		
2018	0.96	100	0.08	100	0.26	90	0.73	100	0.07	100		
DJF	0.49	100	0.07	100	0.08	77	0.45	100	0.07	100		
MAM	1.32	100	0.11	100	0.28	92	1.06	100	0.10	100		
JJA	1.20	100	0.05	100	0.32	100	0.90	100	0.05	100		
SON	0.81	100	0.08	100	0.31	100	0.50	100	0.07	100		

Notation: Red numbers indicate annual or seasonal means based on < 50% data capture.

16

17

18

19 **Table S 6: R²-values for OC versus EC as a function of size fraction and season.**

	Winter	Spring	Summer	Fall
PM₁₀	0.66	0.58	0.51	0.64
PM_{2.5}	0.75	0.69	0.58	0.76

20

Table S 7: Annual mean ($\pm$ SD) relative chemical composition for the period 2001–2018. Unit (%)¹

	OM/PM ₁₀	EC/PM ₁₀	SO ₄ ²⁻ /PM ₁₀	NO ₃ /PM ₁₀	NH ₄ ⁺ /PM ₁₀	SS/PM ₁₀	OM/PM _{2.5}	EC/PM _{2.5}	OM/PM _{10.22}
2001	31±4	2.7±0.4	21	11	6	11	38±5	3.6±0.5	8.9±1.8 ¹⁾²³
2002	26±4	2.1±0.3	20	14	8	10	30±4	2.3±0.3	16±3
2003	29±4	1.6±0.2	23	12	6	11	32±5	2.5±0.4	18±4
2004	27±4	1.9±0.3	19	15	7	14	32±5	2.9±0.4	21±4
2005	25±4	2.4±0.3	20	16	8	13	29±4	3.3±0.5	21±4
2006	26±4	1.8±0.3	20	17	5	12	31±4	3.0±0.4	18±4
2007	27±4	2.9±0.4	15	10	4	14	35±5	4.3±0.6	16±3
2008	25±4	1.7±0.2	14	12	3	18	36±5	3.1±0.4	17±3
2009	26±4	3.7±1.9	15	13	4	13	31±4	2.8±0.4	19±4
2010	34±5	2.4±0.3	17	13	5	11	37±5	3.2±0.5	21±4
2011	25±4	1.7±0.2	14	17	6	16	32±4	3.0±0.4	15±3
2012	22±3	1.8±0.3	17	28	7	17	33±5	3.0±0.4	10±2
2013	29±4	2.0±0.3	15	19	6	23	37±5	3.0±0.4	18±4
2014	28±4	1.6±0.2	18	21	7	20	36±5	2.6±0.4	20±4
2015	26±4	1.9±0.3	16	22	6	28	36±5	3.3±0.5	20±3
2016	32±5	1.5±0.2	14	23	7	23	41±6	2.6±0.4	21±4
2017	38±5	1.5±0.2	19	14	5	27	49±7	2.8±0.4	28±6
2018	34±5	1.6±0.2	14	15	6	19	46±7	2.6±0.4	20±4

¹ 1) Data capture below 50%.

2) Notation: Conversion factors applied OM = OC x 1.9; EC = EC x 1.1

Table S 8: Mean ($\pm$ SD) relative chemical composition of the 3 weekly samples with the highest PM mass concentration (PM_{MAX}) per year for the period 2001–2018.²

	$PM_{10\ MAX}$		OM		EC	SO_4^{2-}		NO_3^-	NH_4^+	SS	$PM_{2.5\ MAX}$		OM		EC
	$\mu g\ m^{-3}$	Season ¹⁾	%	%	%	%	%	%	%	%	$\mu g\ m^{-3}$	Season ¹⁾	%	%	%
2001	12.4 $\pm$ 1.6	234	43 $\pm$ 16	3 $\pm$ 1	14 $\pm$ 10	4 $\pm$ 3	4 $\pm$ 3	3 $\pm$ 3	12.2 $\pm$ 2.4	234	50 $\pm$ 10	5 $\pm$ 2			
2002	23.8 $\pm$ 3.6	223	26 $\pm$ 11	1.8 $\pm$ 0.7	14 $\pm$ 9	13 $\pm$ 14	8 $\pm$ 4	2 $\pm$ 2	18.8 $\pm$ 2.9	223	29 $\pm$ 12	1.7 $\pm$ 0.5			
2003	17.9 $\pm$ 2.7	222	16 $\pm$ 3	1.3 $\pm$ 0.4	17 $\pm$ 3	18 $\pm$ 8	11 $\pm$ 2	8 $\pm$ 3	12.0 $\pm$ 1.9	222	19 $\pm$ 3	1.6 $\pm$ 0.3			
2004	17.0 $\pm$ 6.8	222	23 $\pm$ 9	1.5 $\pm$ 0.4	21 $\pm$ 3	11 $\pm$ 10	8 $\pm$ 4	14 $\pm$ 18	11.6 $\pm$ 6.5	222	27 $\pm$ 9	2.4 $\pm$ 0.8			
2005	16.9 $\pm$ 2.8	244	26 $\pm$ 6	3.4 $\pm$ 1.6	15 $\pm$ 3	20 $\pm$ 3	2 $\pm$ 0	5 $\pm$ 3	12.4 $\pm$ 3.1	224	27 $\pm$ 10	2.9 $\pm$ 0.4			
2006	26.5 $\pm$ 3.0	144	23 $\pm$ 8	4.1 $\pm$ 1.6	19 $\pm$ 7	17 $\pm$ 4	8 $\pm$ 2	7 $\pm$ 5	15.5 $\pm$ 0.9	124	30 $\pm$ 10	2.3 $\pm$ 0.7			
2007	13.8 $\pm$ 3.5	223	40 $\pm$ 2	4.2 $\pm$ 1.5	12 $\pm$ 0	6 $\pm$ 2	4 $\pm$ 1	1 $\pm$ 0	10.7 $\pm$ 3.4	223	48 $\pm$ 2	4.1 $\pm$ 1.5			
2008	12.5 $\pm$ 3.0	222	12 $\pm$ 5	1.5 $\pm$ 0.5	12 $\pm$ 3	20 $\pm$ 7	5 $\pm$ 3	15 $\pm$ 15	7.0 $\pm$ 2.7	223	16 $\pm$ 3	1.9 $\pm$ 0.6			
2009	14.8 $\pm$ 5.7	222	21 $\pm$ 1	1.8 $\pm$ 0.0	12 $\pm$ 5	13 $\pm$ 9	5 $\pm$ 3	3 $\pm$ 2	10.6 $\pm$ 4.1	122	28 $\pm$ 3	2.5 $\pm$ 0.5			
2010	12.0 $\pm$ 2.4	344	21 $\pm$ 6	1.9 $\pm$ 0.6	14 $\pm$ 3	13 $\pm$ 2	3 $\pm$ 1	23 $\pm$ 11	10.7 $\pm$ 3.7	144	42 $\pm$ 9	5.0 $\pm$ 0.6			
2011	16.7 $\pm$ 1.2	224	23 $\pm$ 13	2.0 $\pm$ 1.3	23 $\pm$ 2	17 $\pm$ 4	12 $\pm$ 0	2 $\pm$ 1	12.1 $\pm$ 1.5	224	27 $\pm$ 13	3.0 $\pm$ 1.9			
2012	11.9 $\pm$ 1.7	122	29 $\pm$ 21	3.9 $\pm$ 3.4	10 $\pm$ 9	24 $\pm$ 25	3 $\pm$ 2	1 $\pm$ 0	7.2 $\pm$ 1.4	123	30 $\pm$ 26	3.0 $\pm$ 2.7			
2013	9.4 $\pm$ 0.6	222	12 $\pm$ 4	1.4 $\pm$ 0.5	9 $\pm$ 2	7 $\pm$ 2	8 $\pm$ 0	13 $\pm$ 5	5.7 $\pm$ 0.6	223	30 $\pm$ 9	2.6 $\pm$ 1.0			
2014	17.7 $\pm$ 3.3	224	18 $\pm$ 8	1.5 $\pm$ 0.2	15 $\pm$ 5	25 $\pm$ 6	10 $\pm$ 1	14 $\pm$ 8	8.1 $\pm$ 0.9	122	21 $\pm$ 6	2.5 $\pm$ 0.5			
2015	12.0 $\pm$ 4.0	122	21 $\pm$ 10	2.2 $\pm$ 1.3	12 $\pm$ 7	23 $\pm$ 10	11 $\pm$ 2	12 $\pm$ 12	6.9 $\pm$ 1.7	122	31 $\pm$ 20	3.3 $\pm$ 2.0			
2016	9.2 $\pm$ 0.2	223	30 $\pm$ 22	1.3 $\pm$ 0.3	6 $\pm$ 5	18 $\pm$ 17	7 $\pm$ 7	6 $\pm$ 1	6.7 $\pm$ 1.5	223	37 $\pm$ 25	1.9 $\pm$ 0.3			
2017	9.8 $\pm$ 2.7	114	33 $\pm$ 8	2.3 $\pm$ 1.1	19 $\pm$ 3	14 $\pm$ 7	7 $\pm$ 1	13 $\pm$ 13	6.3 $\pm$ 2.0	112	29 $\pm$ 8	2.7 $\pm$ 0.3			
2018	12.8 $\pm$ 3.1	124	32 $\pm$ 24	1.3 $\pm$ 0.5	11 $\pm$ 3	20 $\pm$ 15	8 $\pm$ 5	5 $\pm$ 1	9.0 $\pm$ 1.0	144	21 $\pm$ 11	1.7 $\pm$ 1.2			

² 1 = DJF; 2 = MAM; 3 = JJA; 4 = SON

Table S 9: Annual and seasonal mean concentrations of TC in PM₁₀, PM_{2.5} and PM_{10-2.5} at Birkenes for 2001–2018 (Unit: $\mu\text{g C m}^{-3}$).

	PM ₁₀		PM _{10-2.5}		PM _{2.5}	
	TC	Capture	TC	Capture	TC	Capture
2001	1.09	63	0.07	48	1.08	63
DJF	0.72	59	0	34	0.75	59
MAM	1.14	92	0.02	65	1.19	92
JJA	2.50	26	0.20	26	2.30	26
SON	0.68	48	0.08	48	0.60	48
2002	1.15	85	0.21	80	1.01	91
DJF	0.65	95	0.05	88	0.60	95
MAM	1.47	92	0.14	92	1.33	92
JJA	1.96	68	0.53	68	1.50	91
SON	0.68	86	0.18	73	0.59	86
2003	1.12	82	0.23	78	0.93	82
DJF	0.94	85	0.05	82	0.95	85
MAM	1.26	86	0.20	76	1.16	86
JJA	1.34	85	0.43	85	0.91	85
SON	0.84	70	0.22	70	0.62	70
2004	0.91	84	0.28	79	0.65	84
DJF	0.65	86	0.09	86	0.56	86
MAM	1.13	86	0.32	66	0.91	86
JJA	1.09	79	0.41	79	0.68	79
SON	0.79	86	0.33	86	0.47	86
2005	0.99	79	0.32	75	0.71	80
DJF	0.56	64	0.06	51	0.64	71
MAM	0.91	85	0.16	85	0.75	85
JJA	1.01	86	0.34	86	0.67	86
SON	1.41	79	0.62	79	0.79	79
2006	1.20	78	0.35	70	0.90	78
DJF	0.91	86	0.08	66	0.88	86
MAM	1.03	85	0.10	72	0.96	85
JJA	1.55	57	0.49	57	1.06	57
SON	1.43	86	0.68	86	0.75	86
2007	0.99	77	0.24	76	0.74	77
DJF	0.76	58	0.09	58	0.67	58
MAM	1.17	86	0.20	86	0.97	86
JJA	1.16	86	0.41	79	0.77	79
SON	0.76	79	0.21	79	0.52	86
2008	0.85	84	0.25	81	0.60	86
DJF	0.47	84	0.05	76	0.43	84
MAM	0.90	86	0.21	79	0.71	86
JJA	1.35	86	0.47	86	0.87	86
SON	0.65	82	0.24	82	0.39	90
2009	0.89	98	0.23	92	0.67	96
DJF	0.68	100	0.05	84	0.58	92
MAM	0.92	92	0.12	85	0.85	92
JJA	1.19	100	0.41	100	0.78	100
SON	0.76	100	0.29	100	0.47	100
2010	1.00	94	0.21	87	0.77	96
DJF	1.09	92	0.06	69	1.03	92
MAM	0.91	85	0.17	85	0.69	100
JJA	1.11	100	0.36	100	0.75	100
SON	0.90	100	0.26	92	0.61	92
2011	0.99	98	0.25	92	0.80	100
DJF	0.71	100	0.09	92	0.64	100
MAM	0.88	92	0.20	85	0.84	100
JJA	1.13	100	0.37	100	0.76	100
SON	1.21	100	0.32	92	0.93	100
2012	0.64	89	0.10	79	0.58	92

	PM ₁₀		PM _{10-2.5}		PM _{2.5}	
	TC	Capture	TC	Capture	TC	Capture
DJF	0.61	100	0.03	92	0.58	100
MAM	0.67	85	0	68	0.73	77
JJA	0.85	70	0.25	56	0.66	100
SON	0.51	100	0.14	92	0.37	92
2013	0.84	98	0.21	92	0.65	98
DJF	0.59	92	0.04	76	0.56	91
MAM	0.89	100	0.17	100	0.72	100
JJA	1.22	100	0.37	100	0.86	100
SON	0.66	100	0.23	92	0.46	100
2014	1.00	100	0.30	94	0.73	96
DJF	0.71	100	0.07	76	0.70	84
MAM	1.02	100	0.24	100	0.78	100
JJA	1.16	100	0.35	100	0.80	100
SON	1.12	100	0.48	100	0.64	100
2015	0.81	98	0.19	88	0.60	88
DJF	0.50	100	0.11	100	0.39	100
MAM	0.70	92	0.10	92	0.60	92
JJA	1.10	100	0.36	63	0.74	63
SON	0.94	100	0.23	96	0.71	96
2016	0.80	100	0.21	94	0.6	100
DJF	0.51	100	0.08	82	0.43	100
MAM	0.90	100	0.22	92	0.71	100
JJA	1.02	100	0.32	100	0.70	100
SON	0.76	100	0.21	100	0.55	100
2017	0.78	94	0.26	78	0.58	94
DJF	0.63	100	0.11	59	0.60	100
MAM	0.70	92	0.14	84	0.58	92
JJA	0.95	86	0.34	86	0.61	86
SON	0.84	100	0.40	83	0.51	100
2018	1.03	100	0.26	90	0.8	100
DJF	0.56	100	0.08	77	0.52	100
MAM	1.43	100	0.30	92	1.06	100
JJA	1.25	100	0.32	92	0.95	100
SON	0.88	100	0.32	100	0.56	100

Notation: Red numbers indicate annual or seasonal means based on < 50% data capture.

30 **Table S 10: Annual and seasonal mean mass concentrations of PM₁₀, PM_{2.5} and PM_{10-2.5} at Birkenes for 2001–2018 (Unit:**
31 **$\mu\text{g m}^{-3}$).**

	PM ₁₀	Capture	PM _{2.5}	Capture	PM _{10-2.5}	Capture
2001	5.8	56	4.6	58	1.7	54
DJF	4.5	59	3.2	59	1.3	59
MAM	6.6	92	5.5	100	1.8	92
JJA	7.9	26	7.3	26	1.3	19
SON	4.9	48	2.9	48	1.9	48
2002	7.5	83	5.7	91	2.2	80
DJF	5.6	88	3.8	95	1.8	88
MAM	11.0	92	8.5	92	2.4	92
JJA	10.0	68	7.3	95	2.9	68
SON	3.9	86	2.8	86	1.5	73
2003	6.7	78	4.8	82	2.4	74
DJF	5.9	76	4.4	85	2.2	76
MAM	9.3	82	6.7	86	3.4	76
JJA	5.9	85	4.4	81	1.7	79
SON	5.4	70	3.5	70	2.3	64
2004	5.7	84	3.4	84	2.4	83
DJF	4.5	86	2.8	86	1.7	86
MAM	8.2	86	5.4	85	3.1	79
JJA	5.7	79	3.3	79	2.4	79
SON	4.3	86	2.0	86	2.3	86
2005	6.5	79	4.0	80	2.6	75
DJF	5.2	64	3.1	71	2.8	51
MAM	6.9	85	4.9	85	1.9	85
JJA	5.5	86	3.4	86	2.1	86
SON	8.0	79	4.4	79	3.6	79
2006	7.8	78	4.7	77	3.4	73
DJF	7.4	86	4.5	79	3.0	79
MAM	6.2	85	4.6	86	2.1	72
JJA	8.3	57	5.4	57	3.0	57
SON	9.5	86	4.5	86	5.0	86
2007	5.8	72	3.3	74	2.5	72
DJF	4.2	38	2.0	38	2.2	38
MAM	7.5	86	4.4	86	3.1	86
JJA	6.1	79	3.5	86	2.5	79
SON	4.4	86	2.4	86	2.0	86
2008	5.7	77	2.8	86	2.7	75
DJF	5.1	84	2.5	90	2.8	84
MAM	7.1	79	4.1	73	2.7	73
JJA	6.1	86	3.1	86	2.9	86
SON	4.0	58	1.9	97	2.1	58
2009	5.8	85	3.6	94	2.3	76
DJF	4.4	96	3	92	1.4	88
MAM	9.4	74	5.3	100	3.4	74
JJA	6.1	71	3.8	92	2.2	63
SON	4.3	99	2.2	92	2.3	83

	PM ₁₀	Capture	PM _{2.5}	Capture	PM _{10-2.5}	Capture
2010	5.1	88	3.4	94	2.2	77
DJF	3.8	100	3.4	87	0.55	64
MAM	5.4	71	3	98	2.3	68
JJA	6.7	90	3.7	100	3.1	91
SON	4.8	92	3.5	92	2.5	85
2011	7.0	98	4.1	100	3.2	96
DJF	5.6	100	3.1	100	2.5	100
MAM	8.2	93	5.5	100	3.9	85
JJA	5.2	100	3.5	100	1.7	100
SON	9.0	100	4.4	100	4.6	100
2012	4.9	89	2.9	92	2.0	85
DJF	4.4	100	2.7	100	1.7	99
MAM	6.7	85	3.2	100	3.3	70
JJA	4.6	70	3.8	70	1.4	70
SON	4.0	100	2.0	100	2.0	100
2013	4.9	92	2.9	86	2.2	84
DJF	3.8	69	2.6	44	1.7	44
MAM	6.4	100	3.7	100	2.8	100
JJA	5.4	100	3.4	100	2.0	100
SON	3.6	100	1.8	100	2.0	92
2014	6.1	98	3.4	96	2.7	94
DJF	5.7	100	3.5	84	2.4	84
MAM	7.2	92	3.7	100	3.3	92
JJA	5.2	100	3.2	100	2.0	100
SON	6.3	100	3.3	100	3.0	100
2015	5.3	98	2.7	88	2.5	88
DJF	5.5	100	2.5	100	31	100
MAM	5.2	100	2.8	100	2.5	100
JJA	5.5	92	2.9	63	1.9	63
SON	5.0	100	2.8	99	2.4	99
2016	4.3	100	2.5	100	1.9	100
DJF	4.1	100	2.1	100	2.0	100
MAM	4.8	100	3.1	100	1.7	100
JJA	4.3	100	2.6	100	1.7	100
SON	4.1	100	2.1	100	2.1	100
2017	3.8	90	2.0	94	1.7	90
DJF	3.7	100	2.2	100	1.4	100
MAM	3.8	92	2.3	92	1.5	92
JJA	3.9	86	2.1	86	1.8	86
SON	3.8	85	1.4	100	2.4	85
2018	5.4	100	3.0	98	2.5	98
DJF	3.8	100	2.4	92	1.4	92
MAM	6.5	100	4.2	100	2.3	100
JJA	5.4	100	3.1	100	2.3	100
SON	6.2	100	2.2	100	3.9	100

Red numbers indicate annual or seasonal means based on < 50% data capture.

Table S 11: Sen slope of annual means and corresponding confidence intervals for significant slopes ($p=0.05$), as well as change in annual mean presented as percentage change per year and as percentage change for the period 2001–2018. Non-significant values in red.

	Slope (% yr ⁻¹)	CI-1	CI-2	Change 2001-2018 (%)
PM₁₀	-2.2	-3.7	-0.7	-38
PM_{2.5}	-4.0	-5.7	-2.2	-69
PM_{10-2.5}	-0.1	-2.2	1.4	-2.4
OC in PM₁₀	0	-1.4	0.9	0
OC in PM_{2.5}	-0.8	-2.8	0.7	-13
OC in PM_{10-2.5}	0.8	-1.7	3.4	13
EC in PM₁₀	-3.9	-5.8	-1.9	-66
EC in PM_{2.5}	-4.2	-6.2	-2.6	-71
TC in PM₁₀	-1.1	-2.0	0.0	-19
TC in PM_{2.5}	-1.5	-3.5	0.0	-26
TC in PM_{10-2.5}	0.0	-2.4	1.9	0
SO₄²⁻	-3.8	-6.1	-1.8	-65
NO₃⁻	0.8	-2.5	4.3	14
NH₄⁺	-2.7	-5.8	0.5	-47
SS	2.2	0.6	4.5	38
Levoglucosan¹⁾	-2.8	-8.8	-0.2	-28

Notation: Trends for levoglucosan are calculated for the period 2008–2018.

Table S 12: Sen slope of seasonal means in % yr⁻¹ and corresponding confidence intervals for significant slopes ($p=0.05$) for 2001–2018. Non-significant values in red.

	DJF		MAM		JJA		SON			
	Slope (% yr ⁻¹)	CI-1	CI-2	Slope (% yr ⁻¹)	CI-1	CI-2	Slope (% yr ⁻¹)	CI-1	CI-2	
PM ₁₀	-1.6	-3.3	-0.1	-3.3	-5.6	-0.9	-2.4	-5.1	-3.3	2.2
PM _{2.5}	-2.4	-4.6	-0.8	-4.4	-7.0	-3.0	-3.0	-7.1	-6.5	0.6
PM _{10-2.5}	0.0	-2.6	3.0	-0.4	-4.2	2.3	-2.3	-4.5	-1.2	4.3
OC in PM ₁₀	-0.2	-2.8	1.4	-1.0	-3.1	1.2	-0.7	-3.2	-1.3	4.0
OC in PM _{2.5}	-0.8	-3.7	1.4	-1.9	-3.7	0.5	-0.6	-2.8	-1.2	4.3
OC in PM _{10-2.5}	3.2	0.0	6.7	3.7	-1.7	7.2	-1.4	-2.5	-2.4	5.4
EC in PM ₁₀	-2.8	-4.9	-0.7	-4.0	-6.0	-1.6	-5.9	-9.8	-9.2	0.0
EC in PM _{2.5}	-3.1	-5.4	-0.7	-4.6	-6.7	-3.0	-4.1	-6.7	-5.5	0.0
TC in PM ₁₀	-1.0	-3.5	0.7	-2.0	-3.4	-0.1	-1.2	-3.6	-2.8	3.5
TC in PM _{2.5}	-1.4	-3.5	0.7	-2.6	-4.2	-0.9	-1.0	-3.6	-2.5	2.7
TC in PM _{10-2.5}	3.4	-0.3	6.7	3.1	-2.6	7.9	-1.7	-3.2	-3.3	4.9
SO ₄ ²⁻	-3.0	-6.1	-0.2	-6.4	-9.0	-3.8	-4.2	-5.9	-5.3	0.7
SS	2.9	-1.4	7.4	1.0	-1.6	3.9	3.7	2.3	0.0	4.8
Levogluconan ¹⁾	-3.3	-15.9	6.7	1.9	-6.7	5.3	-5.7	-18.1	-12.3	7.2

Notation: Trends for levoglucosan are calculated for the period 2008–2018

Table S 13: Sen slope of annual mean ratios and corresponding confidence intervals for significant slopes ($p=0.05$), as well as change in annual mean presented as percentage change per year and as percentage change for the period 2001–2018 (2008–2018 for levoglucosan). Non-significant values in red.

	Slope (% yr ⁻¹)	CI-1	CI-2	Change 2001-2018 (%)
OC_{PM10} to PM₁₀	2.4	0.7	3.4	41
OC_{PM2.5} to PM_{2.5}	3.2	1.7	4.4	55
OC_{PM10-2.5} to PM_{10-2.5}	1.1	-1.3	2.5	17
EC_{PM10} to PM₁₀	-4.5	-7.1	-2.8	-77
EC_{PM2.5} to PM_{2.5}	-3.9	-5.8	-1.9	-66
TC_{PM10} to PM₁₀	1.8	0.3	2.6	30
TC_{PM2.5} to PM_{2.5}	2.6	1.4	3.7	44
TC_{PM10-2.5} to PM_{10-2.5}	0.4	-1.8	1.8	7.1
SO₄²⁻ to PM₁₀	-2.1	-3.4	-0.4	-35
NO₃⁻ to PM₁₀	3.8	0.8	6.3	64
NH₄⁺ to PM₁₀	-0.7	-3.1	2.4	-12
SS to PM₁₀	4.4	3.0	6.7	75
Levoglucosan to OC_{PM10}	-1.8	-10.6	1.8	-18
Levoglucosan to OC_{PM2.5}	-3.6	-9.8	1.3	-36
Levoglucosan to EC_{PM10}	2.8	-3.5	6.5	28
Levoglucosan to EC_{PM2.5}	2.3	-2.2	5.0	24
Levoglucosan to TC_{PM10}	-1.1	-9.0	2.7	-11
Levoglucosan to TC_{PM2.5}	-3.1	-8.1	2.0	-31

Notation: Trends for levoglucosan are calculated for the period 2008–2018.

Table S 14: Sen slope of seasonal mean ratios and corresponding confidence intervals for significant slopes ($p=0.05$), as well as change in annual mean presented as percentage change per year and as percentage change for the period 2001 – 2018 (2008 – 2018 for levoglucosan). Non-significant values in red.

	DJF		MAM		JJA		SON	
	Slope (% yr ⁻¹)		Slope (% yr ⁻¹)		Slope (% yr ⁻¹)		Slope (% yr ⁻¹)	
	CI-1	CI-2	CI-1	CI-2	CI-1	CI-2	CI-1	CI-2
OC _{PM10} to PM ₁₀	1.8	-0.6	1.7	3.7	2.3	3.6	1.7	3.7
OC _{PM2.5} to PM _{2.5}	1.8	-0.4	3.1	5.8	3.9	5.1	3.1	5.8
OC _{PM10-2.5} to PM _{10-2.5}	2.8	-0.9	-0.3	2.6	1.6	3.1	-0.3	2.6
EC _{PM10} to PM ₁₀	-4.6	-7.2	-4.7	-2.7	-7.1	-3.4	-4.7	-2.5
EC _{PM2.5} to PM _{2.5}	-3.9	-5.4	-3.6	-2.5	-4.3	-1.7	-3.6	-1.2
TC _{PM10} to PM ₁₀	0.9	-1.2	1.1	3.2	1.7	2.8	1.1	2.7
TC _{PM2.5} to PM _{2.5}	1.2	-0.6	-0.7	3.3	0.7	1.9	-0.7	1.6
TC _{PM10-2.5} to PM _{10-2.5}	2.2	-1.6	2.4	4.4	3.2	4.1	2.4	4.6
SO ₄ ²⁻ to PM ₁₀	-2.2	-4.0	-1.3	0.1	-1.3	0.0	-1.3	-0.4
NO ₃ ⁻ to PM ₁₀	7.3	3.2	1.2	11.1	4.4	7.1	1.2	4.1
NH ₄ ⁺ to PM ₁₀	2.6	-1.7	-1.4	5.4	-0.6	3.6	-1.4	3.0
SS to PM ₁₀	4.1	-0.3	4.8	7.4	6.2	8.0	4.8	7.6
Levoglucosan to OC _{PM10}	-5.9	-7.9	-4.9	2.5	0.0	0.0	-4.9	0.0
Levoglucosan to OC _{PM2.5}	-2.7	-6.5	-4.2	1.8	0.0	3.7	-4.2	0.0
Levoglucosan to EC _{PM10}	3.1	-6.9	1.1	8.6	8.2	13.6	1.1	5.1
Levoglucosan to EC _{PM2.5}	2.5	-3.7	-0.4	6.4	3.5	10.7	-0.4	5.6
Levoglucosan to TC _{PM10}	-3.5	-7.4	-5.6	3.5	0.0	0.0	-5.6	0.0
Levoglucosan to TC _{PM2.5}	-3.2	-6.4	-2.1	2.1	0.0	7.4	-2.1	1.8

Notation: Trends for levoglucosan are calculated for the period 2008–2018

	Levo- glucosan	Cap	Man- nosan	Cap	Galact- osan	Cap	Levo/ Mann	Cap	Ara- bitol	Cap	Man- nitol	Cap	2-methyl- erythritol	Cap	2-methyl- threitol	Cap	Tre- halose	Cap	Glu- cose
2008	10.90	90	2.68	90	0.91	90	4.7	75											
DJF	10.24	90	2.32	90	0.69	90	4.7	69											
MAM	8.00	86	1.37	86	0.40	86	5.5	82											
JJA	13.40	86	4.48	86	2.00	90	3.4	58											
SON	11.34	97	2.63	100	0.53	97	4.5	89											
2009	10.62	98	1.73	98	0.57	98	5.9	90											
DJF	20.53	92	3.40	92	1.05	92	5.6	83											
MAM	11.84	100	1.84	100	0.69	100	6.6	100											
JJA	1.91	100	0.35	100	0.09	100	5.4	94											
SON	9.08	100	1.49	100	0.50	100	5.4	85											
2010	17.33	100	3.89	100	1.11	100	5.2	85											
DJF	43.04	100	9.39	100	2.85	100	4.9	92											
MAM	10.22	100	2.69	100	0.79	100	4.6	71											
JJA	2.96	100	1.06	100	0.11	100	5.2	84											
SON	13.71	100	2.52	100	0.73	100	5.7	92											
2011	11.64	100	2.33	100	0.83	100	4.7	90											
DJF	14.74	100	3.25	100	1.13	100	4.4	84											
MAM	11.79	100	2.16	100	0.94	100	5.3	92											
JJA	4.00	100	0.81	100	0.16	100	4.8	100											
SON	16.23	100	3.11	100	1.12	100	4.5	85											
2012	10.54	100	2.34	100	0.73	100	4.4	90											
DJF	21.75	100	4.95	100	1.43	100	4.8	100											
MAM	10.67	100	1.96	100	0.61	100	5.5	100											
JJA	1.75	100	0.65	100	0.20	100	3.2	70											
SON	8.07	100	1.82	100	0.69	100	3.8	92											
2013	9.62	100	1.57	100	0.50	100	5.6	92											
DJF	18.21	100	2.88	100	0.85	100	5.3	92											
MAM	10.88	100	1.81	100	0.62	100	6.7	100											
JJA	2.74	100	0.49	100	0.15	100	5.8	92											
SON	6.66	100	1.13	100	0.36	100	5.5	85											
2014	12.78	100	2.99	100	0.60	100	5	96											
DJF	20.14	100	4.72	100	0.92	100	4.2	92											
MAM	12.44	100	2.42	100	0.62	100	5.8	92											
JJA	3.15	100	0.56	100	0.14	100	5.8	100											
SON	16.10	100	4.47	100	0.89	100	4.1	100											
2015	9.36	88	1.36	98	0.43	98	6.4	80											
DJF	9.84	100	1.53	98	0.45	98	6.9	80											
MAM	12.31	92	1.45	92	0.63	92	6.6	61											
JJA	2.26	100	0.44	100	0.07	100	5.6	84											
SON	14.04	100	2.16	100	0.61	100	6.1	97											

	Levo- glucosan	Cap	Man- nosan	Cap	Galact- osan	Cap	Levo/ Mann	Cap	Ara- bitol	Cap	Man- nitol	Cap	2-methyl- erythritol	Cap	2-methyl- threitol	Cap	Tre- halose	Cap	Glu- cose	Cap
2016	7.52	97	1.21	97	0.30	94	5.7	92	4.78	99	3.89	99	0.38	99	0.13	99	3.17	99	5.07	99
DIF	12.54	94	1.87	94	0.47	94	6.4	93	1.70	94	1.09	94	0.02	94	0.01	94	0.76	94	1.47	94
MAM	7.62	100	1.25	100	0.31	100	5.9	100	4.67	100	3.40	100	0.21	100	0.11	100	5.82	100	4.60	100
JJA	2.37	100	0.47	100	0.09	77	5.3	100	8.29	100	6.69	100	1.01	100	0.29	100	3.73	100	9.93	100
SON	7.92	92	1.30	92	0.28	100	5.5	76	4.26	100	4.17	100	0.24	100	0.09	100	2.19	100	4.04	100
2017	8.24	94	1.35	95	0.32	94	5.6	92	5.52	94	5.70	94	0.37	94	0.10	94	3.37	94	5.17	94
DIF	15.77	100	2.49	100	0.65	100	5.6	100	1.16	100	1.40	100	0.01	100	0.01	100	1.03	100	2.03	100
MAM	6.22	92	0.98	92	0.23	92	6.6	84	3.90	92	3.65	92	0.05	92	0.02	92	1.58	92	4.11	92
JJA	2.27	86	0.48	86	0.06	86	4.8	86	9.15	86	8.45	86	1.32	86	0.35	86	4.21	86	7.09	86
SON	7.84	100	1.31	100	0.29	100	5.5	100	8.18	100	9.49	100	0.17	100	0.06	100	6.61	100	7.58	100
2018	9.77	100	1.62	100	0.39	100	6.1	98	5.76	100	5.65	100	0.45	98	0.16	98	2.83	100	4.16	100
DIF	13.50	100	2.21	100	0.60	100	5.9	100	0.72	100	0.94	100	0.01	92	0.01	92	0.60	100	2.74	100
MAM	13.64	100	2.04	100	0.54	100	6.8	100	4.18	100	4.21	100	0.24	100	0.08	100	1.91	100	3.59	100
JJA	1.38	100	0.25	100	0.04	100	5.8	100	8.66	100	7.94	100	1.24	100	0.42	100	2.90	100	4.52	100
SON	10.64	100	2.01	100	0.41	100	5.5	92	9.38	100	9.47	100	0.25	100	0.12	100	5.91	100	5.79	100

54 **Table S 16: Seasonal mean ($\pm$ SD) concentrations of TC_{bb}, OC_{bb} and EC_{bb} in PM₁₀ and PM_{2.5} at Birkenes 2008–2018.**
55 (Unit: $\mu\text{g C m}^{-3}$).

	TC _{bb} PM ₁₀	OC _{bb} PM ₁₀	EC _{bb} PM ₁₀	TC _{bb} PM _{2.5}	OC _{bb} PM _{2.5}	EC _{bb} PM _{2.5}
	TC _{bb} PM ₁₀	OC _{bb} PM ₁₀	EC _{bb} PM ₁₀	TC _{bb} PM _{2.5}	OC _{bb} PM _{2.5}	EC _{bb} PM _{2.5}
2008	0.160$\pm$0.042	0.138$\pm$0.042	0.021$\pm$0.005	0.141 $\pm$ 0.034	0.121 $\pm$ 0.034	0.020 $\pm$ 0.004
DJF	0.150 $\pm$ 0.039	0.130 $\pm$ 0.039	0.020 $\pm$ 0.004	0.133 $\pm$ 0.032	0.114 $\pm$ 0.032	0.019 $\pm$ 0.004
MAM	0.117 $\pm$ 0.031	0.102 $\pm$ 0.030	0.016 $\pm$ 0.003	0.104 $\pm$ 0.025	0.089 $\pm$ 0.025	0.015 $\pm$ 0.003
JJA	0.196 $\pm$ 0.05	0.170 $\pm$ 0.051	0.026 $\pm$ 0.006	0.174 $\pm$ 0.042	0.149 $\pm$ 0.042	0.025 $\pm$ 0.006
SON	0.166 $\pm$ 0.043	0.144 $\pm$ 0.043	0.022 $\pm$ 0.005	0.147 $\pm$ 0.036	0.126 $\pm$ 0.036	0.021 $\pm$ 0.005
2009	0.156$\pm$0.041	0.135$\pm$0.040	0.021$\pm$0.005	0.138$\pm$0.034	0.118$\pm$0.034	0.020$\pm$0.004
DJF	0.301 $\pm$ 0.078	0.261 $\pm$ 0.078	0.040 $\pm$ 0.009	0.266 $\pm$ 0.065	0.228 $\pm$ 0.065	0.038 $\pm$ 0.008
MAM	0.174 $\pm$ 0.045	0.150 $\pm$ 0.045	0.023 $\pm$ 0.005	0.154 $\pm$ 0.037	0.131 $\pm$ 0.037	0.022 $\pm$ 0.005
JJA	0.028 $\pm$ 0.007	0.024 $\pm$ 0.007	0.004 $\pm$ 0.001	0.025 $\pm$ 0.006	0.021 $\pm$ 0.006	0.004 $\pm$ 0.001
SON	0.133 $\pm$ 0.035	0.115 $\pm$ 0.035	0.018 $\pm$ 0.004	0.118 $\pm$ 0.029	0.101 $\pm$ 0.029	0.017 $\pm$ 0.004
2010	0.254$\pm$0.066	0.220$\pm$0.066	0.034$\pm$0.008	0.225$\pm$0.055	0.192$\pm$0.055	0.032$\pm$0.007
DJF	0.631 $\pm$ 0.164	0.547 $\pm$ 0.164	0.084 $\pm$ 0.019	0.558 $\pm$ 0.136	0.478 $\pm$ 0.136	0.080 $\pm$ 0.018
MAM	0.150 $\pm$ 0.039	0.130 $\pm$ 0.039	0.020 $\pm$ 0.004	0.132 $\pm$ 0.032	0.113 $\pm$ 0.032	0.019 $\pm$ 0.004
JJA	0.043 $\pm$ 0.011	0.038 $\pm$ 0.011	0.006 $\pm$ 0.001	0.038 $\pm$ 0.009	0.033 $\pm$ 0.009	0.006 $\pm$ 0.001
SON	0.201 $\pm$ 0.052	0.174 $\pm$ 0.052	0.027 $\pm$ 0.006	0.178 $\pm$ 0.043	0.152 $\pm$ 0.043	0.026 $\pm$ 0.006
2011	0.171$\pm$0.044	0.148$\pm$0.044	0.023$\pm$0.005	0.151$\pm$0.037	0.129$\pm$0.037	0.022$\pm$0.005
DJF	0.216 $\pm$ 0.056	0.187 $\pm$ 0.056	0.029 $\pm$ 0.006	0.191 $\pm$ 0.047	0.164 $\pm$ 0.047	0.027 $\pm$ 0.006
MAM	0.173 $\pm$ 0.045	0.150 $\pm$ 0.045	0.023 $\pm$ 0.005	0.153 $\pm$ 0.037	0.131 $\pm$ 0.037	0.022 $\pm$ 0.005
JJA	0.059 $\pm$ 0.015	0.051 $\pm$ 0.015	0.008 $\pm$ 0.002	0.052 $\pm$ 0.013	0.044 $\pm$ 0.013	0.007 $\pm$ 0.002
SON	0.238 $\pm$ 0.062	0.206 $\pm$ 0.062	0.032 $\pm$ 0.007	0.210 $\pm$ 0.051	0.180 $\pm$ 0.051	0.030 $\pm$ 0.007
2012	0.155$\pm$0.040	0.134$\pm$0.040	0.021$\pm$0.005	0.137$\pm$0.033	0.117$\pm$0.033	0.020$\pm$0.004
DJF	0.319 $\pm$ 0.083	0.276 $\pm$ 0.083	0.043 $\pm$ 0.009	0.282 $\pm$ 0.069	0.241 $\pm$ 0.069	0.040 $\pm$ 0.009
MAM	0.156 $\pm$ 0.041	0.136 $\pm$ 0.041	0.021 $\pm$ 0.005	0.138 $\pm$ 0.034	0.118 $\pm$ 0.034	0.020 $\pm$ 0.004
JJA	0.026 $\pm$ 0.007	0.022 $\pm$ 0.007	0.003 $\pm$ 0.001	0.023 $\pm$ 0.006	0.019 $\pm$ 0.006	0.003 $\pm$ 0.001
SON	0.118 $\pm$ 0.031	0.102 $\pm$ 0.031	0.016 $\pm$ 0.004	0.105 $\pm$ 0.026	0.090 $\pm$ 0.025	0.015 $\pm$ 0.003
2013	0.141$\pm$0.037	0.122$\pm$0.037	0.019$\pm$0.004	0.125$\pm$0.030	0.107$\pm$0.030	0.018$\pm$0.004
DJF	0.267 $\pm$ 0.069	0.231 $\pm$ 0.069	0.036 $\pm$ 0.008	0.236 $\pm$ 0.058	0.202 $\pm$ 0.058	0.034 $\pm$ 0.008
MAM	0.159 $\pm$ 0.041	0.138 $\pm$ 0.041	0.021 $\pm$ 0.005	0.141 $\pm$ 0.034	0.121 $\pm$ 0.034	0.020 $\pm$ 0.004
JJA	0.040 $\pm$ 0.010	0.035 $\pm$ 0.010	0.005 $\pm$ 0.001	0.035 $\pm$ 0.009	0.030 $\pm$ 0.009	0.005 $\pm$ 0.001
SON	0.098 $\pm$ 0.025	0.085 $\pm$ 0.025	0.013 $\pm$ 0.003	0.086 $\pm$ 0.021	0.074 $\pm$ 0.021	0.012 $\pm$ 0.003
2014	0.187$\pm$0.049	0.162$\pm$0.049	0.025$\pm$0.006	0.166$\pm$0.040	0.142$\pm$0.040	0.024$\pm$0.005
DJF	0.295 $\pm$ 0.077	0.256 $\pm$ 0.077	0.040 $\pm$ 0.009	0.261 $\pm$ 0.064	0.224 $\pm$ 0.064	0.037 $\pm$ 0.008
MAM	0.182 $\pm$ 0.047	0.158 $\pm$ 0.047	0.024 $\pm$ 0.005	0.161 $\pm$ 0.039	0.138 $\pm$ 0.039	0.023 $\pm$ 0.005
JJA	0.046 $\pm$ 0.012	0.040 $\pm$ 0.012	0.006 $\pm$ 0.001	0.041 $\pm$ 0.010	0.035 $\pm$ 0.010	0.006 $\pm$ 0.001
SON	0.236 $\pm$ 0.061	0.204 $\pm$ 0.061	0.032 $\pm$ 0.007	0.209 $\pm$ 0.051	0.179 $\pm$ 0.051	0.030 $\pm$ 0.007
2015	0.137$\pm$0.036	0.119$\pm$0.036	0.018$\pm$0.004	0.121$\pm$0.030	0.104$\pm$0.030	0.017$\pm$0.004
DJF	0.144 $\pm$ 0.038	0.125 $\pm$ 0.038	0.019 $\pm$ 0.004	0.128 $\pm$ 0.031	0.109 $\pm$ 0.031	0.018 $\pm$ 0.004
MAM	0.180 $\pm$ 0.047	0.156 $\pm$ 0.047	0.024 $\pm$ 0.005	0.160 $\pm$ 0.039	0.137 $\pm$ 0.039	0.023 $\pm$ 0.005
JJA	0.033 $\pm$ 0.009	0.029 $\pm$ 0.009	0.004 $\pm$ 0.001	0.029 $\pm$ 0.007	0.025 $\pm$ 0.007	0.004 $\pm$ 0.001
SON	0.206 $\pm$ 0.054	0.178 $\pm$ 0.054	0.028 $\pm$ 0.006	0.182 $\pm$ 0.044	0.156 $\pm$ 0.044	0.026 $\pm$ 0.006
2016	0.110$\pm$0.029	0.096$\pm$0.029	0.015$\pm$0.003	0.098$\pm$0.024	0.084$\pm$0.024	0.014$\pm$0.003
DJF	0.184 $\pm$ 0.048	0.159 $\pm$ 0.048	0.025 $\pm$ 0.005	0.162 $\pm$ 0.040	0.139 $\pm$ 0.040	0.023 $\pm$ 0.005
MAM	0.112 $\pm$ 0.029	0.097 $\pm$ 0.029	0.015 $\pm$ 0.003	0.099 $\pm$ 0.024	0.085 $\pm$ 0.024	0.014 $\pm$ 0.003

	TC _{bb} PM ₁₀	OC _{bb} PM ₁₀	EC _{bb} PM ₁₀	TC _{bb} PM _{2.5}	OC _{bb} PM _{2.5}	EC _{bb} PM _{2.5}
JJA	0.035±0.009	0.030±0.009	0.005±0.001	0.031±0.007	0.026±0.007	0.004±0.001
SON	0.116±0.030	0.101±0.030	0.016±0.003	0.103±0.025	0.088±0.025	0.015±0.003
2017	0.121±0.031	0.105±0.031	0.016±0.004	0.107±0.026	0.092±0.026	0.015±0.003
DJF	0.231±0.060	0.200±0.060	0.031±0.007	0.204±0.050	0.175±0.050	0.029±0.007
MAM	0.091±0.024	0.079±0.024	0.012±0.003	0.081±0.020	0.069±0.020	0.012±0.003
JJA	0.033±0.009	0.029±0.009	0.004±0.001	0.029±0.007	0.025±0.007	0.004±0.001
SON	0.115±0.030	0.100±0.030	0.015±0.003	0.102±0.025	0.087±0.025	0.015±0.003
2018	0.143±0.037	0.124±0.037	0.019±0.004	0.127±0.031	0.108±0.031	0.018±0.004
DJF	0.198±0.052	0.171±0.051	0.026±0.006	0.175±0.043	0.150±0.043	0.025±0.006
MAM	0.200±0.052	0.173±0.052	0.027±0.006	0.177±0.043	0.151±0.043	0.025±0.006
JJA	0.020±0.005	0.018±0.005	0.003±0.001	0.018±0.004	0.015±0.004	0.003±0.001

57 **Table S 17: Equations showing the relationship between EC, OC and TC for ambient PM_{2.5} aerosol filter samples**
58 **collected at Birkenes in 2014, obtained by temperature calibrated Quartz and EUSAAR-2 temperature programs.**

EC	$EC_{\text{EUSAAR-2, TOT}} = EC_{\text{QUARTZ, TOT}} \times 1.6118$	$R^2 = 0.876$	n = 50	Eq. S 16
OC	$OC_{\text{EUSAAR-2, TOT}} = OC_{\text{QUARTZ, TOT}} \times 0.8687$	$R^2 = 0.977$	n = 50	Eq. S 17
TC	$TC_{\text{EUSAAR-2, TOT}} = TC_{\text{QUARTZ, TOT}} \times 0.9151$	$R^2 = 0.976$	n = 50	Eq. S 18

59

60