



Measurement report: Observations of long-lived volatile organic compounds from the 2019–2020 Australian wildfires during the COALA campaign

Asher P. Mouat¹, Clare Paton-Walsh³, Jack B. Simmons³, Jhonathan Ramirez-Gamboa³,
David W. T. Griffith³, and Jennifer Kaiser^{1,2}

¹Department of Civil and Environmental Engineering,
Georgia Institute of Technology, Atlanta GA 30332, USA

²Department of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta GA 30332, USA

³School of Earth, Atmospheric, and Life Sciences,
University of Wollongong, Wollongong, NSW 2522, Australia

Correspondence: Asher P. Mouat (amouat3@gatech.edu)

Received: 30 August 2021 – Discussion started: 9 September 2021

Revised: 2 June 2022 – Accepted: 22 June 2022 – Published: 31 August 2022

Abstract. In 2019–2020, Australia experienced its largest wildfire season on record. Smoke covered hundreds of square kilometers across the southeastern coast and reached the site of the COALA-2020 (Characterizing Organics and Aerosol Loading over Australia) field campaign in New South Wales. Using a subset of nighttime observations made by a proton-transfer-reaction time-of-flight mass spectrometer (PTR-ToF-MS), we calculate emission ratios (ERs) and factors (EFs) for 15 volatile organic compounds (VOCs). We restrict our analysis to VOCs with sufficiently long lifetimes to be minimally impacted by oxidation over the ~ 8 h between when the smoke was emitted and when it arrived at the field site. We use oxidized VOC to VOC ratios to assess the total amount of radical oxidation: maleic anhydride / furan to assess OH oxidation, and (cis-2-butenediol + furanone) / furan to assess NO₃ oxidation. We examine time series of O₃ and NO₂ given their closely linked chemistry with wildfire plumes and observe their trends during the smoke event. Then we compare ERs calculated from the freshest portion of the plume to ERs calculated using the entire nighttime period. Finding good agreement between the two, we are able to extend our analysis to VOCs measured in more chemically aged portions of the plume. Our analysis provides ERs and EFs for six compounds not previously reported for temperate forests in Australia: acrolein (a compound with significant health impacts), methyl propanoate, methyl methacrylate, maleic anhydride, benzaldehyde, and creosol. We compare our results with two studies in similar Australian biomes, and two studies focused on US temperate forests. We find over half of our EFs are within a factor of 2.5 relative to those presented in Australian biome studies, with nearly all within a factor of 5, indicating reasonable agreement. For US-focused studies, we find similar results with over half our EFs within a factor of 2.5, and nearly all within a factor of 5, again indicating reasonably good agreement. This suggests that comprehensive field measurements of biomass burning VOC emissions in other regions may be applicable to Australian temperate forests. Finally, we quantify the magnitude attributable to the primary compounds contributing to OH reactivity from this plume, finding results comparable to several US-based wildfire and laboratory studies.

1 Introduction

Wildfire smoke significantly affects atmospheric composition, chemistry, human health, and radiative balance (Andreae and Merlet, 2001; Yokelson et al., 2008; Akagi et al., 2011; Ford et al., 2018; Gregory et al., 2018; Sokolik et al., 2019; Macsween et al., 2020). Wildfire season duration and intensity are predicted to increase in the future, suggesting a growing influence of biomass burning in coming decades (Fairman et al., 2015; Donovan et al., 2017; Abatzoglou et al., 2019). Volatile organic compounds (VOCs) emitted from biomass burning (BBVOCs) are directly harmful to human health and can contribute to the formation of ozone and secondary organic aerosol (SOA) (Akagi et al., 2012; Keywood et al., 2013; Lawson et al., 2015; Sekimoto et al., 2017). Predictions of BBVOC emissions are complicated by the complexity of combustion and fuel characteristics, and model parametrizations are based on a limited number of field observations (Hatch et al., 2015; Sekimoto et al., 2018).

Australia wildfires emit 7%–8% of global biomass burning emissions, producing more volatilized carbon than the United States and Europe, with smoke plumes significantly influencing local and even global air quality (Ito and Penner, 2004; Van Der Werf et al., 2010; Keywood et al., 2013; Lawson et al., 2015). In 2019–2020, Australia experienced its worst wildfire season on record with an estimated 19 million hectares of land destroyed (Filkov et al., 2020). This particular season is now colloquially known as the Black Summer, due to its prolonged intensity and length (October 2019–February 2020). Many of Australia's major cities were blanketed in smoke for weeks at a time, leading to long-term exposure to excessive concentrations of harmful atmospheric compounds (Borchers Arriagada et al., 2020). These fires predominantly affected the temperate forests of the state of New South Wales (NSW), burning the largest land area of anywhere in the country (Davey and Sarre, 2020). Despite the impact on atmospheric composition from Australian fires, its biomes remain understudied, particularly these same NSW forests (Lawson et al., 2015). Given the complexity and variability in biomass burning scenarios and the use of emission factors (EFs, in units of kilograms of VOC emitted per kilogram fuel burnt) to inform air quality models, this can lead to issues in effectively constraining emissions. For example, Lawson et al. (2017) reported a strongly non-linear response in simulated ozone (O_3) when varying biomass burning (BB) EFs, showing the resulting sensitivity from chemical transport models (CTMs). This sparseness of measurements leads to the use of North American EFs (such as those from Burling et al., 2011, or Akagi et al., 2011) to inform CTMs, simulating emissions of geographically separate biomes. Even among similar biomes (for instance, the temperate forests of the US), fuel types differ and thus can influence the speciation of VOCs emitted (Coggon et al., 2016; Hatch et al., 2017; Guérette et al., 2018). Further evidence of this is found in a study by Guérette et al. (2018) showing that

EFs of some VOCs (e.g., formic acid, ethane, monoterpenes, acetonitrile) can be 3–5 times higher than those measured in the US, and attributing this to fuel type.

A complicating factor in deriving EFs from field observations is accounting for the influence of chemical processing. EFs are ideally based on observations close to the fire. When this is not possible, indicators of plume chemical age, such as oxidized VOC (OVOC) to VOC ratios, can be used to diagnose the relative age of a plume. During the day, downwind VOC concentrations are primarily influenced by OH-initiated oxidation. At night, NO_3 -initiated oxidation can significantly influence observed VOC concentrations (Decker et al., 2019; Kodros et al., 2020). There are several methods in existence for assessing daytime oxidation, but fewer are known for the night (De Gouw et al., 2006; Liu et al., 2016; Gregory et al., 2018; Decker et al., 2019). In this work, we use the maleic anhydride-to-furan ratio introduced in Gkatzelis et al. (2020) to assess OH oxidation. We examine the use of a new OVOC/VOC ratio, *cis*-2-butenediol + furanone-to-furan, as an indicator of nighttime oxidation.

To further assess the effects of nighttime transport on biomass burning emissions, we look at the magnitude of OH reactivity measured that results from the compounds which most substantially contribute to it and determine the relative contributions of the resulting chemical groups. Certain categories of BBVOCs like furans or phenols, which are emitted in the combustion process, are important as they enhance OH reactivity and resultingly have high O_3 and SOA forming potential, and are considered to be understudied (Gilman et al., 2015; Hatch et al., 2017). Most wildfire studies are conducted during the daytime, with plume oxidation focused on interactions with the OH radical and O_3 (Liu et al., 2016; Coggon et al., 2019; Palm et al., 2020; Decker et al., 2021; Permar et al., 2021). However, the plume studied here spent a significant amount of time transported under nighttime conditions.

Additionally, we use time series to observe chemical trends in ozone (O_3) and nitrogen dioxide (NO_2) as their emissions and chemical behavior are intimately linked with biomass burning chemistry. O_3 production in wildfire plumes is contingent on initial emissions, local environment, and atmospheric processing during transportation. Wildfire plumes emit significant precursors of O_3 , but there is not a general consensus towards generation or depletion, with various campaigns reporting measurements in either case, especially in the instance of processed, downwind plumes (Verma et al., 2009; Alvarado et al., 2010; Jaffe and Wigder, 2012; Lawson et al., 2015; Brey and Fischer, 2016; Müller et al., 2016). NO_2 is emitted during the combustion process and has a non-linear relationship to O_3 production via reactions with these VOC precursors. However, the NO_2 radical has additional chemical pathways with OH, NO_3 , and phenolic compounds leading to a general NO_x -limited regime (Jaffe and Wigder, 2012; Liang et al., 2022; Robinson et al., 2021). Furthermore, there are again fewer observations for the effect of nighttime

oxidation processes on O₃ production with a recent modeling study conducted by Decker et al. (2019). O₃ production in wildfire smoke remains a significant source of uncertainty in its contribution to the tropospheric O₃ budget (Jaffe and Wigder, 2012; Young et al., 2018; Xu et al., 2021).

Here, we use observations from a proton-transfer-reaction time-of-flight mass spectrometer (PTR-ToF-MS) during the 2019–2020 Australian wildfire season to derive EFs of 15 compounds, including 6 compounds for which there are no previous observations. We examine a subset of smoke-influenced nighttime observations made by a PTR-ToF-MS during the COALA-2020 field campaign. NO₃-initiated oxidation dominated the chemical processing late in the night, as the plume traveled ~ 8 h to the field site from large, highly active fires to the south. We also use co-located Fourier transform infrared (FTIR) measurements of CO₂, CO and CH₄ to derive these EFs for nighttime longer-lived VOCs ($\tau_{\text{BBVOC}+\text{NO}_3} \geq \text{average transport time}$). We compare these results with five related studies, two focused on Australian temperate forests, two focused on US temperature forests, and one reporting EFs used to represent temperate biomes across the globe. We find generally good agreement across several of these studies and discuss potential reasons for discrepancies seen in EFs for selected compounds.

2 Field site and instrument description

2.1 Field site and active fires

The COALA-2020 field site was located in Cataract Scout Park (34.247° S, 150.825° E) at 400 m above sea level, 15 km inland, and 30 km to the northwest of the nearest urban area (Wollongong, NSW). Figure 1 shows the field site relative to the fires active between 1 and 5 February 2020. We use the Suomi VIIRS thermal anomalies product filtering for points at high confidence levels to avoid counting any reflective false positives from plains or urban centers. Also plotted is the normalized difference vegetative index (NDVI), which is determined from measurements aboard the MODIS Terra satellite (Didan, 2021). The fires are primarily located in temperate forests along the southeastern coast, with a small inland group near Canberra. These forests consist of open, tall woodlands made up of *Eucalyptus* species grouped generally as dry sclerophyll.

2.2 PTR-ToF-MS and supporting observations

VOCs were measured using an Ionicon PTR-ToF-MS 4000, which operated with a mass resolution between 2000–3000 FWHM $m \Delta m^{-1}$ and at a mass range spanning $m/z = 18$ –256. The drift tube was held at a temperature of 70 °C, pressure at 2.60 mbar, and an $E/N = 120$ Td (electric field to molecular number density ratio; 1 townsend = 10^{-21} V m²). The instrument was housed in a climate-controlled unit, connected to a 15 m long, 1/4 in. outer diam-

eter (OD) PTFE insulated line attached to a 10 m tall mast, placing inlet height 0.5 m above canopy height. The sample flowed through the inlet at 3 SLPM for a residence time of 2.5 s. Peak separation of 1 min averaged spectra was conducted in Ionicon's PTR-Viewer 3 software.

Calibrations were performed using two VOC cylinders designed by Airgas on 31 January 2020, three days before measuring the smoke event discussed here. A second calibration was performed in the following week with little change in instrument sensitivity. The cylinders contained 17 compounds spanning a mass range of 33–154 Da and are shown in Table S1 in the Supplement. Many of these compounds are reported in the final EFs list – methanol, acetonitrile, acetaldehyde, acrolein, acetone, MVK + MACR, benzene, C₈ aromatics, and C₉ benzenes. All compounds used either do not fragment under these drift tube conditions or have known fragmentary peaks. Instrument zeros were determined using ultra-zero air. Limits of detection (3σ) for calibrated species are also given in Table S1 and range between 5–165 ppt. The raw counts per second (cps) were corrected for instrument transmission, which was determined using a subset of the species in the calibration standards. Corrected cps are then normalized (ncps) to the reagent ion signal ($\text{H}_3\text{O}^+ \text{ccps} \times 10^6 \text{ncps}$) using the methodology described by Sekimoto et al. (2017). For compounds of interest not included in the calibration standards, we use the method described by Sekimoto et al. (2017), which yields uncertainties at 100 %. Table S2 shows all compounds presented in this study alongside whether they were included in the calibration standards and their respective uncertainty.

In addition to the PTR-ToF-MS measurements, we use observations of CO, CO₂, and CH₄ obtained from the collocated FTIR system. Information of this instrument and its setup is provided in Griffith et al. (2012).

3 Observed CO, VOC, and OVOC enhancements

Figure 2 shows the observations of CO and VOCs during a smoky period on 3–4 February 2020. CO and acetonitrile – long-lived tracers associated with wildfires (Coggon et al., 2016) – are used to identify the total period of time during which observations were impacted by smoke. Enhancements in both species started at 17:30 LT on 3 February and lasted until 19:00 LT on 4 February, when wind direction shifted.

We use furan, a short-lived smoke tracer, and its oxidation products to determine which periods of the smoke event represent the least oxidized plume. Furan is highly reactive with OH ($k_{\text{OH}+\text{furan}} = 4.04 \times 10^{-11} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1}$ at 298 K and 1 atm) and NO₃ ($k_{\text{NO}_3+\text{furan}} = 1.36 \times 10^{-12} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1}$ at 298 K and 1 atm). OH-initiated oxidation produces maleic anhydride, which has low reactivity with both OH and NO₃ ($\tau_{\text{OH}} = 3.99 \text{ d}$, $\tau_{\text{NO}_3} = 1.42 \text{ d}$ with $[\text{OH}]_{\text{Avg}} = 2 \times 10^6 \text{ molec. cm}^{-3}$ and $[\text{NO}_3]_{\text{Avg}} = 8 \times 10^7 \text{ molec. cm}^{-3}$, with reaction rate constants from Grosjean




Figure 1. Active fires from 1–5 February 2020 and their proximity to the COALA-2020 field site. Normalized Difference Vegetative Index (NDVI) is plotted at 250 m resolution from the MOD13A1 dataset acquired by measurements via the MODIS Terra satellite. Pixels have been filtered to contain cloud coverage less than 30 % and VI usefulness bits indicating top two tiers of data quality. Fire counts are plotted using the VNP14IMGTDL_NRT data from Suomi VIIRS satellite imaging overlaid with HYSPLIT back trajectories. Each tail represents a trajectory 12 h prior to reaching the site and is colored by its starting time. Circles indicate 1 h intervals moving backwards from the start time.




Figure 2. VOCs and CO on 3–4 February 2020 with the shaded area representing sunset to sunrise. The peak in CO after sunset (start of gray-shaded area) is used to denote the beginning of the smoke event. We limit our analysis to sunrise on the following day. The color labels A–D indicate individual times used to calculate ERs (see Sect. 5.2 in main text). m/z 85 in the bottom time series indicates the sum of furanone and cis-2-butenediol.

and Williams (1992) and Bierbach et al. (1994) and no reported direct emissions). The ratio of maleic anhydride-to-furan therefore provides a relative measure of the plume photochemical age. Using aircraft-based observations of wildfire plumes in the western US, Gkatzelis et al. (2020) found

that maleic anhydride-to-furan ratios below 0.10 indicate the plume has undergone little OH processing.

Nighttime in-plume furan oxidation is dominated by NO_3 , with contributions from O_3 (Decker et al., 2019). While many BBVOCs are highly reactive with NO_3 , there is substantially less research on indicators of NO_3 oxidation. Decker et al. (2019) track NO_3 chemistry using the ratio of total reactive nitrogen (NO_y) to NO_x , and Kodros et al. (2020) examine NO_3 -reacted products such as nitrocatechol and nitrophenol of phenolic compounds (e.g., phenol, catechol, cresol). Measurements of NO_y were not made during this field campaign, and NO_3 products of phenols were subject to high uncertainty due to fragmentation in our PTR-ToF-MS measurement. We therefore examine a new indicator of NO_3 processing using furan's dominant NO_3 products – cis-2-butenediol and furanone (Berndt et al., 1997). Both products are relatively long lived, with lifetimes estimated at $\tau_{\text{cis-2-butenediol}} = 9 \text{ d}$ and $\tau_{\text{furanone}} = 8 \text{ h}$ assuming an average concentration of $[\text{NO}_3] = 8 \times 10^7 \text{ molec. cm}^{-3}$ (O'Dell et al., 2020). Lab-based studies and field campaigns conducted in the US and Australia suggest that furan and furanone EFs are comparable, with study-averaged values for furan ranging from 0.132–0.51 g kg^{-1} and 0.27–0.57 for furanone (Andreae and Merlet, 2001; Akagi et al., 2011; Hatch et al., 2015; Stockwell et al., 2015; Liu et al., 2017; Koss et al., 2018; Selimovic et al., 2018). No furan EFs have been reported for Australian temperate forests and only one furanone EF is reported from Lawson et al. (2015) at a comparable value at

0.57 g kg^{-1} . Additionally, emissions modeled in Decker et al. (2019) from wildfires suggest that furan and furanone are emitted in roughly equal proportions. As such, we operate not on the assumption of negligible OVOC emissions, but that variability in OVOC/VOC ratios are driven by chemical aging. Cis-2-butenediol and furanone are both measured at m/z 85, and from here onwards will be denoted as such.

Figure 2 shows furan enhancements, which begin later on 3 February than acetonitrile, maleic anhydride, and m/z 85 enhancements, indicating a less oxidized plume was being sampled. Maleic anhydride concentrations are high during the initial period of the smoke event, suggesting significant OH-initiated processing throughout the day before the plume reached the site. After sunrise, furan decays faster than CO, and maleic anhydride concentrations begin to rise, again showing the impact of OH-initiated oxidation. Enhancements in m/z 85 are seen when the smoke arrives and vary throughout the night. Just prior to sunrise (04:00–06:15 LT), both OVOC/VOC ratios rapidly decrease (Fig. 3), corresponding with a rise in furan, CO, and acetonitrile. Maleic anhydride/furan drops to 0.05, which is within the lower range of the chemically younger plumes reported by Gkatzelis et al. (2020). The ratio of m/z 85 to furan is around 2.5. While we cannot use this to quantify plume age since the two products are measured as a sum, we note that this period constitutes the lowest ratio throughout the event, with surrounding periods having ratios 1.6–2.8 times greater. We note that at a value of 2.5, this plume has likely undergone significant aging, despite this being the freshest smoke detected during the campaign. Further corroboration of these results, determined via particulate matter (PM) composition, can be found in Simmons et al. (2022). In their study, a ToF-ACSM was employed and observed the ratio of PM_{10} mass fraction at mass-to-charge ratio 44 (f_{44}), where a lower mass fraction indicates a less oxidized plume. A similar decrease at f_{44} in the same timeframe as the m/z 85 and maleic anhydride tracers is noted.

The rapid decreases in OVOC/VOC ratios are unlikely to result from shifts in chemistry alone. Instead, this suggests a shift in meteorological conditions which brings in smoke from a closer source, in agreement with measured wind direction, which shifted from flowing northeast to north at this time. We further investigate plume transport using a back-trajectory model.

4 Plume origin and transport time

We use a HYSPLIT back-trajectory model (Stein et al., 2015) to determine the origin and transport time of the smoke arriving at the site throughout the smoke event. The meteorological input used is the Global Data Assimilation (GDAS) dataset. The model was set to assess trajectories at three different altitudes at 10, 500, and 1500 m above ground level (a.g.l.) to capture plume height. Our period of interest spans




Figure 3. Product-to-reactant ratio for furan oxidation products. Both ratios indicate the period just before sunrise is least oxidized. Again, the color labels A–D indicate individual times used to calculate ERs. m/z 85 indicates the sum of furanone and cis-2-butenediol.

from 17:00 LT on 3 February, just before CO enhancements are seen at the site, to 06:00 LT on 4 February when furan concentrations rapidly decrease. The model was set to calculate a new 12 h trajectory every hour during this time. Back trajectories are shown in Fig. 1. For every hour in the event (each represented by a color), one can track the origin of the sampled air mass 12 h in advance of its arrival.

A shift in trajectories occurred between 17:00 and 18:00 LT on 3 February, corresponding with the arrival of the smoke plume as indicated by observed CO enhancements. Subsequent trajectories originate near the fires located ~ 230 – 375 km from the field site on the southeast coast. The model shows that air masses initially stayed at low altitude and were lofted to ~ 560 m a.g.l. when passing over the active fires ~ 25 km to the south, near Canberra (Fig. S1 in the Supplement). The plume descended to 10 m a.g.l. as it reached the coast. The model suggests smoke sampled later in the evening (between 04:00–06:00 LT on 4 February) spent more time over land compared to previous points in the event. This shift in trajectories and the increasing intensity of fires near Canberra during this time signify possible contributions to the decrease in OVOC to VOC age marker ratios. Further investigation is conducted via HYSPLIT forward trajectories in the supplement (Figs. S2 and S3). In short, during this period, plumes from the Canberra fires were lofted to 2000 m a.g.l. well before crossing with the SE fire plume, which attained a maximum altitude up to 560 m a.g.l. This indicates little influence from the Canberra fires on our measurements. Given that there are two major clusters approximately 70 km apart in the SE, the influence of precipitation and wind speed (Figs. S4–S6) is considered to determine whether combustion conditions were comparable. Both fires experienced similar total precipitation in the month

prior and experienced similar wind speeds during this smoke plume event. As a result, we conclude that combustion conditions are similar and that EFs derived from this plume would be representative of a biome average. Over the entire course of the event, HYSPLIT analysis suggests transport time from the fires to the field site is around 8 h (> 200 km), but potentially shorter for the time frame immediately prior to sunrise.

5 O₃ and NO₂ time series

Detailed time series of O₃ and NO₂ are presented in this section in Fig. 4. Information regarding instrumentation and corresponding setups can be found in Sect. 2.1 of Simmons et al. (2022). Like Fig. 2, a CO time series is provided to outline the general trend of smoke during the event.

A non-smoke-influenced daytime and nighttime average (composed of 8 h averages) was calculated for O₃ and NO₂ concentrations using data from the month of March. Smoke around the continent had been either transported or removed by rain by this time. O₃ was calculated to have a daytime concentration of 24.6 ppb and a nighttime concentration of 19.5 ppb. Respective concentrations were calculated for NO₂ at 2.2 ppb in the day and 3.3 ppb in the evening. Additionally, averages for a larger suite of gas and aerosol phase variables over all smoke events sampled during the COALA-2020 campaign can be found in Simmons et al. (2022).

As stated before, smoke-related enhancements are visible around 17:30 LT in Fig. 4d, with the hours prior being virtually devoid of tracers. Enhancements pick up without a shift in wind direction, with winds at this time traveling to the northwest, consistent with the HYSPLIT trajectories presented in Fig. 2. As the wind approaches a more easterly direction, enhancements in CO are maintained, and concentrations of more reactive BBVOCs begin to increase. O₃ concentration on 3 February reaches a maximum of approximately 25 ppb around 14:00 LT and maintains this level until sunset, when it decreases as biogenic sources are no longer emitting and photolysis is halted. O₃ concentration decreases to a minimum 15.6 ppb and NO₂ decreases down to 0.8 ppb, both around midnight and both below the nighttime monthly average despite enhancements in CO. O₃ has a $R^2 = 0.48$ with CO and, when considering the known transport time of this smoke, indicates transportation rather than local production. Given the comparatively low concentrations of both compounds at this time, it is likely that this plume is depleting these species. This is compounded with the low concentrations of NO₂ in this temperate forest setting and, despite emitting NO_x, wildfire plumes being generally NO_x-limited (Jaffe and Wigder, 2012; Robinson et al., 2021).

Around 03:30, the wind shifts from traveling northwest to west, significantly enhancing O₃, NO₂, CO, and total VOC concentrations, corresponding to the least aged portion discussed in Sect. 3. Sunrise occurs around 06:30 LT coinciding with a steady decline in highly reactive VOC enhancements

(Fig. 2) and NO₂ (Fig. 4c). Liang et al. (2022) found a significant correlation of $R^2 = 0.86$ between NO₂ and maleic anhydride for a transported plume of similar age oxidized in the daytime. The opposite trend is observed in our scenario despite our measurements exhibiting comparable trends from maleic anhydride. The NO_x-limited environment and differences in biogenic VOC (BVOC) quantities arising from the forest setting in this study and the urban setting in theirs are likely responsible for the opposing trends in the NO₂ time series. Maleic anhydride similarly peaks around noon on 4 February, and both its production and the fast depletion of furan indicate that OH chemical pathways generally oxidize this plume faster than NO₃ reaction pathways. While O₃ concentration continues to increase after sunrise, it cannot be stated that this is dominantly due to BBVOC oxidation given the strong source of BVOC emissions from the surrounding forest. Isoprene nitrates sequester NO₂, ultimately leading to O₃ production. The diel cycle of O₃ and isoprene on a non-smoke-affected day strongly correlate to temperature and photoactive radiation. O₃ does achieve a max concentration of 30 ppb at 12:00 LT on 4 February, which is approximately 5.5 ppb above the daytime average and higher than the prior day despite similar temperatures (23.6 °C on 3 February, and 24.5 °C on 4 February). This most likely results from the combination of transported O₃ compounded with enhanced reactivity from the plume plus local, biogenic-related production. The plume is diluted at a consistent rate until 18:00 LT on 4 February when a shift in wind direction significantly reduces CO enhancements and concludes the smoke event.

6 Emission factors

6.1 Species selection

To identify compounds which would be suitable for EF derivation, we compare the list of measured ions with compounds identified in previous literature such as Brill et al. (2014), Hatch et al. (2015), Gilman et al. (2015), Stockwell et al. (2015), Bruns et al. (2017), Koss et al. (2018), and the PTR Library (Pagonis et al., 2019). To corroborate species assignment, we examine correlations of identifiable compounds with CO, acetonitrile, furans, and phenolic compounds, which are well-established smoke tracers. We also examine tracer–tracer relationships, for instance the anti-trend between maleic anhydride and furan resulting from OH oxidation. We exclude compounds with low proton affinities that are known to have humidity-dependent calibration factors (e.g., HCHO, HCN). This results in 150 identified VOCs species measured during the smoke event.

We further filter our VOC list by two criteria. First, VOC + NO₃ reaction rates must be included either in the NIST Chemical Kinetics Database (Manion et al., 2015) or Master Chemical Mechanism (v3.3.1) (Bloss et al., 2005; Jenkin et al., 1997, 2003; Saunders et al., 2003). Second, the




Figure 4. Time series for O₃, NO₂, CO, and wind direction. (a) Wind direction is read as true north is 0° and east is 90°. (b) O₃ trends well with CO until sunrise occurs, wherein BBVOC + OH oxidation combined with biogenic VOC emissions led to daily production. The close trend with CO over nighttime indicates transport rather than local formation. (c) NO₂ also shows a similar trend but upon sunrise begins to negatively correlate with CO and O₃. (d) CO smoke tracer provided as time series reference.

VOC must have a significantly long lifetime against NO₃ oxidation to be minimally impacted over the 8 h transit time from the active fires to the field site ($\tau_{\text{BBVOC}+\text{NO}_3} < 8 \text{ h}$, again assuming $[\text{NO}_3] = 8 \times 10^7 \text{ molec. cm}^{-3}$).

6.2 Calculating emission ratios

An ER is defined here as the slope of a regression of a given VOC to CO (both in units of ppb). Following Guérette et al. (2018), ERs are reported if correlation between a given VOC and CO are well correlated, with $R^2 \geq 0.5$. High correlation minimizes the impact of the choice of regression method (e.g., orthogonal, York) on calculated slopes (Wu and Yu, 2018) and removes the need to account for background corrections (additional discussion of surrounding influential sources can be found in Simmons et al., 2022). We use a reduced major axis regression to determine emission ratios. Given the time component that affects our measurements, it should be noted that compounds with low emission factors and high reactivity are likely to be excluded as they have been reacted away before reaching the site, thus exhibiting an insufficient CO correlation.

We first derive ERs using all data from the “freshest” portion of the plume as determined from OVOC/VOC ratios (marked “D” in Fig. 2). This produces 15 ERs that meet our criteria. We expect this period to provide the most accurate representation of original VOC emissions. We then calculate ERs for more aged portions of the smoke event (Periods A–C, Fig. 2), performing regression analysis on the chemically distinct time periods. The start and end time of each period is determined by visual inspection of VOC/CO behaviors,

which all exhibit similar distinct periods. Figure 5 provides an example of the analysis using acrolein. We average the slopes from each of these lines to derive an average ER for the full smoke event and compare to just the freshest portion of the plume (Period D). We find that using only the freshest smoke compared to using all the data generates very similar results for 9 of the 15 compounds (of which these 9 all have multiple ERs over the evening). Relative differences of the resultant ERs are within 1.5%–47% with two outliers: C₈ aromatics (88%) and C₃ benzenes (212%). Three compounds have only one ER from all four periods (maleic anhydride, benzaldehyde, and creosol) so there is no standard deviation, but the remaining compounds from period D are captured within 1σ of ERs from periods A–D (shown in Fig. S7). Good agreement between methods allows us to extend our analysis beyond the freshest part of the plume and therefore allows us to report ERs for a larger number of compounds. When focusing only on the freshest part of the plume, maleic anhydride and benzaldehyde must be excluded due to insufficient R^2 with CO. All ERs reported here and used in EF calculation use the “average over evening” method and include these compounds. Additionally, only one ER for CO₂ and CH₄ have been calculated using the dataset from periods A–D. Both these compounds are long-lived, and from visual inspection, they do not form distinct time periods like the VOC ERs (shown in Fig. 4). A table with the resultant VOC ERs is also provided in the Supplement (Table S3). We use the CO₂ ER to determine an average modified combustion efficiency with the following equation:

$$\text{MCE} = \frac{\text{ER}_{\text{CO}_2}}{\text{ER}_{\text{CO}_2} + \text{ER}_{\text{CO}}}, \quad (1)$$

where the ER_{CO} is just unity and $\text{ER}_{\text{CO}_2/\text{CO}}$ is 10.82 ppb CO_2 ppb CO^{-1} . This results in an MCE calculation of 0.92, indicating a less efficient, even mixture of smoldering and flaming (Akagi et al., 2011).

6.3 Calculating emission factors

Emission factors are defined as the mass of some trace gas emitted per mass of dry biomass burnt. The most direct way of calculating this quantity is capturing total emissions released from a fire as well as knowing the quantity of fuel burnt. Unless experiments are conducted in a laboratory setting, these quantities are not known. As such, emission factors are calculated according to the carbon mass balance method (Akagi et al., 2011; Selimovic et al., 2018), using CO as the reference gas for the 15 reported species, which produces the following equation:

$$\text{EF}_X = F_{\text{carbon}} \times 1000 \times \text{MM}_X / \text{MM}_C \times \text{ER}_{X/\text{CO}} / \sum \text{ER}_{Y/\text{CO}} \quad (2)$$

where $F_{\text{carbon}} = 0.5$ and is the assumed carbon fractional content of the fuel as used in previous studies (Akagi et al., 2011; Paton-Walsh et al., 2014). MM_X is the molar mass of compound X; MM_C is the molar mass of carbon; $\text{ER}_{X/\text{CO}}$ is the CO ER of X; and $\sum \text{ER}_{Y/\text{CO}}$ is the sum of $\text{ER}_{\text{CO}_2/\text{CO}}$, $\text{ER}_{\text{CH}_4/\text{CO}}$, and $\text{ER}_{\text{CO}/\text{CO}}$. These ERs constitute the major volatilized carbon components of the plume, but the resulting EFs may be overestimated by 1%–2% (Andreae and Merlet, 2001) as this method assumes all volatilized carbon is detected including particulate carbon and VOCs.

EFs derived in this work are presented in Table 1 alongside results from two eastern Australia-based studies by Lawson et al. (2015) and Guérette et al. (2018), two western US-based studies sampling emissions from corresponding temperate fuel types by Liu et al. (2017) and Permar et al. (2021), and one study by Akagi et al. (2011) that provides EFs for general temperate zones. Additionally, Fig. S8 displays these results via scatter plot.

First, in comparison with the Australia-based studies, Guérette et al. (2018) reports EFs notably larger than those presented in this work, with only benzene and C_8 aromatics showing good agreement. Except for these two compounds and C_3 benzenes, Guérette et al. (2018) reports larger EFs than Lawson et al. (2015) and none within agreement. Our results more closely agree with Lawson et al. (2015) with methanol, acetone, and furanone EFs within 1σ , and acetonitrile and acetaldehyde falling within a factor of 2. This agreement is likely due to both this work and Lawson et al. (2015) examining opportunistically intercepted smoke plumes that experienced some processing, whereas Guérette et al. (2018)

sampled near-source, controlled ground burns. Guérette et al. (2018) reports an acetonitrile EF ~ 4.5 times higher than this work and ~ 3 times greater than Lawson et al. (2015) constituting one of the largest disparities. This is attributed to the native and abundant acacias, which are N-fixing species located mainly in forest understories. Their measurements likely had a higher proportion of this foliage constituting the total fuel load due to both proximity to the forest floor and resulting leaf litter. Another of the largest differences is MVK + MACR, which shows a disparity of ~ 6 times this work and 3 times that of Lawson et al. (2015). This is also most likely explained by differences in sampling approach in that proportional contributions of vegetation vary and plumes in Guérette et al. (2018) did not undergo any dilution or photochemical processing.

In comparison with US-based studies, methanol, acetonitrile, acetone, and benzene agree across both studies within 1σ , with acrolein, methyl propanoate, methyl methacrylate, C_3 benzenes, and creosol agreeing very well with values reported by Permar et al. (2021). It should be noted though that within the estimated uncertainties, the value for creosol reported by Permar et al. (2021) is ~ 3.5 times greater than the value in this work, which constitutes another of the largest disparities in this dataset. Additionally, methanol agrees well with the value from Akagi et al. (2011). The EF for m/z 85 in this work is also expectedly larger than both other values presented here at ~ 3 times greater than Permar et al. (2021). This is likely due to the plume sampled in this work undergoing the longest transport of any plumes measured in other studies.

Perhaps an unexpected finding is that EFs derived in this work agree better with observations in the US than the Guérette et al. (2018) study, which was in the same region as the COALA-2020 measurements. It should be noted that all studies except Guérette et al. (2018) are from plumes sampled several kilometers downwind. Differences previously characterized as arising from varying fuel types may actually result from measurement approaches to deriving EFs and proximity to emission source. Agreement across results from this work and from the US-based studies lends credence to the use of newly presented EFs for modeling purposes in temperate Australian forests. Further corroborating this notion is the extremely good agreement (all EFs within uncertainty for all three studies) found between EFs in this work and those presented in Stockwell et al. (2015) and Koss et al. (2018). These results can be seen in the Supplement in Fig. S9.

7 OH reactivity

As this plume has been shown to oxidize faster when exposed to the OH radical as opposed to the NO_3 radical, this indicates that the nighttime transport of this plume would be able to comparably preserve OH reactivity. We investigate this by




Figure 5. Example ER analysis (a) using acrolein, wherein the smoke event is partitioned into four periods over the evening. Average ERs (slopes) from periods A–C agree closely with those in the freshest portion of the plume (D). Panels (b) and (c) show the singular ERs derived for CO₂ and CH₄ using the entire nighttime dataset (A–D).

Table 1. EFs (g kg⁻¹) derived in this work compared to two studies conducted in the same or near temperate Australian forests, two US-based aircraft campaigns sampling western temperate US fuels, and one study reporting EFs across geographically distant temperate forests. Again, *m/z* 85 indicates the sum of furanone and cis-2-butenediol.

Compound	Formula	<i>m/z</i>	Biome location					Temperate forests
			AU		US		Akagi et al. (2011)	
			This work	Guérette et al. (2018)	Lawson et al. (2015)	Liu et al. (2017)		
Methanol	CH ₄ O	33.00	2.01 ± 0.58	3.0 ± 0.5	2.07 ± –	2.45 ± 1.43	33.00	1.93 ± 1.38
Acetonitrile	C ₂ H ₃ N	42.03	0.16 ± 0.03	0.70 ± 0.10	0.25 ± –	0.25 ± 0.13	42.03	–
Acetaldehyde	C ₂ H ₄ O	45.03	0.57 ± 0.20	1.20 ± 0.30	0.92 ± –	1.64 ± 0.52	45.03	–
Acrolein	C ₃ H ₄ O	57.03	0.23 ± 0.08	–	–	–	57.03	–
Acetone	C ₃ H ₆ O	59.05	0.55 ± 0.28	0.80 ± 0.20	0.54 ± –	1.13 ± 0.82	59.05	–
MVK+MACR	C ₄ H ₆ O	71.05	0.18 ± 0.03	1.0 ± 0.30	0.38 ± –	0.33 ± 0.06	71.05	–
Benzene	C ₆ H ₆	79.05	0.25 ± 0.08	0.39 ± 0.07	0.69 ± –	0.43 ± 0.12	79.05	–
<i>m/z</i> 85	C ₄ H ₄ O ₂	85.03	0.83 ± 0.27	–	0.57 ± –	0.39 ± –	85.03	–
Methyl propanoate	C ₄ H ₈ O ₂	89.06	0.07 ± 0.03	–	–	–	89.06	–
Maleic Anhydride	C ₄ H ₂ O ₃	99.00	0.05 ± –	–	–	–	99.00	–
Methyl methacrylate	C ₅ H ₈ O ₂	101.06	0.07 ± 0.04	–	–	–	101.06	–
Benzaldehyde	C ₇ H ₆ O	107.05	0.05 ± –	–	–	–	107.05	–
C ₈ aromatics	C ₈ H ₁₀	107.09	0.08 ± 0.05	0.11 ± 0.03	0.26 ± –	0.15 ± 0.004	107.09	–
C ₃ benzenes	C ₉ H ₁₂	121.10	0.07 ± 0.06	–	0.27 ± –	–	121.10	–
Creosol	C ₈ H ₁₀ O ₂	139.08	0.05 ± –	–	–	–	139.08	–

* Dashes indicate either EF or EF variability not reported in study.

first determining which compounds were most significant in their enhancements and then determining their corresponding OH reactivity.

First, a subset of the PTR-ToF-MS data was created by calculating ERs using the methodology described in Sect. 6.2 over the same nighttime period. However, we did not filter out compounds by their atmospheric lifetime, and any unidentified species were not considered regardless of correlation strength. This means the resulting OH reactivity is likely to be slightly low, but this method ensures reactivity solely from compounds attributable to BB emissions is being gauged. Then, an average for each compound was calculated using the same period for ERs. These nighttime aver-

ages were then compared with their diurnal cycles calculated using data from 1–19 March 2020 (ending date of PTR-ToF-MS sampling ambient air). If a compound's mean over the smokey period is greater than the mean of its diel cycle plus 1σ over the same timeframe, this compound is considered in the transported OH reactivity. Finally, we background correct the nighttime concentrations using the March diurnal cycles and convert to reactivity using Eq. (3):

$$R_{\text{OH}} = \sum [\text{VOC}_i] \cdot A \cdot k_{\text{VOC}_i+\text{OH}}, \quad (3)$$

where [VOC_{*i*}] is the concentration of the *i*th VOC in units of parts per billion, *A* is the conversion factor to molec._{*i*} cm⁻³ (*A* = 2.46 × 10¹⁰ in units of molec._{air} cm⁻³ ppb⁻¹ at 1 atm

and 25 °C), and $k_{\text{VOC}_i} + \text{OH}$ is the OH rate constant for the corresponding VOC_i . Rate constants were again sourced from the same databases as the NO_3 rate constants. The rate constant used for m/z 85 was determined as an average of the constant provided in Koss et al. (2018) ($k_{\text{OH}} = 44.2 \times 10^{-12} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1}$) and Bierbach et al. (1994) ($k_{\text{OH}} = 52.1 \times 10^{-12} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1}$) assuming both compounds contributed equally to signal at this mass peak.

Ultimately, 26 compounds were determined to have the most significant contributions, transporting an average OH reactivity of 5.25 s^{-1} , with a minimum of 3.15 s^{-1} occurring around 03:00 LT on 4 February and a maximum of 9.83 s^{-1} around 20:00 LT on 3 February, shown in Fig. 6. These values are well within range of those seen in nighttime and aged daytime transported plumes by Liang et al. (2022), who measured a total OH reactivity range from approximately $4\text{--}26 \text{ s}^{-1}$. We calculate an OH reactivity from the primary biogenic VOCs (isoprene plus monoterpenes) for further comparison. The maximum biogenic value, achieved around 12:00 LT on 4 February, is 6.35 s^{-1} , and the average biogenic reactivity over the course of the campaign is 5.90 s^{-1} , indicating that the nighttime conditions allowed for the transport of a reactivity quantity that approximately doubled OH reactivity at the COALA-2020 field site. Additionally, there is little variability in the relative contributions to reactivity across these different groups over the course of the smoke event, indicating the plume experienced a consistent oxidation over the course of its travel.

Compounds from the plume have been grouped into four categories to capture their diversity. Expectedly, biogenic emissions contribute the most to total reactivity (attributable dominantly to isoprene), but the furans group is the most reactive with values from $1.24\text{--}3.93 \text{ s}^{-1}$. This group contains various furans (furan, 2-methylfuran, m/z 85, and furfural alcohol) wherein m/z 85 is by far the most significant, contributing up to 69% of the group total. This high m/z 85 presence explains why this group is also the most OH reactive as most furans are largely oxidized by NO_3 during this transport timeframe, except m/z 85, which has a long τ_{NO_3} but a comparatively shorter τ_{OH} . The furan reactivity range is comparable to lab-based values measured in Gilman et al. (2015), which ranged from $1.3\text{--}5.5 \text{ s}^{-1}$. Both these studies find lower furan reactivities than lab measurements made in Koss et al. (2018) at an average reactivity of 14.2 s^{-1} , where furans constitute the third highest reactivity group. Aromatics make up the second most reactive group (range of $0.66\text{--}2.14 \text{ s}^{-1}$) in this study, with dominant contributions from phenol (39%), styrene (33%), and catechol (32%). Catechol's contribution is likely less than this as other studies have revealed that it shares a significant portion of its mass peak with 5-methyl furfural (Stockwell et al., 2015; Koss et al., 2018). Despite their high NO_3 reactivity, phenolic compounds still dominate the overall OH reactivity contributions in this category. These compounds appear across other studies as primary contributors to OH reactivity (Gilman et al.,

2015; Hatch et al., 2017; Koss et al., 2018; Sekimoto et al., 2018; Decker et al., 2019; Liang et al., 2022). Alkenes (range of $0.86\text{--}1.83 \text{ s}^{-1}$) are on par with aromatics, for which their reactivity is largely attributable to propene and butene, followed finally by non-aromatic oxygenates (range of $0.28\text{--}1.87 \text{ s}^{-1}$), which contain compounds like methanol, acetaldehyde, and acetic acid. The comparably low reactivity from this group is unexpected as other studies have shown that the dominant contributions to reactivity come from this group (Gilman et al., 2015; Koss et al., 2018; Liang et al., 2022).

8 Conclusions

EFs were derived for a total of 15 trace gas species via measurements from a PTR-ToF-MS and an FTIR spectrometer, the resulting OH reactivity of the transported plume quantified, and O_3 and NO_2 time series investigated. The COALA-2020 ground-based field campaign opportunistically sampled a sustained biomass burning plume from 3–4 February 2020 during the 2019–2020 wildfire season in New South Wales, Australia. We determined via HYSPLIT trajectories that the most likely pathway traveled by the plume was from a distance ranging from $\sim 230\text{--}375 \text{ km}$ south from fires along the temperate forests of the east coast with contributions from more inland fires near Canberra, Australia. This plume lofted to an altitude of 500 m a.g.l. as it passed over active fires $\sim 8 \text{ h}$ out from the field site, before descending to 10 m a.g.l. while traveling over the ocean and reaching the site at 17:30 LT. All data used in the derivation of EFs were limited from sunset on 3 February to sunrise on 4 February as this period showed the greatest enhancements of reactive BB tracers like furan. Through visual inspection, we partitioned this plume event into four portions and calculated and averaged the individual ERs. We used two age marker ratios derived from furan radical oxidation to determine the freshest portion of the plume and found that ERs from this portion corresponded well with the averaged ERs (within 1σ). Using EFs from the entire evening allowed for the inclusion of three more VOC EFs into this analysis which, for the freshest portion of the plume, did not meet the selection criteria for ERs.

We have further characterized wildfire emissions in Australia's temperate region by providing a more comprehensive suite of biome-averaged VOC EFs. This suite introduces new EFs for acrolein, methyl propanoate, methyl methacrylate, maleic anhydride, benzaldehyde, and creosol. Of particular note is acrolein, which has been shown to be a gas-phase variable posing significant harm to human health (O'Dell et al., 2020; Simmons et al., 2022). When compared with values reported from two Australian studies located in the same or nearby temperate forests, we find mixed agreement with results from Guérette et al. (2018), as only two values are captured within our EF variability, with acetonitrile differing by a factor of ~ 4.5 and MVK + MACR differing by a factor of ~ 6 . However, two compounds are within the range




Figure 6. Selected compounds with significantly high smoke-related enhancements are grouped into categories of varying reactivity based on known reactivity groups, except for the “isoprene + monoterpenes” group, which is the sum of isoprene (m/z 69) and monoterpene (m/z 137) reactivities. This captures every compound included in this OH reactivity calculation.

of variability for Lawson et al. (2015) and two others are well within a factor of 2, which indicates reasonable agreement. Furthermore, comparison with two recent US studies that report data on analogous temperate zones, as well as one report covering global temperate regions, show generally good agreement for 9 of the 15 compounds, with several others within a factor of 2, indicating very good agreement. This closer agreement with these studies, as well as that of Lawson et al. (2015), is likely due to the measurement approach when deriving EFs as both US-based studies were aircraft campaigns, and the Australia-based study intercepted a transported plume much like this work. Guérette et al. (2018) sampled controlled burns on a ground campaign virtually at the emission source. This indicates that variability previously ascribed to differing fuel types may be overshadowed by sampling approach and that comprehensive measurements from US-based studies may be useful for studying Australian biomes. Agreement with both Lawson et al. (2015) and the US-based studies indicates that results here are valid for future use in Australian, biome-specific biomass burning studies. Compounding this is the excellent agreement found between EFs in this study and a comparison of two laboratory, US-based, temperate fuel studies, indicating the potential for lab-based results to be similarly applicable. Chemically comprehensive near-source observations of Australian fuel types are needed to evaluate the importance delineating temperate forest EFs in different regions across the globe.

Probing the OH reactivity of the plume revealed that the nighttime conditions, despite the long transport time, transported a quantity that effectively doubled OH reactivity at the COALA-2020 field site, with contributions arising from expected classes of compounds such as furans (most contribution), aromatics (second), and alkenes (third). m/z 85 contributed most significantly of the furans measured, which is

due to its long NO_3 lifetime but short OH lifetime. Other furans had largely been reacted away before reaching the COALA-2020 field site. Phenol had the largest contribution of the measurable phenolic compounds despite its high NO_3 reactivity. Alkenes and aromatics were found, as a group, to have an on par reactivity and, unexpectedly, non-aromatic oxygenates contributed the least.

Data availability. Data from time periods used for analysis in this work are available from the PANGAEA archive at <https://doi.org/10.1594/PANGAEA.927277> (Mouat et al., 2021a) and <https://doi.org/10.1594/PANGAEA.939407> (Mouat et al., 2021b).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-22-11033-2022-supplement>.

Author contributions. APM conducted PTR-ToF-MS measurements and subsequent data analysis. JBS, CPW, and JRG oversaw the maintenance and in-person operation of the PTR-ToF-MS for much of the COALA-2020 field campaign. CO measurements were provided by DWTG. CPW led the COALA-2020 campaign, whilst JK led PTR-ToF-MS instrument deployment and data analysis. All coauthors have provided substantial input during the process of drafting this work.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. We thank Travis Naylor, Ian Galbally and all the UOW COALA-2020 team for their aid in conducting measurements during the field campaign and all input thereafter. We additionally gratefully acknowledge the NOAA Air Resources Laboratory (ARL) for providing the HYSPLIT transport and dispersion model used for analysis in this publication. We acknowledge the use of data and/or imagery from NASA's Land, Atmosphere Near real-time Capability for EOS (LANCE) system (<https://earthdata.nasa.gov/lance>, last access: 11 March 2022), part of NASA's Earth Observing System Data and Information System (EOSDIS) (Simmons et al., 2022).

Financial support. This research has been supported by the National Science Foundation (grant no. 2016646).

Review statement. This paper was edited by Ivan Kourtchev and reviewed by two anonymous referees.

References

- Abatzoglou, J. T., Williams, A. P., and Barbero, R.: Global Emergence of Anthropogenic Climate Change in Fire Weather Indices, *Geophys. Res. Lett.*, 46, 326–336, <https://doi.org/10.1029/2018GL080959>, 2019.
- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- Akagi, S. K., Craven, J. S., Taylor, J. W., McMeeking, G. R., Yokelson, R. J., Burling, I. R., Urbanski, S. P., Wold, C. E., Seinfeld, J. H., Coe, H., Alvarado, M. J., and Weise, D. R.: Evolution of trace gases and particles emitted by a chaparral fire in California, *Atmos. Chem. Phys.*, 12, 1397–1421, <https://doi.org/10.5194/acp-12-1397-2012>, 2012.
- Alvarado, M. J., Logan, J. A., Mao, J., Apel, E., Riemer, D., Blake, D., Cohen, R. C., Min, K.-E., Perring, A. E., Browne, E. C., Wooldridge, P. J., Diskin, G. S., Sachse, G. W., Fuelberg, H., Sessions, W. R., Harrigan, D. L., Huey, G., Liao, J., Case-Hanks, A., Jimenez, J. L., Cubison, M. J., Vay, S. A., Weinheimer, A. J., Knapp, D. J., Montzka, D. D., Flocke, F. M., Pollack, I. B., Wennberg, P. O., Kurten, A., Crounse, J., Clair, J. M. St., Wisthaler, A., Mikoviny, T., Yantosca, R. M., Carouge, C. C., and Le Sager, P.: Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: an integrated analysis of aircraft and satellite observations, *Atmos. Chem. Phys.*, 10, 9739–9760, <https://doi.org/10.5194/acp-10-9739-2010>, 2010.
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 955–966, <https://doi.org/10.1029/2000GB001382>, 2001.
- Berndt, T., Böge, O., and Rolle, W.: Products of the Gas-Phase Reactions of NO₃ Radicals with Furan and Tetramethylfuran, *Environ. Sci. Technol.*, 31, 1157–1162, <https://doi.org/10.1021/es960669z>, 1997.
- Bierbach, A., Barnes, I., Becker, K. H., and Wiesen, E.: Atmospheric Chemistry of Unsaturated Carbonyls: Butenedial, 4-Oxo-2-pentenal, 3-Hexene-2,5-dione, Maleic Anhydride, 3/Furan-2-one, and 5-Methyl-3H-furan-2-one, *Environ. Sci. Technol.*, 28, 715–729, 1994.
- Bloss, C., Wagner, V., Jenkin, M. E., Volkamer, R., Bloss, W. J., Lee, J. D., Heard, D. E., Wirtz, K., Martin-Reviejo, M., Rea, G., Wenger, J. C., and Pilling, M. J.: Development of a detailed chemical mechanism (MCMv3.1) for the atmospheric oxidation of aromatic hydrocarbons, *Atmos. Chem. Phys.*, 5, 641–664, <https://doi.org/10.5194/acp-5-641-2005>, 2005.
- Borchers Arriagada, N., Palmer, A. J., Bowman, D. M., Morgan, G. G., Jalaludin, B. B., and Johnston, F. H.: Unprecedented smoke-related health burden associated with the 2019–20 bushfires in eastern Australia, *Med. J. Australia*, 213, 282–283, 2020.
- Brey, S. J. and Fischer, E. V.: Smoke in the city: How often and where does smoke impact summertime ozone in the United States?, *Environ. Sci. Technol.*, 50, 1288–1294, 2016.
- Brilli, F., Gioli, B., Ciccioli, P., Zona, D., Loreto, F., Janssens, I. A., and Ceulemans, R.: Proton Transfer Reaction Time-of-Flight Mass Spectrometric (PTR-TOF-MS) determination of volatile organic compounds (VOCs) emitted from a biomass fire developed under stable nocturnal conditions, *Atmos. Environ.*, 97, 54–67, <https://doi.org/10.1016/j.atmosenv.2014.08.007>, 2014.
- Bruns, E. A., Slowik, J. G., El Haddad, I., Kilic, D., Klein, F., Dommen, J., Temime-Roussel, B., Marchand, N., Baltensperger, U., and Prévôt, A. S. H.: Characterization of gas-phase organics using proton transfer reaction time-of-flight mass spectrometry: fresh and aged residential wood combustion emissions, *Atmos. Chem. Phys.*, 17, 705–720, <https://doi.org/10.5194/acp-17-705-2017>, 2017.
- Burling, I. R., Yokelson, R. J., Akagi, S. K., Urbanski, S. P., Wold, C. E., Griffith, D. W. T., Johnson, T. J., Reardon, J., and Weise, D. R.: Airborne and ground-based measurements of the trace gases and particles emitted by prescribed fires in the United States, *Atmos. Chem. Phys.*, 11, 12197–12216, <https://doi.org/10.5194/acp-11-12197-2011>, 2011.
- Coggon, M. M., Veres, P. R., Yuan, B., Koss, A., Warneke, C., Gilman, J. B., Lerner, B. M., Peischl, J., Aikin, K. C., Stockwell, C. E., Hatch, L. E., Ryerson, T. B., Roberts, J. M., Yokelson, R. J., and de Gouw, J. A.: Emissions of nitrogen-containing organic compounds from the burning of herbaceous and arboreal biomass: Fuel composition dependence and the variability of commonly used nitrile tracers, *Geophys. Res. Lett.*, 43, 9903–9912, <https://doi.org/10.1002/2016GL070562>, 2016.
- Coggon, M. M., Lim, C. Y., Koss, A. R., Sekimoto, K., Yuan, B., Gilman, J. B., Hagan, D. H., Selimovic, V., Zarzana, K. J., Brown, S. S., Roberts, J. M., Müller, M., Yokelson, R., Wisthaler, A., Krechmer, J. E., Jimenez, J. L., Cappa, C., Kroll, J. H., de Gouw, J., and Warneke, C.: OH chemistry of non-methane organic gases (NMOGs) emitted from laboratory and ambient biomass burning smoke: evaluating the influence of furans and oxygenated aromatics on ozone and secondary NMOG formation, *Atmos. Chem. Phys.*, 19, 14875–14899, <https://doi.org/10.5194/acp-19-14875-2019>, 2019.
- Davey, S. M. and Sarre, A.: Editorial: the 2019/20 Black Summer bushfires, *Aust. Forestry*, 83, 47–51, <https://doi.org/10.1080/00049158.2020.1769899>, 2020.

- Decker, Z. C. J., Zarzana, K. J., Coggon, M., Min, K.-E., Pollock, I., Ryerson, T. B., Peischl, J., Edwards, P., Dubé, W. P., Markovic, M. Z., Roberts, J. M., Veres, P. R., Graus, M., Warneke, C., de Gouw, J., Hatch, L. E., Barsanti, K. C., and Brown, S. S.: Nighttime Chemical Transformation in Biomass Burning Plumes: A Box Model Analysis Initialized with Aircraft Observations, *Environ. Sci. Technol.*, 53, 2529–2538, <https://doi.org/10.1021/acs.est.8b05359>, 2019.
- Decker, Z. C. J., Robinson, M. A., Barsanti, K. C., Bourgeois, I., Coggon, M. M., DiGangi, J. P., Diskin, G. S., Flocke, F. M., Franchin, A., Fredrickson, C. D., Gkatzelis, G. I., Hall, S. R., Halliday, H., Holmes, C. D., Huey, L. G., Lee, Y. R., Lindaas, J., Middlebrook, A. M., Montzka, D. D., Moore, R., Neuman, J. A., Nowak, J. B., Palm, B. B., Peischl, J., Piel, F., Rickly, P. S., Rollins, A. W., Ryerson, T. B., Schwantes, R. H., Sekimoto, K., Thornhill, L., Thornton, J. A., Tyndall, G. S., Ullmann, K., Van Rooy, P., Veres, P. R., Warneke, C., Washenfelder, R. A., Weinheimer, A. J., Wiggins, E., Winstead, E., Wisthaler, A., Womack, C., and Brown, S. S.: Nighttime and daytime dark oxidation chemistry in wildfire plumes: an observation and model analysis of FIREX-AQ aircraft data, *Atmos. Chem. Phys.*, 21, 16293–16317, <https://doi.org/10.5194/acp-21-16293-2021>, 2021.
- de Gouw, J. A., Warneke, C., Stohl, A., Wollny, A. G., Brock, C. A., Cooper, O. R., Holloway, J. S., Trainer, M., Fehsenfeld, F. C., Atlas, E. L., Donnelly, S. G., Stroud, V., and Lueb, A.: Volatile organic compounds composition of merged and aged forest fire plumes from Alaska and western Canada, *J. Geophys. Res.-Atmos.*, 111, D10303, <https://doi.org/10.1029/2005JD006175>, 2006.
- Didan, K.: MODIS/Terra Vegetation Indices 16-Day L3 Global 250 m SIN Grid V061 [dataset], <https://doi.org/10.5067/MODIS/MOD13Q1.061>, 2021.
- Donovan, V. M., Wonkka, C. L., and Twidwell, D.: Surging wildfire activity in a grassland biome, *Geophys. Res. Lett.*, 44, 5986–5993, 2017.
- Fairman, T. A., Nitschke, C. R., and Bennett, L. T.: Too much, too soon? A review of the effects of increasing wildfire frequency on tree mortality and regeneration in temperate eucalypt forests, *Int. J. Wildland Fire*, 25, 831–848, 2015.
- Filkov, A. I., Ngo, T., Matthews, S., Telfer, S., and Penman, T. D.: Impact of Australia's catastrophic 2019/20 bushfire season on communities and environment. Retrospective analysis and current trends, *Journal of Safety Science and Resilience*, 1, 44–56, <https://doi.org/10.1016/j.jnlsr.2020.06.009>, 2020.
- Ford, B., Val Martin, M., Zelasky, S., Fischer, E., Anenberg, S., Heald, C. L., and Pierce, J.: Future fire impacts on smoke concentrations, visibility, and health in the contiguous United States, *GeoHealth*, 2, 229–247, 2018.
- Gilman, J. B., Lerner, B. M., Kuster, W. C., Goldan, P. D., Warneke, C., Veres, P. R., Roberts, J. M., de Gouw, J. A., Burling, I. R., and Yokelson, R. J.: Biomass burning emissions and potential air quality impacts of volatile organic compounds and other trace gases from fuels common in the US, *Atmos. Chem. Phys.*, 15, 13915–13938, <https://doi.org/10.5194/acp-15-13915-2015>, 2015.
- Gkatzelis, G., Coggon, M. M., Sekimoto, K., Gilman, J., Lamplugh, A., Bourgeois, I., Peischl, J., Ryerson, T. B., Veres, P. R., Neuman, J. A., Womack, C., Brown, S. S., Rollins, A. W., Rickly, P., Bela, M., Schwantes, R., Katich, J. M., Lindaas, J., Jimenez, J. L., Campuzano Jost, P., Guo, H., Nault, B. A., Pagonis, D., Schueneman, M., Day, D. A., Wisthaler, A., Piel, F., Tomsche, L., Mikoviny, T., Hair, J. W., Shingler, T. J., Fenn, M. A., Selimovic, V., Huey, L. G., Ji, Y., Lee, Y. R., Tanner, D., Nowak, J. B., DiGangi, J. P., Halliday, H. S., Diskin, G. S., Fried, A., Weibring, P., Wolfe, G. M., St Clair, J. M., Hannun, R. A., Liao, J., Hanisco, T. F., Travis, K., Roberts, J., Trainer, M., Schwarz, J. P., Crawford, J. H., and Warneke, C.: Non-methane organic and nitrogen emissions from wildfire plumes during FIREX-AQ, AGU Fall Meeting, 1 December 2020, Online, 2020AGUFMA224.0013G, 2020.
- Gregory, R. W., Yayne-abeba, A., Matthew, S. L., and Yu-Mei, H.: Impacts of a large boreal wildfire on ground level atmospheric concentrations of PAHs, VOCs and ozone, *Atmos. Environ.*, 178, 19–30, <https://doi.org/10.1016/j.atmosenv.2018.01.013>, 2018.
- Griffith, D. W. T., Deutscher, N. M., Caldow, C., Kettlewell, G., Riggenbach, M., and Hammer, S.: A Fourier transform infrared trace gas and isotope analyser for atmospheric applications, *Atmos. Meas. Tech.*, 5, 2481–2498, <https://doi.org/10.5194/amt-5-2481-2012>, 2012.
- Grosjean, D. and Williams, E. L.: Environmental persistence of organic compounds estimated from structure-reactivity and linear free-energy relationships. Unsaturated aliphatics, *Atmos. Environ. A-Gen.*, 26, 1395–1405, [https://doi.org/10.1016/0960-1686\(92\)90124-4](https://doi.org/10.1016/0960-1686(92)90124-4), 1992.
- Guérette, E.-A., Paton-Walsh, C., Desservettaz, M., Smith, T. E. L., Volkova, L., Weston, C. J., and Meyer, C. P.: Emissions of trace gases from Australian temperate forest fires: emission factors and dependence on modified combustion efficiency, *Atmos. Chem. Phys.*, 18, 3717–3735, <https://doi.org/10.5194/acp-18-3717-2018>, 2018.
- Hatch, L. E., Luo, W., Pankow, J. F., Yokelson, R. J., Stockwell, C. E., and Barsanti, K. C.: Identification and quantification of gaseous organic compounds emitted from biomass burning using two-dimensional gas chromatography–time-of-flight mass spectrometry, *Atmos. Chem. Phys.*, 15, 1865–1899, <https://doi.org/10.5194/acp-15-1865-2015>, 2015.
- Hatch, L. E., Yokelson, R. J., Stockwell, C. E., Veres, P. R., Simpson, I. J., Blake, D. R., Orlando, J. J., and Barsanti, K. C.: Multi-instrument comparison and compilation of non-methane organic gas emissions from biomass burning and implications for smoke-derived secondary organic aerosol precursors, *Atmos. Chem. Phys.*, 17, 1471–1489, <https://doi.org/10.5194/acp-17-1471-2017>, 2017.
- Ito, A. and Penner, J. E.: Global estimates of biomass burning emissions based on satellite imagery for the year 2000, *J. Geophys. Res.-Atmos.*, 109, D14S05, <https://doi.org/10.1029/2003JD004423>, 2004.
- Jaffe, D. A. and Wigder, N. L.: Ozone production from wildfires: A critical review, *Atmos. Environ.*, 51, 1–10, 2012.
- Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31, 81–104, [https://doi.org/10.1016/S1352-2310\(96\)00105-7](https://doi.org/10.1016/S1352-2310(96)00105-7), 1997.
- Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 181–193, <https://doi.org/10.5194/acp-3-181-2003>, 2003.

- Keywood, M., Kanakidou, M., Stohl, A., Dentener, F., Grassi, G., Meyer, C. P., Torseth, K., Edwards, D., Thompson, A. M., Lohmann, U., and Burrows, J.: Fire in the Air: Biomass Burning Impacts in a Changing Climate, *Crit. Rev. Env. Sci. Tec.*, 43, 40–83, <https://doi.org/10.1080/10643389.2011.604248>, 2013.
- Kodros, J. K., Papanastasiou, D. K., Paglione, M., Masiol, M., Squizzato, S., Florou, K., Skyllakou, K., Katsionoudis, C., Nenes, A., and Pandis, S. N.: Rapid dark aging of biomass burning as an overlooked source of oxidized organic aerosol, *P. Natl. Acad. Sci. USA*, 117, 33028, <https://doi.org/10.1073/pnas.2010365117>, 2020.
- Koss, A. R., Sekimoto, K., Gilman, J. B., Selimovic, V., Coggon, M. M., Zarzana, K. J., Yuan, B., Lerner, B. M., Brown, S. S., Jimenez, J., Krechmer, J., Roberts, J. M., Warneke, C., Yokelson, R. J., and de Gouw, J.: Non-methane organic gas emissions from biomass burning: identification, quantification, and emission factors from PTR-ToF during the FIREX 2016 laboratory experiment, *Atmos. Chem. Phys.*, 18, 3299–3319, <https://doi.org/10.5194/acp-18-3299-2018>, 2018.
- Lawson, S. J., Keywood, M. D., Galbally, I. E., Gras, J. L., Caine, J. M., Cope, M. E., Krummel, P. B., Fraser, P. J., Steele, L. P., Bentley, S. T., Meyer, C. P., Ristovski, Z., and Goldstein, A. H.: Biomass burning emissions of trace gases and particles in marine air at Cape Grim, Tasmania, *Atmos. Chem. Phys.*, 15, 13393–13411, <https://doi.org/10.5194/acp-15-13393-2015>, 2015.
- Lawson, S. J., Cope, M., Lee, S., Galbally, I. E., Ristovski, Z., and Keywood, M. D.: Biomass burning at Cape Grim: exploring photochemistry using multi-scale modelling, *Atmos. Chem. Phys.*, 17, 11707–11726, <https://doi.org/10.5194/acp-17-11707-2017>, 2017.
- Liang, Y., Weber, R. J., Misztal, P. K., Jen, C. N., and Goldstein, A. H.: Aging of Volatile Organic Compounds in October 2017 Northern California Wildfire Plumes, *Environ. Sci. Technol.*, 56, 1557–1567, <https://doi.org/10.1021/acs.est.1c05684>, 2022.
- Liu, X., Zhang, Y., Huey, L. G., Yokelson, R. J., Wang, Y., Jimenez, J. L., Campuzano-Jost, P., Beyersdorf, A. J., Blake, D. R., Choi, Y., St. Clair, J. M., Crounse, J. D., Day, D. A., Diskin, G. S., Fried, A., Hall, S. R., Hanisco, T. F., King, L. E., Meinardi, S., Mikoviny, T., Palm, B. B., Peischl, J., Perring, A. E., Pollack, I. B., Ryerson, T. B., Sachse, G., Schwarz, J. P., Simpson, I. J., Tanner, D. J., Thornhill, K. L., Ullmann, K., Weber, R. J., Wennberg, P. O., Wisthaler, A., Wolfe, G. M., and Ziemba, L. D.: Agricultural fires in the southeastern U.S. during SEAC4RS: Emissions of trace gases and particles and evolution of ozone, reactive nitrogen, and organic aerosol, *J. Geophys. Res.-Atmos.*, 121, 7383–7414, <https://doi.org/10.1002/2016JD025040>, 2016.
- Liu, X., Huey, L. G., Yokelson, R. J., Selimovic, V., Simpson, I. J., Müller, M., Jimenez, J. L., Campuzano-Jost, P., Beyersdorf, A. J., Blake, D. R., Butterfield, Z., Choi, Y., Crounse, J. D., Day, D. A., Diskin, G. S., Dubey, M. K., Fortner, E., Hanisco, T. F., Hu, W., King, L. E., Kleinman, L., Meinardi, S., Mikoviny, T., Onasch, T. B., Palm, B. B., Peischl, J., Pollack, I. B., Ryerson, T. B., Sachse, G. W., Sedlacek, A. J., Shilling, J. E., Springston, S., St. Clair, J. M., Tanner, D. J., Teng, A. P., Wennberg, P. O., Wisthaler, A., and Wolfe, G. M.: Airborne measurements of western U.S. wildfire emissions: Comparison with prescribed burning and air quality implications, *J. Geophys. Res.-Atmos.*, 122, 6108–6129, <https://doi.org/10.1002/2016JD026315>, 2017.
- MacSween, K., Paton-Walsh, C., Roulston, C., Guérette, E.-A., Edwards, G., Reisen, F., Desservettaz, M., Cameron, M., Young, E., and Kubistin, D.: Cumulative firefighter exposure to multiple toxins emitted during prescribed burns in Australia, *Expos. Health*, 12, 721–733, 2020.
- Manion, J. A., Huie, R. E., Levin, R. D., Burgess Jr., D. R., Orkin, V. L., Tsang, W., McGivern, W. S., Hudgens, J. W., Knyazev, V. D., Atkinson, D. B., Chai, E., Tereza, A. M., Lin, C. Y., Allison, T. C., Mallard, W. G., Westly, F., Herron, J. T., Hampson, R. F., and Frizzell, D. H.: NIST Chemical Kinetics Database (2015.09), NIST [data set], <https://kinetics.nist.gov/kinetics/index.jsp> (last access: 11 August 2022), 2015.
- Mouat, A. P., Kaiser, J., Paton-Walsh, C., Ramirez-Gamboa, J., Naylor, T. A., and Simmons, J. B.: Volatile organic compound measurements at Cataract Scout Park, Australia, taken during the COALA-2020 campaign, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.927277>, 2021a.
- Mouat, A. P., Kaiser, J., Paton-Walsh, C., Ramirez-Gamboa, J., Naylor, T. A., and Simmons, J. B.: Additional measurements of volatile organic compounds by PTR-ToF-MS at Cataract Scout Park, Australia, taken during the COALA-2020 campaign, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.939407>, 2021b.
- Müller, M., Anderson, B. E., Beyersdorf, A. J., Crawford, J. H., Diskin, G. S., Eichler, P., Fried, A., Keutsch, F. N., Mikoviny, T., Thornhill, K. L., Walega, J. G., Weinheimer, A. J., Yang, M., Yokelson, R. J., and Wisthaler, A.: In situ measurements and modeling of reactive trace gases in a small biomass burning plume, *Atmos. Chem. Phys.*, 16, 3813–3824, <https://doi.org/10.5194/acp-16-3813-2016>, 2016.
- O'Dell, K., Hornbrook, R. S., Permar, W., Levin, E. J. T., Garofalo, L. A., Apel, E. C., Blake, N. J., Jarnot, A., Pothier, M. A., Farmer, D. K., Hu, L., Campos, T., Ford, B., Pierce, J. R., and Fischer, E. V.: Hazardous Air Pollutants in Fresh and Aged Western US Wildfire Smoke and Implications for Long-Term Exposure, *Environ. Sci. Technol.*, 54, 11838–11847, <https://doi.org/10.1021/acs.est.0c04497>, 2020.
- Pagonis, D., Sekimoto, K., and de Gouw, J.: A Library of Proton-Transfer Reactions of H₃O⁺ Ions Used for Trace Gas Detection, *J. Am. Soc. Mass Spectr.*, 30, 1330–1335, <https://doi.org/10.1007/s13361-019-02209-3>, 2019.
- Palm, B. B., Peng, Q., Fredrickson, C. D., Lee, B. H., Garofalo, L. A., Pothier, M. A., Kreidenweis, S. M., Farmer, D. K., Pokhrel, R. P., and Shen, Y.: Quantification of organic aerosol and brown carbon evolution in fresh wildfire plumes, *P. Natl. Acad. Sci. USA*, 117, 29469–29477, 2020.
- Paton-Walsh, C., Smith, T. E. L., Young, E. L., Griffith, D. W. T., and Guérette, É.-A.: New emission factors for Australian vegetation fires measured using open-path Fourier transform infrared spectroscopy – Part 1: Methods and Australian temperate forest fires, *Atmos. Chem. Phys.*, 14, 11313–11333, <https://doi.org/10.5194/acp-14-11313-2014>, 2014.
- Permar, W., Wang, Q., Selimovic, V., Wielgasz, C., Yokelson, R. J., Hornbrook, R. S., Hills, A. J., Apel, E. C., Ku, I.-T., Zhou, Y., Sive, B. C., Sullivan, A. P., Collett Jr, J. L., Campos, T. L., Palm, B. B., Peng, Q., Thornton, J. A., Garofalo, L. A., Farmer, D. K., Kreidenweis, S. M., Levin, E. J. T., DeMott, P. J., Flocke, F., Fischer, E. V., and Hu, L.: Emissions of Trace Organic Gases From Western U.S. Wildfires Based on WE-CAN Aircraft

- Measurements, *J. Geophys. Res.-Atmos.*, 126, e2020JD033838, <https://doi.org/10.1029/2020JD033838>, 2021.
- Robinson, M. A., Decker, Z. C., Barsanti, K. C., Coggon, M. M., Flocke, F. M., Franchin, A., Fredrickson, C. D., Gilman, J. B., Gkatzelis, G. I., and Holmes, C. D.: Variability and time of day dependence of ozone photochemistry in western wildfire plumes, *Environ. Sci. Technol.*, 55, 10280–10290, 2021.
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, <https://doi.org/10.5194/acp-3-161-2003>, 2003.
- Sekimoto, K., Li, S.-M., Yuan, B., Koss, A., Coggon, M., Warneke, C., and de Gouw, J.: Calculation of the sensitivity of proton-transfer-reaction mass spectrometry (PTR-MS) for organic trace gases using molecular properties, *Int. J. Mass Spectrom.*, 421, 71–94, <https://doi.org/10.1016/j.ijms.2017.04.006>, 2017.
- Sekimoto, K., Koss, A. R., Gilman, J. B., Selimovic, V., Coggon, M. M., Zarzana, K. J., Yuan, B., Lerner, B. M., Brown, S. S., Warneke, C., Yokelson, R. J., Roberts, J. M., and de Gouw, J.: High- and low-temperature pyrolysis profiles describe volatile organic compound emissions from western US wildfire fuels, *Atmos. Chem. Phys.*, 18, 9263–9281, <https://doi.org/10.5194/acp-18-9263-2018>, 2018.
- Selimovic, V., Yokelson, R. J., Warneke, C., Roberts, J. M., de Gouw, J., Reardon, J., and Griffith, D. W. T.: Aerosol optical properties and trace gas emissions by PAX and OP-FTIR for laboratory-simulated western US wildfires during FIREX, *Atmos. Chem. Phys.*, 18, 2929–2948, <https://doi.org/10.5194/acp-18-2929-2018>, 2018.
- Simmons, J. B., Paton-Walsh, C., Mouat, A. P., Kaiser, J., Humphries, R. S., Keyword, M., Griffith, D. W. T., Sutresna, A., Naylor, T., and Ramirez-Gamboa, J.: Bushfire smoke plume composition and toxicological assessment from the 2019–2020 Australian Black Summer, *Air Qual. Atmos. Hlth.*, accepted, 2022.
- Sokolik, I., Soja, A., DeMott, P., and Winker, D.: Progress and challenges in quantifying wildfire smoke emissions, their properties, transport, and atmospheric impacts, *J. Geophys. Res.-Atmos.*, 124, 13005–13025, 2019.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *B. Am. Meteorol. Soc.*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- Stockwell, C. E., Veres, P. R., Williams, J., and Yokelson, R. J.: Characterization of biomass burning emissions from cooking fires, peat, crop residue, and other fuels with high-resolution proton-transfer-reaction time-of-flight mass spectrometry, *Atmos. Chem. Phys.*, 15, 845–865, <https://doi.org/10.5194/acp-15-845-2015>, 2015.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.
- Verma, S., Worden, J., Pierce, B., Jones, D. B. A., Al-Saadi, J., Boersma, F., Bowman, K., Eldering, A., Fisher, B., Jourdain, L., Kulawik, S., and Worden, H.: Ozone production in boreal fire smoke plumes using observations from the Tropospheric Emission Spectrometer and the Ozone Monitoring Instrument, *J. Geophys. Res.-Atmos.*, 114, D02303, <https://doi.org/10.1029/2008JD010108>, 2009.
- Wu, C. and Yu, J. Z.: Evaluation of linear regression techniques for atmospheric applications: the importance of appropriate weighting, *Atmos. Meas. Tech.*, 11, 1233–1250, <https://doi.org/10.5194/amt-11-1233-2018>, 2018.
- Xu, L., Crouse, J. D., Vasquez, K. T., Allen, H., Wennberg, P. O., Bourgeois, I., Brown, S. S., Campuzano-Jost, P., Coggon, M. M., Crawford, J. H., DiGangi, J. P., Diskin, G. S., Fried, A., Gargulinski, E. M., Gilman, J. B., Gkatzelis, G. I., Guo, H., Hair, J. W., Hall, S. R., Halliday, H. A., Hanisco, T. F., Hannun, R. A., Holmes, C. D., Huey, L. G., Jimenez, J. L., Lamplugh, A., Lee, Y. R., Liao, J., Lindaas, J., Neuman, J. A., Nowak, J. B., Peischl, J., Peterson, D. A., Piel, F., Richter, D., Rickly, P. S., Robinson, M. A., Rollins, A. W., Ryerson, T. B., Sekimoto, K., Selimovic, V., Shingler, T., Soja, A. J., Clair, J. M. S., Tanner, D. J., Ullmann, K., Veres, P. R., Walega, J., Warneke, C., Washenfelder, R. A., Weibring, P., Wisthaler, A., Wolfe, G. M., Womack, C. C., and Yokelson, R. J.: Ozone chemistry in western U.S. wildfire plumes, *Sci. Adv.*, 7, eabl3648, <https://doi.org/10.1126/sciadv.abl3648>, 2021.
- Yokelson, R. J., Christian, T. J., Karl, T. G., and Guenther, A.: The tropical forest and fire emissions experiment: laboratory fire measurements and synthesis of campaign data, *Atmos. Chem. Phys.*, 8, 3509–3527, <https://doi.org/10.5194/acp-8-3509-2008>, 2008.
- Young, P. J., Naik, V., Fiore, A. M., Gaudel, A., Guo, J., Lin, M. Y., Neu, J. L., Parrish, D. D., Rieder, H. E., Schnell, J. L., Tilmes, S., Wild, O., Zhang, L., Ziemke, J., Brandt, J., Delcloo, A., Doherty, R. M., Geels, C., Hegglin, M. I., Hu, L., Im, U., Kumar, R., Luhar, A., Murray, L., Plummer, D., Rodriguez, J., Saiz-Lopez, A., Schultz, M. G., Woodhouse, M. T., and Zeng, G.: Tropospheric Ozone Assessment Report: Assessment of global-scale model performance for global and regional ozone distributions, variability, and trends, *Elementa*, 6, 10, <https://doi.org/10.1525/elementa.265>, 2018.