



Ice microphysical processes in the dendritic growth layer: a statistical analysis combining multi-frequency and polarimetric Doppler cloud radar observations

Leonie von Terzi¹, José Dias Neto², Davide Ori¹, Alexander Myagkov³, and Stefan Kneifel^{1,4}

¹Institute of Geophysics and Meteorology, University of Cologne, Cologne, Germany

²Department of Geosciences and Remote Sensing, Delft University of Technology, Delft, the Netherlands

³Radiometer Physics GmbH, Meckenheim, Germany

⁴Meteorological Institute, Ludwig-Maximilians University Munich, Germany

Correspondence: Leonie von Terzi (lterzi@uni-koeln.de)

Received: 7 April 2022 – Discussion started: 11 April 2022

Revised: 24 August 2022 – Accepted: 25 August 2022 – Published: 13 September 2022

Abstract. The dendritic growth layer (DGL), defined as the temperature region between -20 and -10 °C, plays an important role for ice depositional growth, aggregation and potentially secondary ice processes. The DGL has been found in the past to exhibit specific observational signatures in polarimetric and vertically pointing radar observations. However, consistent conclusions about their physical interpretation have often not been reached.

In this study, we exploit a unique 3-months dataset of mid-latitude winter clouds observed with vertically pointing triple-frequency (X-, Ka-, W-band) and polarimetric W-band Doppler radars. In addition to standard radar moments, we also analyse the multi-wavelength and polarimetric Doppler spectra. New variables, such as the maximum of the spectral differential reflectivity (ZDR) ($sZDR_{max}$), allows us to analyse the ZDR signal of asymmetric ice particles independent of the presence of low ZDR producing aggregates. This unique dataset enables us to investigate correlations between enhanced aggregation and evolution of small ice particles in the DGL. For this, the multi-frequency observations are used to classify all profiles according to their maximum average aggregate size within the DGL. The strong correlation between aggregate class and specific differential phase shift (KDP) confirms the expected link between ice particle concentration and aggregation. Interestingly, no correlation between aggregation class and $sZDR_{max}$ is visible. This indicates that aggregation is rather independent of the aspect ratio and density of ice crystals. A distinct reduction of mean Doppler velocity in the DGL is found to be strongest for cases with largest aggregate sizes. Analyses of spectral edge velocities suggest that the reduction is the combined result of the formation of new ice particles with low fall velocity and a weak updraft. It appears most likely that this updraft is the result of latent heat released by enhanced depositional growth. Clearly, the strongest correlations of aggregate class with other variables are found inside the DGL. Surprisingly, no correlation between aggregate class and concentration or aspect ratio of particles falling from above into the DGL could be found. Only a weak correlation between the mean particle size falling into the DGL and maximum aggregate size within the DGL is apparent. In addition to the correlation analysis, the dataset also allows study of the evolution of radar variables as a function of temperature. We find the ice particle concentration continuously increasing from -18 °C towards the bottom of the DGL. Aggregation increases more rapidly from -15 °C towards warmer temperatures. Surprisingly, KDP and $sZDR_{max}$ are not reduced by the intensifying aggregation below -15 °C but rather reach their maximum values in the lower half of the DGL. Also below the DGL, KDP and $sZDR_{max}$ remain enhanced until -4 °C. Only there, additional aggregation appears to deplete ice crystals and therefore reduce KDP and $sZDR_{max}$.

The simultaneous increase of aggregation and particle concentration inside the DGL necessitates a source mechanism for new ice crystals. As primary ice nucleation is expected to decrease towards warmer temperatures,

secondary ice processes are a likely explanation for the increase in ice particle concentration. Previous laboratory experiments strongly point towards ice collisional fragmentation as a possible mechanism for new particle generation. The presence of an updraft in the temperature region of maximum depositional growth might also suggest an important positive feedback mechanism between ice microphysics and dynamics which might further enhance ice particle growth in the DGL.

1 Introduction

Recent space-borne analyses underline once more the importance of understanding ice growth processes in clouds as more than 70 % of global precipitation is found to be generated via the ice phase (Heymsfield et al., 2020; Field and Heymsfield, 2015; Mülmenstädt et al., 2015). The dendritic growth layer (DGL), usually located between -20 and -10 °C, is known to play an important role for the growth and evolution of ice and snow in clouds. The reasons for its importance are manifold: the difference in saturation vapour pressure between ice and liquid reaches a maximum at -12 °C which favours depositional growth by the Wegener–Bergeron–Findeisen (WBF) process (Korolev, 2007). In addition, the particular plate-like shapes that particles grow into in the DGL lead to a distinct maximum in the depositional growth rate at -15 °C, where dendritic particles can exceed 1.5 mm in size within 10 min growth time at liquid water saturation (Takahashi, 2014). The fragile structure of those particles has also been found by airborne in situ observations (Schwarzenboeck et al., 2009; Hobbs and Rangno, 1998, 1990, 1985) and laboratory experiments (Takahashi et al., 1995; Griggs and Choulaton, 1986; Vardiman, 1978) to favour collisional breakup. This secondary ice process (SIP) gains increasing attention by the scientific community (e.g. Georgakaki et al., 2022; Phillips et al., 2018) as an important process which could explain the discrepancy between number of ice nucleating particles (INPs) and ice particle number concentration (IPNC) (Kanji et al., 2017). Unlike for example the Hallett–Mossop rime splintering process (Field et al., 2017; Hallett and Mossop, 1974), ice collisional fragmentation could provide new secondary ice particles over a wide temperature range. Finally, ice particles which are multiplied in number by SIP and grow rapidly in the DGL have been found to also aggregate very efficiently in the DGL (Lamb and Verlinde, 2011). Especially the branched structure of dendrites appears to be responsible for enhancing the stickiness of the crystals and thus favouring the formation of aggregates (Connolly et al., 2012). The various ice growth signatures in the DGL which can be observed for example with radars could be linked to the intensity of surface precipitation (Trömel et al., 2019, and references therein). This highlights again the importance to properly understand the interplay of microphysical processes in the DGL in order to accurately model surface precipitation.

Dual-polarisation radar observations are a powerful tool to observe the result of several of the aforementioned growth processes in the DGL. The abundance of plate-like particles in the DGL leads to vertically distinct layers of enhanced differential reflectivity (ZDR) and propagational differential phase shift (KDP) close to the -15 °C region (Griffin et al., 2018; Schrom and Kumjian, 2016; Schrom et al., 2015; Moisseev et al., 2015; Thompson et al., 2014; Bechini et al., 2013; Andrić et al., 2013; Kennedy and Rutledge, 2011; Trapp et al., 2001, among others). Interestingly, the ZDR layer appears at slightly higher altitudes as the KDP enhancement. ZDR is independent of the particle concentration but increases strongly with particle density and aspect ratio (Kumjian, 2013). However, the formation of aggregates leads to a decrease of ZDR despite the presence of asymmetric crystals which explains its layered structure. KDP is strongly related to particle concentration and, in contrast to ZDR, is not reflectivity-weighted, and thus not strongly influenced by the presence of large aggregates. While the general connection of ZDR and KDP layers, and intensive plate-like growth and subsequent aggregation is widely accepted, a definite conclusion on the reasons for the distinct vertical structure has not yet been reached (e.g. Schrom and Kumjian, 2016; Schrom et al., 2015; Moisseev et al., 2015).

Polarimetric information from the DGL is mostly lacking in vertically pointing radar observations. Similar to slant measuring radars, vertically pointing radars commonly observe a rapid increase in the radar reflectivity factor (Z_e , henceforth called reflectivity) in the DGL, in particular at -15 °C (e.g. Schrom and Kumjian, 2016; Zawadzki, 2013). The Doppler spectra collected with zenith-pointing radars revealed two distinct features in the DGL: first, the Doppler spectra often reveal an additional slow secondary mode in the DGL. In case of low turbulence and weak vertical air motions, the Doppler velocities can be related to the particles' terminal fall velocity. Apparently, the second spectral mode indicates the formation of new ice particles that often increase in fall speed and eventually merge with the main aggregate mode falling from higher altitudes. Second, the mean Doppler velocity (MDV) often reveals a slight but temporally very persistent reduction in the DGL. Various explanations for those features have been presented in the literature (see for example discussion presented in Schrom and Kumjian, 2016; Schrom et al., 2015). Zawadzki (2013) argues that vertical air motion at -15 °C is necessary to enhance supersaturation which enables the nucleation and subsequent growth

of plate-like particles. Other authors assign the MDV reduction simply to the evolution of a new secondary mode in the spectrum (e.g. Oue et al., 2018; Moisseev et al., 2015, 2009; Shupe et al., 2004; Zawadzki et al., 2001; Field, 2000, and references therein). The explanations for the origin of a new, slow ice particle mode include sedimentation of ice particles into the DGL from higher altitudes (Moisseev et al., 2015), enhanced primary nucleation due to upward air motion (Zawadzki, 2013) and secondary ice particle formation (e.g. Kennedy and Rutledge, 2011). Also, buoyancy-driven upward motion due to latent heat release of rapidly growing ice particles by water vapour deposition has been discussed as a potential reason for the decrease of MDV (Schrom and Kumjian, 2016).

An increasing number of ground-based sites are equipped with polarimetric and multi-frequency cloud radars. The use of higher frequencies does not only substantially increase the backscattered signal, especially of small ice particles scattering in the Rayleigh regime, but it also reduces observational limitations in some polarimetric variables. KDP is the range derivative of the differential phase shift and generally affected by high measurement noise. As KDP is inversely proportional to the wavelength, it can be more reliably estimated at shorter wavelength, also for small concentrations of asymmetric ice particles (Bringi et al., 2001). Moreover, even particles whose reflectivity values are below the radar detection level will cause some differential phase shift. Hence, KDP is sensitive also to the presence of extremely small, asymmetric ice crystals such as those expected to be produced by SIP (secondary ice process). The elevation-dependence of polarimetric cloud radar observations allows to infer shape, orientation and apparent density of ice crystals (e.g. Myagkov et al., 2016a; Matrosov et al., 2012). In addition, most polarimetric cloud radars provide polarimetric Doppler spectra which allow assigning of the polarimetric signatures to specific Doppler velocities. Most previous studies used linear depolarisation ratio (LDR) spectra from vertically pointing radar observations to investigate the evolution of columnar and needle particles between -10 and 0°C (e.g. Giangrande et al., 2016; Oue et al., 2015). Spectral ZDR from an S-band radar system has been used by Spek et al. (2008) to retrieve particle size distribution of aggregates and plates. Pfitzenmaier et al. (2018) analysed spectral LDR from a zenith-pointing Ka-band and spectral ZDR from a slant-viewing S-band radar to study ice particle growth along fall streaks.

Aggregation in the DGL can be only indirectly detected by radar polarimetry as a reduction of for example ZDR and concurrent increase of radar reflectivity (Kumjian, 2013). In contrast, the increase in mean particle size can be well observed as an increasing reflectivity difference in multi-frequency cloud radar observations (e.g. Kneifel et al., 2011; Liao et al., 2008, 2005; Matrosov, 1992). Ice particles increasing in size begin to scatter less radiation back relative to particles that can be still approximated by Rayleigh scattering (usually valid if particles size \ll wavelength). As this de-

viation happens first at the shorter wavelength, the logarithmic reflectivity difference (also called dual wavelength ratio, DWR) increases with the mean size of the particle size distribution (PSD). Also, DWR can be resolved spectrally, which allows to constrain particle scattering models (Kneifel et al., 2016), to retrieve the particle size distribution (Mróz et al., 2021; Barrett et al., 2019) or to separate attenuation and differential scattering effects (e.g. Li and Moisseev, 2019; Tridon and Battaglia, 2015).

The majority of previous radar studies on the DGL focus their analysis on a number of case studies. A more statistical investigation is presented by Trömel et al. (2019), where X-band radar observations of 52 stratiform precipitation cases obtained close to Bonn, Germany, were analysed using quasi-vertical profiles (QVP, Ryzhkov et al., 2016). They found a correlation of KDP and Ze in the DGL, and were able to link signatures in the DGL to surface precipitation. Similarly, in a statistical analysis of 27 d of C-band observations close to the city of Turin, northern Italy, Bechini et al. (2013) linked enhanced KDP in the DGL to an enhanced Ze at the surface. Schneebeli et al. (2013) analysed a dataset of polarimetric X-band radar observations of clouds ranging in temperature between -30 to 0°C collected in the Swiss Alps. Interestingly, they were unable to find a distinct KDP maximum in the DGL but rather a continuous increase of KDP and Ze towards warmer temperatures related to a general increase of the ice water content (IWC).

Only a few studies attempted to combine different radar approaches for studying the DGL. Oue et al. (2018) used vertically pointing and slant-viewing polarimetric cloud radars to study the DGL in Arctic clouds. By combining Doppler spectra of a vertically pointing Ka-band radar with slant polarimetric observations, they were able to assign the increasing ZDR signatures in the DGL with the slow, secondary mode in the reflectivity Doppler spectra. A similar correlation of spectral bi-modalities and polarimetric signatures in the DGL have also been identified in mid-latitude clouds (Moisseev et al., 2015).

In this study, we present an in-depth analysis of vertically pointing triple-frequency (X-, Ka-, W-band) Doppler spectra combined with spectral polarimetric observations from a W-band cloud radar operated at a fixed 30° elevation angle. To our knowledge, such a combined multi-frequency analysis including spectral polarimetric observations obtained at W-band with a simultaneous transmit simultaneous receive (STSR) mode radar have not been presented so far. The 3-months dataset of winter clouds observed at a mid-latitude European site close to Cologne, Germany, are described in Sect. 2.1. A combined view with the various radar observables on the DGL is illustrated for a case study in Sect. 3. The case study description provides an overview of typical radar signatures which have been observed in the DGL in previous studies, and an introduction to new observables based on spectral polarimetry and multi-frequency observations. In Sect. 4, a statistical analysis is presented, which aims

to connect polarimetric and spectral signatures dominated by newly generated ice crystals to the maximum aggregate size reached in the DGL. The section also includes a spectral analysis aimed to disentangle the contributions of upward air motion and secondary ice particle mode on the MDV reduction observed in the DGL. In Sect. 5.1, we summarise the vertical evolution of the various radar variables with a special focus on the temperature level where changes in the different variables are most pronounced. Profiles from laboratory experiments are added to this conceptual picture to allow an in-depth discussion of the most likely evolution of microphysical processes in the DGL. In Sect. 5.2, the role of sedimenting particles from higher altitudes and especially cloud top temperature on the signatures in the DGL are briefly discussed. The main findings of our statistical analysis are summarised in Sect. 6.

2 Data and methods

2.1 TRIPEX-pol campaign

The results presented in this study are based on a multi-month dataset obtained during the campaign “TRIPLE-frequency and Polarimetric radar Experiment for improving process observation of winter precipitation” (TRIPEX-pol). The campaign took place from November 2018 until January 2019 at the Jülich ObservatorY for Cloud Evolution Core Facility (JOYCE-CF, Löhnert et al., 2015, 50°54′31″N, 6°24′49″E; 111 m a.s.l.) located ca. 40 km west of Cologne, Germany. Similar to a previous winter campaign (TRIPEX, Dias Neto et al., 2019), radar Doppler spectra and moments were continuously collected from a combination of vertically pointing triple-frequency (X-, Ka- and W-band) radars. The main difference from the earlier TRIPEX campaign is an extension of the observational capabilities by two additional radars: a new X-band radar with better sensitivity and the possibility to record Doppler spectra, and a scanning polarimetric Doppler W-band radar (Table 1). The vertically pointing and the scanning W-band radars are both frequency modulated continuous wave (FMCW) systems manufactured by Radiometer Physics GmbH (Myagkov et al., 2020; Küchler et al., 2017). The X- and Ka-band systems are pulsed radar systems manufactured by Metek GmbH (Mróz et al., 2021; Görzdorf et al., 2015). All four radar systems were installed at the same roof platform within horizontal distances of less than 20 m. The resolution in range and time were adjusted to allow a very close radar volume matching (Table 1). The polarimetric W-band radar was measuring at 30° constant elevation (CEL) for intervals of 5 min towards West. In between the CEL measurements, the radar was performing single range height indicator (RHI, from 30 to 150° elevation) and plan position indicator (PPI, at 85° elevation) scans intended for wind profiling. Auxiliary instruments at JOYCE including rain gauges, microwave radiometers and Doppler wind lidars provide additional information about the

atmospheric state and precipitation on the surface (for further details, see Löhnert et al., 2015). The combination of various remote-sensing instruments also allows the continuous generation of Cloudnet classification and categorisation products (Illingworth et al., 2007). Besides in situ and remote-sensing observations, Cloudnet products also incorporate thermodynamic and wind information for JOYCE-CF extracted from analysis fields provided by the European Centre for Medium-Range Weather Forecasts (ECMWF) Integrated Forecast System (IFS) model.

2.2 Processing of the vertically pointing radar data

The dataset from the three vertically pointing radars was processed in three levels. Level 0 contains the regridded and cleaned Doppler spectra. Level 1 contains the radar moments calculated from the level 0 dataset (Sect. 2.2.1). The level 2 processing follows closely the method presented in Dias Neto et al. (2019). It includes corrections for radar-specific calibration offsets and gas, liquid and ice attenuation (Sects. 2.2.2 and 2.2.3). Key methods of the processing steps will be discussed in the following subsections.

2.2.1 Doppler spectra processing

Despite the similarity of the radar resolutions in space and time (see Table 1), the measured Doppler spectra of each radar had to be regridded to a reference time–height grid. For the reference grid, we chose a temporal resolution of 4 s and a range resolution of 36 m. The original data were matched to the reference grid using the method of nearest neighbours but only considering data points with a maximum displacement of ± 17 m in range and ± 2 s in time.

The centre frequency of the vertically pointing W-band radar had to be slightly changed from 94.00 to 94.12 GHz in order to avoid interference with the W-band polarimetric radar. This change caused some spectral artefacts which are caused by the spectral impurities of the used chirp generator. The level of impurities is considerably lower when the radar operates at the default centre frequency. Also, some weak “side lobes” appeared in the X-band Doppler spectra when stronger signals were present close to the ground.

In order to identify the spectral region with “true” atmospheric signal, we selected the Ka-band radar as reference. The Ka-band radar provides the highest sensitivity of all radars, and its Doppler spectra showed no artefacts. Due to the different heights of the lowest usable range gates of the radars, return signals below 400 m altitude were omitted. The Ka-band Doppler spectra were used to derive a spectral mask for each range gate and time step. For this, we first identified the spectral edges by subtracting the noise floor using the common method by Hildebrand and Sekhon (1974) and then locating the outermost spectral bins which exceed the noise level by 3 dB. Our spectral mask is defined by the Doppler velocity bins identified by this method to contain real sig-

Table 1. Technical specifications of the four radars that were deployed during the TRIPEX-pol campaign. The W-band and the W-band pol radar are FMCW radars, therefore the range resolution, Doppler velocity resolution and the Nyquist range vary for the different chirps. The values in this table are valid for the lowest chirp region (W-band: 215–1475 m, W-band pol: 107–715 m). The full chirp tables for both W-band radars are provided in Appendix A.

Specifications	X-band	Ka-band	W-band	W-band pol
Frequency [GHz]	9.4	35.5	94.1	94.0
Polarimetry	single-pol	LDR	LDR	STSR
Number of spectral averages	10	20	13	28
Half-power beam width [°]	1.0	0.6	0.6	0.6
Range resolution [m]	36.0	36.0	36.0	35.8
Temporal resolution [s]	2	2	3	7
Sensitivity at 1 km [dBz], 2 s integration time	−50	−63	−58	−58
Maximum range [km]	12	15	16	16
Doppler velocity resolution [m s^{-1}]	0.038	0.04	0.04	0.05
Nyquist range [m s^{-1}]	± 78	± 20	± 10	± 6

nal. This spectral mask was then applied to filter the Doppler spectra of the other two vertically pointing radars. This filtering could unfortunately not remove all artefacts, as the W-band artefacts did also sometimes overlap with the Doppler spectrum from real atmospheric targets.

From the regridded and filtered Doppler spectra, the common radar moments are derived including equivalent radar reflectivity factor (Z_e), mean Doppler velocity (MDV), spectrum width and skewness. The Ka-band spectra were also used to derive the fast-falling edge and slow-falling edge of the Doppler velocity of each spectrum. Those spectral edge velocities were derived in a similar way as the spectral mask. In case of strong atmospheric signals, spectral leakages might cause biases in the spectral edge velocity estimate. We mitigate this effect by neglecting all spectral lines which are lower than 40 dBz with respect to the maximum spectral line. Examples of the derived spectral edge velocities are shown in Fig. 3d where they are overlaid to the original spectra.

2.2.2 Evaluation of radar reflectivity calibration and antenna pointing

The reflectivity calibration of all four radars was evaluated using the drop size distributions (DSDs) measured during rainfall periods by the PARSIVEL optical disdrometer (Löffler-Mang and Joss, 2000) which was installed at JOYCE-CF in close vicinity to the radars (Fig. 1). The method is identical to the approach described in Dias Neto et al. (2019). The DSDs are used to calculate the Z_e distribution for each radar frequency and rainfall event. This distribution is then compared to the measured Z_e distribution at the lowest usable range gate (more details and discussion of uncertainties are provided in Dias Neto et al., 2019). The offsets estimated with this method for the three radars are 0 dBz for the X-band, an underestimation of 3 dBz for Ka-band and an overestimation of 2 dBz for the W-band. We applied the

disdrometer-based method to 21 rainfall cases and found no systematic temporal drifts of the estimated offsets. The W-band radar data obtained during the TRIPEX-pol campaign have also been used in Myagkov et al. (2020) to evaluate different calibration methods including also disdrometer-based methods. Our estimated bias of 2 dBz lies within their estimated values of 0.5 to 2.1 dBz for the vertically pointing W-band radar. For the polarimetric W-band radar, they found an underestimation of 0.7 ± 0.7 dBz.

Accurate zenith pointing is crucial for the analysis of Doppler spectra and MDV in order to avoid velocity biases induced by horizontal wind. The absolute pointing of the Ka-band radar has been evaluated using sun-tracking scans (e.g. Muth et al., 2012). The pointing accuracy during the campaign was found to be better than $\pm 0.1^\circ$ in elevation and azimuth. For the non-scanning X- and W-band radar, the pointing could only be evaluated relative to the absolute calibrated Ka-band radar. Following the approach shown in Kneifel et al. (2016), the pointing of the X- and W-band radars has been evaluated in relation to the Ka-band radar. For this, the difference in mean Doppler velocity (MDV) between X- and Ka-band (Ka- and W-band) has been analysed in dependency of the horizontal wind speed and direction obtained from Cloudnet. The analysis of the MDV differences obtained during TRIPEX-pol indicated that the misalignment between X- and Ka-band as well as W- and Ka-band did not exceed 0.1° .

2.2.3 Attenuation correction and relative DWR calibration

At cloud radar frequencies, atmospheric gases and hydrometeors cause attenuation which generally increases with frequency. Similar to Dias Neto et al. (2019), we first corrected the Z_e profiles for the estimated attenuation by gases. The gas attenuation profiles were calculated with the Passive and Active Microwave TRAnsfer model (PAMTRA, Mech




Figure 1. Measurement setup on the roof platform of JOYCE-CF in Jülich, western Germany (Löhnert et al., 2015). The four radars measuring during the TRIPEX-pol campaign were mounted in close vicinity to ensure optimal volume matching. A description of the radar setup can be found in Table 1 and Sect. 2. The radar measurement setup is complemented with a microwave radiometer HATPRO (in front of the W-band radar at the far right of the picture), a PARSIVEL (behind the W-band radar) and a Pluvio precipitation gauge on the far left of the picture. Information about further instrumentation on the JOYCE-CF roof platform is given in Sect. 2 and in Löhnert et al. (2015).

et al., 2020) which takes into account contributions by nitrogen, oxygen and water vapour. Profiles of temperature, humidity and pressure provided by Cloudnet were used as input for PAMTRA. Estimating the vertical profile of attenuation by liquid and frozen hydrometeors is challenging as accurate profiles of hydrometeor mass content and size distributions are required. As profile information of liquid and ice are unavailable, we only estimate the total path integrated attenuation following the approach presented in Dias Neto et al. (2019).

This method leverages on the fact that small ice particles can be assumed to be Rayleigh scatterers for which the radar reflectivity factor,

$$Ze(\lambda) = \frac{\lambda^4}{\pi^5 |K_\lambda|^2} \int \sigma_\lambda(D) N(D) dD, \quad (1)$$

is independent of the wavelength λ , if we assume a constant dielectric factor $|K_\lambda|^2$ (e.g. Kneifel et al., 2015; Hogan et al., 2000). $\sigma_\lambda(D)$ is the backscattering cross section of a particle with maximum size D and $N(D)$ is the particle size distribution. The dual-wavelength ratio,

$$DWR_{\lambda_1, \lambda_2} = Ze_{\lambda_1} - Ze_{\lambda_2}, \quad (2)$$

is the difference in reflectivity in logarithmic units at two wavelengths λ_1, λ_2 (usually with $\lambda_1 > \lambda_2$). With the particle size getting closer to the wavelength, the backscattering cross section increases less than expected from the Rayleigh approximation due to the destructive interference of electromagnetic waves scattered by various parts of the particle (see Fig. C1a). This deviation from the Rayleigh scattering behaviour starts first at the smallest wavelength. As a result, increasing DWRs can be attributed to larger mean particle sizes in the radar volume. If the mean particle size becomes large enough, also the largest wavelength would transition to the non-Rayleigh regime and the DWR will increase slower, eventually reaching a saturation point (Mason et al., 2019).

The total DWR measured under real conditions at a certain range can be written as the sum,

$$DWR_{\lambda_1, \lambda_2} = DWR_{\text{scat}} + DWR_{\text{hard}} + DWR_{\text{att}}. \quad (3)$$

DWR_{scat} is due to differential scattering of particles. Constant hardware-related offset DWR_{hard} might originate for example from radar miscalibration or differential radome attenuation. Differential attenuation causes a propagational component DWR_{att} which accumulates with increasing range. The major contributions to total attenuation are due to rain, the melting layer and supercooled liquid water. Ice and snow also contribute to W-band attenuation but as shown by Tridon et al. (2020), an ice water path larger than 1 kg m^{-2} is needed to cause 1 dBz two-way attenuation.

As proposed originally by Hogan et al. (2000) and recently evaluated by Tridon et al. (2020), a reflectivity threshold can be used to identify cloud regions where DWR_{scat} is negligible. The remaining DWR can then be attributed to the sum of DWR_{att} and DWR_{hard} . The X-band profiles would be least affected by attenuation but the X-band sensitivity is often too low to capture Rayleigh regions at high altitudes. Therefore, we use Ze profiles from the Ka-band radar as our reference which have been corrected for gas attenuation. Ka-band reflectivities between $-30 \text{ dBz} < Ze_{\text{Ka}} < -10 \text{ dBz}$ and between $-15 \text{ dBz} < Ze_{\text{Ka}} < 0 \text{ dBz}$ are used for estimating the non-scattering DWR components for W-Ka and X-Ka, respectively. To exclude partially melted particles, we additionally restricted the cloud regions used for the relative DWR calibration to be at least 1 km above the melting layer. Following the approach in Dias Neto et al. (2019), we also exclude profiles where the number of valid measurements within a 15 min time window is less than 300. Further, regions for which the variance of the DWRs exceeds 2 dB^2 , or where the correlation between Ze measurements from the reference radar (Ka) and one of the other radars (X, W) is

less than 0.7, are discarded. The estimated relative DWR offset for a moving time window of 15 min is then applied to the entire profile. As we expect the major contributions to the total attenuation from the rain part and the melting layer, this approach appears to be justified for the ice part of the cloud. In case of additional attenuation in the ice part, for example due to layers of supercooled liquid, our approach would cause an overestimation of the true attenuation, and hence our processed DWR_{scat} values would underestimate the real DWR_{scat} below the attenuation layer (even returning negative values). For convenience, in the following, we use the radar bands (X, Ka, W) instead of the wavelengths as indices (e.g. DWR_{KaW}). The DWRs have also been derived spectrally (sDWRs) by calculating the difference of the logarithmic power in each Doppler spectral bin. Identical corrections and relative calibrations as used for the DWR are applied for sDWR.

2.3 Processing of the polarimetric radar data

The polarimetric W-band radar observations were collected at constant 30° elevation and a fixed azimuth angle of 235° for 5 min time periods. The azimuth direction is close to the main wind direction where most cloud systems have been advected from during the campaign (south–west to north–west according to wind information from Cloudnet). In order to minimise time-lag differences related to different observation volumes of the slanted polarimetric radar and vertically pointed systems, we average all measurements over 5 min.

The polarimetric radar moments and the polarimetric Doppler spectra have been projected to the height above ground, and then regridded to the same observations. At a maximum range of the polarimetric radar of 16 km (see also Table 1), the height above ground is 8 km, and the maximum horizontal distance between the vertically pointing radar and the polarimetric radar is $\cos(30^\circ) \times 16 \text{ km} = 13.86 \text{ km}$. In order to reduce the noise of the specific differential phase shift (KDP), we first smoothed the differential phase shift ϕ_{dp} over five range gates (corresponding to 180 m) using a moving window mean and then averaged it over 5 min before calculating KDP as half of the discrete range derivative of ϕ_{dp} :

$$\text{KDP} = \frac{\Delta\phi_{\text{dp}}}{2 \times \Delta r}, \quad (4)$$

with Δr being the distance between the range gates.

The ZDR is defined as

$$\text{ZDR} = 10 \log_{10} \left(\frac{Z_{\text{eH}}}{Z_{\text{eV}}} \right), \quad (5)$$

with the radar reflectivity at horizontal polarisation (Z_{eH}) and vertical polarisation (Z_{eV}). The ZDR values are dominated by the particles contributing largest to the reflectivity. In addition to ZDR, we also derived the spectral ZDR (sZDR) as the logarithmic difference of horizontal and vertical power in each Doppler spectral bin. The maximum in

sZDR called $sZDR_{\text{max}}$ indicates the presence of high ZDR producing ice particles within the radar volume even in cases where the ZDR is for example lowered by the presence of low ZDR producing aggregates. The quality of polarimetric measurements strongly depends on the signal-to-noise ratio (SNR). Variance in ZDR, ϕ_{dp} and ρ_{hv} drastically increases at very low SNR (Eq. 6.122 and Sect. 6.5 in Bringi and Chandrasekar, 2001). Therefore, in order to use only high quality data for the following analysis, we omit all polarimetric observations when SNR (or in case of sZDR, spectral SNR in a spectral bin) is less than 10 dB.

2.4 Evaluation of the temperature information from Cloudnet

In Sects. 4 and 5.2, we statistically analyse different radar variables with respect to their temperature-dependent behaviour. As no regular radio soundings are available at JOYCE-CF, we use the temperature information included in the Cloudnet products. We evaluated the temperature information from Cloudnet with the temperature measured by 27 radiosondes which were launched throughout the course of the TRIPEX–pol campaign for all heights below 8 km (see Chap. 3.3 in Dias Neto, 2021). The analysis showed a good agreement between the temperature information from Cloudnet and the measured temperature from the radiosondes. The mean difference between the Cloudnet and radiosonde is 0.2°C , with a standard deviation of the mean difference of 1.1°C . The correlation between the Cloudnet temperature and measured temperature is 0.9. Further, the root mean square difference was found to be $\text{RMSD} \approx 1.1^\circ\text{C}$.

2.5 Description of the dataset

Most clouds and precipitation events that occurred during the 89 d of the campaign were caused by mid-latitude frontal systems which are common during wintertime at JOYCE-CF. According to the Cloudnet classification, ice and mixed phase clouds were present 46.6 % of the time (1029 h). Rainfall was observed during 9.2 % (202.8 h) of the time causing a total accumulation of 152.7 mm of rainfall measured by the Pluvio weighing gauge installed at JOYCE-CF. On three days, snowfall was reaching the ground. A total of 18.6 h of snowfall produced a total liquid equivalent accumulation of 10.8 mm. The coldest temperature of -7.0°C was observed on 24 January 2019, one of the three days with snowfall on the ground. The warmest temperature, reaching 16.7°C , during the campaign was observed on 24 November 2018.

3 Snowfall case study: 30 January 2019

With the following snowfall case study, we aim to provide an impression of the data quality and illustrate the complementary information in spectral multi-frequency and polarimetric observations. Common observational features of the

DGL visible in standard radar moments and in new spectral variables will be discussed. The case study selected shows more pronounced signals than the average profiles discussed later in the statistical analysis. However, the case is not an exceptional event and similar profiles of radar variables can be found frequently at JOYCE-CF during similar winter cases.

On 30 January 2019, a frontal system passed over JOYCE-CF and caused snowfall reaching the surface with a liquid equivalent accumulation of 6.6 mm. During the entire day, temperatures at the surface remained below freezing, ranging between -2.4 and -0.4 °C. The various radar observables from zenith pointing and slant polarimetric observations are displayed in Fig. 2. The cloud system produced snowfall reaching the ground mainly between 05:00 and 19:00 UTC. The overall cloud structure observed in zenith is very similar to the radar measurements at 30° elevation. This similarity was regularly observed for the 3-months time period probably due to the predominantly large-scale structure of the winter precipitation observed during the campaign.

The MDV (Fig. 2b) throughout the case are found to be larger than -1.5 m s⁻¹ which indicates unrimed or only slightly rimed particles (Kneifel and Moisseev, 2020). Here we use the convention that negative (MDV) velocities correspond to motion towards the ground. Faster falling particles therefore have smaller (more negative) values than slower falling particles. When plotting the DWR in the triple-frequency space (DWR_{KaW} against DWR_{XKa}, not shown), we also find a “hook shape” which has been previously found to indicate predominance of unrimed aggregates (Kneifel et al., 2015). Certainly, the particles observed on the surface are not necessarily representative for the particles sampled with the radars due to impacts of advection and further growth processes during sedimentation towards the surface. Still, the snowfall reaching the ground was mostly comprised of unrimed or only slightly rimed crystals and aggregates. Visual observation on the ground at the site between 09:00 and 10:00 UTC (Fig. B1) reveals the presence of stellar and dendritic crystals reaching up to 4 mm in size mixed with unrimed aggregates with maximum sizes up to 10 mm.

The combined radar observations reveal several features which have been reported and discussed in previous literature related to the DGL (Ori et al., 2020; Barrett et al., 2019; Griffin et al., 2018; Oue et al., 2018; Schrom et al., 2015; Moisseev et al., 2015; Andrić et al., 2013; Bechini et al., 2013, among others). The Ze values (Fig. 2a) rapidly increase at the -15 °C temperature level most likely due to an increase in the mean particle size as indicated by DWR_{KaW} (Fig. 2c). A layer of enhanced $sZDR_{max}$ (Fig. 2d) values up to 4 dB at -15 °C indicates a rapid generation of asymmetric particles. Note that at 30° elevation, ZDR is expected to be in general smaller than ZDR measured at lower elevation angles which have often been used in previous studies where data from lower frequency scanning radar systems have been analysed. According to Myagkov et al. (2016b), such values of differential reflectivity at these temperatures correspond to

horizontally aligned strongly oblate (plate-like) ice particles. For the following analysis, we define the aspect ratio similar to Takahashi et al. (1991) as the ratio of the *a* and *c* axis of an ice crystal. Plate-like particles for example then have aspect ratios larger than 1.

The decrease of $sZDR_{max}$ from 4 to 2 dB towards lower layers indicates a change in particle properties (i.e. aspect ratios become closer to unity and/or apparent ice density becomes smaller) of the strong ZDR producing particles found at -15 °C. Also the KDP (Fig. 2e) shows an immediate increase at -15 °C with values of up to 3 ° km⁻¹. Interestingly, both KDP and $sZDR_{max}$ remain enhanced down to the surface despite the ongoing aggregation indicated by increasing DWR_{KaW} towards the ground.

Additional insights into the vertical evolution of particle populations and their contribution to radar moments can be gained from the analysis of vertical profiles of Doppler spectra (Fig. 3d–f). The first aggregates which produce a noticeable DWR_{KaW} signal (see Fig. 3e) appear at -18 °C on the fast edge of the Doppler spectra (ca. -1 to -1.2 m s⁻¹). Interestingly, the spectral velocity where we find the largest sZe values at this temperature is at slightly lower velocities (-0.8 to -1 m s⁻¹). This could indicate that the number concentration of the aggregates producing the enhanced $sDWR_{KaW}$ region is still low. At around the same temperature level, we also find a secondary mode in sZe and a broadening of $sDWR_{KaW}$ (indicated by magenta square in Fig. 3d, e). At its first appearance, the secondary mode shows initial spectral velocities close to -0.3 m s⁻¹ which indicates small ice particles rather than supercooled liquid droplets which typically produce a narrow spectral mode around 0 m s⁻¹ (e.g. Kalesse et al., 2016; Shupe et al., 2004). The spectral asymmetry introduced by the secondary mode leads to a rapid change in the Doppler spectral skewness (Fig. 3b). Starting at -18 °C, the skewness increases from values close to unity (symmetrical spectra) to 1.4 at -17 °C (Doppler spectra are skewed towards the slower falling side). The secondary mode appears alongside polarimetric signatures. KDP starts to increase at -20 °C, reaching a maximum of 2.1 ° km⁻¹ at -15 °C, correlating well with the increase and the maximum of $sZDR$ of around 4 dB. The enhanced KDP and $sZDR$ values indicate an increase in concentration and aspect ratio of small ice crystals.

For temperatures warmer than -15 °C, the fall velocities, sZe and $sDWR_{KaW}$ of the secondary mode increase until they merge with the main mode at around -12 °C. Interestingly, at this temperature we also see a distinct slow-down of the MDV from approx. -0.9 m s⁻¹ at -17 °C to -0.6 m s⁻¹ at -12 °C. Looking at the spectrogram for this particular time, it appears as if the slow-down in MDV is the result of the new mode and a general shift of the entire spectrum towards larger values. The latter is compatible with the effect of upward air motion. This slow-down of the entire Ze spectrum is illustrated in Fig. B2a) which presents a zoomed view into the temperature region -16 to -10 °C. The main mode con-




Figure 2. Snowfall event occurring on 30 January 2019 at JOYCE. From the vertically pointing radars, the (a) Z_e and (b) MDV at Ka-band, and the (c) DWR_{KaW} are shown as time–height plots. From polarimetric observations at W-band and 30° elevation angle (mapped to height above ground), the (d) maximum spectral ZDR $sZDR_{max}$ and (e) KDP are presented. In (a–e), the dashed red lines depict the -20 , -15 and -10 °C isotherms. Impressions of dendrites and aggregates sampled on the ground between 09:00 and 10:00 UTC are provided in Fig. B1.

tributes more power to the spectrum than the newly formed secondary mode. Therefore, shifts of the main mode towards slower or faster velocities dominate changes in MDV. Hence, the slow-down of the main spectral mode at -12.5 °C reduces the MDV at this temperature.

Just slightly below the temperature level where the secondary mode merges with the main mode, we find the largest $sDWR_{KaW}$ values of up to 10 dB at spectral velocity bins between -1 and -1.5 $m\ s^{-1}$. At temperatures around -10 °C, the $sZDR_{max}$ values increase again. The maximum in $sZDR_{max}$ at around -8 °C roughly coincides with the appearance of a weak secondary mode in the sZ_e and an increase in KDP. The new particle mode as well as the enhanced $sZDR_{max}$ and KDP values mostly disappear at temperatures higher than -5 °C while sZ_e and $sDWR_{KaW}$ remain enhanced.

The signatures found in this case study are largely in agreement with radar signatures reported in previous studies about particle growth and aggregation in the DGL (Griffin et al., 2018; Schrom and Kumjian, 2016; Moisseev et al., 2015; Schrom et al., 2015; Andrić et al., 2013; Bechini et al., 2013, among others). However, some differences are also found, especially when comparing our results to studies that

analysed lower frequency polarimetric radar observations. Those observations frequently revealed layers of enhanced KDP and ZDR at the -15 °C level. One reason for the less layered appearance of KDP and ZDR in our case could be related to the higher frequency used for polarimetric observations in this study. As KDP is inversely proportional to the radar wavelength (e.g. Bringi et al., 2001), we are able to observe KDP signals from relatively small particle concentrations which are difficult to detect by low-frequency polarimetric radars. For example, a KDP signal of $1^\circ\ km^{-1}$ observed at W-band would only be $0.1^\circ\ km^{-1}$ at X-band. Also, the SNR is much higher for cloud radars since the maximum distance measured is smaller than for typical weather radars. As a result, only the maximum of the KDP enhancement close to the -15 °C level might be detectable by low-frequency radars and regions with enhanced particle concentrations, but KDP values below the detection limit might be missed. Another reason for the less layered appearance of KDP and ZDR might be an underrepresentation of cases with strong forcing conditions during TRIPEX–pol. More intense vertical air motions are expected to result in a larger concentration of particles and abundance of dendrites, that is expected to lead to stronger aggregation and more intense de-




Figure 3. Radar profiles and spectrograms from the snowfall event occurring on 30 January 2019 at JOYCE at 13:30 UTC. Vertical profiles of (a) Ze and MDV, (b) skewness from the Ka-band radar, and (c) KDP and ZDR from the W-band radar are shown. Spectrograms (i.e. vertical evolution of Doppler spectra) of (d) Ze from the Ka-band are shown together with the derived spectral edge velocities (red lines). Panel (e) shows spectrograms of DWR_{KaW} and (f) ZDR from the W-band radar. The magenta squares in (d–f) depict the region where a secondary spectral mode is visible. An example of a single Doppler spectrum, where the second mode is visible, is also provided in Fig. B2.

pletion of ice crystals (e.g. Moisseev et al., 2015; Schrom et al., 2015). Radars operating at longer wavelengths can clearly detect these cases with large concentrations of ice crystals at -15°C but might miss cases with weaker forcing due to sensitivity limits. The differences in sensitivity between W-band and lower frequency radars might cause a selection bias of low-frequency radars with respect to stronger depositional growth and aggregation cases. A more detailed discussion of the expected similarities and differences of ZDR and KDP at X-band and W-band is provided in Appendix C.

In the following sections, we will apply our analysis to all ice-containing clouds included in our dataset. As this case study illustrated, the combination of observations from the slant (Fig. 2d–e) and the zenith direction (Fig. 2a–c) appears to be reasonable, especially when applying an additional temporal averaging over 5 min to the profiles. This will allow us to link polarimetric and multi-frequency zenith observations in order to better understand which radar variables are connected to different intensities of aggregation in the DGL.

4 Profile classification by the mean aggregate size in the DGL using DWR_{KaW}

Aggregation becomes particularly active in the DGL, causing rapid changes in radar quantities sensitive to the mean size, such as Ze or DWR (see also Fig. 2). As a growing aggre-

gate will deviate from Rayleigh scattering first at the shortest wavelength, we use DWR_{KaW} as our most sensitive measure for the onset of aggregation. In order to exclude multi-layer or sublimation cases, we require the radar profiles to be continuous within the DGL. Following the approach presented in Dias Neto (2021), these profiles were sorted into three classes according to their maximum DWR_{KaW} value reached within the DGL (Table 2). Assuming inverse exponential PSDs combined with particle and scattering properties of dendritic aggregates (Ori et al., 2021), we find the three DWR_{KaW} classes representing mean mass diameters (D_0) ranging from 1 to 6 mm. However, the D_0 estimation likely represents only a lower limit, especially for DWR_{KaW} class 3. As is shown in Mason et al. (2019), the shape of the PSD influences the shape of the triple-frequency signatures, and by extension also the DWR measurements. A narrow PSD with a large D_0 might account for the same DWR_{KaW} as a more wide PSD with a smaller D_0 . The DWR_{KaW} classes were chosen such that there is a similar number of profiles within each DWR_{KaW} class. All profiles with DWR_{KaW} values exceeding 9.5 dB are excluded as they are most likely related to partially rimed aggregates or due to an insufficient correction of W-band attenuation.

Table 2. Definition of maximum DWR_{KaW} intervals within the DGL (i.e. temperature region between -20 and -10 °C) used to classify radar profiles according to the particles' maximum mean mass diameters D_0 in the DGL. The last column denotes the number of available radar profiles with continuous observations within the DGL.

max (DWR_{KaW}) [dB]	class number	approx. D_0 [mm]	number of profiles
0–1.5	1	< 0.75	222
1.5–4.0	2	0.75–1.5	223
4.0–9.5	3	1.5–6	190

4.1 Relation of vertically pointing radar variables to the mean aggregate size in the DGL

After classifying all vertically pointing profiles according to their maximum DWR_{KaW} in the DGL, we can now investigate how other radar moments evolve as a function of temperature for the different classes (Fig. 4). The profiles of DWR_{KaW} (Fig. 4b) reveal that the maximum D_0 is reached at the lower end of the DGL (-10 °C) with the strongest DWR_{KaW} increase found at temperatures warmer than -15 °C. This is also the temperature where we find DWR_{XKa} (Fig. 4c) to rapidly increase reaching values of 2 dB at -10 °C for the highest DWR_{KaW} class. Both DWRs only slightly change from -10 to -5 °C. From -5 °C towards the melting layer we find an additional increase in the DWRs, especially in DWR_{XKa} . This is in agreement with previous DWR studies (Ori et al., 2020; Dias Neto et al., 2019) and in situ observations (Lawson et al., 1998). The most common explanation for this second aggregation layer is an increasing thickness of the quasi-liquid layer of the ice surface causing the sticking efficiency to rapidly increase at $T > -5$ °C coupled with enhanced depositional growth at around -5 °C (Lamb and Verlinde, 2011).

Interestingly, small differences in the Ze median and quantile profiles between the aggregation classes are already visible at temperatures lower than -20 °C (Fig. 4a). At the top of the DGL, we find a 3 dB difference in Ka-band reflectivity medians between the different aggregation classes which increase up to 10 dBz at the bottom of the DGL. Between -20 and -15 °C, the slopes of the Ze medians are relatively similar. From -15 °C to the bottom of the DGL, we find a relatively constant Ze for class 1, an unchanged linear increase for class 2 and a more rapidly increasing Ze curve for class 3. Unlike the DWRs, the Ze medians remain relatively constant or even decrease for temperatures between -10 and 0 °C. This behaviour might be related to non-Rayleigh scattering effects at Ka-band (also indicated by increasing DWR_{XKa}) and/or a result of microphysics. For example, a simultaneous increase in mean particle size and a decreasing number concentration caused by aggregation might have a compensating effect on Ze. The quantiles of the distribution in general fol-

low the Ze median throughout the shown temperature range. At the top of the DGL, the spread between the 25th and 75th percentile is 11 dB (10, 12.2 dB) for aggregation classes 1 (2, 3). This spread remains similar throughout the DGL. Only at warmer temperatures the spread increases, reaching 16.6 dB (24.4, 30 dB) at 0 °C.

The most intriguing signature, however, is found for the MDV in the DGL (Fig. 4d). The magnitude of MDV increases (in an absolute sense) from ca. -0.6 m s $^{-1}$ at -30 °C to -0.9 m s $^{-1}$ at -18 °C. This continuous increase is expected due to evolving particle size related to depositional growth and aggregation. Unlike Ze and DWR_{KaW} , the MDVs are only slightly more negative for the larger DWR_{KaW} classes. Also, the quantiles of the distributions reveal a larger overlap in most regions. When the temperature increases above -16 °C, the largest DWR_{KaW} class shows a pronounced reduction of the MDV reaching a local minimum of -0.8 m s $^{-1}$ at -13 °C. This slow-down is also visible in the vertical evolution of the quantiles. The lower (upper) quantile is reduced from -1.1 m s $^{-1}$ (-0.83 m s $^{-1}$) at -16 °C to -1.03 m s $^{-1}$ (-0.63 m s $^{-1}$) at -13 °C, resulting in a slow-down of 0.075 m s $^{-1}$ (0.15 m s $^{-1}$). At the bottom of the DGL, the MDV values increase to only slightly larger values (-0.8 to -1.1 m s $^{-1}$) as compared to the top of the DGL. This “slow-down” of the MDV in the DGL appears to increase with DWR_{KaW} class. Different explanations for this slow-down in the DGL have been discussed in literature. A common explanation for this effect is the occurrence of a new and slower ice mode in the Doppler spectrum (similar to the mode shown in Fig. 2f) which would cause a reduction in the MDV (e.g. Schrom and Kumjian, 2016). An alternative explanation proposes the existence of a frequently occurring updraft at -15 °C produced by large-scale lifting (Zawadzki, 2013). A third explanation assumes that the latent heat released by enhanced depositional growth of ice particles in the DGL will locally increase buoyancy and eventually cause upward air motion (Schrom and Kumjian, 2016; Zawadzki, 2013). Obviously, if only the MDV is considered, it is impossible to disentangle vertical air motion and microphysical effects. In the following section, we will extend the analysis to the Doppler spectra in order to better estimate the relative contributions of microphysics and upward motion on the observed MDV slow-down in the DGL.

4.2 Estimating the contribution of vertical air motion and new particle mode to the MDV reduction in the DGL

The observed slow-down of the MDV in the DGL could be solely caused by the appearance of a slow, secondary particle mode or entirely due to an upward air motion. Both effects might even be connected to each other, for example if a large-scale updraft is locally enhancing relative humidity which might then trigger nucleation of new ice particles. Alternatively, rapid depositional growth of ice particles




Figure 4. Median (solid line) and quantiles (shading) of (a) Ze at Ka-band, (b) DWR_{KaW} , (c) DWR_{XKa} and (d) MDV (Ka-band) profiles stratified with temperature and classified into classes of maximum DWR_{KaW} in the DGL (see class definition in Table 2). The temperature region of the DGL is indicated by the dashed red lines. Only profiles which are continuous in the DGL are considered.

in the DGL might release latent heat which could cause a buoyancy-driven upward air motion.

Vertical air motion can be derived from Doppler spectra if the terminal velocity of a spectral mode or distinct spectral feature is well-known. In mixed-phase clouds, the presence of a narrow spectral peak due to small supercooled liquid water droplets can sometimes be used to infer vertical air motion, assuming that the terminal velocity of the droplets is negligible (e.g. Zhu et al., 2021; Shupe et al., 2004). Unfortunately, a supercooled liquid peak does not appear in all ice clouds and is often only occurring in a relatively narrow height region of the cloud. However, vertical air motion is commonly assumed to impact all particles in the radar volume in the same way. As a result, the spectrum will be shifted to higher or lower Doppler velocities but without changing its shape. In contrast, a new particle mode will only affect the slow edge of the spectrum while the fast-falling particles will be mostly unaffected. This slow-down effect due to new particle formation will be detectable as soon as the new mode overcomes the noise threshold even if no distinct secondary peak can be identified. Therefore, if the MDV slow-down is solely an effect of vertical air motion, we expect the slow and the fast spectral edge to show an identical reduction in Doppler velocity. If the slow-down is caused by new particles only, the slow edge should decrease while the fast edge should remain constant or increase.

The median and quantile profiles of slow and fast spectral edges separated for the three DWR_{KaW} classes are shown in Fig. 5. We can clearly see that the slow-down on the slow edge is larger than on the fast edge for all classes. This implies that the MDV slow-down is indeed a combination of vertical air motion and the formation of a new, slower spectral mode. The reduction on the fast edge is almost only visible for the largest DWR_{KaW} class while the reduction on the slow edge is also noticeable for the lower DWR_{KaW} classes. On the fast edge, the differences in the median velocity profiles between class 1 and 3 are around 0.2 to 0.3 $m\ s^{-1}$ throughout the entire temperature region, while on the slow edge the differences are negligibly small. When focusing on

the largest DWR_{KaW} class and zooming into the temperature region of the DGL (Fig. 5b), we can roughly estimate the velocity reduction at the two spectral edges. We assume that the spectral edges without new particle mode or updraft would remain constant or increase towards warmer temperatures. For each spectral edge profile, we search for the level where we find the velocities indicating the onset of a slow-down (vertical red lines in Fig. 5b). If we subtract the velocities at this level from the values of the strongest reduction found in the DGL, we obtain a lower limit estimate of the real slow-down. For the slow edge, we find a reduction of 0.28 $m\ s^{-1}$ between -18 and -14 $^{\circ}C$. On the fast edge, the reduction is almost a factor of 3 smaller with a reduction of 0.1 $m\ s^{-1}$ between -16 and -14 $^{\circ}C$. The total slow-down in the MDV is 0.18 $m\ s^{-1}$ between -16 and -14 $^{\circ}C$, which is less than for the slow edge simply due to the stronger contribution of the larger (faster) particles to the MDV. A Kolmogorov–Smirnov two-sample test revealed that the slow-down on the slow-falling edge is statistically significant, while it is not significant for the fast edge.

In case of no updraft, we would expect the fast edge velocity to continuously increase, similarly as for temperatures colder than -16 $^{\circ}C$. So even if the Kolmogorov–Smirnov test indicates that the slow-down on the fast edge is not statistically significant, we argue that the persistent stagnation of the fall velocities over the temperature range from between -16 and -14 $^{\circ}C$ strongly points towards the presence of an updraft. The presence of an updraft can also be detected at the slow edge velocities. The median values reduce at -14 $^{\circ}C$ to almost 0 $m\ s^{-1}$; the quantiles indicated by the shaded areas in Fig. 5b show even upward (positive) velocities which strongly indicates an upward air motion. The presence of an updraft was also found by Dias Neto (2021) for a similar winter dataset collected at JOYCE-CF. Our estimated updraft velocity represents, however, only a lower boundary of the true updraft speed as the terminal velocity of the particles at the spectral edges are unknown.




Figure 5. (a) Derived Doppler velocities of the spectral edges classified into DWR_{KaW} classes (see also Table 2). Solid lines indicate the median and the shading denotes the quantiles of the distribution. The dashed red lines indicate the location of the DGL. (b) Zoomed view into the DGL region but only including the spectral edge velocities (green) and MDV (black) of class 3. The vertical dashed red lines in (b) visualise the points in the spectral edges and MDV which have been used to estimate the maximum velocity reduction (see details in the text).

4.3 Investigation of the small particle mode

The spectral analysis in the previous section clearly indicated the formation of a new, slow falling ice particle mode in the DGL. As polarimetric variables are known to be particularly sensitive to newly formed ice crystals with asymmetric shape, we also sort the polarimetric profiles according to the DWR_{KaW} classification in order to analyse their relation to the aggregate class.

Interestingly, the median ZDR profiles do not reveal a clear maximum of ZDR at -15°C (Fig. 6a) despite the expected presence of dendrites with very low aspect ratios. Between -30 and 0°C we find the ZDR profiles to be increasingly shifted to smaller values for higher aggregate class. The spread between the quantiles is rather large compared to the relative difference between the aggregate classes. The differences between the lower and upper quantile varies between 0.3 and 0.5 dB, with a similar magnitude for all classes and throughout the entire analysed temperature region. This indicates that even if the medians and quantiles are shifted towards smaller values for higher aggregate class, there is a large variability within each class. From the ZDR profiles alone it is very difficult to tell whether the shift towards small values is caused by a change of small ice particle properties or by varying number, size or density of aggregates.

As already illustrated in the case study analysis (Fig. 2h), the high ZDR producing particles usually populate at lower Doppler velocities and are thus well separated from larger, low ZDR producing aggregates. As a result, the $s\text{ZDR}_{\text{max}}$ (Fig. 6b) is mostly unaffected by the presence of aggregates and shows the ZDR signature of the most ZDR producing particles present in the radar volume. Increasing first at -18°C from 1.0 dB, $s\text{ZDR}_{\text{max}}$ of aggregate class 2 and 3 reaches a maximum within the bottom half of the DGL (1.5 and 1.7 dB for class 2 and 3, respectively). Below the DGL, $s\text{ZDR}_{\text{max}}$ remains relatively constant between 1 and 1.5 dB

down to -3°C . Further down, $s\text{ZDR}_{\text{max}}$ drops to values of 0.8 to 1 dB, which are similar to the values found at temperatures colder than -20°C . Overall, the $s\text{ZDR}_{\text{max}}$ in the DGL is only weakly dependent on aggregate class, with only slightly larger values (ca. 0.5 dB) for aggregate class 2 and 3 compared to class 1. This seems to indicate that the properties impacting ZDR, such as aspect ratio or ice density of the dendrites and other plate-like particles growing in the DGL, are overall relatively similar for aggregate classes 2 and 3. However, the position of the maxima within the DGL of aggregate class 2 and 3 differs. Class 3 reaches its maximum slightly below -15°C , while class 2 reaches its maximum at the bottom of the DGL. This might indicate that the $s\text{ZDR}_{\text{max}}$ producing particles of class 2 continue to grow and increase their aspect ratio and/or density throughout the entire DGL, while the particles in class 3 reach their largest aspect ratio already slightly below -15°C .

Most notably, when comparing ZDR and $s\text{ZDR}_{\text{max}}$, the $s\text{ZDR}_{\text{max}}$ profiles lack the shift towards higher values with lower aggregate class. As both variables are independent of particle concentration, it appears most likely that the ZDR shift is related to on average larger aggregates throughout the profile rather than less ZDR producing ice crystals. This interpretation is consistent with the DWR_{KaW} profiles in Fig. 4b which show overall larger mean aggregate sizes between -30 and 0°C for increasing aggregate classes. The maximum size of aggregates in the DGL appears to be correlated with the mean size of particles above the DGL. For temperatures colder than -18°C , the spread between the lower and upper quantile of all classes is approximately 1 dB. At -18°C , which is the temperature level where the medians of class 2 and 3 start to increase, we also find an increasing spread between the quantiles of class 2 and 3 reaching values up to 3.5 dB (4 dB) for class 3 (class 2) in the lower half of the DGL. Below the DGL, the spread slightly reduces again, un-




Figure 6. Same as in Fig. 4, but showing profiles of W-band (a) ZDR, (b) $sZDR_{\max}$, (c) KDP and (d) Ka-band skewness (for interpretation of the skewness sign, see text). Note that the polarimetric variables have been obtained at 30° elevation and the skewness from the zenith Doppler spectra.

til approximately 1.5 dB at 0°C . In general, the upper quantile follows the course of the median. This large spread of the quantiles within the DGL indicates that many profiles included in the dataset show only a rather weak or completely absent increase in $sZDR_{\max}$ within the DGL. This is most likely also the explanation why the median of $sZDR_{\max}$ is substantially smaller than the $sZDR_{\max}$ profile presented in Fig. 3.

Similar to $sZDR_{\max}$, the median KDP profiles (Fig. 6c) above the DGL show only low values (0.25°km^{-1}) and no clear difference between the aggregate classes. This implies that the particles falling from aloft into the DGL are not only similar in terms of their properties impacting ZDR, but also in terms of average concentration. All median KDP profiles start to increase at -18°C which is again very similar to the behaviour of $sZDR_{\max}$. The KDP values within the DGL show a stronger increase for larger aggregate classes as observed for $sZDR_{\max}$. Interestingly, the KDP profile observed for aggregate class 3 is on average not peaking directly at -15°C , but rather linearly increasing from -18°C towards -12°C where it reaches a maximum of 0.7°km^{-1} . The KDP profiles remain enhanced from the bottom of the DGL down to -3°C where their values suddenly drop to nearly 0°km^{-1} . The rapid decrease of KDP and $sZDR_{\max}$ at temperatures larger than -3°C is correlated with the increase of DWR_{XKa} (Fig. 4c) in this region. It appears to be quite likely that the second intensified aggregation zone close to 0°C is mainly responsible for the reduction of high ZDR producing ice particles. Similar to $sZDR_{\max}$, the spread between the upper and lower quantile is small (within 0.5°km^{-1}) above the DGL. At -18°C the spread especially of class 3 increases to 1.1°km^{-1} at -12°C . The spread reduces again at temperatures warmer than -5°C to 0.65°km^{-1} at 0°C . Again, the distributions show a large overlap between the different classes. Also for the largest aggregate class, we find profiles with very weak increase of KDP. On the other side, the upper quantiles also clearly reveal a higher likelihood for extreme KDP and $sZDR_{\max}$ values for higher aggregate classes.

In order to complement the picture of the small particle evolution, in Fig. 6d we also included the skewness of the non-polarimetric Ka-band Doppler spectra recorded in zenith. Asymmetric broadening of the spectra on the fast edge (negative skewness) or on the slow edge (positive skewness) can be very well detected in the skewness profile (see also example profile in Fig. 3). The formation of new, small ice particles can be expected to result in positive skewness values even if the spectra do not reveal a separated spectral mode. Above the DGL, all median skewness profiles are close to 0 indicating on average symmetrical spectra. Similar to KDP, the skewness values increase most rapidly at -18°C . We also find clearly larger skewness values (up to 0.2) for higher aggregate classes. For the largest aggregate class, the skewness values also remain close to 0.2 down to the -5°C level. This similarity to the signature found in KDP is even more surprising as we can expect processes such as riming (broadening on fast spectral edge) to decrease skewness, and hence potentially masking the signature of the small particle mode. Also, when looking at the 75th percentiles (shaded areas), we find, similar to $sZDR_{\max}$ and KDP, distinct maxima at -15 and -5°C .

Comparing the results of the statistical analysis to the case study presented in Sect. 3, we note the smaller values of KDP, $sZDR_{\max}$ and skewness found in the statistics. In the case study, $sZDR_{\max}$ values of up to 4 dB at around -15°C were reached, alongside a maximum KDP of approx. 2°km^{-1} and a skewness of 1.3. However, in the statistics we classify the profiles with the maximum DWR_{KaW} . In Fig. 2, one can see that we do not always find an increase in KDP or $sZDR_{\max}$ for increasing DWR_{KaW} . For example, at 06:30 UTC, a strong increase in KDP below -15°C coincides with an enhanced $sZDR_{\max}$ and DWR_{KaW} . At later periods of the day, for example at 18:00 UTC, we see enhanced DWR_{KaW} without enhanced KDP and $sZDR_{\max}$. Those examples explain why the medians in our statistical analysis are shifted to smaller values.

Summarising the results of this section, we can say that our statistics reveal that the aggregation in the DGL is corre-

lated to growth and evolution of asymmetric particles within the DGL. Aggregation in the DGL appears to be stronger if larger aggregates fall already from higher levels into the DGL. Most notably, signatures related to crystal growth or aggregation that evolved in the DGL appear to persist to lower layers until -5°C is reached. In the following section, we will discuss our main findings together with laboratory results, and attempt a physical interpretation of the potentially involved processes.

5 Discussion

5.1 Interpretation of the temperature dependence of ice particles, aggregates and vertical air motion in the DGL

The most striking features visible in the radar data of the DGL are rapid changes of vertical gradients within a relatively small temperature range, and distinct maxima at specific temperatures. In order to simplify their interpretation, we schematically show the radar profiles for the highest aggregation class in Fig. 7, together with related results from previous laboratory studies. Although we are lacking in situ observations from inside the cloud for our dataset, we base our interpretation on well-established relations between microphysical processes and distinct radar signatures. For example, aggregation can be clearly associated with an increase in DWR (e.g. Ori et al., 2020; Barrett et al., 2019; Dias Neto et al., 2019; Mason et al., 2019), whereas plate-like particle growth is strongly linked to enhanced ZDR and KDP (e.g. Moisseev et al., 2015; Schrom et al., 2015).

According to Takahashi (2014), hereafter “TKH14”, the increase in mass and size of plate-like particles is strongest between -16 and -13°C with a local maximum at -15°C for stellar particles. When growing at constant temperature and high supersaturation, they develop distinct habits ranging from sectors (SEC), broad branched particles (BB), stellars (STEL), dendrites (DEN) and fern-like dendrites (FERN) (indicated on the left in Fig. 7). This shape dependency on temperature was also confirmed by airborne in situ observations (Bailey and Hallett, 2009). As mentioned in (Bailey and Hallett, 2009, among others), the shape of the particles does not only depend on temperature, but also on the supersaturation that the particle experiences during growth. During the TRIPEX-pol campaign, we did not have sufficient relative humidity information. In the following, we therefore only focus our interpretation on the observed temperature-dependent growth of ice particles.

The plate-like particle growth connected to increasing $s\text{ZDR}_{\text{max}}$ values seems to start at -20°C . This temperature roughly coincides with the temperature level where the aspect ratio of crystals observed in the laboratory starts to deviate from unity (see Fig. 2 in Takahashi et al., 1991, hereafter TKH91). Initially, the increase in $s\text{ZDR}_{\text{max}}$ is relatively moderate down to -15°C . At this temperature level,

$s\text{ZDR}_{\text{max}}$ strongly increases reaching its maximum values between -13 and -14°C . As long as the particles grow in the plate-like growth regime, we can expect a certain correlation of aspect ratio and size. Regions showing enhanced $s\text{ZDR}_{\text{max}}$ values are also likely connected to larger sized crystals. Interestingly, the maximum in $s\text{ZDR}_{\text{max}}$ is found at slightly warmer temperatures than -15°C where the maximum size and aspect ratio is found in TKH14 and TKH91. A simple explanation might be the fact that unlike the particles grown in the experiments by TKH14 and TKH91, particles in real clouds are sedimenting into different temperature regions while growing. Once the sedimenting particles reach the -15°C level, they grow most efficiently by vapour deposition as indicated by the strong gradient found in the $s\text{ZDR}_{\text{max}}$ profile. The most favourable growth region (indicated in the curve of maximum particle size found in TKH14) extends from -15°C to almost -13°C which is the location where $s\text{ZDR}_{\text{max}}$ indicates the largest particles with most extreme aspect ratios. But why are $s\text{ZDR}_{\text{max}}$ values not increasing further towards the bottom of the DGL? Scattering calculations (Myagkov et al., 2016b; Hogan et al., 2000) indicate that ZDR is only slowly increasing once a certain aspect ratio is reached (see also Fig. C1b). The $s\text{ZDR}_{\text{max}}$ values can therefore be expected to reach a certain saturation value once the particle grew into a shape with very low aspect ratio. Further, in case of dendritic growth, the effective density of the particle decreases with size, leading to a saturation of ZDR. With increasing size, also the particles' cross sectional area strongly increases which makes it more likely for the particle to collide with another crystal or aggregate. At a certain point, increasing aggregation (more likely for larger crystals) might counteract the general increase in crystal size leading to the observed slight decrease in $s\text{ZDR}_{\text{max}}$ towards -10°C . The increasing DWR_{KaW} values in the DGL appear to be consistent with this interpretation. The $s\text{ZDR}_{\text{max}}$ and DWR_{KaW} most rapidly increase both at -15°C but unlike $s\text{ZDR}_{\text{max}}$, the DWR_{KaW} continues to rise throughout the DGL. This effect might also be responsible for the different vertical location of the maximum in $s\text{ZDR}_{\text{max}}$ for aggregate class 2 and 3 (Fig. 6b). More intense aggregation (class 3) might consume the largest dendrites earlier, which results in a $s\text{ZDR}_{\text{max}}$ maximum closer to the -15°C level.

The temperature region of maximum crystal growth found in TKH14 also roughly coincides with the temperature region where the upward air motion is found to be largest. This indicates that the weak upward air motion might indeed be a result of the latent heat released by the intensified depositional growth. If we look again at the spectral edge velocities in Fig. 5b, we see that the slow edge velocity begins to decrease already at a temperature of -18°C which is almost 2°C colder than the temperature where the fast edge is affected by the updraft. At -18°C , the number of new particles appears to increase, which can be also seen in the KDP profile. The latent heat release due to the increasing number and more favourable growth conditions starting at -16°C




Figure 7. Schematic plot combining the main features found in the radar profile statistics (coloured lines) discussed in Sect. 4 and results from laboratory experiments (black lines) in the DGL. The leftmost column denotes temperature regions where specific crystal types are growing (according to TKH91 and TKH14). The abbreviations used in this study and the ones according to the classification of Kikuchi et al. (2013) used in TKH14 denote thick hexagonal plates (TPL, P1b), hexagonal plates (PL, P1a), sectors (SEC, P2a), broad branched (BB, P2b), stellars (STEL, P3a), dendrites (DEN, P3b) and fern-like dendrites (FERN, P3c). (a) The maximum crystal size and (b) aspect ratio (defined as ratio of the maximum dimensions along the a and c axis) which is reached in the experiments by TKH14 (their Table 1) and TKH91 (their Fig. 2) at constant temperature and liquid water saturation after 10 min growth time. The solid line indicates the temperature region in which the experiment was conducted and the dashed line shows the interpolation to warmer/colder temperatures which the fitted function to the measurement predicts. (c) Number of fragments collected after the collision of two ice spheres at different temperatures (fit to values shown in Fig. 14 in TKH95). Median profiles of observed (d) $sZDR_{\max}$ (blue), (e) DWR_{KaW} (orange) and (f) KDP (green) for the largest aggregate class shown in Figs. 4 and 6. Note that the original profiles have been reduced to the main features such as maximum/minimum or strongest vertical change in order to simplify the discussion. The grey shaded area denotes the temperature region where the spectral edge analysis (Sect. 4.2) indicated upward air motion possibly related to latent heat release. The titles in (d–f) indicate the most common particle properties to which the polarimetric variables are related (a more detailed discussion is provided in the text).

might finally be sufficient to also increase buoyancy enough to sustain upward air motion. The updraft might even cause a positive feedback, as its presence enables the particles to grow in the favourable growth region longer. This would further enhance their mass uptake by deposition and increase the latent heat release.

The reverse explanation that an updraft produces local enhancement of supersaturation leading to subsequent nucleation and depositional growth can not be entirely ruled out by the observations. However, it appears rather unlikely that a synoptic or small-scale dynamical feature would be statistically prevalent in this narrow temperature region, despite the large number and variety of cloud and weather conditions included in our statistics. We speculate that one reason why such a latent heat-driven updraft has not been detected so far in numerical weather prediction models might be simply related to the fact that most models do not include an explicit habit prediction scheme which is probably needed to reproduce the intensive growth rate found in the laboratory. However, when simulating a heavy rainfall case, Lee and Baik (2018) found that simulations with a bin microphysics schemes reveal intense latent heat release due to depositional growth. This latent heat release is sufficient to cause an updraft and a positive feedback mechanism. The latent heat release in bulk schemes was found to be substantially weaker.

The profile of KDP, with its sudden increase at -18°C and its nearly linear increase towards -12°C , appears to be more challenging to be explained by updraft and depositional growth features. If depositional growth alone was responsible for the increase in KDP, we would expect the profile to have a similar shape as $sZDR_{\max}$. KDP is well-known to be closely related to the concentration of asymmetric particles (Kumjian, 2013). Primary ice nucleation appears to be rather unlikely as an explanation for increasing KDP in the DGL. The activation of INPs is expected to decrease with warmer temperatures (Kanji et al., 2017) which is opposite to the KDP signature found in our dataset. The increase of KDP towards the bottom of the DGL is even more surprising as ongoing aggregation (increasing DWR_{KaW} values) should reduce the number concentration of KDP producing particles, therefore reducing KDP or at the least keeping KDP constant.

Single scattering computations for realistically shaped crystals and aggregates show that KDP is by approximately a factor of 3 larger for dendrites than for same-sized aggregates (Appendix C). Therefore, the contribution of aggregates to total KDP can not be completely neglected. A simple calculation example for the bottom of the DGL using DWR , Z_e and KDP values for class 3 revealed that aggregates might contribute 1/3 to the total KDP (Appendix C). KDP at W-band can be interpreted in a similar way as commonly done

for lower frequency radars. Unlike for Ze and ZDR, KDP seems to not be affected by non-Rayleigh scattering effects. It should be noted that this behaviour is expected to change for increasing size parameters (e.g. at radar frequencies higher than W-band).

Secondary ice processes (SIP, Korolev and Leisner, 2020; Field et al., 2017) appear to be a more likely explanation for the observed increase in particle concentration. Several in situ studies (e.g. Rangno and Hobbs, 2001; Hobbs and Rangno, 1998, 1990, 1985) have reported fragments of stellar and dendrites in the DGL. The most likely SIP explaining this effect appears to be fragmentation of ice particles when colliding with each other. Unfortunately, this process has so far only been studied in three laboratory studies (Takahashi et al., 1995; Griggs and Choulaton, 1986; Vardiman, 1978). The number of fragments ejected when colliding two ice spheres at constant temperature and water saturation has been found by Takahashi et al. (1995), hereafter TKH95, to be particularly enhanced inside the DGL between -18 and -12 °C. Although the collisional energy from two cm-sized ice spheres used in TKH95 is quite unrealistic for the scenarios observed in our cases, the rapid growth of fragile dendritic arms on the ice spheres observed by TKH95 indicates that such structures are likely to also grow on aggregates which we found to be sedimenting from above into the DGL. It appears quite probable that collision or even touching of two aggregates might already cause fragmentation of the delicate structures growing on their surfaces. Griggs and Choulaton (1986) also reports that dendrite crystals can fragment without collisions simply due to air drag. Recent modelling studies suggest that ice collisional fragmentation can elevate the ice number concentration by three orders of magnitude (Georgakaki et al., 2022).

Clearly, more insights into the competing effects of aggregation and potential particle generation by ice fragmentation need to be explored, for example using model simulations which include habit-dependent growth and also recent formulations of ice fragmentation (Phillips et al., 2018). Also, new laboratory studies constraining the physical basis of ice fragmentation are clearly needed. Future measurement campaigns should combine in-cloud in situ measurements with multi-frequency and polarimetric radar observations in order to further constrain the hypothesised ice microphysical processes.

5.2 Dependency of DWR and polarimetric quantities on cloud top temperature

Previous studies (Trömel et al., 2019; Griffin et al., 2018; Oue et al., 2018) found evidence that the cloud top temperature (CTT) is correlated with polarimetric radar features within the DGL. Griffin et al. (2018) analysed five winter storms at S-band and found an increase of the 80th percentile of KDP values inside the DGL with colder CTT. Similar dependencies were also found for a large set of mid-latitude

clouds (Trömel et al., 2019) and case studies of Arctic mixed-phase clouds (Oue et al., 2018). Griffin et al. (2018) explained the high KDP in the DGL with a high number of irregular crystals or nearly isometric aggregates falling from cloud top. At colder CTT, the primary ice production can be expected to be larger due to the temperature dependence of INP activation (DeMott et al., 2010; Kanji et al., 2017). At $T < -37$ °C, also homogeneous ice nucleation can further increase the number of ice particles. Although we found no clear correlation of aggregation in the DGL with particle concentration or habit of particles sedimenting from aloft in Sect. 4.3, we sorted our profiles with respect to CTT in order to allow a direct comparison with previous studies.

We determined cloud top height using the Ka-band radar because it is the most sensitive radar and it is less affected by signal attenuation compared to the W-band. The temperature information is taken again from the model analysis implemented in Cloudnet. All profiles with continuous values from cloud top down to -10 °C level (to avoid multi-layer cloud cases) were sorted into three 10 °C-wide CTT regimes ranging from -60 to -30 °C (Fig. 8).

Focusing first on the region above the DGL (-30 to -20 °C), the DWR_{KaW} profiles for the different CTT (Fig. 8b) show rather similar values around 0 dB. Also, KDP values for different CTT are relatively similar above the DGL, varying between 0.1 and 0.4 °km $^{-1}$, and showing maximum differences between the CTT regimes of less than 0.2 °km $^{-1}$ (Fig. 8e). A slightly larger separation with CTT regime can be found for $sZDR_{max}$ and ZDR (Fig. 8f, d). Colder CTT seem to be connected to slightly larger ZDR values. The maximum differences in ZDR and $sZDR_{max}$ for the three CTT regimes are both around 0.25 dB. Also Ze at Ka-band (Fig. 8a) shows larger values for colder CTT. KDP, Ze and ZDR show a larger spread between the lower and upper quantile. For Ze, the spread between the quantiles of class 1 (2, 3) reaches approximately 8.75 dB (11.25, 13.7 dB). The quantiles of all classes show large overlap. Similarly, the quantiles of KDP are mostly overlapping and spreading 1.2 °km $^{-1}$. ZDR shows the largest spread relative to the magnitude of the median of 0.5 dB (0.7, 0.8 dB) above the DGL. However, the upper and lower quantiles of the classes differ similar to the medians.

From these observations, we can conclude that the concentration of particles (KDP) and average size of aggregates or polycrystals (DWR_{KaW}) above the DGL does not substantially vary for different CTT. The similarity of the CTT dependency in ZDR and $sZDR_{max}$ rather indicates that the three CTT regimes lead to different shapes of ZDR producing crystals. This might simply be related to different habits that the particles grow into after being nucleated at different CTT. Particles falling from lower CTT have also had more time to grow by deposition, and can hence reach larger sizes and potentially more asymmetric shapes.

Considering that the analysis was performed for the same cases as analysed in Sect. 4, the median profiles appear to




Figure 8. Dependency of (a) Z_e , (b) DWR_{KaW} , (c) MDV, (d) ZDR (e) KDP and (f) $sZDR_{max}$ on cloud top temperature (CTT). The profiles of each radar observable were stratified with temperature and classified into CTT classes. As in Fig. 4, the solid lines depict the median of the distribution in each CTT class. The shaded area indicates the quantiles of the distribution.

be much less dependent on CTT as compared to sorting them with respect to the maximum DWR_{KaW} in the DGL. Except for a clear slow-down in MDV, the signatures inside the DGL, for example the steep increase in $sZDR_{max}$ at -15°C , are much less pronounced or completely absent. Still, certain weak dependencies on CTT can be found in some variables shown in Fig. 8. The total ZDR for profiles starting at colder CTT decreases in the DGL. For the coldest CTT class, ZDR decreases from 0.6 dB at -20°C to 0.4 dB at 10°C . In contrast, for the warmest CTT class, the ZDR profile stays rather constant at 0.4 dB. The KDP values increase stronger for colder CTT regimes. For CTT between -60 and -50°C , KDP increases from around 0.4°km^{-1} at -20°C to 0.7°km^{-1} at -10°C . In contrast, for the warmest CTT class, KDP increases only from 0.2°km^{-1} at -20°C to 0.35°km^{-1} at -10°C . The MDV slow-down is less pronounced for colder CTT (0.03 m s^{-1} for the coldest, 0.11 m s^{-1} for the warmest CTT class). Below -15°C , the particles fall faster for colder CTT (-0.97 m s^{-1} for the coldest CTT class, -0.85 m s^{-1} for the warmer CTT classes). Interestingly, the DWR_{KaW} profiles show no evidence of a strong dependence of aggregation in the DGL on CTT regime. They only increase from 0 dB at -20°C to 1–1.5 dB at -10°C . Interestingly, also $sZDR_{max}$ lacks a clear maximum in the DGL and increases much less than when sorting the profiles with maximum DWR_{KaW} . Only the median profile of the warmest CTT class increases slightly from 0.95 dB at -20°C to 1.3 dB at -10°C . The upper quantile of the

distribution, however, shows a stronger increase for the two warmest CTT classes, reaching 2.29 and 1.88 dB at the bottom of the DGL for class 2 and 3, respectively.

The results obtained by sorting the profiles according to their CTT reveal, similar to Sect. 4, that the main changes of particle concentration, crystal shape and aggregate size take place within the DGL. The role of particles sedimenting from upper layers appear to be small regardless of whether we sort by CTT or maximum DWR_{KaW} . This is in agreement with Dias Neto (2021), who found no clear dependency of aggregation strength on CTT. However, why do we still find larger KDP values for colder CTT if the influence of particles sedimenting from above is weak? Deeper clouds might simply overall provide a more favourable environment (for example larger updrafts, larger super-saturation) for ice particle growth. This might also lead to higher super-saturation inside the DGL. As a result, we expect primary ice nucleation and secondary ice processes to be more enhanced in such an environment. A more intense depositional growth of delicate dendritic structures will most likely also impact number and size of fragments caused by ice collisional fragmentation. As discussed in the previous section, there is growing evidence that fragmentation is a potential source for enhancing particle number concentration in the DGL. The dependence of KDP on CTT found in our study and in previous work might therefore be less related to the larger nucleation rates expected for colder CTT. The sorting by CTT might simply result into a separation of cloud regimes with more or less favourable

growth conditions. This aspect should be further investigated with future campaign datasets that include a large number of reliable humidity profiles (e.g. from frequent radio soundings).

6 Summary and conclusions

A statistical analysis of 3 months of ground-based, triple-frequency (X-, Ka-, W-band) Doppler radar observations combined with polarimetric Doppler W-band observations was conducted at the JOYCE-CF site in order to better understand growth signatures and related processes in the DGL. Similar to previous studies, we find rapid aggregation taking place in the DGL in combination with the formation of a new ice particle mode, most likely associated to dendritic particles.

After classifying the profiles with respect to their maximum average particle size (maximum DWR_{KaW}), we found a substantial reduction of the MDV in the DGL, which is strongest for the highest aggregate class. An analysis of the spectral edge velocities revealed that part of the reduction is due to a new mode of slow falling ice particles, that first appears in the spectra at $-18\text{ }^{\circ}\text{C}$. In addition, an updraft in the order of 0.1 m s^{-1} is revealed by the fast spectral edge velocity with a maximum reached at $-14\text{ }^{\circ}\text{C}$. As suggested by previous studies, it appears most likely that the updraft is a result of latent heat release due to enhanced depositional growth in the DGL.

Clearly, processes in the DGL are strongly tied to temperature. After combining the main signatures revealed by spectral multi-frequency and polarimetric observations with recent laboratory experiments, we derived the following interpretation of particle evolution within the DGL:

- The concentration of particles indicated by KDP continuously increases from -18 to $-12\text{ }^{\circ}\text{C}$. This increase in concentration seems not to be strongly affected by the particle concentration falling from above. Only for the 75 % percentile we find a distinct maximum of KDP at $-12\text{ }^{\circ}\text{C}$, similar to previous studies using lower frequency radars. The temperature where KDP first increases roughly coincides with the temperature where laboratory studies found an increase in the numbers of fragments ejected due to particle collisions. Such an SIP could potentially compensate the loss of particles due to aggregation within the DGL and hence explain the continuous increase of KDP.
- The maximum spectral ZDR ($sZDR_{max}$) indicates that the aspect ratio of dendritic particles strongly increases at $-15\text{ }^{\circ}\text{C}$, coinciding with the temperature of maximum growth rate and aspect ratio measured in the laboratory. Slight temperature shifts between the radar observations and laboratory results can be most likely assigned to particle sedimentation while growing. Similar

as for KDP, no strong difference is found in $sZDR_{max}$ for particles sedimenting into the DGL from above. However, DWR_{KaW} and ZDR indicate that aggregates sedimenting from higher altitudes into the DGL are larger for cases with enhanced aggregation in the DGL.

- The temperature region where the analysis of the spectral edge velocity indicated an updraft coincides with the region of strongest depositional growth and increase in mean aggregate size. This updraft, potentially connected to latent heat release, might cause a positive feedback as it would enhance the residence time of small particles in the favourable growth zone.

Sorting the profiles with respect to cloud top temperature revealed only slight differences in ice particle shape but nearly negligible differences in concentration or mean size for particles entering the DGL from above. The strongest change in concentration, aspect ratio and mean aggregate size is again observed within the DGL. This highlights the importance of processes taking place inside the DGL for evolution of particle concentration and size. Larger aspect ratios and sizes of ice particles falling from above into the DGL and the generally stronger increase of KDP in the DGL for colder CTT might be simply explained by the overall more favourable growth conditions expected for deeper cloud systems, such as higher super-saturation.

Our statistical analysis further revealed that KDP and $sZDR_{max}$ values remain enhanced after leaving the DGL down to $-3\text{ }^{\circ}\text{C}$ where their values rapidly decrease towards the melting layer. Other SIP being active at warmer temperatures might be a potential source for new ice particles that maintain the high KDP values. Intensified aggregation at temperatures warmer than $-3\text{ }^{\circ}\text{C}$ is the most likely explanation for the rapid decrease of polarimetric variables with concurrent increase of DWR. This increase in aggregation might be explained by the strongly increasing thickness of a quasi-liquid layer (QLL) on ice surfaces, which is expected to increase the sticking efficiency of all ice particles.

This study clearly demonstrates the added value of combining different radar approaches including Doppler spectral analysis, high-frequency radar polarimetry and multi-frequency observations for ice microphysical studies. A statistical analysis as presented in this work can provide robust estimates of potential correlations between different radar variables and their specific temperature dependency. Unlike case studies, such a statistical approach also provides an estimation of the natural variability of observables involved. Statistically-based observational process signatures are very useful for evaluating and improving microphysical schemes in weather prediction models (e.g. Karrer et al., 2021; Ori et al., 2020). They are also urgently needed as constraint for recent model developments such as habit-dependent growth (Jensen et al., 2017; Sulia and Kumjian, 2017; Harrington et al., 2013; Hashino and Tripoli, 2007) and Lagrangian Monte Carlo models where the particle history can be traced

(Grabowski et al., 2019; Brdar and Seifert, 2018). Signatures in remote-sensing datasets can also guide new laboratory studies which are inevitably needed to improve our process understanding and microphysical parameterisations in models. Such laboratory studies in addition to in situ measurements or Monte Carlo modelling studies could also provide unique opportunities to validate our findings and the hypothesised ice microphysical processes of this study.

Appendix A: Chirp tables of the vertically pointing W-band and slant polarimetric W-band radar.

The range resolution, number of spectral averages, Doppler velocity resolution and Nyquist range vary with range for the two FMCW W-band radars. This is due to different chirp settings being defined for different range gate regions. The details of the chirp settings applied during TRIPEX-pol are summarised in Tables A1 and A2.

Table A1. Chirp table for the vertical W-band radar installed during the TRIPEX-pol campaign.

Specifications	Chirp 1	Chirp 2	Chirp 3	Chirp 4
Range [m]	215–1475	1482–3986	3999–8151	8165–11998
Range resolution [m]	36.0	12.8	12.8	12.8
Number of spectral averages	13	13	15	11
Doppler velocity resolution [m s^{-1}]	0.04	0.027	0.028	0.029
Nyquist range [m s^{-1}]	± 10.26	± 6.85	± 3.41	± 1.81

Table A2. Chirp table for the polarimetric W-band radar installed during the TRIPEX-pol campaign.

Specifications	Chirp 1	Chirp 2	Chirp 3
Range [m]	107–715	751–5902	5938–17994
Range resolution [m]	35.8	35.8	35.8
Number of spectral averages	28	56	104
Doppler velocity resolution [m s^{-1}]	0.05	0.042	0.039
Nyquist range [m s^{-1}]	± 6.35	± 4.98	± 2.66

Appendix B: Snow particle observations on the surface and zoomed view on Doppler spectra observed on 30 January 2019 at JOYCE-CF

On 30 January 2018, snowfall was observed on the ground at JOYCE-CF. Between 09:00 and 10:00 UTC, large dendritic crystals and aggregates could be photographed (Fig. B1). The pictures are meant to complement the remote-sensing observations from the case study presented in Sect. 3.

For a better visibility of the second spectral mode described in Sect. 3, we show a zoomed view of the spectral Z_e in the DGL (Fig. B2a) and a single Doppler spectrum extracted close to the -14°C level where the second, slow falling mode can be recognised (Fig. B2b).




Figure B1. Example photographs of dendritic crystals and aggregates reaching the surface during the snowfall event occurring on 30 January 2019 at JOYCE. The sample pictures were taken at 09:00 UTC. The long ticks on the scale denote 1 cm while the short ticks denote 1 mm.




Figure B2. Zoom into the spectrogram shown in Fig. 3: (a) spectral Z_e from the Ka-band radar, the magenta line depicts the position of the maximum of the main spectral mode. As in Fig. 3d, the red lines indicate the derived spectral edge velocities. The blue line in (a) shows the temperature from which the single spectrum in (b) was taken.

Appendix C: Comparability of W-band and X-band polarimetric observations and estimation of the contribution of aggregates to KDP

The majority of polarimetric observations in the DGL have been obtained by lower frequency systems (e.g. S-, C-, X-band). Only during recent years, an increasing number of higher frequency polarimetric cloud radars (mainly Ka- and W-band) became available. The use of higher frequencies has a number of advantages, such as larger KDP (increasing with λ^{-1}) for a given particle population. However, also non-Rayleigh scattering effects become increasingly important at higher frequencies which can make the interpretation of high-frequency polarimetric observations more challenging.

Certainly, an in-depth discussion of the differences in polarimetric observations at various frequencies is out of the scope of this study. However, a comparison of the size and frequency dependence of some key radar variables, such as Ze, ZDR and KDP, for single particles can help to understand which variables are more or less affected by non-Rayleigh scattering effects. Figure C1 shows those three variables which were derived using a recent scattering database by Lu et al. (2016) which also contains scattering properties needed for polarimetric quantities. As our focus in this study is on the DGL, we focused on horizontally aligned branched planar crystals and aggregates of stellars (HD-P1d) at an elevation angle of 30° (consistent with our observations during Tripex-pol). In addition to W-band, values are calculated for X-band as this is the lowest frequency included in the database.

Up to 1 mm particle size, the single-particle Ze at X- and W-band are almost identical as expected from Rayleigh theory (Fig. C1a). The slight differences between crystals and aggregates are most likely due to their different mass-size relations. No specific resonance effects are visible at X-band up to 10 mm size, while at W-band we find the first distinct minimum at 3 mm which is close to the wavelength (3.3 mm). As a result, the Ze at W-band is lower than at X-band for particles larger than 1 mm which is the reason for increasing DWR at larger mean size.

Similar resonance phenomena can also be found in ZDR at very similar particle sizes (Fig. C1b). ZDR at X- and W-band are very similar up to 1 mm size with larger values (3 dB) for crystals and smaller values (1.3 dB) for aggregates owing to their lower density and less extreme aspect ratio. The ZDR values at X-band remain relatively constant over the entire size range. At W-band we find, in addition to a strong minimum at 3 mm, two distinct maxima at 2 and 4–5 mm. Similar but overall less extreme resonance phenomena can also be found for the aggregates at W-band. The ZDR at W-band from the aggregates seems also to increase with particle size.

Despite the large values in ZDR, which can be reached at specific diameters due to resonance phenomena at W-band (between 0 and 9 dB), the differences in total ZDR between

X- and W-band can be expected to be relatively small as the extreme values are likely to cancel out when integrating over a PSD (Matrosov, 2021). Even for spectrally resolved ZDR, resonance effects might be difficult to detect as various ice particle sizes are likely to fall into the same Doppler velocity bin due to natural variability in particle shape and orientation. Matrosov (2021) measured ZDR at W- and Ka-band in Arctic clouds and found that the ZDR differences are slightly increasing with ZDR but, on average, the differences are found to be less than 0.5 dB.

Most interesting for the interpretation of the results of our study is the comparison of single-particle KDP shown in Fig. C1c. One can see that after scaling KDP with λ , there are only very small differences found between X- and W-band. Especially, no resonance phenomena as observed for Ze and ZDR can be found for KDP at any size and for both particle types. This is in agreement with Lu et al. (2015), who showed that simulated KDP at cloud radar wavelengths does not exhibit resonance phenomena. Also, measured KDP values at Ka- and W-band reveal only the expected wavelength scaling (Matrosov, 2021). Also the strong increase of KDP with particle size is remarkably similar at X- and W-band (Fig. C1c). As expected, the KDP for crystals is much larger (up to 1 order of magnitude at 4 mm) than for aggregates. However, as shown in the following simple calculation, the contribution of aggregates to the total KDP can usually not be neglected.

In the observational statistics (Fig. 7), we saw that KDP and DWR continuously increase towards the bottom of the DGL. Can those enhanced KDP values maybe entirely be explained by the contribution from aggregates as for example suggested by Moisseev et al. (2015)? We try to shed light on this question with the following simple calculation.

For simplicity, we assume an inverse exponential size distribution for the aggregates of the form,

$$N(D) = N_0 \cdot \exp(-\Lambda D), \quad (\text{C1})$$

with the slope parameter Λ in m^{-1} , the intercept parameter N_0 in m^{-4} and the particle size D . It appears reasonable to assume that at the bottom of the DGL, DWR_{KaW} and Ze are dominated by the contributions from the aggregates. Using the scattering properties shown in Fig. C1, we can directly estimate Λ to be $(2.25 \times 10^{-3})^{-1} \text{m}^{-1}$ for the maximum DWR_{KaW} of 4.3 dB observed at -12°C for the largest aggregate class. With this Λ , we need to assume N_0 to be $5.6 \times 10^4 \text{m}^{-4}$ in order to match the Ka-band Ze of 10.2 dBz at -12°C .

The W-band KDP caused by this aggregate distribution is $0.28^\circ \text{km}^{-1}$ which is roughly one-third of the observed KDP. If we repeat the same calculation with the 75th percentile of the radar variables measured at -12°C , we obtain for a DWR_{KaW} of 6 dB, and Ze of 13 dBz, a KDP of 1.5°km^{-1} . The aggregates contribute still $0.32^\circ \text{km}^{-1}$ (20%) to the total KDP. This high contribution of aggregates to the observed KDP is most likely an upper limit, since we assume the

aggregates to be perfectly horizontally aligned. The naturally occurring tumbling and fluttering of the particles within clouds would reduce the KDP (and ZDR) produced by aggregates.

Unfortunately, we have less constraints on the PSD for the small, presumably dendritic particles at the -12°C level. For the case study shown in Fig. 2, we observed on-the-ground dendrites reaching up to 5 mm size (see example in Fig. B1). Cloud chamber experiments by TH14 show that particle sizes of 1–1.5 mm are reached in the DGL temperature regime after a growth time of 10 min. In order to produce the remaining KDP signal for the median KDP value at -12°C , a concentration of 2500 dendrites per m^3 with 1 mm or 120 m^{-3} with a size of 5 mm would be needed. For the 75th percentile of the KDP observed at -12°C , the concentration would increase to 4150 and 200 m^{-3} , respectively. For comparison, the expected number of ice nucleating particles at -12°C ranges between 1000 and 2000 m^{-3} (e.g. DeMott et al., 2010).




Figure C1. Scattering properties of single particles for X- and W-band calculated with dendrites (branched planar crystals, blue and orange crosses) and aggregates of stellar crystals (HD-P1d, light blue and yellow circles) from the scattering database presented in Lu et al. (2016). The calculation of vertically pointing Ze for a single particle at X- and W-band is shown in (a), the ZDR at X- and W-band and 30° elevation in (b), and the single particle KDP at X- and W-band in (c). KDP was scaled with the wavelength λ to allow a better comparison of the diameter-dependent behaviour at X- and W-band.

Code and data availability. The quality processed level 2 dataset is available on Zenodo under the <https://doi.org/10.5281/zenodo.5025636>, (von Terzi, 2021). The dataset that was used for this statistical analysis (5 min temporal average containing all polarimetric variables and the zenith variables, and variables derived from the Doppler spectra) is available under <https://doi.org/10.5281/zenodo.5025636>, (von Terzi, 2021). Due to the large size, the full level-0 dataset, containing the Doppler spectra of the four radars, is only available on request. The code used to reprocess, classify and plot is available in <https://doi.org/10.5281/zenodo.7050251>, (von Terzi, 2022).

Author contributions. Data analysis, post-processing and generation of figures was performed by LvT with contributions from JDN and AM. LvT and SK conceptualised the methods and interpretation. LvT and DO performed the scattering calculations. LvT and SK prepared the paper with contributions from all co-authors.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Special issue statement. This article is part of the special issue "Fusion of radar polarimetry and numerical atmospheric modelling towards an improved understanding of cloud and precipitation processes (ACP/AMT/GMD inter-journal SI)". It is not associated with a conference.

Acknowledgements. Contributions by José Dias Neto, Davide Ori and Stefan Kneifel were funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under grant KN 1112/2-1 and KN 1112/2-2 as part of the Emmy-Noether Group "Optimal combination of Polarimetric and Triple Frequency radar techniques for Improving Microphysical process understanding of cold clouds" (OPTIMIce). The TRIPEX-pol campaign and work provided by Leonie von Terzi have been supported by the DFG Priority Program SPP2115 "Fusion of Radar Polarimetry and Numerical Atmospheric Modelling Towards an Improved Understanding of Cloud and Precipitation Processes" (PROM) under grant PROM-IMPRINT (project number 408011764). We thank the Regional Computing Center of the University of Cologne (RRZK) for providing computing time on the DFG-funded (Funding number: INST 216/512/IFUGG) High Performance Computing (HPC) system CHEOPS as well as their support. Leonie von Terzi and José Dias Neto also acknowledge support from the Graduate School of Geosciences of the University of Cologne. The authors are indebted to staff of the University of Cologne, research center Jülich, and RPG, especially Birger Bohn, Rainer Haseneder-Lind, Pavel Krobot, Bernhard Pospichal, Avdulah Saljihi, and Kai Schmidt for their help with the installation of the W-band radars, and

Kai Schmidt for the preparation of the scanning polarimetric W-band radar for the TRIPEX-pol campaign.

Financial support. This research has been supported by the Deutsche Forschungsgemeinschaft (grant nos. 408011764, KN 1112/2-1, and KN 1112/2-2).

Review statement. This paper was edited by Matthew Lebsock and reviewed by two anonymous referees.

References

- Andrić, J., Kumjian, M. R., Zrnić, D. S., Straka, J. M., and Melnikov, V. M.: Polarimetric signatures above the melting layer in winter storms: An observational and modeling study, *J. Appl. Meteorol. Clim.*, 52, 682–700, 2013.
- Bailey, M. P. and Hallett, J.: A comprehensive habit diagram for atmospheric ice crystals: Confirmation from the laboratory, AIRS II, and other field studies, *J. Atmos. Sci.*, 66, 2888–2899, 2009.
- Barrett, A. I., Westbrook, C. D., Nicol, J. C., and Stein, T. H. M.: Rapid ice aggregation process revealed through triple-wavelength Doppler spectrum radar analysis, *Atmos. Chem. Phys.*, 19, 5753–5769, <https://doi.org/10.5194/acp-19-5753-2019>, 2019.
- Bechini, R., Baldini, L., and Chandrasekar, V.: Polarimetric radar observations in the ice region of precipitating clouds at C-band and X-band radar frequencies, *J. Appl. Meteorol. Clim.*, 52, 1147–1169, 2013.
- Brdar, S. and Seifert, A.: McSnow: A Monte-Carlo particle model for riming and aggregation of ice particles in a multidimensional microphysical phase space, *J. Adv. Model. Earth Sy.*, 10, 187–206, 2018.
- Bringi, V. N. and Chandrasekar, V.: Polarimetric Doppler Weather Radar, Cambridge University Press, ISBN: 0-521-62384-7, 2001.
- Bringi, V. N., Keenan, T., and Chandrasekar, V.: Correcting C-band radar reflectivity and differential reflectivity data for rain attenuation: A self-consistent method with constraints, *IEEE T. Geosci. Remote*, 39, 1906–1915, 2001.
- Connolly, P. J., Emersic, C., and Field, P. R.: A laboratory investigation into the aggregation efficiency of small ice crystals, *Atmos. Chem. Phys.*, 12, 2055–2076, <https://doi.org/10.5194/acp-12-2055-2012>, 2012.
- DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M., Eidhammer, T., and Rogers, D.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *P. Natl. Acad. Sci. USA*, 107, 11217–11222, 2010.
- Dias Neto, J.: Investigating aggregation in ice and snow clouds using novel combination of triple-frequency cloud radars and radar Doppler spectra, Ph.D. thesis, Universität zu Köln, Cologne, <https://kups.ub.uni-koeln.de/53405/> (last access: 6 September 2022), 2021.
- Dias Neto, J., Kneifel, S., Ori, D., Trömel, S., Handwerker, J., Bohn, B., Hermes, N., Mühlbauer, K., Lenefer, M., and Simmer, C.: The TRIPLE-frequency and Polarimetric radar Experi-

- ment for improving process observations of winter precipitation, *Earth Syst. Sci. Data*, 11, 845–863, <https://doi.org/10.5194/essd-11-845-2019>, 2019.
- Field, P. R.: Bimodal ice spectra in frontal clouds, *Q. J. Roy. Meteor. Soc.*, 126, 379–392, 2000.
- Field, P. R. and Heymsfield, A.: Importance of snow to global precipitation, *Geophys. Res. Lett.*, 42, 9512–9520, 2015.
- Field, P. R., Lawson, R. P., Brown, P. R., Lloyd, G., Westbrook, C., Moisseev, D., Miltenberger, A., Nenes, A., Blyth, A., Choulaton, T., Connolly, P., Buehl, J., Crosier, J., Cui, Z., Dearden, C., DeMott, P., Flossmann, A., Heymsfield, A., Huang, Y., Kalesse, H., Kanji, Z. A., Korolev, A., Kirchgaessner, A., Lasher-Trapp, S., Leisner, T., McFarquhar, G., Philips, V., Stith, J., and Sullivan, S.: Secondary ice production: Current state of the science and recommendations for the future, *Meteor. Mon.*, 58, 7–1, 2017.
- Georgakaki, P., Sotiropoulou, G., Vignon, É., Billault-Roux, A.-C., Berne, A., and Nenes, A.: Secondary ice production processes in wintertime alpine mixed-phase clouds, *Atmos. Chem. Phys.*, 22, 1965–1988, <https://doi.org/10.5194/acp-22-1965-2022>, 2022.
- Giangrande, S. E., Toto, T., Bansemmer, A., Kumjian, M. R., Mishra, S., and Ryzhkov, A. V.: Insights into riming and aggregation processes as revealed by aircraft, radar, and disdrometer observations for a 27 April 2011 widespread precipitation event, *J. Geophys. Res.-Atmos.*, 121, 5846–5863, 2016.
- Görsdorf, U., Lehmann, V., Bauer-Pfundstein, M., Peters, G., Vavriv, D., Vinogradov, V., and Volkov, V.: A 35-GHz polarimetric Doppler radar for long-term observations of cloud parameters – Description of system and data processing, *J. Atmos. Ocean. Tech.*, 32, 675–690, 2015.
- Grabowski, W. W., Morrison, H., Shima, S.-I., Abade, G. C., Dziekan, P., and Pawlowska, H.: Modeling of cloud microphysics: Can we do better?, *B. Am. Meteorol. Soc.*, 100, 655–672, 2019.
- Griffin, E. M., Schuur, T. J., and Ryzhkov, A. V.: A polarimetric analysis of ice microphysical processes in snow, using quasi-vertical profiles, *J. Appl. Meteorol. Clim.*, 57, 31–50, 2018.
- Griggs, D. and Choulaton, T.: A laboratory study of secondary ice particle production by the fragmentation of rime and vapour-grown ice crystals, *Q. J. Roy. Meteor. Soc.*, 112, 149–163, 1986.
- Hallett, J. and Mossop, S.: Production of secondary ice particles during the riming process, *Nature*, 249, 26–28, 1974.
- Harrington, J. Y., Sulia, K., and Morrison, H.: A method for adaptive habit prediction in bulk microphysical models. Part I: Theoretical development, *J. Atmos. Sci.*, 70, 349–364, 2013.
- Hashino, T. and Tripoli, G.: The Spectral Ice Habit Prediction System (SHIPS). Part I: Model description and simulation of the vapor deposition process, *J. Atmos. Sci.*, 64, 2210–2237, 2007.
- Heymsfield, A. J., Schmitt, C., Bansemmer, A., Gettelman, A., Field, P. R., and Liu, C.: Contributions of the liquid and ice phases to global surface precipitation: Observations and global climate modeling, *J. Atmos. Sci.*, 77, 2629–2648, 2020.
- Hildebrand, P. H. and Sekhon, R.: Objective determination of the noise level in Doppler spectra, *J. Appl. Meteorol.*, 13, 808–811, 1974.
- Hobbs, P. V. and Rangno, A. L.: Ice particle concentrations in clouds, *J. Atmos. Sci.*, 42, 2523–2549, 1985.
- Hobbs, P. V. and Rangno, A. L.: Rapid development of high ice particle concentrations in small polar maritime cumuliform clouds, *J. Atmos. Sci.*, 47, 2710–2722, 1990.
- Hobbs, P. V. and Rangno, A. L.: Microstructures of low and middle-level clouds over the Beaufort Sea, *Q. J. Roy. Meteor. Soc.*, 124, 2035–2071, 1998.
- Hogan, R. J., Illingworth, A. J., and Sauvageot, H.: Measuring crystal size in cirrus using 35- and 94-GHz radars, *J. Atmos. Ocean. Tech.*, 17, 27–37, 2000.
- Illingworth, A., Hogan, R., O’connor, E., Bouniol, D., Brooks, M., Delanoë, J., Donovan, D., Eastment, J., Gaussiat, N., Goddard, J. W. F., Haeffelin, M., Klein Baltin, H., Krasnov, O. A., Pelon, J., Piriou, J.-M., Protat, A., Russchenberg, J., Seifert, A., Tompkins, A. M., van Zadelhoff, G.-J., Vinit, F., Willén, U., Wilson, D. R., and Wrench, C. L.: Cloudnet: Continuous evaluation of cloud profiles in seven operational models using ground-based observations, *B. Am. Meteorol. Soc.*, 88, 883–898, 2007.
- Jensen, A. A., Harrington, J. Y., Morrison, H., and Milbrandt, J. A.: Predicting ice shape evolution in a bulk microphysics model, *J. Atmos. Sci.*, 74, 2081–2104, 2017.
- Kalesse, H., Szyrmer, W., Kneifel, S., Kollias, P., and Luke, E.: Fingerprints of a riming event on cloud radar Doppler spectra: observations and modeling, *Atmos. Chem. Phys.*, 16, 2997–3012, <https://doi.org/10.5194/acp-16-2997-2016>, 2016.
- Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Krämer, M.: Overview of ice nucleating particles, *Meteor. Mon.*, 58, 1–1, 2017.
- Karrer, M., Seifert, A., Ori, D., and Kneifel, S.: Improving the representation of aggregation in a two-moment microphysical scheme with statistics of multi-frequency Doppler radar observations, *Atmos. Chem. Phys.*, 21, 17133–17166, <https://doi.org/10.5194/acp-21-17133-2021>, 2021.
- Kennedy, P. C. and Rutledge, S. A.: S-band dual-polarization radar observations of winter storms, *J. Appl. Meteorol. Clim.*, 50, 844–858, 2011.
- Kikuchi, K., Kameda, T., Higuchi, K., and Yamashita, A.: A global classification of snow crystals, ice crystals, and solid precipitation based on observations from middle latitudes to polar regions, *Atmos. Res.*, 132, 460–472, 2013.
- Kneifel, S. and Moisseev, D.: Long-term statistics of riming in nonconvective clouds derived from ground-based Doppler cloud radar observations, *J. Atmos. Sci.*, 77, 3495–3508, 2020.
- Kneifel, S., Kulie, M., and Bennartz, R.: A triple-frequency approach to retrieve microphysical snowfall parameters, *J. Geophys. Res.-Atmos.*, 116, D11203, <https://doi.org/10.1029/2010JD015430>, 2011.
- Kneifel, S., von Lerber, A., Tiira, J., Moisseev, D., Kollias, P., and Leinonen, J.: Observed relations between snowfall microphysics and triple-frequency radar measurements, *J. Geophys. Res.-Atmos.*, 120, 6034–6055, 2015.
- Kneifel, S., Kollias, P., Battaglia, A., Leinonen, J., Maahn, M., Kalesse, H., and Tridon, F.: First observations of triple-frequency radar Doppler spectra in snowfall: Interpretation and applications, *Geophys. Res. Lett.*, 43, 2225–2233, 2016.
- Korolev, A.: Limitations of the Wegener–Bergeron–Findeisen mechanism in the evolution of mixed-phase clouds, *J. Atmos. Sci.*, 64, 3372–3375, 2007.
- Korolev, A. and Leisner, T.: Review of experimental studies of secondary ice production, *Atmos. Chem. Phys.*, 20, 11767–11797, <https://doi.org/10.5194/acp-20-11767-2020>, 2020.
- Küchler, N., Kneifel, S., Löhnert, U., Kollias, P., Czekala, H., and Rose, T.: A W-band radar–radiometer system for accurate and

- continuous monitoring of clouds and precipitation, *J. Atmos. Ocean. Tech.*, 34, 2375–2392, 2017.
- Kumjian, M. R.: Principles and Applications of Dual-Polarization Weather Radar. Part I: Description of the Polarimetric Radar Variables., *Journal of Operational Meteorology*, 1, 226–242, <https://doi.org/10.15191/nwajom.2013.0119>, 2013.
- Lamb, D. and Verlinde, J.: *Physics and chemistry of clouds*, Cambridge University Press, ISBN: 9780511976377, 2011.
- Lawson, R. P., Stewart, R. E., and Angus, L. J.: Observations and numerical simulations of the origin and development of very large snowflakes, *J. Atmos. Sci.*, 55, 3209–3229, 1998.
- Lee, H. and Baik, J.-J.: A comparative study of bin and bulk cloud microphysics schemes in simulating a heavy precipitation case, *Atmosphere*, 9, 475, <https://doi.org/10.1175/JTECH1808.1>, 2018.
- Liao, L., Meneghini, R., Iguchi, T., and Detwiler, A.: Use of dual-wavelength radar for snow parameter estimates, *J. Atmos. Ocean. Tech.*, 22, 1494–1506, 2005.
- Liao, L., Meneghini, R., Tian, L., and Heymsfield, G. M.: Retrieval of snow and rain from combined X-and W-band airborne radar measurements, *IEEE T. Geosci. Remote*, 46, 1514–1524, 2008.
- Li, H. and Moiseev, D.: Melting layer attenuation at Ka-and W-bands as derived from multifrequency radar Doppler spectra observations, *J. Geophys. Res.-Atmos.*, 124, 9520–9533, 2019.
- Löffler-Mang, M. and Joss, J.: An optical disdrometer for measuring size and velocity of hydrometeors, *J. Atmos. Ocean. Techn.*, 17, 130–139, 2000.
- Löhnert, U., Schween, J., Acquistapace, C., Ebell, K., Maahn, M., Barrera-Verdejo, M., Hirsikko, A., Bohn, B., Knaps, A., O’connor, E., Simmer, C., Wahner, A., and Crewell, S.: JOYCE: Jülich observatory for cloud evolution, *B. Am. Meteorol. Soc.*, 96, 1157–1174, 2015.
- Lu, Y., Aydin, K., Clothiaux, E. E., and Verlinde, J.: Retrieving cloud ice water content using millimeter-and centimeter-wavelength radar polarimetric observables, *J. Appl. Meteorol. Clim.*, 54, 596–604, 2015.
- Lu, Y., Jiang, Z., Aydin, K., Verlinde, J., Clothiaux, E. E., and Botta, G.: A polarimetric scattering database for non-spherical ice particles at microwave wavelengths, *Atmos. Meas. Tech.*, 9, 5119–5134, <https://doi.org/10.5194/amt-9-5119-2016>, 2016.
- Mason, S. L., Hogan, R. J., Westbrook, C. D., Kneifel, S., Moiseev, D., and von Terzi, L.: The importance of particle size distribution and internal structure for triple-frequency radar retrievals of the morphology of snow, *Atmos. Meas. Tech.*, 12, 4993–5018, <https://doi.org/10.5194/amt-12-4993-2019>, 2019.
- Matrosov, S. Y.: Radar reflectivity in snowfall, *IEEE T. Geosci. Remote*, 30, 454–461, 1992.
- Matrosov, S. Y., Mace, G. G., Marchand, R., Shupe, M. D., Hallar, A. G., and McCubbin, I. B.: Observations of ice crystal habits with a scanning polarimetric W-band radar at slant linear depolarization ratio mode, *J. Atmos. Ocean. Tech.*, 29, 989–1008, 2012.
- Matrosov, S. Y.: Polarimetric radar variables in snowfall at Ka and W-band frequency bands: A comparative analysis, *J. Atmos. Ocean. Techn.*, 38, 91–101, 2021.
- Mech, M., Maahn, M., Kneifel, S., Ori, D., Orlandi, E., Kollias, P., Schemann, V., and Crewell, S.: PAMTRA 1.0: the Passive and Active Microwave radiative TRANSfer tool for simulating radiometer and radar measurements of the cloudy atmosphere, *Geosci. Model Dev.*, 13, 4229–4251, <https://doi.org/10.5194/gmd-13-4229-2020>, 2020.
- Moiseev, D., Saltikoff, E., and Leskinen, M.: Dual-polarization weather radar observations of snow growth processes, in: 34th Conference on Radar Meteorology, Williamsburg, VA, 5–9 October 2009, 13B.2, https://ams.confex.com/ams/34Radar/techprogram/paper_156123.htm (last access: 5 September 2020), 2009.
- Moiseev, D. N., Lautaportti, S., Tyynela, J., and Lim, S.: Dual-polarization radar signatures in snowstorms: Role of snowflake aggregation, *J. Geophys. Res.-Atmos.*, 120, 12644–12655, 2015.
- Mróz, K., Battaglia, A., Kneifel, S., von Terzi, L., Karrer, M., and Ori, D.: Linking rain into ice microphysics across the melting layer in stratiform rain: a closure study, *Atmos. Meas. Tech.*, 14, 511–529, <https://doi.org/10.5194/amt-14-511-2021>, 2021..
- Mülmenstädt, J., Sourdeval, O., Delanoë, J., and Quaas, J.: Frequency of occurrence of rain from liquid-, mixed-, and ice-phase clouds derived from A-Train satellite retrievals, *Geophys. Res. Lett.*, 42, 6502–6509, 2015.
- Muth, X., Schneebeli, M., and Berne, A.: A sun-tracking method to improve the pointing accuracy of weather radar, *Atmos. Meas. Tech.*, 5, 547–555, <https://doi.org/10.5194/amt-5-547-2012>, 2012.
- Myagkov, A., Seifert, P., Bauer-Pfundstein, M., and Wandinger, U.: Cloud radar with hybrid mode towards estimation of shape and orientation of ice crystals, *Atmos. Meas. Tech.*, 9, 469–489, <https://doi.org/10.5194/amt-9-469-2016>, 2016a.
- Myagkov, A., Seifert, P., Wandinger, U., Bühl, J., and Engelmann, R.: Relationship between temperature and apparent shape of pristine ice crystals derived from polarimetric cloud radar observations during the ACCEPT campaign, *Atmos. Meas. Tech.*, 9, 3739–3754, <https://doi.org/10.5194/amt-9-3739-2016>, 2016b.
- Myagkov, A., Kneifel, S., and Rose, T.: Evaluation of the reflectivity calibration of W-band radars based on observations in rain, *Atmos. Meas. Tech.*, 13, 5799–5825, <https://doi.org/10.5194/amt-13-5799-2020>, 2020.
- Ori, D., Schemann, V., Karrer, M., Dias Neto, J., von Terzi, L., Seifert, A., and Kneifel, S.: Evaluation of ice particle growth in ICON using statistics of multi-frequency Doppler cloud radar observations, *Q. J. Roy. Meteor. Soc.*, 146, 3830–3849, 2020.
- Ori, D., von Terzi, L., Karrer, M., and Kneifel, S.: snowScatt 1.0: consistent model of microphysical and scattering properties of rimed and unrimed snowflakes based on the self-similar Rayleigh–Gans approximation, *Geosci. Model Dev.*, 14, 1511–1531, <https://doi.org/10.5194/gmd-14-1511-2021>, 2021.
- Oue, M., Kumjian, M. R., Lu, Y., Verlinde, J., Aydin, K., and Clothiaux, E. E.: Linear depolarization ratios of columnar ice crystals in a deep precipitating system over the Arctic observed by zenith-pointing Ka-band Doppler radar, *J. Appl. Meteorol. Clim.*, 54, 1060–1068, 2015.
- Oue, M., Kollias, P., Ryzhkov, A., and Luke, E. P.: Toward exploring the synergy between cloud radar polarimetry and Doppler spectral analysis in deep cold precipitating systems in the Arctic, *J. Geophys. Res.-Atmos.*, 123, 2797–2815, 2018.
- Pfützenmaier, L., Unal, C. M. H., Dufournet, Y., and Russchenberg, H. W. J.: Observing ice particle growth along fall streaks in mixed-phase clouds using spectral polarimetric radar data, *Atmos. Chem. Phys.*, 18, 7843–7862, <https://doi.org/10.5194/acp-18-7843-2018>, 2018.

- Phillips, V. T., Patade, S., Gutierrez, J., and Bansemer, A.: Secondary ice production by fragmentation of freezing drops: Formulation and theory, *J. Atmos. Sci.*, 75, 3031–3070, 2018.
- Rangno, A. L. and Hobbs, P. V.: Ice particles in stratiform clouds in the Arctic and possible mechanisms for the production of high ice concentrations, *J. Geophys. Res.-Atmos.*, 106, 15065–15075, 2001.
- Ryzhkov, A., Zhang, P., Reeves, H., Kumjian, M., Tschallener, T., Trömel, S., and Simmer, C.: Quasi-vertical profiles—A new way to look at polarimetric radar data, *J. Atmos. Ocean. Tech.*, 33, 551–562, 2016.
- Schneebeil, M., Dawes, N., Lehning, M., and Berne, A.: High-resolution vertical profiles of X-band polarimetric radar observables during snowfall in the Swiss Alps, *J. Appl. Meteorol. Clim.*, 52, 378–394, 2013.
- Schrom, R. S. and Kumjian, M. R.: Connecting microphysical processes in Colorado winter storms with vertical profiles of radar observations, *J. Appl. Meteorol. Clim.*, 55, 1771–1787, 2016.
- Schrom, R. S., Kumjian, M. R., and Lu, Y.: Polarimetric radar signatures of dendritic growth zones within Colorado winter storms, *J. Appl. Meteorol. Clim.*, 54, 2365–2388, 2015.
- Schwarzenboeck, A., Shcherbakov, V., Lefevre, R., Gayet, J.-F., Pointin, Y., and Duroure, C.: Indications for stellar-crystal fragmentation in Arctic clouds, *Atmos. Res.*, 92, 220–228, 2009.
- Shupe, M. D., Kollias, P., Matrosov, S. Y., and Schneider, T. L.: Deriving mixed-phase cloud properties from Doppler radar spectra, *J. Atmos. Ocean. Tech.*, 21, 660–670, 2004.
- Spek, A., Unal, C., Moisseev, D., Russchenberg, H., Chandrasekar, V., and Dufournet, Y.: A new technique to categorize and retrieve the microphysical properties of ice particles above the melting layer using radar dual-polarization spectral analysis, *J. Atmos. Ocean. Tech.*, 25, 482–497, 2008.
- Sulia, K. J. and Kumjian, M. R.: Simulated polarimetric fields of ice vapor growth using the adaptive habit model. Part I: Large-eddy simulations, *Mon. Weather Rev.*, 145, 2281–2302, 2017.
- Takahashi, T.: Influence of liquid water content and temperature on the form and growth of branched planar snow crystals in a cloud, *J. Atmos. Sci.*, 71, 4127–4142, 2014.
- Takahashi, T., Endoh, T., Wakahama, G., and Fukuta, N.: Vapor diffusional growth of free-falling snow crystals between -3 and -23 °C, *J. Meteorol. Soc. Jpn.*, 69, 15–30, 1991.
- Takahashi, T., Nagao, Y., and Kushiyama, Y.: Possible high ice particle production during graupel–graupel collisions, *J. Atmos. Sci.*, 52, 4523–4527, 1995.
- Thompson, E. J., Rutledge, S. A., Dolan, B., Chandrasekar, V., and Cheong, B. L.: A dual-polarization radar hydrometeor classification algorithm for winter precipitation, *J. Atmos. Ocean. Tech.*, 31, 1457–1481, 2014.
- Trapp, R. J., Schultz, D. M., Ryzhkov, A. V., and Holle, R. L.: Multiscale structure and evolution of an Oklahoma winter precipitation event, *Mon. Weather Rev.*, 129, 486–501, 2001.
- Tridon, F. and Battaglia, A.: Dual-frequency radar Doppler spectral retrieval of rain drop size distributions and entangled dynamics variables, *J. Geophys. Res.-Atmos.*, 120, 5585–5601, 2015.
- Tridon, F., Battaglia, A., and Kneifel, S.: Estimating total attenuation using Rayleigh targets at cloud top: applications in multilayer and mixed-phase clouds observed by ground-based multifrequency radars, *Atmos. Meas. Tech.*, 13, 5065–5085, <https://doi.org/10.5194/amt-13-5065-2020>, 2020.
- Trömel, S., Ryzhkov, A. V., Hickman, B., Mühlbauer, K., and Simmer, C.: Polarimetric radar variables in the layers of melting and dendritic growth at X band – Implications for a nowcasting strategy in stratiform rain, *J. Appl. Meteorol. Clim.*, 58, 2497–2522, 2019.
- Vardiman, L.: The generation of secondary ice particles in clouds by crystal–crystal collision, *J. Atmos. Sci.*, 35, 2168–2180, 1978.
- von Terzi, L.: TRIPEX-pol level-2 data (Version LV2), Zenodo [data set], <https://doi.org/10.5281/zenodo.5025636>, 2021.
- von Terzi, L.: OPTIMICE-team/DGL-analysis-ACP-2022: Analysis of the dendritic growth layer for von Terzi et al. ACP, 2022 (v1.0), Zenodo [code], <https://doi.org/10.5281/zenodo.7050251>, 2022.
- Zawadzki, I.: Observations of snow growth by a vertically pointing radar, in: 36th Conference on Radar Meteorology, Breckenridge Colorado, 16–20 September 2013, 11A.6, <https://ams.confex.com/ams/36Radar/webprogram/Paper229071.html> (last access: 9 September 2022) 2013.
- Zawadzki, I., Fabry, F., and Szyrmer, W.: Observations of supercooled water and secondary ice generation by a vertically pointing X-band Doppler radar, *Atmos. Res.*, 59, 343–359, 2001.
- Zhu, Z., Kollias, P., Yang, F., and Luke, E.: On the Estimation of In-Cloud Vertical Air Motion Using Radar Doppler Spectra, *Geophys. Res. Lett.*, 48, e2020GL090682, <https://doi.org/10.1029/2020GL090682>, 2021.