



Supplement of

Ammonium adduct chemical ionization to investigate anthropogenic oxygenated gas-phase organic compounds in urban air

Peeyush Khare et al.

Correspondence to: Jordan E. Krechmer (jordan.krechmer@bruker.com) and Drew R. Gentner (drew.gentner@yale.edu)

The copyright of individual parts of the supplement might differ from the article licence.

Table of Contents

Figures

- Fig. S1** Aerial visualization of the sampling location at the ASRC (City University of New York).
Fig. S2 Regional map showing radial distances of 10, 20 and 30 kilometers from the sampling site.
Fig. S3 Stability of ammonium-parent ion adducts as a function of voltage and IMR temperatures.
Fig. S4 Benzene-to-toluene ratio over the sampling period.
Fig. S5 Average ambient mass spectra collected using Vocus CI-ToF with NH₄⁺ as the reagent ion.
Fig. S6 Sensitivity analysis for CI-ToF response factors as a function of isomer.
Fig. S7 Comparison between measured concentrations of select analytes by the NH₄⁺-ToF and PTR-ToF instruments over the sampling period.
Fig. S8 Instrument response to ethanol during on-field calibration periods.
Fig. S9 Wind speed and direction at the sampling site during the measurement period.
Fig. S10 Potential chemical losses of different chemical species due to OH oxidation.
Fig. S11 Geometric mean ion abundances of all measured ions averaged over the duration of the sampling campaign.
Fig. S12 Geometric mean of absolute concentrations averaged over the entire duration of the sampling campaign.
Fig. S13 Linear regression slopes vs geometric mean concentration ratios with C₃H₆O.
Fig. S14 Image plot of concentration-based linear correlation coefficients.
Fig. S15 Image plot of ion abundance-based linear correlation coefficients.
Fig. S16 List of measured ion formulas sorted by ions with the highest counts of concentration-based correlation coefficients exceeding 0.9.
Fig. S17 The complete list of measured ion formulas sorted by ions with the highest counts of ion abundance-based correlation coefficients exceeding 0.9.
Fig. S18 Concentration ratios versus annual emissions inventories (compared to acetone) labelled as likely isomers.
Fig. S19 Comparison of correlation coefficients against select tracer compounds using ion abundances of all target species.
Fig. S20 Examples of observed strong linear correlations between C₇H₈O and select other species.

Tables

- Table S1** Chemical composition of the analytical standard cylinder used for mass calibration of select species.
Table S2 Mass response factors of select measured ions.
Table S3 Geometric mean ion abundances measured for all calibrated and non-calibrated ions.
Table S4 Relative importance of different ionization pathways for select oxygenated organic compounds
Table S5 The limits of detection for different calibrated ion formulas.
Table S6 Emissions contributions to select major species in New York City.
Table S7 List of the calibrated ion formulas measured, their possible chemical isomers and examples of uses.
Table S8 List of calibrated ions and their enhancement ratios with tracer compounds with uncertainties over the sampling period.
-

S1. Detailed description of correlations observed between trends in measured ions

To supplement the discussion in the main text and accompany correlation-related figures S9-12, here we use both concentration and ion abundance-based linear regressions to compare correlations between oxygenated species that potentially have similar or co-located sources. Only ions with the strongest correlations ranging from 0.85 to 1 were considered here to reduce the effect of meteorological influences on linear regressions. Compounds are also identified whose dynamics in the atmosphere could be important from an emissions or air quality perspective and warrant further investigation in future studies. Given that a wide range of possible uses exist for many of these compounds, mentions of specific sources are not intended to exclude other potential contributing source types, but rather to identify co-varying compounds (and shared uses) where potentially useful to the reader.

Glycols and glycol ethers

$C_2H_6O_2$, a prominent ion in this study identified primarily as ethylene glycol, showed very good correlations ($r > 0.9$) with 19 other ions, and was most strongly correlated ($r > 0.95$) with C_3H_9NO (e.g. aminopropanol). Most of these were volatile C_xH_yO ions with carbon numbers ≤ 6 including C_3H_6O (e.g. acetone), C_4H_6O (e.g. MVK), $C_6H_{10}O$ (e.g. cyclohexanone) and C_5H_8O (e.g. cyclopentanone), but also $C_{10}H_{12}O$ (e.g. benzyl acetone) and among others. The remaining ones were largely ester-containing $C_xH_yO_2$ ions. These compounds are widely used as solvents, with some also likely constituents of personal care products. This suggested that solvents could be an important contributor of $C_2H_6O_2$ in ambient environments.

Glycol ether-related ions generally did not show very strong correlations with other ions. With the exception of $C_8H_{18}O_2$ (e.g. EGHE) and $C_8H_{18}O_3$ (e.g. dipropylene glycol butyl ether or DGBE), almost all ions in this category were correlated with $r \leq 0.9$. EGHE is used in many consumer and commercial products while DGBE is a popular solvent. The strongest correlations ($r > 0.95$) of $C_8H_{18}O_2$ were with ions representing menthol, linalool and benzyl acetate, which was consistent with the popular use of EGHE in fragrance-related products. EGHE also showed good correlations ($r > 0.9$) with 29 other ions that were mainly constituted by C_xH_yO and $C_xH_yO_4$ ion groups such as C_7H_8O , $C_{11}H_{20}O$, $C_{14}H_{22}O$, $C_{16}H_{30}O_4$, $C_{10}H_{22}O_4$ and $C_9H_{18}O_4$. In the case of DGBE, no ions showed $r > 0.95$ but 17 ions were well-correlated ($r > 0.9$). The highest of these were ions containing EGHE, benzyl alcohol and some oxy-terpenoids such as menthol and linalool.

$C_6H_{12}O_3$ likely included propylene glycol methyl ether acetate (PGMEA) and 2-ethoxyethyl acetate that are widely used in coatings, printing inks and cleaning products. However, only 3 other ions correlated well ($r > 0.9$) with $C_6H_{12}O_3$. These included $C_7H_{14}O_3$ (e.g. ethyl-3-ethoxypropionate), $C_8H_{16}O_3$ (e.g. butoxyethyl acetate) and $C_6H_{12}O$ (e.g. methyl butyl ketone)

while another 33 ions showed an $r > 0.85$. The highest among these remaining ones were C_xH_yO ions containing acetone, MVK and benzyl acetone.

Select personal care product-related compounds

Personal care products also likely contributed to glycol ether-related ion measurements. For example, $C_8H_{10}O_2$ (e.g. phenoxyethanol) was well-correlated ($r > 0.9$) with C_7H_8O , $C_{10}H_{12}O$, $C_{10}H_{18}O$ and $C_{10}H_{20}O$ ions, which represent benzyl alcohol, benzyl acetone, menthol and linalool. This was consistent with two major contributing isomers to $C_8H_{10}O_2$ being phenoxyethanol and 1,4 dimethoxybenzene that are used in personal care products. The isomer contributions were further substantiated by $C_8H_{10}O_2$ concentration peaks that coincided with morning and evening commute hours. Among other species in this category, $C_{10}H_{22}O_4$ (e.g. triethylene glycol butyl ether or TEGBE) most strongly correlated with $C_{14}H_{22}O$ (e.g. α -isomethyl ionone) ($r > 0.95$) that is primarily used in perfumes and other cosmetic products. Other major correlations included ions containing EGHE and benzyl acetate ($r > 0.9$).

Carbonyls

Ions representing carbonyls also showed good correlations with a wide range of ions. C_3H_6O (e.g. acetone) exhibited strong correlations ($r > 0.95$) with C_3H_9NO (e.g. aminopropanol), $C_6H_{10}O$ (e.g. cyclohexanone) and $C_8H_{14}O$ (6-methyl-5-hepten-2-one). However, it was also well-correlated ($r > 0.9$) with 30 more ions that were dominated by C_xH_yO compounds with $C_{\#}$'s ranging from 4 to 12 such as MVK, MEK, cyclopentanone, and benzyl acetone. In addition, $C_xH_yO_2$ ions related to cyclohexyl acetate, ethyl acetoacetate, ethylene glycol and butyl acrylate were also well-correlated with acetone. $C_6H_{10}O$ (e.g. cyclohexanone) correlated strongly with $C_4H_6O_2$, C_5H_8O and $C_8H_{14}O_2$ ions consistent with the popular use of cyclohexanone as a general solvent. $C_6H_{10}O$ also correlated well with $C_8H_{16}O$, $C_{10}H_{12}O$ and $C_6H_{10}O_2$ ions that are related to personal care products.

C_4H_8O (e.g. MEK) exhibited particularly strong correlations ($r > 0.95$) with ions prevalent in oxygenated solvents suggesting connections to solvent-related sources. There likely were smaller contributions from personal care products as evidenced by good correlations with C_3H_9NO , $C_6H_{10}O$, $C_6H_{12}O$ and $C_8H_{14}O$. Other carbonyl-related ions (e.g. $C_6H_{12}O$) were mainly strongly correlated with compounds used in fragrances and cosmetic products. C_7H_6O (e.g. benzaldehyde) showed strongest correlation with C_8H_8O (e.g. acetophenone) which is used in solvents and fragrances but also found in automobile exhaust. It also varied strongly with $C_6H_{12}O$, $C_8H_{16}O$, $C_{10}H_{12}O$ and $C_9H_{10}O_2$ that suggest solvents and cosmetics to be major sources. Overall, this supports large contributions of solvent-related sources to carbonyl-related ions.

Some variability was observed in the number of species with which the oxy-terpenoid-related ions were strongly correlated, suggestive of multiple sources types. For example, $C_{10}H_{20}O$ (e.g. menthol) showed good correlations ($r > 0.9$) with 41 ions. Of these, 14 ions were strongly

correlated ($r > 0.95$) and included glycol ethers, carbonyls, esters and alcohols. $C_{10}H_{18}O$ correlated nearly perfectly with $C_{10}H_{20}O$. On the other hand, $C_{10}H_{16}O$ (e.g. camphor) was well-correlated ($r > 0.9$) with only four ions that included other major oxy-terpenoid-related ions, i.e. $C_{10}H_{20}O$ (e.g. menthol), $C_{10}H_{18}O$ (e.g. linalool), $C_9H_{12}O_2$ (e.g. phenoxy propanol) and $C_{12}H_{20}O_2$ (e.g. geranyl acetate). Correlations with $C_{10}H_{30}O_5Si_5$ (e.g. D5 siloxane) were also around 0.9. Most of these compounds (except D5) can be found in cleaning products, which likely contributed to $C_{10}H_{18}O$ ion signal since it strongly varied with the same ions but also with $C_{11}H_{22}O_2$ and $C_{19}H_{18}O_2$ (e.g. FAMES) and $C_{10}H_{20}O_2$ (e.g. hydroxycitronellal).

On the contrary, no ions correlated very strongly ($r > 0.95$) with D5 siloxane but 13 ions showed $r > 0.9$. Menthol- and linalool-containing ions showed the best correlations with D5 ($r \sim 0.93$) followed by several $C_xH_yO_2$ ions such as $C_{12}H_{20}O_2$ (e.g. α -terpinyl acetate) and $C_9H_{10}O_2$ (e.g. benzyl acetate), which are used in cosmetics and other fragrance-related products. C_7H_8O (e.g. benzyl alcohol) concentrations varied very similarly to several ions containing oxy-terpenoids, monoterpenes, D5 siloxane and acetates that are found in cosmetic products.

Fragrances and related solvents likely also contributed to $C_7H_{14}O_2$ (e.g. amyl acetate) since it showed very strong correlations with $C_4H_6O_2$, $C_8H_{14}O_2$, C_7H_8O , $C_8H_{16}O$ and $C_{10}H_{20}O$ ions. The highest correlations ($r > 0.96$) were observed with $C_8H_{16}O$ (e.g. octanal) and C_7H_8O (e.g. benzyl alcohol) suggesting that fragrance-based products were likely important contributors of $C_7H_{14}O_2$. Similarly, $C_8H_{16}O_2$ (e.g. caprylic acid) also exhibited similar variations as fragrance-related esters and solvents, and correlated strongly with $C_7H_{14}O_2$ suggesting possible similar sources.

$C_{12}H_{24}O_2$ (e.g. lauric acid) was reasonably correlated ($r \sim 0.9$) with $C_8H_{14}O_2$, $C_{10}H_{18}O_2$, $C_8H_{16}O_4$ and $C_8H_{16}O$ that are prominently used in fragrances. Among other semivolatile ions, $C_{18}H_{34}O_2$ (e.g. oleic acid), did not show appreciable correlation with other ions potentially due to variations in emissions pathways, sources and gas-to-particle partitioning effects.

Nitrogen-containing species

Among the nitrogen-containing ions measured, C_2H_7NO (e.g. monoethanolamine) correlated strongly with $C_{10}H_{30}O_5Si_5$ (e.g. D5) except the period when it was influenced by biomass burning. C_2H_7NO also correlated very strongly ($r > 0.9$) with $C_4H_8O_2$ (e.g. ethyl acetate) and with some solvents. Unlike C_2H_7NO , $C_4H_{11}NO_2$ (e.g. diethanolamine) did not show appreciable correlation with tracers of personal care products such as D5 siloxane. This could be due to diethanolamine being primarily used as feedstock for the production of diethanolamides and diethanolamine salts of fatty acids that are used in cosmetic products. Triethanolamine showed very weak correlations ($r < 0.85$) with majority of the measured ions. Its timeseries varied somewhat similarly with only some aldehyde- and ester-containing ions that are prevalent in fragrance-related products (e.g. phenoxyethyl isobutyrate, pentanal). This suggested that triethanolamine is also likely emitted

from personal care products where it sees common usage, though with potential partitioning limitations.

Fatty acid methyl esters

FAME-containing ions strongly correlated with each other suggesting overlap in their sources in urban environments. $C_9H_{18}O_2$ (e.g. methyl octanoate) also correlated strongly with ions containing menthol, octanal, benzyl acetone, hydroxycitronellal and esters that are prevalent in personal care products. Some correlation was also observed with ions containing carbonyl solvents that are used in both consumer and other commercial products suggesting multiple source contributions to $C_9H_{18}O_2$. The same ions formed the best set of correlations for $C_{11}H_{22}O_2$ (e.g. methyl decanoate).

Texanol and TXIB

$C_{12}H_{24}O_3$ (e.g. texanol) which is a tracer for water-based paints. $C_{16}H_{30}O_4$ represents TXIB that is found in a variety of architectural products (Gkatzelis et al., 2021). They did not correlate significantly with each other given some variations in their use. In fact, $C_{12}H_{24}O_3$ correlated considerably ($r > 0.85$) only with the $C_8H_{18}O_2$ ion related to EGHE which is also found in water-based coatings among other wider uses. On the other hand, $C_{16}H_{30}O_4$ showed good correlation with a wider range of compounds and the concentrations of $C_{16}H_{30}O_4$ also exceeded $C_{12}H_{24}O_3$ throughout the measurement period. The strongest similarities in its timeseries were observed with ions containing glycol ethers prevalent in paint-related products, but aromatic esters (e.g. $C_9H_{10}O_2$; benzyl acetate) and some oxy-terpenoids-related ions (e.g. $C_{10}H_{20}O$; menthol, $C_{10}H_{18}O$; linalool) showed good correlation with $C_{16}H_{30}O_4$.

$C_3H_8O_3$ (e.g. glycerol) showed particularly strong correlation with $C_3H_6O_2$ that is related to methyl acetate, propanoic acid and/or ethyl formate. $C_8H_8O_3$ (e.g. methyl paraben) varied similarly as $C_9H_{12}O_2$ (e.g. phenoxy propanol), $C_5H_6O_2$ (e.g. furfuryl alcohol) and C_7H_8O (e.g. benzyl alcohol) that are used in personal care products (except furfuryl alcohol). It also correlated with oxy-terpenoids found in fragrance-related products.



Figure S1. Aerial visualization (© Google Maps) of the sampling location at the ASRC (City University of New York), with view from NNW. The instrument inlet pointed toward downtown Manhattan shown in the distance in the top-right of the image.




Figure S2. Regional map showing radial distances of 10, 20 and 30 kilometers from the sampling site (starred) at the ASRC (City University of New York). Credits: <https://www.calcmaps.com/map-radius/>




Figure S3. Stability of ammonium-parent ion adducts as a function of voltage and IMR temperature for select compounds including highly oxygenated species.




Figure S4. Benzene to toluene ratios over the sampling period where a ratio of 1.8:1 was used as a filter to remove major biomass burning influences. Note: Biomass burning emissions span a range of benzene-to-toluene ratios with some variance (Sheu et al., 2020) and the filter is set to be inclusive of the average ratio of 1.9:1 in Koss et al (Koss et al., 2018). We further note that the enhancement in the benzene-to-toluene ratio during 1/26 – 1/30 may be partly influenced by more aged background air masses coming from predominantly west-northwest direction (See discussion in Section 3.2).




Figure S5. Average ambient air mass spectra collected using Vocus CI-ToF with NH_4^+ as the reagent ion.




Figure S6. Sensitivity analysis of CI-ToF response factors comparing isomer pairs with variations in instrument sensitivity to NH_4^+ ionization (using calibration standards); shown as variations in simulated concentrations of select ion formulas in response to the changes in the magnitude of the mass calibration factor dependent on relative contribution of each isomer in the pair (x-axis).




Figure S7. A comparison between measured concentrations of select analytes by NH₄-ToF and co-located PTR-ToF (Stony Brook University) instruments over the sampling period. The spread observed in comparisons of C₁₀H₁₆ and C₄H₈O could largely be attributed to variations within relative responses to isomers with the different ionization schemes of the two instruments.




Figure S8. Instrument response to ethanol during on-field calibration periods.




Figure S9. Wind speed and direction at the sampling site during the measurement period. The missing data for the on-site weather station included a period of high wind speeds and periodic heavy rain in the afternoon (until 4:30 LT) followed by calm conditions with winds from South-Northeast between 12-7 PM and West-Northwest thereafter as shown by regional data from the LaGuardia weather station.




Figure S10. Potential chemical losses due to OH oxidation, shown as fraction remaining with distance after wintertime daytime atmospheric oxidation at different wind speeds for the chemical species listed in table S2. Note: An aerial distance of 10–15 kilometers from the sampling site at ASRC, CUNY covers the commercial areas of New York City, as well as, parts of New Jersey as shown in Figure S2. Daytime monoterpene and sesquiterpene losses may vary between specific isomers, but are shown here as α -pinene and α -cedrene, respectively.




Figure S11. Geometric mean ion abundances of all measured ions averaged over the duration of the sampling campaign.




Figure S12. Geometric mean of absolute concentrations (i.e. without 5th percentile background subtraction) of mass calibrated ions averaged over the entire duration of the sampling campaign and color coded by their annual emissions as estimated by the VCPy inventory.




Figure S13. Concentration ratios of measured analytes relative to acetone as determined by linear regression slopes (y-axis) compared to background-subtracted geometric mean concentration ratios (x-axis) and colored by correlation coefficient.




Figure S14. Image-based representation of the various degrees of concentration-based linear correlations between mass calibrated compound pairs.




Figure S15. Image-based representation of the various degrees of ion abundance-based linear correlations between all measured ions.




Figure S16. List of measured formulas sorted by ions with the highest counts of concentration-based large correlation coefficients exceeding 0.9. The number of poor correlations (i.e. < 0.5) for each compound is also shown.




Figure S17. The complete list of targeted ion formulas sorted by ions with the highest counts of ion abundance-based correlation coefficients exceeding 0.9. The number of poor correlations (i.e. < 0.5) for each compound is also shown.




Figure S18. Concentration ratios versus annual emissions inventories (compared to acetone), labelled as likely isomers. Note: Supporting figure for Figure 4 showing same data.




Figure S19. Comparison of correlation coefficients against select tracer compounds using ion abundances of all target species. Nearly 35% of the measured ions correlated highly ($0.9 < r < 1$) with benzyl alcohol's ion.




Figure S20. Examples of observed strong linear correlations between C_7H_8O (e.g. benzyl alcohol) and, (a) $C_9H_{10}O_2$ (e.g. benzyl acetate) (b) $C_4H_{10}O_2$ (e.g. 1-methoxy-2-propanol) and (c) $C_8H_{10}O_2$ (e.g. phenoxyethanol).

Table S1. Chemical composition of the analytical standard cylinder (Apel-Riemer Environmental) used for mass calibration of select species.

| Multicomponent calibration mixture in nitrogen | |
|---|---|
| Compound | Concentration (ppb), (uncertainty: 5%) |
| Ethanol | 981 |
| Acetonitrile | 993 |
| Acetone | 989 |
| Acrylonitrile | 987 |
| Isoprene | 954 |
| Methyl vinyl ketone | 997 |
| Methyl ethyl ketone | 994 |
| Benzene | 989 |
| o-Xylene | 1003 |
| a-Pinene | 1000 |
| 1,2,4-Trimethylbenzene | 973 |
| Octamethylcyclotetrasiloxane | 968 |
| Decamethylcyclopentasiloxane | 962 |
| b-Caryophyllene | 99.5 |

Table S2. Mass response factors (and standard deviations) for mass calibration of select measured ions (using analytical standards; e.g. AccuStandard, Sigma-Aldrich). Standard deviations are determined based on linear regressions with multi-point LCS calibration curves. Cylinder-based standard calibrations are shown in italics (i.e. Table S1).

| Analytical standard | Formula | Res. fac. (avg) | Res. fac. (st. dev) |
|---------------------------------------|--|-----------------|---------------------|
| Glycerol | C ₃ H ₈ O ₃ | 2617 | 923 |
| <i>D5-siloxane</i> | C ₁₀ H ₃₀ O ₅ Si ₅ | 2200 | 110 |
| Triethylene glycol | C ₆ H ₁₄ O ₄ | 1876.504 | 71 |
| Ethyl acetoacetate | C ₆ H ₁₀ O ₃ | 1548.422 | 112 |
| Furfural | C ₅ H ₄ O ₂ | 1522 | 64 |
| Ethylene glycol hexyl ether | C ₈ H ₁₈ O ₂ | 1320.445 | 40 |
| Cyclohexanone | C ₆ H ₁₀ O | 1042.756 | 69 |
| Hydroxycitronellal | C ₁₀ H ₂₀ O ₂ | 1007.586 | 63 |
| Texanol | C ₁₂ H ₂₄ O ₃ | 995.1243 | 65 |
| Acetophenone | C ₈ H ₈ O | 937.9794 | 20 |
| Benzyl acetone | C ₁₀ H ₁₂ O | 761.2294 | 86 |
| Pentadioic acid, dimethyl ester | C ₇ H ₁₂ O ₄ | 737.1291 | 60 |
| Propylene glycol methyl ether acetate | C ₆ H ₁₂ O ₃ | 679.011 | 55 |
| 1-Methoxy-2-propanol | C ₄ H ₁₀ O ₂ | 645.9141 | 69 |
| n-Butyl acetate | C ₆ H ₁₂ O ₂ | 521.7066 | 34 |
| 2-Butoxyethanol | C ₆ H ₁₄ O ₂ | 495.2152 | 40 |
| 2-(2-Butoxyethoxy)ethanol | C ₈ H ₁₈ O ₃ | 450.3138 | 184 |
| 6-Methyl 5-hepten-2-one | C ₈ H ₁₄ O | 424.3785 | 11 |
| Propyl paraben | C ₁₀ H ₁₂ O ₃ | 389.7354 | 74 |
| Maltol | C ₆ H ₆ O ₃ | 380.425 | 37 |
| Dimethyl phthalate | C ₁₀ H ₁₀ O ₄ | 366.3753 | 23 |
| Cyclopentanone | C ₅ H ₈ O | 355.6696 | 9 |
| <i>Methyl ethyl ketone</i> | C ₄ H ₈ O | 350 | 18 |
| 2-Hexanone | C ₆ H ₁₂ O | 345.8968 | 18 |
| Phenoxy ethanol | C ₈ H ₁₀ O ₂ | 345.677 | 119 |
| Methyl pyrrolidone | C ₅ H ₉ NO | 331.8216 | 41 |
| Benzyl alcohol | C ₇ H ₈ O | 329.976 | 23 |
| Ethyl paraben | C ₉ H ₁₀ O ₃ | 323.4804 | 73 |
| Dipropylene glycol | C ₆ H ₁₄ O ₃ | 303.4106 | 43 |
| 2-Methoxy ethanol | C ₃ H ₈ O ₂ | 295.6414 | 30 |
| Methyl methacrylate | C ₅ H ₈ O ₂ | 260 | 13 |
| <i>Acetone</i> | C ₃ H ₆ O | 251 | 13 |
| <i>MVK</i> | C ₄ H ₆ O | 250 | 12 |
| Butyl paraben | C ₁₁ H ₁₄ O ₃ | 238.0273 | 43 |
| Isopropyl acetate | C ₅ H ₁₀ O ₂ | 209.0689 | 23 |
| Benzaldehyde | C ₇ H ₆ O | 203.1 | 14 |
| Methyl paraben | C ₈ H ₈ O ₃ | 199.2943 | 53 |
| Menthol | C ₁₀ H ₂₀ O | 180.1444 | 11 |
| <i>D4-siloxane</i> | C ₈ H ₂₄ O ₄ Si ₄ | 170.36 | 52.26 |
| Diethylene glycol | C ₄ H ₁₀ O ₃ | 160.6576 | 28 |
| Eugenol | C ₁₀ H ₁₂ O ₂ | 148.3496 | 7 |
| Diethyl phthalate | C ₁₂ H ₁₄ O ₄ | 140.7306 | 8 |
| TXIB | C ₁₆ H ₃₀ O ₄ | 131.0738 | 14 |
| Dipropylene glycol propyl ether | C ₉ H ₂₀ O ₃ | 118.3448 | 15 |
| <i>Acetonitrile</i> | C ₂ H ₃ N | 115 | 6 |
| Methyl acetate | C ₃ H ₆ O ₂ | 113.0 | 6 |
| Cyclohexyl acetate | C ₈ H ₁₄ O ₂ | 112.9 | 6 |
| Benzyl acetate | C ₉ H ₁₀ O ₂ | 104.4707 | 10 |
| 2-Phenoxyethyl isobutyrate | C ₁₂ H ₁₆ O ₃ | 100.8308 | 9 |
| Butyric acid | C ₄ H ₈ O ₂ | 100.464 | 19 |
| Methyl benzoate | C ₈ H ₈ O ₂ | 81.46912 | 6 |
| <i>Monoterpenes</i> | C ₁₀ H ₁₆ | 70 | 3 |
| Benzoic acid | C ₇ H ₆ O ₂ | 57.82037 | 4 |
| Octanoic acid | C ₈ H ₁₆ O ₂ | 56.12189 | 4 |
| Methyl octanoate | C ₉ H ₁₈ O ₂ | 38.58139 | 3 |
| Vinyl acetate | C ₄ H ₆ O ₂ | 28 | 2 |
| <i>Sesquiterpenes</i> | C ₁₅ H ₂₄ | 27 | 1 |
| Ethylene glycol | C ₂ H ₆ O ₂ | 11.48517 | 1 |

Note: In some cases, response factors were calculated by averaging the responses for isomer standards. These include C₈H₁₈O₂ (EGHE and 1,2 octanediol), C₆H₁₂O₃ (PGMEA and 2-ethoxyethyl acetate), C₈H₁₈O₃ (2-(2-butoxyethoxy) ethanol and diethylene glycol butyl ether), C₆H₁₂O (2-hexanone and 4-methyl-2-pentanone), C₅H₁₀O₂ (Isopropyl acetate and n-propyl acetate) and C₄H₈O₂ (Butyric acid and ethyl acetate).

Table S3. Geometric mean ion abundances (ions s⁻¹) and standard deviations measured for all calibrated and non-calibrated ions.

| Ion formula | Mean abundance ± standard deviation | Ion formula | Mean abundance ± standard deviation | Ion formula | Mean abundance ± standard deviation |
|--|-------------------------------------|---|-------------------------------------|---|-------------------------------------|
| C ₃ H ₆ O | 239.76±183.96 | C ₆ H ₈ O | 6.46±3.06 | C ₄ H ₁₁ NO ₂ | 2.62±1.11 |
| C ₄ H ₈ O ₂ | 68.26±66.72 | C ₅ H ₁₀ O ₃ | 6.29±3.34 | C ₈ H ₁₀ O | 2.57±0.82 |
| C ₆ H ₁₂ O ₂ | 53.77±71.95 | C ₉ H ₁₈ O | 6.25±2.82 | C ₁₄ H ₂₂ O | 2.52±0.74 |
| C ₃ H ₆ O ₂ | 49.2±42.65 | C ₉ H ₁₄ O | 6.13±2.57 | C ₄ H ₇ NO | 2.51±0.49 |
| C ₄ H ₈ O | 43.98±28.61 | C ₅ H ₈ O ₄ | 6.05±2.1 | C ₁₀ H ₁₀ O ₄ | 2.42±0.54 |
| C ₂ H ₄ O ₂ | 36.06±21.25 | C ₃ H ₇ NO ₂ | 5.92±2.61 | C ₁₀ H ₂₂ O ₄ | 2.4±0.74 |
| C ₁₀ H ₃₀ O ₅ Si ₅ | 35.12±33.31 | C ₈ H ₁₄ O | 5.74±2.96 | C ₁₀ H ₁₂ O ₂ | 2.38±0.76 |
| C ₃ H ₈ O | 29.66±19.76 | C ₄ H ₄ O ₄ | 5.59±1.59 | C ₁₈ H ₃₆ O ₂ | 2.33±0.16 |
| C ₂ H ₃ N | 28.91±12.8 | C ₁₀ H ₂₀ O | 5.57±3.22 | C ₇ H ₆ O ₃ | 2.32±0.22 |
| C ₅ H ₈ O ₂ | 28.07±31.59 | C ₇ H ₁₀ O | 5.35±3.01 | C ₁₀ H ₂₂ O ₃ | 2.31±0.69 |
| C ₂ H ₆ O ₂ | 27.99±41.6 | C ₄ H ₁₀ O ₃ | 5.11±2.77 | C ₁₂ H ₂₄ O | 2.31±0.73 |
| C ₂ H ₅ NO | 27.5±5.14 | C ₈ H ₁₆ O ₂ | 4.89±2.65 | C ₁₆ H ₃₀ O ₄ | 2.3±0.57 |
| C ₃ H ₈ O ₂ | 24.32±14.97 | C ₈ H ₁₄ O ₂ | 4.85±2.3 | C ₈ H ₁₈ O | 2.29±0.73 |
| C ₅ H ₁₀ O ₂ | 23.79±22.06 | C ₇ H ₈ O | 4.7±2.05 | C ₁₇ H ₃₄ O ₂ | 2.2±0.44 |
| C ₄ H ₆ O ₂ | 22.68±11.09 | C ₆ H ₁₀ O ₄ | 4.62±2.15 | C ₆ H ₁₄ O ₄ | 2.17±0.65 |
| C ₅ H ₁₀ O | 21.12±9.91 | C ₄ H ₉ NO | 4.35±3.29 | C ₆ H ₁₅ NO | 2.12±0.9 |
| C ₆ H ₁₂ O | 20.24±14.68 | C ₇ H ₁₆ O ₂ | 4.34±3.08 | C ₈ H ₁₄ O ₄ | 2.11±0.51 |
| C ₅ H ₄ O ₂ | 19.93±9.62 | C ₆ H ₆ O ₃ | 4.18±0.97 | C ₈ H ₁₆ O ₃ | 2.11±0.66 |
| C ₄ H ₆ O ₃ | 19.25±9.42 | C ₆ H ₆ O | 4.07±0.94 | C ₁₂ H ₁₄ O ₄ | 2.11±0.42 |
| C ₁₀ H ₁₆ O | 17.72±8.34 | C ₉ H ₁₀ O ₂ | 4.07±2.01 | C ₆ H ₇ NO ₂ | 2.08±0.26 |
| C ₂ H ₆ O | 16.29±6.17 | C ₅ H ₁₂ O ₂ | 3.97±1.83 | C ₁₁ H ₂₂ O ₂ | 2.07±0.55 |
| C ₃ H ₆ O ₃ | 15.28±7.64 | C ₈ H ₂₄ O ₄ Si ₄ | 3.91±0.6 | C ₁₄ H ₂₀ O | 2.05±0.58 |
| C ₇ H ₆ O | 15.03±7.09 | C ₉ H ₁₈ O ₄ | 3.64±1.86 | C ₁₅ H ₂₂ O ₂ | 1.97±0.4 |
| C ₄ H ₆ O | 13.89±8.85 | C ₉ H ₁₆ O ₂ | 3.61±1.38 | C ₁₁ H ₂₆ O ₅ Si | 1.94±0.5 |
| C ₆ H ₁₀ O ₂ | 13.75±9.21 | C ₇ H ₁₆ O ₃ | 3.57±1.82 | C ₉ H ₁₀ O ₃ | 1.93±0.34 |
| C ₇ H ₁₄ O | 13.47±9.28 | C ₈ H ₁₈ O ₃ | 3.46±1.73 | C ₁₈ H ₃₄ O ₂ | 1.93±0.69 |
| C ₃ H ₄ O | 13.13±3.82 | C ₇ H ₆ O ₂ | 3.42±0.5 | C ₉ H ₂₀ O ₃ | 1.91±0.47 |
| C ₆ H ₁₀ O | 12.39±6.54 | C ₄ H ₁₁ NO | 3.38±1.66 | C ₁₃ H ₂₄ | 1.87±0.29 |
| C ₆ H ₁₄ O ₂ | 11.62±9.17 | C ₇ H ₁₂ O ₃ | 3.38±1.29 | C ₁₈ H ₂₆ O | 1.84±0.34 |
| C ₆ H ₁₄ O ₃ | 10.87±8.58 | C ₁₁ H ₂₀ O | 3.35±1.64 | C ₁₁ H ₁₄ O ₂ | 1.81±0.38 |
| CH ₆ O ₅ Si | 10.85±4.6 | C ₅ H ₁₂ O ₃ | 3.34±1.34 | C ₁₆ H ₃₄ O | 1.8±0.21 |
| C ₅ H ₈ O | 10.56±5.52 | C ₃ H ₈ O ₃ | 3.31±1.37 | C ₁₄ H ₂₈ O ₂ | 1.78±0.24 |
| C ₁₀ H ₁₆ | 10.55±6.75 | C ₈ H ₁₈ O ₂ | 3.27±1.39 | C ₁₂ H ₂₄ O ₂ | 1.78±0.27 |
| C ₈ H ₈ O | 9.82±5.9 | C ₁₀ H ₁₈ O ₂ | 3.23±1.19 | C ₁₀ H ₁₂ O ₃ | 1.74±0.42 |
| C ₅ H ₆ O ₂ | 9.1±3.23 | C ₁₀ H ₁₂ O | 3.19±1.43 | C ₈ H ₈ O ₄ | 1.72±0.27 |
| C ₂ H ₄ O | 8.67±1.77 | C ₁₂ H ₂₀ O ₂ | 3.14±1.19 | C ₁₃ H ₂₂ O | 1.71±0.33 |
| C ₁₀ H ₁₈ O | 8.58±5.34 | C ₈ H ₁₀ O ₂ | 3±1.07 | C ₁₃ H ₁₀ O | 1.69±0.39 |
| C ₆ H ₁₂ O ₃ | 8.45±7.3 | C ₈ H ₄ O ₃ | 2.98±0.92 | C ₆ H ₁₅ NO ₃ | 1.66±0.41 |
| C ₄ H ₁₀ O ₂ | 8.29±5.17 | C ₉ H ₁₈ O ₂ | 2.96±0.91 | C ₁₂ H ₁₆ O ₃ | 1.65±0.22 |
| C ₃ H ₉ NO | 7.62±3.67 | C ₇ H ₁₄ O ₃ | 2.94±1.41 | C ₁₈ H ₃₂ O ₂ | 1.62±0.46 |
| C ₈ H ₈ O ₂ | 7.49±1.25 | C ₅ H ₉ NO | 2.85±1.07 | C ₈ H ₁₆ O ₄ | 1.62±0.27 |
| C ₁₀ H ₁₆ O ₂ | 7.41±4.34 | C ₉ H ₁₄ O ₃ | 2.82±0.97 | C ₉ H ₂₀ O ₄ | 1.58±0.32 |
| C ₅ H ₆ O | 7.22±2.94 | C ₉ H ₁₂ O ₂ | 2.81±1 | C ₉ H ₁₉ NO ₄ | 1.57±0.51 |
| C ₈ H ₁₆ O | 6.92±3.2 | C ₁₆ H ₃₂ O ₂ | 2.75±0.43 | C ₁₅ H ₂₄ O | 1.56±0.24 |
| C ₆ H ₁₀ O ₃ | 6.8±3.59 | C ₈ H ₈ O ₃ | 2.69±0.8 | C ₁₁ H ₁₆ O ₂ | 1.53±0.26 |
| C ₆ H ₆ O ₂ | 6.79±3.34 | C ₁₀ H ₂₀ O ₂ | 2.68±0.84 | C ₁₂ H ₁₈ O ₃ | 1.5±0.23 |
| C ₇ H ₁₂ O ₂ | 6.71±3.96 | C ₁₀ H ₁₄ O | 2.66±0.65 | C ₆ H ₁₀ O ₆ | 1.46±0.23 |
| C ₇ H ₁₄ O ₂ | 6.69±3.85 | C ₂ H ₇ NO | 2.65±1.42 | C ₁₁ H ₁₄ O ₃ | 1.4±0.27 |
| C ₄ H ₁₀ O | 6.63±3.77 | C ₇ H ₁₂ O ₄ | 2.62±0.84 | C ₁₁ H ₂₂ O ₄ | 1.25±0.17 |
| C ₁₂ H ₂₄ O ₃ | 6.58±3.95 | C ₁₁ H ₂₂ O | 2.62±0.99 | | |

Table S4. Relative importance of different ionization pathways for select oxygenated organic compounds during Vocus-NH₄⁺ ionization.

| Standard compound | [M+NH ₄] ⁺ | [M+H] ⁺ | [M] ⁺ |
|-----------------------------------|-----------------------------------|--------------------|------------------|
| Methyl benzoate | 100% | 0% | 0% |
| Cyclohexyl acetate | 100% | 0% | 0% |
| 2-Butoxy ethanol | 100% | 0% | 0% |
| Glycerol | 100% | 0% | 0% |
| Ethylene glycol hexyl ether | 99.8% | 0.2% | 0% |
| 6-methyl-5-hepten-2-one | 63% | 37% | 0% |
| 2-Hexanone | 100% | 0% | 0% |
| PGMEA | 88% | 12% | 0% |
| Pentanedioic acid dimethyl ester | 86% | 14% | 0% |
| 1-Methoxy-2-propanol | 96% | 1% | 3% |
| Di(propylene glycol) propyl ether | 97% | 3% | 0% |
| Benzyl alcohol | 100% | 0% | 0% |

Note: The presence of an unsaturated double bond (C=C) in 6-methyl-5-hepten-2-one results in significant ionization via the proton-transfer ionization pathway (Xu et al., 2022).

Table S5. Limits of detection (in parts per trillion, ppt) for different mass calibrated ion formulas calculated using 1 minute averages of zero periods that occurred 79 times over a 24-hour sampling period.

| Compound formula, i | Probable compounds, i | Limit of detection, ppt |
|--|---|-------------------------|
| C ₄ H ₆ O ₂ | Vinyl acetate, Methyl acrylate | 228 |
| C ₃ H ₆ O | Acetone | 203 |
| C ₂ H ₆ O ₂ | Ethylene glycol | 161 |
| C ₄ H ₈ O ₂ | Ethyl acetate, Butyric acid | 75 |
| C ₂ H ₃ N | Acetonitrile | 65 |
| C ₃ H ₆ O ₂ | Methyl acetate, Propionic acid, Hydroxyacetone, Ethyl formate | 32 |
| C ₈ H ₈ O ₂ | Methyl benzoate | 31 |
| C ₄ H ₆ O | MVK, MACR | 26 |
| C ₉ H ₁₈ O ₂ | Methyl octanoate | 19 |
| C ₇ H ₆ O ₂ | Benzoic acid | 19 |
| C ₄ H ₈ O | MEK, THF, Cyclopropyl carbinol | 18 |
| C ₁₅ H ₂₄ | Sesquiterpenes | 18 |
| C ₇ H ₆ O | Benzaldehyde | 16 |
| C ₁₀ H ₁₆ | Monoterpenes | 14 |
| C ₄ H ₁₀ O ₃ | Diethylene glycol | 11 |
| C ₅ H ₈ O ₂ | Methyl methacrylate | 11 |
| C ₃ H ₈ O ₂ | 2-Methoxy ethanol, propylene glycol | 7.6 |
| C ₅ H ₁₀ O ₂ | Isopropyl acetate, n-propyl acetate | 7.4 |
| C ₈ H ₂₄ O ₄ Si ₄ | D4-siloxane | 6.9 |
| C ₈ H ₁₆ O ₂ | Caprylic acid (i.e., Octanoic acid) | 6.3 |
| C ₉ H ₁₀ O ₂ | Benzyl acetate | 6.2 |
| C ₆ H ₁₂ O | 2-Hexanone, 4-Methyl-2-pentanone | 4.7 |
| C ₅ H ₈ O | Cyclopentanone | 4.3 |
| C ₆ H ₆ O ₃ | Maltol | 4.1 |
| C ₁₀ H ₃₀ O ₅ Si ₅ | D5-siloxane | 4.0 |
| C ₈ H ₁₄ O ₂ | Cyclohexyl acetate | 4.0 |
| C ₇ H ₈ O | Benzyl alcohol | 3.7 |
| C ₆ H ₁₄ O ₃ | Dipropylene glycol | 3.6 |
| C ₅ H ₄ O ₂ | Furfural | 3.3 |
| C ₁₀ H ₁₂ O ₂ | Eugenol | 3.1 |
| C ₁₂ H ₁₆ O ₃ | 2-Phenoxyethyl isobutyrate | 3.1 |
| C ₈ H ₈ O ₃ | Methyl paraben | 2.9 |
| C ₁₆ H ₃₀ O ₄ | TXIB | 2.9 |
| C ₁₂ H ₁₄ O ₄ | Diethyl phthalate | 2.9 |
| C ₆ H ₁₂ O ₂ | n-Butyl acetate | 2.8 |
| C ₁₀ H ₂₀ O | Menthol | 2.6 |
| C ₁₀ H ₁₀ O ₄ | Dimethyl phthalate | 1.7 |
| C ₉ H ₂₀ O ₃ | Dipropylene glycol propyl ether | 1.7 |
| C ₉ H ₁₀ O ₃ | Ethyl paraben | 1.5 |
| C ₄ H ₁₀ O ₂ | 1-Methoxy-2-propanol | 1.5 |
| C ₈ H ₈ O | Acetophenone | 1.3 |
| C ₁₁ H ₁₄ O ₃ | Butyl paraben | 1.2 |
| C ₈ H ₁₀ O ₂ | Phenoxy ethanol | 1.2 |
| C ₈ H ₁₄ O | 6-Methyl 5-hepten-2-one | 1.1 |
| C ₁₀ H ₁₂ O ₃ | Propyl paraben | 1.1 |
| C ₆ H ₁₀ O | Cyclohexanone | 1.0 |
| C ₅ H ₉ NO | Methyl pyrrolidone | 0.9 |
| C ₆ H ₁₂ O ₃ | Propylene glycol methyl ether acetate | 0.9 |
| C ₁₂ H ₂₄ O ₃ | Texanol | 0.9 |
| C ₃ H ₈ O ₃ | Glycerol | 0.8 |
| C ₈ H ₁₈ O ₃ | 2-(2-Butoxyethoxy)ethanol, DGBE | 0.8 |
| C ₆ H ₁₄ O ₂ | 2-Butoxyethanol, 1-propoxy-2-propanol | 0.8 |
| C ₇ H ₁₂ O ₄ | Pentadiolic acid, dimethyl ester | 0.7 |
| C ₆ H ₁₀ O ₃ | Ethyl acetoacetate | 0.6 |
| C ₁₀ H ₂₀ O ₂ | Hydroxycitronellal | 0.6 |
| C ₁₀ H ₁₂ O | Benzyl acetone | 0.5 |
| C ₆ H ₁₄ O ₄ | Triethylene glycol | 0.5 |
| C ₈ H ₁₈ O ₂ | Ethylene glycol hexyl ether | 0.2 |

Note: While the hourly measurements used in this study will utilize a greater number of points for averaging, the ion signal averages over longer (> 1 min) durations are not expected to yield significant differences in the detection limits (Bertram et al., 2011; Novak, Vermeuel, & Bertram, 2020).

Table S6. Emissions contributions to select major species in New York City calculated using VCPy, anthropogenic emissions from the National Emissions Inventory (NEI) and SPECIATE databases. **Note:** Fractions of contribution are calculated using total emissions of all isomers in each source category. This table includes all table 1 compounds with emissions in the NEI inventory and were calibrated in this study.

| Compounds | % Contributions from individual source types [†] | | | | |
|---|---|---------|----------|-------|-----------------|
| | Point | On-road | Non-road | VCPs | Other non-point |
| Ethylene glycol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Acetone | 0.1 | 0.2 | 0.4 | 99.0 | 0.3 |
| Ethyl acetate, Butyric acid | 0.0 | 0.0 | 0.0 | 92.5 | 7.5 |
| Methyl acetate, Propionic acid, Hydroxyacetone, Ethyl formate | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Monoterpenes (e.g., limonene, α -Pinene) | 0.0 | 0.2 | 0.0 | 99.8 | 0.0 |
| MEK, THF, Cyclopropyl carbinol | 0.1 | 6.6 | 4.8 | 72.0 | 16.5 |
| Isopropyl acetate, n-propyl acetate | 0.0 | 0.0 | 0.0 | 82.3 | 17.7 |
| Methyl methacrylate | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Butyl acetate | 0.0 | 0.0 | 0.0 | 92.9 | 7.1 |
| 2-Methoxy ethanol, propylene glycol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| 2-Hexanone, 4-Methyl-2-pentanone | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Dipropylene glycol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| 2-Butoxyethanol*, 1-propoxy-2-propanol | 0.0 | 0.0 | 0.0 | 98.6 | 1.4 |
| D4 siloxane | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| D5 siloxane | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Benzyl alcohol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| 1-Methoxy-2-propanol | 0.1 | 0.0 | 0.0 | 95.7 | 4.3 |
| Cyclohexanone | 0.0 | 0.0 | 0.0 | 99.9 | 0.1 |
| PGMEA, 2-Ethoxyethyl acetate | 0.0 | 0.0 | 0.0 | 97.0 | 3.0 |
| Methyl pyrrolidone | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Phenoxyethanol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| 2-(2-Butoxyethoxy)ethanol, DGBE | 0.0 | 0.0 | 0.0 | 99.4 | 0.6 |
| Dimethyl phthalate | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Benzaldehyde | 0.0 | 36 | 58 | 4.5 | 0.6 |
| Texanol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Pentadioic acid, dimethyl ester | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Ethylene glycol hexyl ether, 1,2-Octanediol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Glycerol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Triethylene glycol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Diethylene glycol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Menthol | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |
| Dipropylene glycol propyl ether | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 |

[†]Source types are defined as per the National Emissions Inventory (NEI) framework. Point sources include industrial and non-industrial commercial facilities; Other non-point sources include cooking, residential heating, commercial combustion, asphalt paving and other commercial and consumer solvent use; On-road sources are primarily heavy- and light-duty vehicles driven on gasoline, diesel and other fuels; Non-road sources include off-road vehicles and equipment that use gasoline, diesel or other fuels, and, VCPs are volatile chemical products used in both commercial and consumer applications.

Table S7. List of the calibrated ion formulas measured using the NH_4^+ -ToF instrument, their possible chemical names and examples of known uses. Listed examples (for calibrated compounds and other potential isomers) are not intended to be comprehensive, but provide examples of common uses that can aid in the interpretation of observations.

| Mass calibrated ions | Analytical standards used | Other likely contributing isomers | Example uses/sources of all listed isomers |
|---|---|---|--|
| $\text{C}_{10}\text{H}_{30}\text{O}_5\text{Si}_5$ | D5-siloxane | | Cosmetic products, antiperspirants |
| $\text{C}_6\text{H}_{14}\text{O}_4$ | Triethylene glycol | | Automobile fluids, lubricants, solvents |
| $\text{C}_6\text{H}_{10}\text{O}_3$ | Ethyl acetoacetate | | |
| $\text{C}_5\text{H}_4\text{O}_2$ | Furfural | | Feedstock chemical, specialty solvent, wood off-gassing, biomass burning |
| $\text{C}_8\text{H}_{18}\text{O}_2$ | Ethylene glycol hexyl ether; 1,2 Octanediol | | |
| $\text{C}_6\text{H}_{10}\text{O}$ | Cyclohexanone | | Solvent for paints, degreasers, polymers, etc. |
| $\text{C}_{10}\text{H}_{20}\text{O}_2$ | Hydroxycitronellal | Octyl acetate, p-Menthane-3,8-diol | Fragrances, solvents, insect repellants |
| $\text{C}_{12}\text{H}_{24}\text{O}_3$ | Texanol | | Paint solvent, other solvents |
| $\text{C}_8\text{H}_8\text{O}$ | Acetophenone | 2-methyl benzaldehyde, phenyl acetaldehyde | Automobile exhaust, fragrances |
| $\text{C}_{10}\text{H}_{12}\text{O}$ | Benzyl acetone | Anethole, Estragole, Cuminaldehyde | Perfumery, cosmetic products |
| $\text{C}_7\text{H}_{12}\text{O}_4$ | Pentadioic acid, dimethyl ester | Diethyl malonate | Fragrances |
| $\text{C}_6\text{H}_{12}\text{O}_3$ | PGMEA; 2-Ethoxyethyl acetate | | Coatings, printing inks, deicing formulations |
| $\text{C}_4\text{H}_{10}\text{O}_2$ | 1-Methoxy-2-propanol | 2-Ethoxyethanol, 1,4 Butanediol | Industrial solvent |
| $\text{C}_6\text{H}_{12}\text{O}_2$ | Butyl acetate | Propyl propanoate, Methyl pentanoate, Ethyl butyrate, Diacetone alcohol | Fragrances, paints, solvents |
| $\text{C}_6\text{H}_{14}\text{O}_2$ | 2-Butoxyethanol | 2-Methyl-2,4-pentanediol | Solvents, cosmetic products, coatings, surfactant stabilizer |
| $\text{C}_3\text{H}_8\text{O}_2$ | 2-Methoxy ethanol; Propylene glycol | Dimethoxymethane | Solvent for varnishes, dyes, wood stains |
| $\text{C}_8\text{H}_{18}\text{O}_3$ | 2-(2-Butoxyethoxy)ethanol; DGBE | | Paints, inks, surface cleaners, varnishes |
| $\text{C}_8\text{H}_{14}\text{O}$ | 6-Methyl 5-hepten-2-one | Filbertone | Skin oil oxidation product, fragrances |
| $\text{C}_{10}\text{H}_{12}\text{O}_3$ | Propyl paraben | | Cosmetics |
| $\text{C}_6\text{H}_6\text{O}_3$ | Maltol | Pyrogallol, Hydroxymethylfurfural | Dyes, other industrial processing, food additive |
| $\text{C}_{10}\text{H}_{10}\text{O}_4$ | Dimethyl phthalate | | Insect repellents, lacquer coatings, plastics |
| $\text{C}_5\text{H}_8\text{O}$ | Cyclopentanone | 3-Penten-2-one | Precursor to fragrances, flavoring agent |
| $\text{C}_4\text{H}_8\text{O}$ | Methyl ethyl ketone | Tetrahydrofuran | Paints, dyes, adhesive remover, degreasers, solvent |
| $\text{C}_6\text{H}_{12}\text{O}$ | 2-Hexanone; 4-Methyl-2-pentanone | Methyl isobutyl ketone | Solvent |
| $\text{C}_8\text{H}_{10}\text{O}_2$ | Phenoxy ethanol | Anisyl alcohol, 1,4-Dimethoxybenzene | Fragrances, soaps |
| $\text{C}_5\text{H}_9\text{NO}$ | Methyl pyrrolidone | | Industrial solvent |
| $\text{C}_7\text{H}_8\text{O}$ | Benzyl alcohol | Anisole | General solvent |
| $\text{C}_9\text{H}_{10}\text{O}_3$ | Ethyl paraben | Ethyl salicylate | Cosmetics |
| $\text{C}_6\text{H}_{14}\text{O}_3$ | Dipropylene glycol | 2-(2-Ethoxyethoxy)ethanol | Industrial solvent, paints, wood stains, textile inks, brake fluids |
| $\text{C}_5\text{H}_8\text{O}_2$ | Methyl methacrylate | Acetylpropionyl, 2-Methyltetrahydrofuran-3-one | Solvents for paints, inks and lacquers, coffee aroma |
| $\text{C}_3\text{H}_6\text{O}$ | Acetone | Propionaldehyde | Wide-use solvent, disinfectant |
| $\text{C}_4\text{H}_6\text{O}$ | Methyl vinyl ketone | Methacrolein, Crotonaldehyde | Chemical precursor |
| $\text{C}_{11}\text{H}_{14}\text{O}_3$ | Butyl paraben | | Antioxidant, cosmetic preservatives |
| $\text{C}_5\text{H}_{10}\text{O}_2$ | Isopropyl acetate; n-Propyl acetate | Isobutyl formate, Methyl isobutyrate | Fragrances, industrial solvent |

| | | | |
|---|---------------------------------|---|---|
| C ₇ H ₆ O | Benzaldehyde | | Soaps, dyes, fragrances |
| C ₈ H ₈ O ₃ | Methyl paraben | | Cosmetics, personal care products |
| C ₁₀ H ₂₀ O | Menthol | Citronellal, Decanal, Rhodinol | Fragrances, insect repellants, food flavoring |
| C ₄ H ₁₀ O ₃ | Diethylene glycol | | Solvent for resins, dyes, oils |
| C ₁₀ H ₁₂ O ₂ | Eugenol | Chavibetol, Propyl benzoate, Hinokitiol | Fragrances, cosmetic preservatives, food flavoring, consumer products |
| C ₁₂ H ₁₄ O ₄ | Diethyl phthalate | | Plasticizer in consumer products |
| C ₁₆ H ₃₀ O ₄ | TXIB | | Inks, coatings, lacquers |
| C ₉ H ₂₀ O ₃ | Dipropylene glycol propyl ether | | Industrial solvent, inks, coatings |
| C ₂ H ₃ N | Acetonitrile | | Industrial solvent, chemical precursor, combustion by-product |
| C ₃ H ₆ O ₂ | Methyl acetate; Propionic acid | Hydroxyacetone, Ethyl formate | Industrial solvent, naturally occurring |
| C ₈ H ₁₄ O ₂ | Cyclohexyl acetate | | Flavoring agent, fragrances |
| C ₉ H ₁₀ O ₂ | Benzyl acetate | Ethyl benzoate, Acetanisole | Fragrances, food flavoring |
| C ₁₂ H ₁₆ O ₃ | 2-Phenoxyethyl isobutyrate | | |
| C ₄ H ₈ O ₂ | Butyric acid; Ethyl acetate | Methyl propionate, Acetoin | Industrial solvent, coatings, fragrances, food flavoring |
| C ₈ H ₈ O ₂ | Methyl benzoate | Phenylacetic acid | Fragrances |
| C ₁₀ H ₁₆ | Monoterpenes | | Widely-used fragrances, Food flavoring, biogenic sources |
| C ₇ H ₆ O ₂ | Benzoic acid | | Fragrances, dyes, insect repellents |
| C ₈ H ₁₆ O ₂ | Octanoic acid | Hexyl acetate | Disinfectants, skin products, industrial solvent |
| C ₈ H ₂₄ O ₄ Si ₄ | D4-siloxane | | Adhesives, silicone rubbers, gels |
| C ₉ H ₁₈ O ₂ | Methyl octanoate | Heptyl acetate, Pentyl butyrate | Fragrances, cigarette additive |
| C ₁₅ H ₂₄ | Sesquiterpenes | | Fragrances, biogenic sources |
| C ₂ H ₆ O ₂ | Ethylene glycol | Methoxymethanol | Antifreeze, coolant, de-icing agent, paints, printing inks |
| C ₄ H ₆ O ₂ | Vinyl acetate; Methyl acrylate | γ-Butyrolactone, Methacrylic acid, Diacetyl | Monomer, precursor/feedstock, nail primers, fragrances |
| C ₃ H ₈ O ₃ | Glycerol | | Personal care products, foods |

Table S8. List of ions calibrated with authentic standards (Table S2), probable contributing isomers, background subtracted geometric mean concentrations (with standard deviations), annual emissions in each inventory, and measured mean concentration enhancement ratios (with standard deviations over the measurement period and linear correlation coefficients) with acetone and other tracers.

| Compound formula, i | Probable compounds, i | Geo. mean concentration, ppt, i | Emissions, kg yr ⁻¹ , VCPy+, FIVE-VCP | Ratios to tracer compounds ($\Delta\text{mol}/\Delta\text{mol}$) [†] | | | |
|---|--|---------------------------------|--|---|--------------------------------------|-------------------------------------|--|
| | | | | $\Delta i/\Delta\text{Benzene}$ (r) | $\Delta i^*1000/\Delta\text{CO}$ (r) | $\Delta i/\Delta\text{Acetone}$ (r) | $\Delta i/\Delta\text{Benzyl alcohol}$ (r) |
| C ₂ H ₆ O ₂ | Ethylene glycol | 2437±3622 | 361511, 236310 | 1.1E+01±2.5E+01 (0.79) | 9.1E+00±2.0E+01 (0.83) | 2.8E+00±6.4E+00 (0.95) | 3.0E+02±6.5E+02 (0.88) |
| C ₃ H ₆ O | Acetone* | 977±783 | 1360720, 1587220 | 3.8E+00±6.7E+00 (0.83) | 3.3E+00±5.2E+00 (0.87) | -- | 1.1E+02±1.6E+02 (0.92) |
| C ₄ H ₆ O ₂ | Methyl acrylate*, Diacyetyl* | 810±396 | 1905, 4638 | 2.1E+00±3.6E+00 (0.82) | 1.8E+00±2.7E+00 (0.89) | 5.6E-01±8.9E-01 (0.95) | 5.9E+01±8.6E+01 (0.94) |
| C ₄ H ₈ O ₂ | Ethyl acetate*, Butyric acid | 679±664 | 30225, 293 | 2.8E+00±5.3E+00 (0.72) | 2.3E+00±4.1E+00 (0.73) | 7.2E-01±1.3E+00 (0.73) | 7.6E+01±1.3E+02 (0.67) |
| C ₃ H ₆ O ₂ | Methyl acetate*, Propionic acid, Hydroxyacetone, Ethyl formate | 435±377 | 50747, 103808 | 1.7E+00±3.1E+00 (0.64) | 1.5E+00±2.4E+00 (0.65) | 4.5E-01±7.9E-01 (0.76) | 4.8E+01±7.7E+01 (0.7) |
| C ₂ H ₃ N | Acetonitrile | 246±102 | | 8.5E-01±1.2E+00 (0.32) | 7.2E-01±9.1E-01 (0.24) | 2.2E-01±3.0E-01 (0.35) | 2.3E+01±2.8E+01 (0.33) |
| C ₁₀ H ₁₆ | Monoterpenes (e.g., limonene*, α -Pinene*) | 156±105 | 60327, 15516 | 5.1E-01±9.0E-01 (0.79) | 4.3E-01±6.9E-01 (0.87) | 1.3E-01±2.3E-01 (0.85) | 1.4E+01±2.2E+01 (0.94) |
| C ₄ H ₈ O | MEK, THF, Cyclopropyl carbinol* | 126±82 | 57457, 277556 | 4.3E-01±7.3E-01 (0.79) | 3.7E-01±5.6E-01 (0.84) | 1.1E-01±1.8E-01 (0.93) | 1.2E+01±1.8E+01 (0.85) |
| C ₅ H ₁₀ O ₂ | Isopropyl acetate*, n-propyl acetate* | 114±106 | 3457, 5289 | 4.4E-01±8.3E-01 (0.61) | 3.7E-01±6.5E-01 (0.69) | 1.1E-01±2.1E-01 (0.69) | 1.2E+01±2.1E+01 (0.58) |
| C ₅ H ₈ O ₂ | Methyl methacrylate* | 108±121 | 1102, - | 4.1E-01±8.8E-01 (0.45) | 3.5E-01±7.0E-01 (0.37) | 1.1E-01±2.2E-01 (0.5) | 1.1E+01±2.2E+01 (0.41) |
| C ₆ H ₁₂ O ₂ | Butyl acetate* | 103±138 | 80120, 56862 | 4.9E-01±1.0E+00 (0.76) | 4.1E-01±8.1E-01 (0.77) | 1.3E-01±2.6E-01 (0.87) | 1.3E+01±2.6E+01 (0.83) |
| C ₈ H ₈ O ₂ | Methyl benzoate* | 92±15 | | 1.1E-01±1.6E-01 (0.72) | 9.1E-02±1.2E-01 (0.75) | 2.8E-02±4.1E-02 (0.78) | 3.0E+00±3.8E+00 (0.79) |
| C ₈ H ₁₆ O ₂ | Caprylic acid* (i.e., Octanoic acid) | 87±47 | 5281, - | 2.5E-01±4.2E-01 (0.81) | 2.1E-01±3.2E-01 (0.92) | 6.5E-02±1.1E-01 (0.92) | 6.9E+00±1.0E+01 (0.95) |
| C ₃ H ₈ O ₂ | 2-Methoxy ethanol, propylene glycol* | 82±51 | 240692, - | 2.9E-01±4.7E-01 (0.71) | 2.4E-01±3.6E-01 (0.71) | 7.5E-01±1.2E-01 (0.85) | 7.9E+00±1.1E+01 (0.77) |
| C ₉ H ₁₈ O ₂ | Methyl octanoate, Nonanoic acid* | 77±24 | | 1.4E-01±2.3E-01 (0.79) | 1.2E-01±1.7E-01 (0.9) | 3.7E-02±5.7E-02 (0.9) | 3.9E+00±5.4E+00 (0.94) |
| C ₇ H ₆ O | Benzaldehyde* | 76±37 | 3156, 14833 | 2.1E-01±3.4E-01 (0.83) | 1.8E-01±2.6E-01 (0.88) | 5.4E-02±8.6E-02 (0.88) | 5.7E+00±8.2E+00 (0.93) |
| C ₁₅ H ₂₄ | Sesquiterpenes (e.g., β - Caryophyllene) | 70±11 | | 7.3E-02±1.2E-01 (0.83) | 6.2E-02±8.6E-02 (0.83) | 1.9E-02±2.9E-02 (0.78) | 2.0E+00±2.7E+00 (0.7) |
| C ₆ H ₁₂ O | 2-Hexanone*, 4-Methyl- 2-pentanone | 59±42 | 6162, 14990 | 2.0E-01±3.6E-01 (0.83) | 1.7E-01±2.8E-01 (0.84) | 5.3E-02±9.0E-02 (0.92) | 5.6E+00±8.8E+00 (0.91) |
| C ₇ H ₆ O ₂ | Benzoic acid* | 59±9 | | 5.8E-02±9.0E-02 (0.48) | 4.9E-02±6.7E-02 (0.39) | 1.5E-02±2.2E-02 (0.4) | 1.6E+00±2.1E+00 (0.45) |
| C ₄ H ₆ O | MVK, MACR | 58±39 | | 1.9E-01±3.3E-01 (0.83) | 1.6E-01±2.5E-01 (0.87) | 4.9E-02±8.3E-02 (0.94) | 5.1E+00±8.0E+00 (0.94) |
| C ₈ H ₁₄ O ₂ | Cyclohexyl acetate | 43±20 | | 1.2E-01±2.0E-01 (0.81) | 1.0E-01±1.5E-01 (0.89) | 3.2E-02±4.9E-02 (0.95) | 3.4E+00±4.7E+00 (0.95) |
| C ₉ H ₁₀ O ₂ | Benzyl acetate | 39±19 | 7, - | 1.0E-01±1.7E-01 (0.82) | 8.8E-02±1.3E-01 (0.89) | 2.7E-02±4.4E-02 (0.87) | 2.9E+00±4.2E+00 (0.95) |
| C ₆ H ₁₄ O ₃ | Dipropylene glycol | 36±28 | 41085, 105732 | 1.4E-01±2.4E-01 (0.65) | 1.2E-01±1.9E-01 (0.71) | 3.6E-02±6.1E-02 (0.7) | 3.8E+00±5.9E+00 (0.8) |
| C ₄ H ₁₀ O ₃ | Diethylene glycol | 32±17 | 7026, 110939 | 8.9E-02±1.5E-01 (0.84) | 7.5E-02±1.2E-01 (0.87) | 2.3E-02±3.8E-02 (0.91) | 2.4E+00±3.7E+00 (0.92) |
| C ₁₀ H ₂₀ O | Menthol, Decanal* | 31±18 | 971, 0.05 | 9.4E-02±1.6E-01 (0.77) | 7.9E-02±1.2E-01 (0.89) | 2.4E-02±4.0E-02 (0.9) | 2.6E+00±3.8E+00 (0.96) |
| C ₅ H ₈ O | Cyclopentanone | 30±16 | | 8.4E-02±1.4E-01 (0.84) | 7.1E-02±1.1E-01 (0.9) | 2.2E-02±3.5E-02 (0.95) | 2.3E+00±3.4E+00 (0.95) |
| C ₆ H ₁₄ O ₂ | 2-Butoxyethanol*, 1- propoxy-2-propanol* | 23±19 | 109288, 72125 | 8.9E-02±1.6E-01 (0.8) | 7.5E-02±1.2E-01 (0.87) | 2.3E-02±3.9E-02 (0.91) | 2.4E+00±3.8E+00 (0.9) |
| C ₈ H ₂₄ O ₄ Si ₄ | D4 siloxane* | 23±3 | 12872, 92707 | 2.3E-02±3.6E-02 (0.38) | 2.0E-02±2.7E-02 (0.48) | 6.0E-03±8.9E-03 (0.48) | 6.4E-01±8.5E-01 (0.59) |
| C ₁₆ H ₃₀ O ₄ | TXIB* | 18±4 | - , 2264 | 2.6E-02±4.2E-02 (0.73) | 2.2E-02±3.2E-02 (0.83) | 6.8E-03±1.1E-02 (0.75) | 7.2E-01±1.0E+00 (0.86) |

| | | | | | | | |
|--|--|-------|----------------|---------------------------|---------------------------|---------------------------|---------------------------|
| C ₁₀ H ₁₂ O ₂ | Eugenol | 16±5 | 45, - | 3.1E-02±4.9E-02 (0.82) | 2.6E-02±3.7E-02 (0.85) | 7.9E-03±1.2E-02 (0.91) | 8.4E-01±1.2E+00 (0.92) |
| C ₉ H ₂₀ O ₃ | Dipropylene glycol propyl ether | 16±4 | 4150, 5966 | 2.3E-02±3.8E-02 (0.65) | 2.0E-02±2.9E-02 (0.71) | 6.1E-03±9.5E-03 (0.62) | 6.4E-01±9.1E-01 (0.73) |
| C ₁₂ H ₁₆ O ₃ | 2-Phenoxyethyl isobutyrate | 16±2 | | 1.6E-02±2.4E-02 (0.73) | 1.3E-02±1.8E-02 (0.76) | 4.1E-03±5.9E-03 (0.79) | 4.4E-01±5.6E-01 (0.83) |
| C ₁₀ H ₃₀ O ₅ Si ₅ | D5 siloxane* | 16±15 | 272778, 323982 | 6.7E-02±1.2E-01 (0.7) | 5.7E-02±9.6E-02 (0.82) | 1.7E-02±3.1E-02 (0.82) | 1.8E+00±3.0E+00 (0.9) |
| C ₁₂ H ₁₄ O ₄ | Diethyl phthalate | 15±3 | 17138, - | 2.3E-02±3.4E-02 (0.64) | 1.9E-02±2.5E-02 (0.7) | 5.9E-03±8.4E-03 (0.65) | 6.2E-01±7.8E-01 (0.71) |
| C ₇ H ₈ O | Benzyl alcohol | 14±6 | 22898, 20791 | 3.6E-02±5.9E-02 (0.85) | 3.1E-02±4.5E-02 (0.92) | 9.5E-03±1.5E-02 (0.92) | -- |
| C ₈ H ₁₄ O | 6-Methyl 5-hepten-2-one | 14±7 | | 4.1E-02±6.6E-02 (0.81) | 3.4E-02±5.0E-02 (0.89) | 1.1E-02±1.6E-02 (0.96) | 1.1E+00±1.6E+00 (0.96) |
| C ₈ H ₈ O ₃ | Methyl paraben | 14±4 | | 2.4E-02±3.9E-02 (0.83) | 2.1E-02±2.9E-02 (0.86) | 6.3E-03±9.7E-03 (0.83) | 6.7E-01±9.2E-01 (0.87) |
| C ₄ H ₁₀ O ₂ | 1-Methoxy-2-propanol | 13±8 | 3558, 2182 | 4.1E-02±7.1E-02 (0.78) | 3.5E-02±5.4E-02 (0.85) | 1.1E-02±1.8E-02 (0.89) | 1.1E+00±1.7E+00 (0.89) |
| C ₅ H ₄ O ₂ | Furfural* | 13±6 | - , 0.01 | 3.4E-02±5.7E-02 (0.71) | 2.9E-02±4.3E-02 (0.62) | 8.8E-03±1.4E-02 (0.56) | 9.3E-01±1.4E+00 (0.66) |
| C ₆ H ₁₀ O | Cyclohexanone | 12±6 | 384, 96838 | 3.6E-02±5.9E-02 (0.84) | 3.0E-02±4.5E-02 (0.91) | 9.4E-03±1.5E-02 (0.96) | 9.9E-01±1.4E+00 (0.92) |
| C ₆ H ₁₂ O ₃ | PGMEA, 2-Ethoxyethyl acetate | 12±11 | 10327, 7450 | 4.7E-02±8.8E-02 (0.78) | 4.0E-02±6.8E-02 (0.76) | 1.2E-02±2.2E-02 (0.9) | 1.3E+00±2.2E+00 (0.86) |
| C ₆ H ₆ O ₃ | Maltol | 11±3 | | 1.3E-02±2.3E-02 (0.59) | 1.1E-02±1.7E-02 (0.44) | 3.4E-03±5.6E-03 (0.42) | 3.6E-01±5.4E-01 (0.49) |
| C ₈ H ₈ O | Acetophenone* | 10±6 | 4, - | 3.2E-02±5.5E-02 (0.81) | 2.7E-02±4.2E-02 (0.85) | 8.4E-03±1.4E-02 (0.89) | 8.8E-01±1.3E+00 (0.9) |
| C ₅ H ₉ NO | Methyl pyrrolidone | 9±3 | 12749, 14015 | 1.9E-02±3.1E-02 (0.72) | 1.6E-02±2.3E-02 (0.78) | 5.0E-03±7.7E-03 (0.77) | 5.3E-01±7.4E-01 (0.78) |
| C ₈ H ₁₀ O ₂ | Phenoxyethanol* | 9±3 | 9851, 0.23 | 1.7E-02±2.9E-02 (0.78) | 1.5E-02±2.2E-02 (0.84) | 4.5E-03±7.2E-03 (0.86) | 4.8E-01±6.9E-01 (0.91) |
| C ₈ H ₁₈ O ₃ | 2-(2-Butoxyethoxy)ethanol, DGBE | 8±4 | 48681, 62011 | 2.1E-02±3.5E-02 (0.85) | 1.8E-02±2.7E-02 (0.91) | 5.4E-03±8.7E-03 (0.89) | 5.7E-01±8.4E-01 (0.94) |
| C ₁₀ H ₁₀ O ₄ | Dimethyl phthalate | 7±1 | 70, - | 9.1E-03±1.5E-02 (0.62) | 7.7E-03±1.1E-02 (0.62) | 2.4E-03±3.6E-03 (0.55) | 2.5E-01±3.4E-01 (0.65) |
| C ₁₂ H ₂₄ O ₃ | Texanol* | 7±4 | 267615, 179276 | 2.0E-02±3.5E-02 (0.57) | 1.7E-02±2.7E-02 (0.74) | 5.3E-03±8.7E-03 (0.67) | 5.6E-01±8.4E-01 (0.74) |
| C ₉ H ₁₀ O ₃ | Ethyl paraben | 6±1 | | 7.0E-03±1.1E-02 (0.84) | 5.9E-03±8.2E-03 (0.84) | 1.8E-03±2.7E-03 (0.85) | 1.9E-01±2.6E-01 (0.9) |
| C ₁₁ H ₁₄ O ₃ | Butyl paraben | 6±1 | | 8.5E-03±1.3E-02 (0.71) | 7.2E-03±9.4E-03 (0.74) | 2.2E-03±3.1E-03 (0.8) | 2.3E-01±2.9E-01 (0.76) |
| C ₆ H ₁₀ O ₃ | Ethyl acetoacetate | 4±2 | | 1.3E-02±2.2E-02 (0.85) | 1.1E-02±1.6E-02 (0.87) | 3.4E-03±5.4E-03 (0.93) | 3.6E-01±5.2E-01 (0.91) |
| C ₁₀ H ₁₂ O | Benzyl acetone | 4±2 | | 1.0E-02±1.7E-02 (0.85) | 8.5E-03±1.3E-02 (0.91) | 2.6E-03±4.2E-03 (0.94) | 2.8E-01±4.1E-01 (0.97) |
| C ₇ H ₁₂ O ₄ | Pentadioic acid, dimethyl ester | 4±1 | 4942, 25606 | 7.2E-03±1.1E-02 (0.8) | 6.1E-03±8.5E-03 (0.84) | 1.9E-03±2.8E-03 (0.87) | 2.0E-01±2.7E-01 (0.89) |
| C ₁₀ H ₁₂ O ₃ | Propyl paraben | 4±1 | | 6.3E-03±1.0E-02 (0.54) | 5.3E-03±7.7E-03 (0.46) | 1.6E-03±2.5E-03 (0.42) | 1.7E-01±2.4E-01 (0.51) |
| C ₁₀ H ₂₀ O ₂ | Hydroxycitronellal | 3±1 | | 5.3E-03±8.4E-03 (0.78) | 4.5E-03±6.3E-03 (0.88) | 1.4E-03±2.1E-03 (0.92) | 1.5E-01±2.0E-01 (0.95) |
| C ₈ H ₁₈ O ₂ | Ethylene glycol hexyl ether*, 1,2-Octanediol | 2±1 | 15836, 7749 | 5.8E-03±9.6E-03 (0.8) | 4.9E-03±7.3E-03 (0.88) | 1.5E-03±2.4E-03 (0.87) | 1.6E-01±2.3E-01 (0.94) |
| C ₃ H ₈ O ₃ | Glycerol | 1±0.5 | 148441, 949405 | 3.3E-04±4.9E-03 (0.64) | 1.6E-03±3.7E-03 (0.65) | 5.3E-04±1.2E-03 (0.74) | 6.3E-02±1.2E-01 (0.73) |
| C ₆ H ₁₄ O ₄ | Triethylene glycol | 1±0.3 | 1718, 955 | 2.1E-03±3.4E-03 (0.47) | 1.8E-03±2.6E-03 (0.45) | 5.5E-04±8.4E-04 (0.4) | 5.8E-02±8.0E-02 (0.51) |

References

- Bertram, T. H., Kimmel, J. R., Crisp, T. A., Ryder, O. S., Yatavelli, R. L. N., Thornton, J. A., ... Worsnop, D. R. (2011). A field-deployable, chemical ionization time-of-flight mass spectrometer. *Atmospheric Measurement Techniques*, 4(7), 1471–1479. <https://doi.org/10.5194/AMT-4-1471-2011>
- Gkatzelis, G. I., Coggon, M. M., McDonald, B. C., Peischl, J., Aikin, K. C., Gilman, J. B., ... Warneke, C. (2021). Identifying Volatile Chemical Product Tracer Compounds in U.S. Cities. *Environmental Science and Technology*, 55(1), 188–199. https://doi.org/10.1021/ACS.EST.0C05467/SUPPL_FILE/ES0C05467_SI_001.PDF
- Koss, A. R., Sekimoto, K., Gilman, J. B., Selimovic, V., Coggon, M. M., Zarzana, K. J., ... De Gouw, J. (2018). Non-methane organic gas emissions from biomass burning: Identification, quantification, and emission factors from PTR-ToF during the FIREX 2016 laboratory experiment. *Atmospheric Chemistry and Physics*, 18(5), 3299–3319. <https://doi.org/10.5194/ACP-18-3299-2018>
- Novak, G. A., Vermeuel, M. P., & Bertram, T. H. (2020). Simultaneous detection of ozone and nitrogen dioxide by oxygen anion chemical ionization mass spectrometry: A fast-time-response sensor suitable for eddy covariance measurements. *Atmospheric Measurement Techniques*, 13(4), 1887–1907. <https://doi.org/10.5194/AMT-13-1887-2020>
- Sheu, R., Stönnner, C., Ditto, J. C., Klüpfel, T., Williams, J., & Gentner, D. R. (2020). Human transport of thirdhand tobacco smoke: A prominent source of hazardous air pollutants into indoor nonsmoking environments. *Science Advances*, 6(10), eaay4109. <https://doi.org/10.1126/sciadv.aay4109>
- Xu, L., Coggon, M. M., Stockwell, C. E., Gilman, J. B., Robinson, M. A., Breitenlechner, M., ... Warneke, C. (2022). A Chemical Ionization Mass Spectrometry Utilizing Ammonium Ions (NH₄⁺ CIMS) for Measurements of Organic Compounds in the Atmosphere. *Aerosol Measurement Techniques Discussions*. <https://doi.org/10.5194/amt-2022-228>