



## Supplement of

## Measurement report: Carbonyl sulfide production during dimethyl sulfide oxidation in the atmospheric simulation chamber SAPHIR

Marc von Hobe et al.

*Correspondence to:* Marc von Hobe (m.von.hobe@fz-juelich.de)

The copyright of individual parts of the supplement might differ from the article licence.



Figure S1 Additional parameters in Experiment I, with time in hours after the first DMS injection and the period with the UV lamps turned on indicated by the coloured bar on the top of panel a. In panel a) k<sub>OH</sub> and radical concentrations are shown with the position of the labels indicating which axis refers to which species. In panel b), NO and NO<sub>2</sub> mixing ratios are given, together with raw PTRMS signals at m/z ratios that are potentially related to relevant products and intermediates (m/z 77: CH<sub>3</sub>SCHO; m/z 79: DMSO; m/z 95: DMSO2 or CH<sub>3</sub>SCH<sub>2</sub>OOH; see Section 2.3 of the article). In panel c), temperature and humidity in the SAPHIR chamber are shown.



Figure S2 Additional parameters in Experiment II, with time in hours after the first DMS injection and the period with the UV lamps turned on indicated by the coloured bar on the top of panel a. In panel a) k<sub>OH</sub> and radical concentrations are shown with the position of the labels indicating which axis refers to which species. In panel b), NO and NO<sub>2</sub> mixing ratios are given, together with raw PTRMS signals at m/z ratios that are potentially related to relevant products and intermediates (m/z 77: CH<sub>3</sub>SCHO; m/z 79: DMSO; m/z 95: DMSO2 or CH<sub>3</sub>SCH<sub>2</sub>OOH; see Section 2.3 of the article). In panel c), temperature and humidity in the SAPHIR chamber are shown.



650

655

**Figure S3** Additional parameters in Experiment III, with time in hours after the first DMS injection and the period with the roof being open indicated by the coloured bar on the top of panel a. In panel a)  $k_{OH}$  and radical concentrations are shown with the position of the labels indicating which axis refers to which species. In panel b), NO and NO<sub>2</sub> mixing ratios are given, together with raw PTRMS signals at m/z ratios that are potentially related to relevant products and intermediates (m/z 77: CH<sub>3</sub>SCHO; m/z 79: DMSO; m/z 95: DMSO2 or CH<sub>3</sub>SCH<sub>2</sub>OOH; see Section 2.3 of the article). In panel c), temperature and

humidity in the SAPHIR chamber are shown.



**Figure S4** Additional parameters in Experiment IV, with time in hours after the first DMS injection and the periods with the UV lamps turned on and the roof being open indicated by the coloured bar on the top of panel a. In panel a) *k*<sub>OH</sub> and radical concentrations are shown with the position of the labels indicating which axis refers to which species. In panel b), NO and NO<sub>2</sub> mixing ratios are given, together with raw PTRMS signals at m/z ratios that are potentially related to relevant products and intermediates (m/z 77: CH<sub>3</sub>SCHO; m/z 79: DMSO; m/z 95: DMSO2 or CH<sub>3</sub>SCH<sub>2</sub>OOH; see Section 2.3 of the article). In panel c), temperature and humidity in the SAPHIR chamber are shown.



**Figure S5** Comparison of observed (black) to expected DMS removal rate (orange) in Experiment IV for  $k_{\text{DMS+OH}}$  from Barnes et al. (2006, and references therin) and OH observed by the DOAS instrument. Also shown is the theoretical DMS removal rate for just the H abstraction channel of the DMS + OH reaction (blue, also with rate constants from Barnes et al., 2006) which represents an upper limit for the HPMTF production rate.





**Figure S6** Sulfur budget of measured sulfur gases (panels a - d correspond to Experiments I – IV respectively). The black line indicates the total sulfur expected in the SAPHIR chamber accounting for DMS additions and chamber loss by dilution. In Experiments II – IV, DMS and two main product sulfur gases, SO<sub>2</sub> and OCS, are measured, and together make up between 70 and 80 % of the total sulfur in the chamber (based on injected DMS and the chamber dilution rate). We expect the remaining part of the sulfur budget to be made up by other product gases (e.g. dimethyl sulfoxide, DMSO, as the major product gas of the OH addition channel) and intermediates including HPMTF and TPA.



**Figure S7** Observed decrease in CO2 mixing ratios and comparison to the dilution rates calculated from the flow into the SAPHIR chamber.