



Supplement of

Active thermokarst regions contain rich sources of ice-nucleating particles

Kevin R. Barry et al.

Correspondence to: Kevin R. Barry (kevin.barry@colostate.edu)


The copyright of individual parts of the supplement might differ from the article licence.

5 Supplement

10


<i>Source</i>	<i>Mean INP Concentration (-15 °C)</i>
<i>Aerosol</i>	$4.4 \cdot 10^{-2} \text{ L}^{-1}$
<i>TKL</i>	$1.2 \cdot 10^5 \text{ mL}^{-1}$
<i>Lagoon</i>	$3.1 \cdot 10^4 \text{ mL}^{-1}$
<i>Seawater</i>	$1.7 \cdot 10^4 \text{ mL}^{-1}$
<i>Active Layer</i>	$3.6 \cdot 10^8 \text{ g}^{-1}$
<i>Permafrost</i>	$1.1 \cdot 10^8 \text{ g}^{-1}$
<i>Sediment</i>	$3.2 \cdot 10^7 \text{ g}^{-1}$
<i>Vegetation</i>	$2.0 \cdot 10^6 \text{ g}^{-1}$
<i>Ice Wedge</i>	$1.7 \cdot 10^5 \text{ g}^{-1}$

Table S1: Average INP concentration at -15 °C for all aerosol and potential source samples. TKL=Thermokarst lake.




15

Figure S1: Complete INP-temperature spectra for all aerosol samples, colored by environment collected (SW=seawater; DOE=fixed site; TKL=thermokarst lake). 95% confidence intervals are plotted (any confidence intervals overlapping with 0 are not shown).




20

Figure S2: A heatmap indicating results of statistical testing for downwind versus upwind aerosol INP concentrations. Purple represents an increase over upwind at 95% confidence, while gray represents no difference at 95% confidence.



- 25 **Figure S3:** Complete INP-temperature spectra for all water samples (thermokarst lake water: green; lagoon: light blue; seawater: dark blue). 95% confidence intervals are plotted (any confidence intervals overlapping with 0 are not shown).



- 30 **Figure S4:** Complete INP-temperature spectra for terrestrial samples (permafrost: gray; sediment: salmon; vegetation: dark green). 95% confidence intervals are plotted (any confidence intervals overlapping with 0 are not shown).




Figure S5: Complete INP-temperature spectra from permafrost cores collected on September 11 (left) and September 17 (right). The different colors represent depth of sample taken in relation to the top of the permafrost core. 95% confidence intervals are plotted (any confidence intervals overlapping with 0 are not shown).





Figure S6: INP concentration per g at -15 °C for active layer (maroon) and ice wedge (bright green) samples. The size of the markers corresponds to the INP concentration.



45 **Figure S7:** Complete INP-temperature spectra for terrestrial samples (active layer: maroon; ice wedge: bright green). 95% confidence intervals are plotted (any confidence intervals overlapping with 0 are not shown).