



Large simulated future changes in the nitrate radical under the CMIP6 SSP scenarios: implications for oxidation chemistry

Scott Archer-Nicholls^{1,a}, Rachel Allen¹, Nathan L. Abraham^{1,2}, Paul T. Griffiths^{1,2}, and Alex T. Archibald^{1,2}

¹Department of Chemistry, University of Cambridge, Cambridge, CB2 1EW, UK

²NCAS-Climate, University of Cambridge, Cambridge, CB2 1EW, UK

^anow at: IT Services, University of Manchester, Manchester, M13 9PL, UK

Correspondence: Alex T. Archibald (ata27@cam.ac.uk)

Received: 7 October 2022 – Discussion started: 16 November 2022

Revised: 6 April 2023 – Accepted: 9 April 2023 – Published: 24 May 2023

Abstract. The nitrate radical (NO_3) plays an important role in the chemistry of the lower troposphere, acting as the principle oxidant during the night together with ozone. Previous model simulations suggest that the levels of NO_3 have increased dramatically since the preindustrial period. Here, we show projections of the evolution of the NO_3 radical from 1850–2100 using the United Kingdom Earth System Model (UKESM1) under the Coupled Model Intercomparison Project Phase 6 (CMIP6) shared socioeconomic pathway (SSP) scenarios. Our model results highlight diverse trajectories for NO_3 , with some scenarios and regions undergoing rapid growth of NO_3 to unprecedented levels over the course of the 21st century and others seeing sharp declines. The local increases in NO_3 (up to 40 ppt above the preindustrial base line) are driven not only by local changes in emissions of nitrogen oxides but have an important climate component, with NO_3 being favoured in warmer future climates. The changes in NO_3 lead to changes in the oxidation of important secondary organic aerosol precursors, with potential impacts on particulate matter pollution regionally and globally. This work highlights the potential for substantial future growth in NO_3 and the need to better understand the formation of secondary organic aerosol (SOA) from NO_3 to accurately predict future air quality and climate implications.

1 Introduction

Whilst nitrogen is ubiquitous in the atmosphere, the majority of gaseous nitrogen-containing molecules are chemically inert. Reactive nitrogen species (NO_y) make up a much smaller fraction but encompass a diverse set of molecules that play a paramount role in the chemistry of the atmosphere. Nitrogen oxide (NO) and nitrogen dioxide (NO_2), collectively known as NO_x , are essential for the catalytic formation of ozone (O_3) in the troposphere, a key air pollutant and greenhouse gas (Monks et al., 2015). The reaction of NO_2 with O_3 produces the nitrate radical (NO_3):



During the daytime, NO_3 can undergo rapid photolysis or reaction with NO, resulting in a very short lifetime, typically in the order of seconds (Wayne et al., 1991). However, at nighttime, in regions of high NO_x , NO_3 is able to persist and become the major oxidant of volatile organic compounds (VOCs) (the hydroxyl radical (OH) and ozone (O_3) dominate this oxidation during the daytime), acting as the most important oxidant during the night (Brown and Stutz, 2012). NO_3 also undergoes a reversible reaction with NO_2 , forming

a thermal equilibrium with dinitrogen pentoxide (N_2O_5):



$$K_{\text{eq}} = \frac{[\text{NO}_2][\text{NO}_3]}{[\text{N}_2\text{O}_5]}. \quad (\text{R3})$$

N_2O_5 has a short lifetime with respect to decomposition at typical atmospheric boundary layer temperatures. As temperature increases, the rate of decomposition of N_2O_5 increases resulting in a greater fraction of reactive nitrogen in the form of NO_3 . This temperature dependence in the equilibrium between NO_2 , NO_3 and N_2O_5 is further driven by the extreme temperature dependence in the formation of NO_3 through Reaction (R1). Due to their tight chemical coupling, NO_3 and N_2O_5 have been termed the N_xO_y family ($\text{N}_x\text{O}_y = \text{NO}_3 + \text{N}_2\text{O}_5$) (Stone et al., 2014). In this sense, sinks of N_2O_5 lead to corresponding indirect loss of NO_3 . Loss of N_xO_y has important implications for tropospheric ozone as it is an important nighttime reservoir of NO_x and O_x ($\text{O} + \text{O}_3 + \text{NO}_2 + \text{others}$) (Archibald et al., 2020a). In the daytime, any remaining N_xO_y is converted back to NO_x ; therefore nighttime sinks of N_xO_y reduce the levels of NO_x available to photochemically form ozone in the troposphere. The heterogeneous hydrolysis of N_2O_5 , which occurs readily on aerosol surfaces, is one of the major nighttime sinks of reactive nitrogen in the troposphere (e.g. Riemer et al., 2003).

NO_3 reactions with alkenes proceed rapidly via the addition of NO_3 to the double bond (Wayne et al., 1991; Brown and Stutz, 2012). As a result, the oxidation of biogenic volatile organic compounds (BVOCs), in particular terpenes, which are emitted in vast quantities by the world's forests (Sakulyanontvittaya et al., 2008), is sensitive to the levels of NO_3 present (Ng et al., 2017). BVOC oxidation by NO_3 can lead to the formation of secondary organic aerosol (SOA) (Fry et al., 2009; Ng et al., 2017), fine aerosols that have implications for the planetary radiation budget and human health (Pöschl, 2005). The extent that NO_3 -initiated chemistry contributes to SOA formation (relative to O_3 and OH-initiated production) depends on the molecular structure of the BVOC (and therefore their rate coefficients for reaction with NO_3), the rate of their emissions and the levels of NO_3 present.

Ng et al. (2017) highlight that there are significant gaps in our understanding of the NO_3 -initiated oxidation of BVOCs. BVOC oxidation chemistry has been studied for many decades (e.g. Brewer et al., 1984); however there are relatively few mechanistic studies of the products of NO_3 -initiated oxidation (e.g. Boyd et al., 2015; Fry et al., 2009; Faxon et al., 2018; Ehn et al., 2017) compared to OH and O_3 (e.g. Atkinson and Arey, 2003; McGillen et al., 2020). Nonetheless there is mounting evidence that NO_3 oxidation could be a significant contributor to SOA formation. Kiendler-Scharr et al. (2016) highlight the ubiquity of organic nitrates in submicron aerosol collected in European nighttime urban conditions, whilst Hamilton et al. (2021) re-

cently found NO_3 oxidation of isoprene to be a significant source of SOA production in Beijing. Recent studies show that highly oxidised molecules (HOMs), formed from the auto-oxidation and accretion of monoterpene oxidation products, are capable of nucleating new aerosol particles without sulfate seeds (Ehn et al., 2014; Tröstl et al., 2016; Bianchi et al., 2019) with important implications for large-scale feedbacks between the climate and the biosphere (Scott et al., 2017; Gordon et al., 2017). Recently, Zhao et al. (2021) have shown that HOMs from NO_3 -initiated oxidation of isoprene may contribute to a significant fraction of the isoprene SOA yield. But these complex processes are so far lacking from Earth system and global chemistry-transport models.

Oxidation of BVOCs is dependent on changes to the oxidant budget (with respect to OH, O_3 and NO_3) and to the changes in the emissions of BVOCs themselves. Our understanding of the processes which control emissions of BVOCs highlights that these have important dependences on climatic parameters including CO_2 concentrations (Arneth et al., 2007), land use, temperature, humidity and precipitation (Arneth et al., 2008). Whilst BVOC emissions models are reasonably well constrained in the present day, past and future trends are less certain, with some models indicating large increases in BVOC emissions in the future (Pacifico et al., 2012) and others small to no increases (Hantson et al., 2017).

Estimates of changes in the levels of NO_3 from the preindustrial period to the present day indicate significant increases, on the order of 100 %–1000 % (Khan et al., 2015), driven by changes in emissions of NO_x . The formation of NO_3 is also dependent on ozone burdens which are likely to change in the future (Archibald et al., 2020a; Griffiths et al., 2021). In addition, the fraction of N_xO_y in the form of NO_3 increases with temperature, while the main sinks are determined by the amount of N_2O_5 which decreases with temperature. Ng et al. (2017) postulated that NO_3 would increase with increasing NO_x emissions, but given the complex chemical processes at work NO_3 is likely to vary nonlinearly with NO_x emissions. To our knowledge, no study has investigated how the NO_3 radical may change in the future under changing NO_x emissions and a changing climate or how these changes will affect BVOC oxidation and SOA formation given changing terpene emissions.

This study presents results on the historic and future evolution of NO_3 using simulations from 1850–2100 made with the United Kingdom Earth System Model (UKESM1) (Sellar et al., 2019; Archibald et al., 2020b). The simulations, based on the Coupled Model Intercomparison Project Phase 6 (CMIP6) Historic and ScenarioMIP scenarios (O'Neill et al., 2016), indicate that there are substantial changes in NO_3 simulated in the future at both the global mean and regional level, depending on the emission scenarios and resulting climate considered. Using these simulations we show that future trends in NO_3 have an impact on the regional oxidation of BVOCs and as a result the SOA budget and burden.

Taking a particular focus on changes occurring over South Asia, a region with severe present-day air quality issues and large variation in how emissions will evolve in this region under different scenarios, we highlight an important role for enhanced levels of NO₃-initiated BVOC oxidation in the future.

2 Methods

2.1 Description of UKESM1




The model simulations make use of the new UK Earth System Model (UKESM1) (Sellar et al., 2019). This is a fully coupled Earth system model using the Global Atmosphere 7.1/Global Land 7.0 (GA7.1/GL7.1; Walters et al., 2019). The dynamical model is coupled with the United Kingdom Chemistry and Aerosol model (UKCA), using the StratTrop mechanism for gas-phase chemistry (Archibald et al., 2020b) and the two-moment GLOMAP-mode aerosol scheme (Mulcahy et al., 2020). UKCA and UKESM1 are designed to quantify the response of the Earth system to forcings and simulate nascent feedbacks that exist in the coupled chemistry–aerosol–climate system. As such, pragmatic decisions have been made to ensure that as complete a representation as possible of many processes is included in the model. For the chemistry of N₂O₅, this includes a fairly basic description of its chemistry in the gas phase and its loss to aerosol surfaces (heterogeneous chemistry) is simplified, with a fixed uptake coefficient being used ($\gamma = 0.1$). Jones et al. (2021) suggest that the simplified treatment of heterogeneous uptake of N₂O₅ leads to an overestimate of the loss of N₂O₅ based on a comparison of modelled and measured HNO₃ across the USA. McDuffie et al. (2018) have determined a median $\gamma = 0.0143$ (range: 2×10^{-5} to 0.1751) based on constrained modelling of aircraft observations. The inclusion of this median uptake coefficient would be likely to lead to an increase in the atmospheric mixing ratio of N_xO_y, and thus our results can be seen as lower bounds on the potential changes in [NO₃]. UKCA does not include any in-particle chemistry for N₂O₅ and also lacks aerosol nitrate in the present implementation, similar to some other CMIP6 class models. The omission of aerosol nitrate is a weakness in the model, but recent work including this process (Jones et al., 2021) suggests that the effects of it are small on N_xO_y.

The experimental setup is that of the simulations conducted as part of the Coupled Model Intercomparison Project Phase 6 (CMIP6; Eyring et al., 2016) DECK and ScenarioMIP experiments. Historical emissions are from the Community Emissions Data System (CEDS; Hoesly et al., 2018). Future emissions progress along one of four benchmark shared socioeconomic pathways (SSPs; Gidden et al., 2019). The analysis focuses on four representative pathways: SSP1-2.6, SSP2-4.5, SSP3-7.0 and SSP5-8.5. Each SSP has different assumptions about the amount of emissions of air pollutant precursors and climate forcings (Gidden et al., 2019).

We perform our analyses focused on 5-year periods (pre-industrial, PI: 1850–1854; present day, PD: 2010–2014; end of century: 2090–2094). These were conducted by performing new simulations using re-start files from the start dumps from the core CMIP6 simulations contributed by UKESM1. It was necessary to re-run the CMIP6 simulations with UKESM1 in order to (i) provide high temporal-resolution output with additional diagnostics for the NO₃ reactions which did not exist in the CMIP6 runs, needed to assess the oxidation of Monoterp and isoprene, and (ii) include a correction to the NO₃ + Monoterp reaction rate coefficient.

2.2 Description of relevant chemical reactions

The representation of BVOC chemistry in the StratTrop chemical mechanism used in UKESM1 is similar to other Earth system models in being very simplified. Isoprene is treated as an individual compound undergoing reactions with OH, O₃ and NO₃, while monoterpenes are represented by a surrogate species, Monoterp, which can undergo oxidation via OH, NO₃ or O₃ with rate coefficients equal to the equivalent oxidation reactions of α -pinene to form an inert species, SEC_ORG, which irreversibly condenses to form SOA (Mann et al., 2010). This simplification has been shown to broadly capture the relation between BVOC emissions, SOA formation and the climate impacts well (Scott et al., 2017; Mulcahy et al., 2020).



where F is some factor between 0 and 1 representing the yield of SOA production from Monoterp. The assumed yield of SEC_ORG from Monoterp is 0.13 but is doubled to 0.26 in order to account for the missing production from isoprene oxidation (Mann et al., 2010; Mulcahy et al., 2020). The rate coefficients k_{OH} , k_{O_3} and k_{NO_3} are equal to the rate coefficients for the reactions between α -pinene and OH, O₃ and NO₃, respectively, from Atkinson et al. (1989). Due to an error, the reaction rate coefficient activation energy for the NO₃ + Monoterp reaction was incorrect in the original version of UKESM1 and has been corrected for these experiments. The rate coefficient used in all previous studies using StratTrop with the GLOMAP aerosol was

$$k_{\text{NO}_3} = 1.19 \times 10^{-12} \times e^{(925/T)} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1}, \quad (\text{R4})$$

whereas the correct form is

$$k_{\text{NO}_3}^* = 1.19 \times 10^{-12} \times e^{(490/T)} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1}. \quad (\text{R5})$$

At $T = 298 \text{ K}$, this difference in activation energy results in k_{NO_3} being 4.3 times faster than $k_{\text{NO}_3}^*$. This correction results in a smaller fraction of Monoterp being oxidised by NO₃ compared to OH and O₃, partly compensated for by an increase in NO₃ burden (see Supplement further details).




Figure 1. Comparison of vertical profile of nighttime concentrations of O_3 (a), NO_2 (b), NO_3 (c) and the $\text{NO}_3 : \text{N}_2\text{O}_5$ ratio (d) from the RONOCO July 2010 flight campaign and 2010–2014 July nighttime values from UKESM1 simulations from grid cells corresponding to flight regions. Lines show means, and error bars show standard deviation.

3 Results

3.1 Comparison of present-day modelled and observed NO_3

Our analysis begins by evaluating the performance of UKESM1 against aircraft observations and pre-compiled ground-based observations presented in Khan et al. (2015). Figure 1 shows the comparison of UKESM1 modelled vertical profiles of O_3 , NO_2 , NO_3 and $\text{NO}_3 : \text{N}_2\text{O}_5$ against observations made in the eastern UK, over the North Sea, as part of the RONOCO summer campaign that took place in July 2010 (Stone et al., 2014). UKESM1 output is not constrained by meteorological forcing, so we sample the model during July 2010–2014, outputting model data once every 27 h and sampling data between 21:00 and 03:00 LT to generate a nighttime climatology for July, which is plotted in orange in Fig. 1 with the error bars reflecting the variability (1σ) in the model

climatology. In general the model simulates the vertical profile of the RONOCO observations well. There is a positive bias in modelled O_3 and a negative bias in modelled NO_2 (consistent with previous analyses with UKCA the chemistry component of UKESM1; e.g. Archibald et al., 2020b). These biases partly offset each other; the vertical profile of the rate of production of NO_3 (Reaction R1) is in good qualitative agreement between the model and observations, with maxima found above the surface below 1 km, but the absolute magnitude of the NO_3 production rate is underestimated in the model by a factor of ~ 2 in this region. One reason for the model disagreement is likely the resolution, whereby UKESM1 is unable to simulate the fine plumes of NO_2 that drive NO_3 production that were observed during the RONOCO campaign (Stone et al., 2014). In addition to the poor horizontal resolution, models like UKESM1 also suffer from biases in vertical resolution and mixing. The simulation of boundary layer height in models like UKESM1 is difficult,

and Fig. 1 suggests that the model boundary layer height is much lower than the observations. This would make sense if in the model there is a significant land fraction in the grid boxes being analysed, as is the case.

Table 1 compares results from the Historical simulation of UKESM1 against the STOCHEM-CRI model data from Khan et al. (2015) and a synthesis of surface observations; see Khan et al. (2015) for details and references for observational data. Our model data are averages from the 2010–2014 period corresponding to the months of each campaign and are in good qualitative and quantitative agreement with the observations (which come from a range of different techniques and cover a range of different dates over the period of ca. the 1990s–2010s; see Khan et al. (2015) for details). The largest disagreement in Table 1 is at Guangzhou, China, where both models predict much lower levels of NO_3 than were observed using a long-path DOAS (differential optical absorption spectroscopy) instrument (Li et al., 2012). Despite being of low horizontal and vertical resolution and in light of the caveats already discussed through the analysis of Fig. 1, we find that UKESM1 reproduces well current observations of NO_3 suggesting it is a suitable tool for simulating past and future changes in the NO_3 burden and distribution.

3.2 Modelled past and future changes in O_3 , NO_x and NO_3

Our analysis focuses on 5-year periods from the simulations, comparing present-day (PD; 2010–2014) and preindustrial (PI; 1850–1854) periods with end of century (2090–2094) predictions under four shared socioeconomic pathways (SSPs) as used for CMIP6. Data for these 5-year periods were collected by re-running the UKESM1 CMIP6 simulations with additional diagnostics for analysis and corrected NO_3 + Monoterp reaction rate coefficient (see the Methods section for details). The future emission pathways we studied, described in more detail in Gidden et al. (2019), are in short the following: SSP1-2.6, a 2 °C world with a focus on sustainability, economic growth and connectivity and low population growth driven by low-carbon technology and energy efficiency; SSP2-4.5, a middle-of-the-road scenario with slower convergence of income levels, greater population growth and reliance on fossil fuels further into the century; SSP3-7.0, a regional rivalry scenario with increased inequality and high population growth in low- to middle-income countries; and SSP5-8.5, similar to SSP1 in terms of population and economic growth but driven by increasing unabated use of energy and fossil fuels. Although SSP5-8.5 leads to the greatest increase in long-lived greenhouse gas emissions, and therefore global warming, by the end of the century, SSP3-7.0 has the highest emissions of short-lived air pollutants (Rao et al., 2017). Emissions of BVOCs also change between these scenarios as they are interactive in UKESM1 (Sellar et al., 2019). BVOC emissions are higher in the PI compared to the PD due to changing land use, reducing forested area

over the 20th century. In the future SSP scenarios, BVOC emissions are predicted to increase as temperatures increase, in spite of the increasing CO_2 levels (which, in the absence of other changes, would cause BVOC emissions to decrease; Arneth et al., 2008).

The results from the UKESM1 simulations show that large changes to monthly average O_3 , NO_2 and NO_3 levels have taken place in the lowest kilometre of the atmosphere from the PI to the PD (Fig. 2a–c). The region of the lowest kilometre is analysed here, rather than the surface level, because nighttime O_3 and NO_3 levels tend to be low at the surface due to rapid reaction with fresh emissions of NO and the low nighttime boundary layer. In both observations and the model, peak NO_3 tends to occur a few hundred metres above the surface (e.g. Fish et al., 1999).

The main driver for the changes between the PI and PD periods is attributed to the increase in anthropogenic emissions, with emissions of NO_x increasing by orders of magnitude over this period. The largest increases in NO_2 and O_3 occur over populated regions (Fig. 2a and b). The changes in O_3 simulated with UKESM1 are in good agreement with other CMIP6 models (Griffiths et al., 2021) and show an increase in the tropospheric burden of around 40 % from the PI–PD. NO_3 is similarly much higher in the PD compared to the PI (Fig. 2c), an increase in tropospheric burden of approximately 75 %, in good agreement with previous global modelling studies (Khan et al., 2015) and with observations (see Sect. 2.1). To first order, NO_3 is proportional to $\text{O}_3 \times \text{NO}_2$ as these species drive its production, but there are some important deviations from this relationship. Firstly, concentrations are lower at high latitudes because colder climates mean more NO_3 is partitioned into N_2O_5 . Changes in NO_3 are also suppressed in regions with high BVOC emissions, such as the south-east USA, South America and South-East Asia/South China, due to the high reactivity with these species. The greatest increases are over the Middle East and the Indian subcontinent – regions with high local temperatures, NO_x emissions and O_3 as well as relatively low BVOC emissions.

The simulated future changes in O_3 , NO_2 and NO_3 in the lowest 1 km depend greatly on the future SSP scenario. Focusing on the change between the future and the PD, the SSP1-2.6 scenario shows large reductions in O_3 , NO_2 and NO_3 across the world, especially in the Northern Hemisphere (Fig. 2d–f). These reductions are similar to but smaller than the increases in NO_3 from the PI to the PD, leading to a tropospheric NO_3 burden under SSP1-2.6 that is 26 % lower than the PD burden. In SSP2-4.5 (Fig. 2g–i) NO_2 decreases in most regions, although to a lesser degree in some regions such as India. However, reductions in O_3 are muted, with similar concentrations across most of the Northern Hemisphere, likely due to competing trends from changes to stratosphere-to-troposphere transport of ozone (driven by increases in the Brewer–Dobson circulation under climate change), lower NO_x emissions (decreasing ozone),

Table 1. Monthly average values from of NO₃ mixing ratios (ppt) from UKESM1 compared with a review of NO₃ measurements and STOCHEM model output adapted from Table 3 in Khan et al. (2015).

Site	Start	Observed (ppt)	STOCHEM (ppt)	UKESM1 (ppt)
Lindenberg	Feb–Sep 1998	4.6	6.9	8.0
Tanus	May 2008	30 ± 20	6.9	6.9
Jerusalem	Jul 2005–Sep 2007	27 ± 43	9.2	11.2
Houston	Aug–Sep 2006	0–149	3.3	11.0
Shanghai	Aug–Oct 2011	16 ± 9	16.6	8.4
Guangzhou	Jul 2006	21.8 ± 1.8	4.7	4.4
Izu Oshima	Jun 2004	3	4.4	7.5
Mace Head	Jul–Aug 1996	5	3.8	3.1
Canary Islands	May 1994	8 ± 3	5	5.7
Finokalia	Jun 2001–Sep 2003	4.2 ± 2.3	16	16.9
Weybourne	Oct–Nov 2004	6	5.5	4.2
East Point	Jul–Aug 2005	13.1	1.6	1.4
Schauinsland	Aug 1990	5.8	12	6.3




Figure 2. Changes in O₃, NO₂ and NO₃ mixing ratios averaged over lowest 1 km of the atmosphere, showing difference between the present-day (2010–2014) and the preindustrial period (1850–1854) (a–c), SSP1-2.6 (2090–2094) – PD (d–f), SSP2-4.5 (2090–2094) – PD (g–i), SSP3-7.0 (2090–2094) – PD (j–l), and SSP5-8.5 (2090–2094) – PD (m–o).

and higher temperatures and BVOC emissions (increasing ozone) (e.g. Archibald et al., 2020a). Changes in NO_3 are spatially variable, with small reductions in North America and China but increasing levels in northern Europe, India and parts of Africa. In the SSP3-7.0 scenario (Fig. 2j–l), which simulates the greatest increase in emissions of short-lived pollutants, near-surface O_3 is predicted to increase over most populated regions, particularly in South Asia and West Africa. In contrast, NO_2 shows diverging trends, decreasing in North America, Europe and China but increasing considerably in India, the Middle East and West Africa. Over most parts of the land and oceans NO_3 is predicted to not change significantly, but specific regions show dramatic increases: by over 10 ppt in West Africa, the Middle East and India (Fig. 2l). These large increases in NO_3 are of a similar order to the PI–PD changes shown in Fig. 2c, effectively doubling near-surface NO_3 concentrations in the future over many populated regions, whilst O_3 increases are only in the order of 50 % greater than the PD. In SSP5-8.5 (Fig. 2m–o), the warmest future scenario, the increases in O_3 in Asia are small and NO_2 is predicted to remain similar to the present day in India whilst decreasing significantly in China. However, there is still a considerable increase in NO_3 over India as well as West Africa and Europe predicted under the SSP5-8.5 scenario (in the order of 10–20 ppt increases above PD levels). These variations in trends between scenarios show that it is not sufficient to assume that lower NO_x emissions in the future will result in reduced NO_3 . Rather, it is a complicated outcome also depending on other changes in the chemical environment and climate.

To investigate the role of changes in climate on NO_3 , through changes in temperature over the lowest 1 km of the atmosphere, the logarithm of K_{eq} (Reaction R3) is plotted against the corresponding temperature values in Fig. 3, using data from each of the scenarios. This shows the strong dependence of K_{eq} , in accordance with the Van 't Hoff relationship. As the climate warms the thermal equilibrium shifts such that more N_xO_y is found as NO_3 , increasing the amount of NO_3 available for oxidation and reducing the sink due to the N_2O_5 heterogeneous reactions. UKESM1 simulates levels of warming by the end of the 21st century that are at the higher end of the CMIP6 multi-model mean but within the ensemble spread (Tebaldi et al., 2021).

3.3 Changes to BVOC oxidation

Figure 4 shows vertically integrated modelled oxidation fluxes for isoprene and Monoterp (the lumped monoterpene species in UKESM1) averaged across the globe and over southern Asia (SA; defined as 5° N, 50° E to 35° N, 95° E, based on the source–receptor region used by the Task Force on Hemispheric Transport of Air Pollution, TF HTAP, <https://www.htap.org>, last access: 1 March 2022; plots for the other TF HTAP regions are included in the Supplement). The modelled oxidation flux is calculated at each model chemistry




Figure 3. Van 't Hoff isochore of the thermal equilibrium between NO_3 and N_2O_5 (a) and change in temperature in UKESM1 simulations from 1850 to 2100 in four representative SSP scenarios and (b) changes in temperature from the historical simulation from 1850 to 2014 and along SSP scenarios from 2014 to 2100.

time step (1 h), thereby taking into account the concentrations of the BVOC, oxidants and the temperature-dependent rate coefficients for their reactions, which are then averaged over each month for analysis. The fluxes are further averaged over the lowest 1 km of the atmosphere to reflect boundary layer oxidation in Fig. 4. In the PI atmosphere, NO_3 -initiated oxidation accounts for less than 1 % of isoprene and 4 % of Monoterp, due to the low NO_x emissions in this period (Fig. 2b). This fraction of oxidation increases to 3 % and 13 % for isoprene and Monoterp, respectively, for the present day (i.e. a factor of 3 increase). UKESM1 predicts that monoterpene emissions increase in a warming climate in all scenarios, particularly in SSP3-7.0 and SSP5-8.5. Across the whole troposphere, the relative amount via NO_3 stays the same in SSP3-7.0 relative to the PD, although the absolute amount increases as BVOC emissions increase. However, over the South Asian region the fraction of Monoterp oxidised by NO_3 increases from 13 % to 15 %. In SSP1-2.6, the Monoterp oxidation fraction by NO_3 decreases in all regions evaluated.

Focusing on Monoterp oxidation, which in UKESM1 leads to the formation of SOA, Fig. 5 shows the vertical profiles of the oxidation of this compound under PD conditions (Fig. 5a and c) and the SSP3-7.0 future scenario (Fig. 5b and d). Similar figures for the other SSP scenarios are included in the Supplement. As with Fig. 4, Fig. 5 shows that the future scenario results in an increase in oxidation of Monoterp (an increase in SOA production in the model) for all oxidants. The vertical profiles show a sharp drop in the oxidation flux away from the surface as a result of the relatively short lifetime of the Monoterp species, which is emitted in the model at the surface and decays away in the vertical rapidly. Focusing on the near surface, where the rate of SOA production via Monoterp oxidation is highest, Fig. 5 shows that over southern Asia UKESM1 pre-




Figure 4. Average oxidation rates in lower 1 km of atmosphere for isoprene (a, b) and Monoterp (c, d), averaged across the whole globe (a, c) and over the South Asia region (b, d).




Figure 5. Vertical profile of Monoterp oxidation via OH, O₃ and NO₃. Panels (a) and (c) show the model-calculated present-day global and South Asian oxidation fluxes, respectively. Panels (b) and (d) show these fluxes at the end of the 21st century under the SSP3-7.0 scenario. The future scenario results in increases in all oxidation fluxes, with the NO₃-initiated oxidation flux increasing by a factor of 2 compared with the present-day oxidation rates in South Asia.

dicts an increase in near-surface NO_3 oxidation of Monoterp of a factor of 2 under the SSP3-7.0 scenario. This implies potential for enhanced production of organic nitrogen containing aerosols under future climate and emission scenarios, compounds whose health and climatic effects have been hitherto less well studied but which observational evidence suggests are ubiquitous in the atmosphere (Kiendler-Scharr et al., 2016).

4 Discussion

Whilst we have demonstrated large potential changes in NO_3 in the future under the SSP scenarios, we acknowledge that there are some limitations to the state-of-the-art CMIP6 models, like UKESM1. Faithful simulation of NO_3 requires models which capture a wide range of processes as discussed in Ng et al. (2017). These include the ability to simulate faithfully the diurnal variability in the boundary layer, which has been shown to be a challenge for these types of models (for example in East Asia; Yue et al. (2021) have shown that CMIP6 models cannot reproduce observed trends in the boundary layer height), and simulating the diurnal emissions of important NO_3 sinks, like isoprene (Cao et al., 2021) – which almost all interactive chemistry CMIP6 models do – and NO_3 sources, like NO_x , which no CMIP6 models do. Further work is required to improve the representation of these key emissions sources and processes in global chemistry–climate models. In addition, the simplistic treatment of SOA in UKESM1 prevents us from being able to explore how the composition of SOA may change under the changing climate and emission scenarios that have been explored for CMIP6. Many previous studies have highlighted how the formation of SOA is highly sensitive to the conditions it is being formed under and how the composition of SOA will change concomitantly (e.g. Hoyle et al., 2011; Pye et al., 2010; Schwantes et al., 2019). These feedbacks between changes in SOA formation mechanism and SOA composition are being explored through future work in UKESM1.

5 Conclusions

In this work we have demonstrated the potential pathways for future evolution of NO_3 , the most important oxidant at night, in the lower troposphere. Analysis of our model simulations against historic measurements highlights that in spite of the relatively low resolution of UKESM1, it is able to capture the magnitude and variability in observations of NO_3 and its precursors (O_3 and NO_2). We have assessed four different future scenarios which span a wide range of possible NO_x and VOC emission pathways and levels of climate change, simulated using UKESM1. This is, to our knowledge, the first assessment of future simulations of NO_3 , and this work highlights the potential for significant increases in this major nighttime

oxidant. In particular we have shown that, depending on the emission scenario, regions of southern Asia are an area of particular interest with potential increases in NO_3 to unprecedented levels, more than double their present-day values. We also demonstrate the importance of climate change for NO_3 and show that the ratio of $\text{NO}_3 : \text{N}_2\text{O}_5$ is predicted to increase under the higher climate forcing future scenarios. The impacts of an increase in NO_3 -dominated BVOC oxidation are, as yet, uncertain. An increasing body of literature is examining the mechanistic pathways through which BVOCs and NO_3 react and the impacts of $\text{BVOC} + \text{NO}_3$ -derived SOA. He et al. (2021) have shown through detailed laboratory experiments using cavity-enhanced absorption spectroscopy that some of the organic nitrates in $\text{BVOC} + \text{NO}_3$ -derived SOA may serve as atmosphere-stable NO_x sinks or reservoirs and will absorb and scatter incoming solar radiation during the daytime leading to an anthropogenic radiative forcing component (given that NO_3 is primarily an anthropogenic species).

The potential increases in NO_3 are shown to be important in the context of enhancing the oxidation of BVOCs in the atmosphere. Due to the poorer understanding of SOA formation from NO_3 oxidation (Ng et al., 2017) than from OH or O_3 oxidation, these findings highlight the need for further lab and field studies to better understand the SOA-forming potential of BVOC oxidation initiated via NO_3 and the incorporation of these data into global chemistry and Earth system models. Our simulated changes in NO_3 are likely dependent on the following:

1. model structure; more work investigating the impacts of vertical and horizontal resolution is required. Our estimates are that low-resolution Earth system models like UKESM1 will underestimate the peak concentrations of NO_3 owing to poor vertical mixing.
2. emissions, including soil NO_x emissions, which are highly uncertain still, and the diurnal and hebdomadal variation in anthropogenic ones – as with most CMIP6 models the emissions in UKESM1 are prescribed without diurnal variations, which are likely to impact NO_3 more than other oxidants. Here we estimate that including vertical profiles of emissions in Earth system models will increase the lifetime of NO_x and hence the rate of NO_3 production, but detailed studies are needed to assess this highly non-linear system.
3. chemical processes, with simplifications in the treatment of N_2O_5 heterogeneous chemistry being a potentially key area (as discussed by Archibald et al. (2020a) in the context of ozone budgets). As eluded to in Sect. 4, a key missing component in UKESM1 is the state dependence of SOA formation. Whilst previous studies have used chamber studies to derive conditional state dependence for the yield of SOA as a function of NO_x (e.g. Pye et al., 2010), future work should more holis-

tically treat the suite of low-volatility SOA precursors and their formation in the gas phase and coupling to the aerosol phase, as discussed by, e.g., Schwantes et al. (2019).

- the chemical mechanism used; Archer-Nicholls et al. (2021) and Weber et al. (2021) show that the NO_y budget is highly sensitive to the use of a more comprehensive chemical mechanism in the UKESM1 framework.

In this work we assessed the impacts of changing k_{NO_3} (see the Supplement S1 and S2 for details). We found that it is very difficult to predict how even a relatively simple change, like changing the rate constant for $\text{BVOC} + \text{NO}_3$ reactions, will modify the output of an Earth system model. Decreasing k_{NO_3} results in vary spatially variable changes in NO_3 , NO_2 and O_3 . In spite of the large change in k_{NO_3} the changes in the fluxes of $\text{BVOC} + \text{NO}_3$ oxidation were much more muted. Dedicated uncertainty analysis looking at points (1) to (4) above are needed. Finally, almost all CMIP6 models that have simulated interactive chemistry have simulated changes in NO_3 , and we encourage the modelling community to undertake a multi-model analysis similar to those focused on OH (Stevenson et al., 2020) and O_3 (Griffiths et al., 2021). Further multi-model analyses are required to constrain the level of model uncertainty in predictions of NO_3 and the potential impacts this could have on future air quality and climate. Co-ordination and partnership between the NO_3 observational community and the modelling community could then allow the future changes in NO_3 to be observationally constrained.

Code availability. Due to intellectual property rights restrictions, we cannot provide the source code or documentation papers for the Unified Model (UM). The Met Office Unified Model is available for use under licence. A number of research organisations and national meteorological services use the UM in collaboration with the Met Office to undertake basic atmospheric process research, produce forecasts, develop the UM code, and build and evaluate Earth system models. For further information on how to apply for a licence, see <http://www.metoffice.gov.uk/research/modelling-systems/unified-model> (last access: 1 October 2021).

Data availability. The UM data used to produce the figures are available from the Centre of Environmental Data Analysis (CEDA) (see Supplement for details; Archer-Nicholls et al., 2023). Data from five experiments are included in the archive: ScenarioMIP SSP1-2.6 (ssp126), SSP2-4.5 (ssp245), SSP3-7.0 (ssp370), SSP5-8.5 (ssp585) experiments and the CMIP Historical experiment (hist) (<https://doi.org/10.5285/b70e6ae10a9f463d88819eb981cd4d0f>; Archer-Nicholls et al., 2023). For the ScenarioMIP experiments, data are archived for the years 2090–2094. For the CMIP Historical experiment, data are archived for the years 1850–1854. The data comprise monthly mean output from both environmental variables, tracer mass mixing ratio and chemical reaction tendencies in moles per grid box per second. Other relevant data may be found in the

CMIP6 archive for CMIP and ScenarioMIP hosted at the Centre for Environmental Data Analysis (CEDA) and at the Earth System Federation Grid.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-23-5801-2023-supplement>.

Author contributions. ATA designed the research and supervised the analysis and led the revision of the paper. SAN and RA led the analysis. NLA and SAN performed the re-runs of UKESM1. PTG contributed to the analysis. All authors contributed to writing the paper.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. The authors would like to thank the UK Met Office Clean Air SPF and the JWCRP UKCA project. UKCA is a collaborative programme supported by NCAS and the Met Office, and we wish to acknowledge them for their continuing support of the project. This work used JASMIN, the UK collaborative data analysis facility.

Financial support. This research has been supported by the Natural Environment Research Council (grant no. NE/P016383/1).

Review statement. This paper was edited by Kostas Tsigaridis and reviewed by two anonymous referees.

References

- Archer-Nicholls, S., Abraham, N. L., Shin, Y. M., Weber, J., Russo, M. R., Lowe, D., Utembe, S. R., O'Connor, F. M., Kerridge, B., Latter, B., Siddans, R., Jenkin, M., Wild, O., and Archibald, A. T.: The Common Representative Intermediates Mechanism Version 2 in the United Kingdom Chemistry and Aerosols Model, *J. Adv. Model. Earth Sy.*, 13, 1–37, <https://doi.org/10.1029/2020MS002420>, 2021.
- Archer-Nicholls, S., Allen, R., Shin, Y., Abraham, N. L., Griffiths, P., and Archibald, A.: UKESM1 diagnostics for CMIP6 ScenarioMIP and CMIP historical experiments, NERC EDS Centre for Environmental Data Analysis [data set], <https://doi.org/10.5285/b70e6ae10a9f463d88819eb981cd4d0f>, 2023.
- Archibald, A. T., Neu, J. L., Elshorbany, Y. F., Cooper, O. R., Young, P. J., Akiyoshi, H., Cox, R. A., Coyle, M.,

- Derwent, R. G., Deushi, M., and Finco, A.: Tropospheric Ozone Assessment Report: A critical review of changes in the tropospheric ozone burden and budget from 1850 to 2100, *Elementa: Science of the Anthropocene*, 8, 1–53, <https://doi.org/10.1525/elementa.2020.034>, 2020a.
- Archibald, A. T., O'Connor, F. M., Abraham, N. L., Archer-Nicholls, S., Chipperfield, M. P., Dalvi, M., Folberth, G. A., Denison, F., Dhomse, S. S., Griffiths, P. T., Hardacre, C., Hewitt, A. J., Hill, R. S., Johnson, C. E., Keeble, J., Köhler, M. O., Morgenstern, O., Mulcahy, J. P., Ordóñez, C., Pope, R. J., Rumbold, S. T., Russo, M. R., Savage, N. H., Sellar, A., Stringer, M., Turnock, S. T., Wild, O., and Zeng, G.: Description and evaluation of the UKCA stratosphere–troposphere chemistry scheme (Strat-Trop v1.0) implemented in UKESM1, *Geosci. Model Dev.*, 13, 1223–1266, <https://doi.org/10.5194/gmd-13-1223-2020>, 2020b.
- Arneth, A., Niinemets, Ü., Pressley, S., Bäck, J., Hari, P., Karl, T., Noe, S., Prentice, I. C., Serça, D., Hickler, T., Wolf, A., and Smith, B.: Process-based estimates of terrestrial ecosystem isoprene emissions: incorporating the effects of a direct CO₂–isoprene interaction, *Atmos. Chem. Phys.*, 7, 31–53, <https://doi.org/10.5194/acp-7-31-2007>, 2007.
- Arneth, A., Monson, R. K., Schurgers, G., Niinemets, Ü., and Palmer, P. I.: Why are estimates of global terrestrial isoprene emissions so similar (and why is this not so for monoterpenes)?, *Atmos. Chem. Phys.*, 8, 4605–4620, <https://doi.org/10.5194/acp-8-4605-2008>, 2008.
- Atkinson, R. and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: A review, *Atmos. Environ.*, 37, 197–219, [https://doi.org/10.1016/S1352-2310\(03\)00391-1](https://doi.org/10.1016/S1352-2310(03)00391-1), 2003.
- Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, R. F., Kerr, J. A., and Troe, J.: Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry: Supplement III, IUPAC Subcommittee on Gas Kinetic Data Evaluation for Atmospheric Chemistry, *J. Phys. Chem. Ref. Data*, 18, 881–1097, <https://doi.org/10.1063/1.555832>, 1989.
- Bianchi, F., Kurtén, T., Riva, M., Mohr, C., Rissanen, M. P., Roldin, P., Berndt, T., Crounse, J. D., Wennberg, P. O., Mentel, T. F., Wildt, J., Junninen, H., Jokinen, T., Kulmala, M., Worsnop, D. R., Thornton, J. A., Donahue, N., Kjaergaard, H. G., and Ehn, M.: Highly Oxygenated Organic Molecules (HOM) from Gas-Phase Autoxidation Involving Peroxy Radicals: A Key Contributor to Atmospheric Aerosol, *Chem. Rev.*, 119, 3472–3509, <https://doi.org/10.1021/acs.chemrev.8b00395>, 2019.
- Boyd, C. M., Sanchez, J., Xu, L., Eugene, A. J., Nah, T., Tuet, W. Y., Guzman, M. I., and Ng, N. L.: Secondary organic aerosol formation from the β -pinene + NO₃ system: effect of humidity and peroxy radical fate, *Atmos. Chem. Phys.*, 15, 7497–7522, <https://doi.org/10.5194/acp-15-7497-2015>, 2015.
- Brewer, D. A., Ogliaruso, M. A., Augustsson, T. R., and Levine, J. S.: The oxidation of isoprene in the troposphere: Mechanism and model calculations, *Atmos. Environ.* (1967), 18, 2723–2744, [https://doi.org/10.1016/0004-6981\(84\)90338-X](https://doi.org/10.1016/0004-6981(84)90338-X), 1984.
- Brown, S. S. and Stutz, J.: Nighttime radical observations and chemistry, *Chem. Soc. Rev.*, 41, 6405–6447, 2012.
- Cao, Y., Yue, X., Liao, H., Yang, Y., Zhu, J., Chen, L., Tian, C., Lei, Y., Zhou, H., and Ma, Y.: Ensemble projection of global isoprene emissions by the end of 21st century using CMIP6 models, *Atmos. Environ.*, 267, 118766, <https://doi.org/10.1016/j.atmosenv.2021.118766>, 2021.
- Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506, 476–479, <https://doi.org/10.1038/nature13032>, 2014.
- Ehn, M., Berndt, T., Wildt, J., and Mentel, T.: Highly oxygenated molecules from atmospheric autoxidation of hydrocarbons: a prominent challenge for chemical kinetics studies, *Int. J. Chem. Kinet.*, 49, 821–831, 2017.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Faxon, C., Hammes, J., Le Breton, M., Pathak, R. K., and Hallquist, M.: Characterization of organic nitrate constituents of secondary organic aerosol (SOA) from nitrate-radical-initiated oxidation of limonene using high-resolution chemical ionization mass spectrometry, *Atmos. Chem. Phys.*, 18, 5467–5481, <https://doi.org/10.5194/acp-18-5467-2018>, 2018.
- Fish, D. J., Shallcross, D. E., and Jones, R. L.: The vertical distribution of NO₃ in the atmospheric boundary layer, *Atmos. Environ.*, 33, 687–691, [https://doi.org/10.1016/S1352-2310\(98\)00332-X](https://doi.org/10.1016/S1352-2310(98)00332-X), 1999.
- Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Wooldridge, P. J., Brown, S. S., Fuchs, H., Dubé, W., Mensah, A., dal Maso, M., Tillmann, R., Dorn, H.-P., Brauers, T., and Cohen, R. C.: Organic nitrate and secondary organic aerosol yield from NO₃ oxidation of β -pinene evaluated using a gas-phase kinetics/aerosol partitioning model, *Atmos. Chem. Phys.*, 9, 1431–1449, <https://doi.org/10.5194/acp-9-1431-2009>, 2009.
- Gidden, M. J., Riahi, K., Smith, S. J., Fujimori, S., Luderer, G., Kriegler, E., van Vuuren, D. P., van den Berg, M., Feng, L., Klein, D., Calvin, K., Doelman, J. C., Frank, S., Fricko, O., Harmsen, M., Hasegawa, T., Havlik, P., Hilaire, J., Hoesly, R., Horing, J., Popp, A., Stehfest, E., and Takahashi, K.: Global emissions pathways under different socioeconomic scenarios for use in CMIP6: a dataset of harmonized emissions trajectories through the end of the century, *Geosci. Model Dev.*, 12, 1443–1475, <https://doi.org/10.5194/gmd-12-1443-2019>, 2019.
- Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., Kürten, A., and Carslaw, K. S.: Causes and importance of new particle formation in the present-day and preindustrial atmospheres, *J. Geophys. Res.-Atmos.*, 122, 8739–8760, <https://doi.org/10.1002/2017JD026844>, 2017.
- Griffiths, P. T., Murray, L. T., Zeng, G., Shin, Y. M., Abraham, N. L., Archibald, A. T., Deushi, M., Emmons, L. K., Galbally, I. E., Hassler, B., Horowitz, L. W., Keeble, J., Liu, J., Moeini, O., Naik, V., O'Connor, F. M., Oshima, N., Tarasick, D., Tilmes, S., Turnock, S. T., Wild, O., Young, P. J., and Zanis, P.: Tropospheric ozone in CMIP6 simulations, *Atmos. Chem. Phys.*, 21, 4187–4218, <https://doi.org/10.5194/acp-21-4187-2021>, 2021.

- Hamilton, J. F., Bryant, D. J., Edwards, P. M., Ouyang, B., Bannan, T. J., Mehra, A., Mayhew, A. W., Hopkins, J. R., Dunmore, R. E., Squires, F. A., Lee, J. D., Newland, M. J., Worrall, S. D., Bacak, A., Coe, H., Percival, C., Whalley, L. K., Heard, D. E., Slater, E. J., Jones, R. L., Cui, T., Surratt, J. D., Reeves, C. E., Mills, G. P., Grimmond, S., Sun, Y., Xu, W., Shi, Z., and Rickard, A. R.: Key Role of NO_3 Radicals in the Production of Isoprene Nitrates and Nitrooxyorganosulfates in Beijing, *Environ. Sci. Technol.*, 55, 842–853, <https://doi.org/10.1021/acs.est.0c05689>, 2021.
- Hantson, S., Knorr, W., Schurgers, G., Pugh, T. A. M., and Arneth, A.: Global isoprene and monoterpene emissions under changing climate, vegetation, CO_2 and land use, *Atmos. Environ.*, 155, 35–45, <https://doi.org/10.1016/j.atmosenv.2017.02.010>, 2017.
- He, Q., Tomaz, S., Li, C., Zhu, M., Meidan, D., Riva, M., Laskin, A., Brown, S. S., George, C., Wang, X., and Rudich, Y.: Optical properties of secondary organic aerosol produced by nitrate radical oxidation of biogenic volatile organic compounds, *Environ. Sci. Technol.*, 55, 2878–2889, <https://doi.org/10.1021/acs.est.0c06838>, 2021.
- Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.
- Hoyle, C. R., Boy, M., Donahue, N. M., Fry, J. L., Glasius, M., Guenther, A., Hallar, A. G., Huff Hartz, K., Petters, M. D., Petäjä, T., Rosenoern, T., and Sullivan, A. P.: A review of the anthropogenic influence on biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 11, 321–343, <https://doi.org/10.5194/acp-11-321-2011>, 2011.
- Jones, A. C., Hill, A., Remy, S., Abraham, N. L., Dalvi, M., Hardacre, C., Hewitt, A. J., Johnson, B., Mulcahy, J. P., and Turnock, S. T.: Exploring the sensitivity of atmospheric nitrate concentrations to nitric acid uptake rate using the Met Office's Unified Model, *Atmos. Chem. Phys.*, 21, 15901–15927, <https://doi.org/10.5194/acp-21-15901-2021>, 2021.
- Khan, M. A. H., Cooke, M. C., Utembe, S. R., Archibald, A. T., Derwent, R. G., Xiao, P., Percival, C. J., Jenkin, M. E., Morris, W. C., and Shallcross, D. E.: Global modeling of the nitrate radical (NO_3) for present and pre-industrial scenarios, *Atmos. Res.*, 164–165, 347–357, <https://doi.org/10.1016/j.atmosres.2015.06.006>, 2015.
- Kiendler-Scharr, A., Mensah, A. A., Friese, E., Topping, D., Nemitz, E., Prévôt, A. S., Äijälä, M., Allan, J., Canonaco, F., Canagaratna, M., and Carbone, S.: Ubiquity of organic nitrates from nighttime chemistry in the European submicron aerosol, *Geophys. Res. Lett.*, 43, 7735–7744, <https://doi.org/10.1002/2016GL069239>, 2016.
- Li, S., Liu, W., Xie, P., Qin, M., and Yang, Y.: Observation of nitrate radical in the nocturnal boundary layer during a summer field campaign in pearl river delta China, *Terr. Atmos. Ocean. Sci.*, 23, 39–48, [https://doi.org/10.3319/TAO.2011.07.26.01\(A\)](https://doi.org/10.3319/TAO.2011.07.26.01(A)), 2012.
- Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., Pickering, S. J., and Johnson, C. E.: Description and evaluation of GLOMAP-mode: a modal global aerosol microphysics model for the UKCA composition-climate model, *Geosci. Model Dev.*, 3, 519–551, <https://doi.org/10.5194/gmd-3-519-2010>, 2010.
- McDuffie, E. E., Fibiger, D. L., Dubé, W. P., Lopez-Hilfiker, F., Lee, B. H., Thornton, J. A., Shah, V., Jaeglé, L., Guo, H., Weber, R. J., Reeves, J. M., Weinheimer, A. J., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., Dibb, J. E., Veres, P., Ebben, C., Sparks, T. L., Wooldridge, P. J., Cohen, R. C., Hornbrook, R. S., Apel, E. C., Campos, T., Hall, S. R., Ullmann, K., and Brown, S. S.: Heterogeneous N_2O_5 uptake during winter: Aircraft measurements during the 2015 WINTER campaign and critical evaluation of current parameterizations, *J. Geophys. Res.-Atmos.*, 123, 4345–4372, <https://doi.org/10.1002/2018JD028336>, 2018.
- McGillen, M. R., Carter, W. P. L., Mellouki, A., Orlando, J. J., Picquet-Varraut, B., and Wallington, T. J.: Database for the kinetics of the gas-phase atmospheric reactions of organic compounds, *Earth Syst. Sci. Data*, 12, 1203–1216, <https://doi.org/10.5194/essd-12-1203-2020>, 2020.
- Monks, P. S., Archibald, A. T., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G. E., Stevenson, D. S., Tarasova, O., Thouret, V., von Schneidemesser, E., Sommariva, R., Wild, O., and Williams, M. L.: Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, *Atmos. Chem. Phys.*, 15, 8889–8973, <https://doi.org/10.5194/acp-15-8889-2015>, 2015.
- Mulcahy, J. P., Johnson, C., Jones, C. G., Povey, A. C., Scott, C. E., Sellar, A., Turnock, S. T., Woodhouse, M. T., Abraham, N. L., Andrews, M. B., Bellouin, N., Browne, J., Carslaw, K. S., Dalvi, M., Folberth, G. A., Glover, M., Grosvenor, D. P., Hardacre, C., Hill, R., Johnson, B., Jones, A., Kipling, Z., Mann, G., Mollard, J., O'Connor, F. M., Palmieri, J., Reddington, C., Rumbold, S. T., Richardson, M., Schutgens, N. A. J., Stier, P., Stringer, M., Tang, Y., Walton, J., Woodward, S., and Yool, A.: Description and evaluation of aerosol in UKESM1 and HadGEM3-GC3.1 CMIP6 historical simulations, *Geosci. Model Dev.*, 13, 6383–6423, <https://doi.org/10.5194/gmd-13-6383-2020>, 2020.
- Ng, N. L., Brown, S. S., Archibald, A. T., Atlas, E., Cohen, R. C., Crowley, J. N., Day, D. A., Donahue, N. M., Fry, J. L., Fuchs, H., Griffin, R. J., Guzman, M. I., Herrmann, H., Hodzic, A., Iinuma, Y., Jimenez, J. L., Kiendler-Scharr, A., Lee, B. H., Lueken, D. J., Mao, J., McLaren, R., Mutzel, A., Osthoff, H. D., Ouyang, B., Picquet-Varraut, B., Platt, U., Pye, H. O. T., Rudich, Y., Schwantes, R. H., Shiraiwa, M., Stutz, J., Thornton, J. A., Tilgner, A., Williams, B. J., and Zaveri, R. A.: Nitrate radicals and biogenic volatile organic compounds: oxidation, mechanisms, and organic aerosol, *Atmos. Chem. Phys.*, 17, 2103–2162, <https://doi.org/10.5194/acp-17-2103-2017>, 2017.
- O'Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.-F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6, *Geosci. Model Dev.*, 9, 3461–3482, <https://doi.org/10.5194/gmd-9-3461-2016>, 2016.
- Pacifico, F., Folberth, G. A., Jones, C. D., Harrison, S. P., and Collins, W. J.: Sensitivity of biogenic isoprene emissions to past, present, and future environmental conditions and implications for atmospheric chemistry, *J. Geophys. Res.-Atmos.*, 117, D22302, <https://doi.org/10.1029/2012JD018276>, 2012.

- Pöschl, U.: Atmospheric aerosols: composition, transformation, climate and health effects, *Angew. Chem. Int. Edit.*, 44, 7520–7540, <https://doi.org/10.1002/anie.200501122>, 2005.
- Pye, H. O. T., Chan, A. W. H., Barkley, M. P., and Seinfeld, J. H.: Global modeling of organic aerosol: the importance of reactive nitrogen (NO_x and NO_3), *Atmos. Chem. Phys.*, 10, 11261–11276, <https://doi.org/10.5194/acp-10-11261-2010>, 2010.
- Rao, S., Klimont, Z., Smith, S. J., Van Dingenen, R., Dentener, F., Bouwman, L., Riahi, K., Amann, M., Bodirsky, B. L., van Vuuren, D. P., Aleluia Reis, L., Calvin, K., Drouet, L., Fricko, O., Fujimori, S., Gernaat, D., Havlik, P., Harmsen, M., Hasegawa, T., and Tavoni, M.: Future air pollution in the Shared Socio-economic Pathways, *Glob. Environ. Chang.*, 42, 346–358, <https://doi.org/10.1016/j.gloenvcha.2016.05.012>, 2017.
- Riemer, N., Vogel, H., Vogel, B., Schell, B., Ackermann, I., Kessler, C., and Hass, H.: Impact of the heterogeneous hydrolysis of N_2O_5 on chemistry and nitrate aerosol formation in the lower troposphere under photochemical conditions, *J. Geophys. Res.*, 108, 4144, <https://doi.org/10.1029/2002JD002436>, 2003.
- Sakulyanontvittaya, T., Duhl, T., Wiedinmyer, C., Helmig, D., Matsunaga, S., Potosnak, M., Milford, J., and Guenther, A.: Monoterpene and sesquiterpene emission estimates for the United States, *Environ. Sci. Technol.*, 42, 1623–1629, <https://doi.org/10.1021/es702274e>, 2008.
- Schwantes, R. H., Charan, S. M., Bates, K. H., Huang, Y., Nguyen, T. B., Mai, H., Kong, W., Flagan, R. C., and Seinfeld, J. H.: Low-volatility compounds contribute significantly to isoprene secondary organic aerosol (SOA) under high- NO_x conditions, *Atmos. Chem. Phys.*, 19, 7255–7278, <https://doi.org/10.5194/acp-19-7255-2019>, 2019.
- Schwantes, R. H., Emmons, L. K., Orlando, J. J., Barth, M. C., Tyn-dall, G. S., Hall, S. R., Ullmann, K., St. Clair, J. M., Blake, D. R., Wisthaler, A., and Bui, T. P. V.: Comprehensive isoprene and terpene gas-phase chemistry improves simulated surface ozone in the southeastern US, *Atmos. Chem. Phys.*, 20, 3739–3776, <https://doi.org/10.5194/acp-20-3739-2020>, 2020.
- Scott, C. E., Arnold, S. R., Monks, S. A., Asmi, A., Paasonen, P., and Spracklen, D. V.: Substantial large-scale feedbacks between natural aerosols and climate, *Nat. Geosci.*, 11, 1–6, <https://doi.org/10.1038/s41561-017-0020-5>, 2017.
- Sellar, A. A., Jones, C. G., Mulcahy, J., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., Mora, L., Kuhlbrodt, T., Rumbold, S., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., and Zerroukat, M.: UKESM1: Description and evaluation of the UK Earth System Model, *J. Adv. Model. Earth Sy.*, 11, 4513–4558, <https://doi.org/10.1029/2019ms001739>, 2019.
- Stevenson, D. S., Zhao, A., Naik, V., O'Connor, F. M., Tilmes, S., Zeng, G., Murray, L. T., Collins, W. J., Griffiths, P. T., Shim, S., Horowitz, L. W., Sentman, L. T., and Emmons, L.: Trends in global tropospheric hydroxyl radical and methane lifetime since 1850 from AerChemMIP, *Atmos. Chem. Phys.*, 20, 12905–12920, <https://doi.org/10.5194/acp-20-12905-2020>, 2020.
- Stone, D., Evans, M. J., Walker, H., Ingham, T., Vaughan, S., Ouyang, B., Kennedy, O. J., McLeod, M. W., Jones, R. L., Hopkins, J., Punjabi, S., Lidster, R., Hamilton, J. F., Lee, J. D., Lewis, A. C., Carpenter, L. J., Forster, G., Oram, D. E., Reeves, C. E., Bauguitte, S., Morgan, W., Coe, H., Aruffo, E., Dari-Salisbury, C., Giammaria, F., Di Carlo, P., and Heard, D. E.: Radical chemistry at night: comparisons between observed and modelled HO_x , NO_3 and N_2O_5 during the RONOCO project, *Atmos. Chem. Phys.*, 14, 1299–1321, <https://doi.org/10.5194/acp-14-1299-2014>, 2014.
- Tebaldi, C., Debeire, K., Eyring, V., Fischer, E., Fyfe, J., Friedlingstein, P., Knutti, R., Lowe, J., O'Neill, B., Sanderson, B., van Vuuren, D., Riahi, K., Meinshausen, M., Nicholls, Z., Tokarska, K. B., Hurtt, G., Kriegl, E., Lamarque, J.-F., Meehl, G., Moss, R., Bauer, S. E., Boucher, O., Brovkin, V., Byun, Y.-H., Dix, M., Gualdi, S., Guo, H., John, J. G., Kharin, S., Kim, Y., Koshiro, T., Ma, L., Olivie, D., Panickal, S., Qiao, F., Rong, X., Rosenbloom, N., Schupfner, M., Séférian, R., Sellar, A., Semmler, T., Shi, X., Song, Z., Steger, C., Stouffer, R., Swart, N., Tachiiri, K., Tang, Q., Tatebe, H., Voldoire, A., Volodin, E., Wyser, K., Xin, X., Yang, S., Yu, Y., and Ziehn, T.: Climate model projections from the Scenario Model Intercomparison Project (ScenarioMIP) of CMIP6, *Earth Syst. Dynam.*, 12, 253–293, <https://doi.org/10.5194/esd-12-253-2021>, 2021.
- Tröstl, J., Chuang, W. K., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C., Bianchi, F., Wagner, R., Simon, M., Lehtipalo, K., Williamson, C., Craven, J. S., Duplissy, J., Adamov, A., Almeida, J., Bernhammer, A.-K., Breitenlechner, M., and Baltensperger, U.: The role of low-volatility organic compounds in initial particle growth in the atmosphere, *Nature*, 533, 527–531, <https://doi.org/10.1038/nature18271>, 2016.
- Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosa-Mas, C. E., Hjorth, J., Bras, G. L., Moortgat, G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The Nitrate Radical: Physics, Chemistry, and the Atmosphere, *Atmos. Environ.*, 25A, 1–203, [https://doi.org/10.1016/0960-1686\(91\)90192-A](https://doi.org/10.1016/0960-1686(91)90192-A), 1991.
- Walters, D., Baran, A. J., Boutle, I., Brooks, M., Earnshaw, P., Edwards, J., Furtado, K., Hill, P., Lock, A., Mannes, J., Morcrette, C., Mulcahy, J., Sanchez, C., Smith, C., Stratton, R., Tennant, W., Tomassini, L., Van Weverberg, K., Vosper, S., Willett, M., Browse, J., Bushell, A., Carslaw, K., Dalvi, M., Essery, R., Gedney, N., Hardiman, S., Johnson, B., Johnson, C., Jones, A., Jones, C., Mann, G., Milton, S., Rumbold, H., Sellar, A., Ujiie, M., Whittall, M., Williams, K., and Zerroukat, M.: The Met Office Unified Model Global Atmosphere 7.0/7.1 and JULES Global Land 7.0 configurations, *Geosci. Model Dev.*, 12, 1909–1963, <https://doi.org/10.5194/gmd-12-1909-2019>, 2019.
- Weber, J., Archer-Nicholls, S., Abraham, N. L., Shin, Y. M., Bannan, T. J., Percival, C. J., Bacak, A., Artaxo, P., Jenkin, M., Khan, M. A. H., Shallcross, D. E., Schwantes, R. H., Williams, J., and Archibald, A. T.: Improvements to the representation of BVOC chemistry–climate interactions in UKCA (v11.5) with the CRI-Strat 2 mechanism: incorporation and evaluation, *Geosci. Model Dev.*, 14, 5239–5268, <https://doi.org/10.5194/gmd-14-5239-2021>, 2021.
- Yue, M., Wang, M., Guo, J., Zhang, H., Dong, X., and Liu, Y.: Long-Term Trend Comparison of Planetary Boundary Layer Height in Observations and CMIP6 Models over China, *J. Climate*, 34, 8237–8256, 2021.
- Zhao, D., Pullinen, I., Fuchs, H., Schrade, S., Wu, R., Acir, I.-H., Tillmann, R., Rohrer, F., Wildt, J., Guo, Y., Kiendler-Scharr, A., Wahner, A., Kang, S., Vereecken, L., and Mentel, T. F.: Highly oxygenated organic molecule (HOM) formation in the isoprene oxidation by NO_3 radical, *Atmos. Chem. Phys.*, 21, 9681–9704, <https://doi.org/10.5194/acp-21-9681-2021>, 2021.