



Supplement of

Convective organization and 3D structure of tropical cloud systems deduced from synergistic A-Train observations and machine learning

Claudia J. Stubenrauch et al.

Correspondence to: Claudia J. Stubenrauch (claudia.stubenrauch@lmd.ipsl.fr)

The copyright of individual parts of the supplement might differ from the article licence.

Artificial Neural Network predictions and evaluation




Figure S1: Box-and-whisker plots of $Z_{top}(\text{predicted}) - Z_{top}(\text{observed})$ and $DZ(\text{predicted}) - DZ(\text{observed})$ in km, for Cb, Ci, thin Ci, mid- and lowlevel clouds separately over ocean (left) and over land (right). The DZ results are shown for two developed models: The first model includes the predicted Z_{top} as input parameter (middle panels), the second does not (bottom panels). The boxes show the quartiles of the data while the whiskers extend to show the rest of the distribution.




Figure S2: Normalized distributions of $DZ(\text{predicted}) - DZ(\text{observed})$, for Cb over ocean and over land. Compared are Cbs for which $DZ > 10$ km is predicted (left) and those for which $DZ < 10$ km is predicted. The prediction models have been applied to 20% of the collocated data and are compared with the results derived from CloudSat-lidar 2B GEOPROF data.




Figure S3: Annual cycle of I_{org} , ROME, ABCOP, COP, mean area and total area estimated from convective areas defined by different variables. These areas are built from grid cells covered by at least 90% UT clouds, with rain rate indicator > 2 , $T_{\text{clld}} < 230 \text{ K}$ & $\epsilon_{\text{clld}} > 0.95$, or using the 2% largest rain rate indicator. The latter leads to a constant total area of convection. Monthly statistics of UT clouds averaged over four observation times from 2008 to 2018.

35




Figure S4: Geographical maps of intense precipitation (indicated by rain rate indicator > 2) occurrence, separately for the four seasons of boreal winter, spring, summer and autumn. Monthly statistics averaged over four observation times from 2008 to 2018.

40






Figure S5: Annual cycle of anvil vertical extent. Monthly statistics averaged over four observation times from 2008 to 2018.

45



50 **Figure S6: Scatter distributions of the 12 month running mean UT radiative heating rate and I_{org} anomalies, the latter obtained by using intense precipitation as proxy to define convective areas, separately for two locations in the tropics: 0N-140E (Warm Pool, left) and 5S-170W (Central Pacific, right). Linear regression fits are also shown, with correlation coefficients. Monthly statistics of UT clouds averaged over four observation times from 2008 to 2018.**

55



60 **Figure S7: Change in radiative heating rates with Oceanic Niño index (ONI). The troposphere is divided into three layers: upper troposphere (100-200 hPa), mid troposphere (200-600 hPa), and low troposphere (600-900 hPa). Monthly statistics from 2008 to 2018.**




Figure S8: Time series of deseasonalized monthly anomalies of I_{org} , ROME, COP and ABCOP, using different proxies to define the convective areas.