



Water activity and surface tension of aqueous ammonium sulfate and D-glucose aerosol nanoparticles

Eugene F. Mikhailov¹, Sergey S. Vlasenko¹, and Alexei A. Kiselev²

¹Department of Atmospheric Physics, Saint Petersburg State University, St Petersburg, 199034, Russia

²Atmospheric Aerosol Research Department, Institute for Meteorology and Climate Research, Karlsruhe Institute of Technology (KIT), Hermann-von-Helmholtz Platz 1, 76344 Eggenstein-Leopoldshafen, Germany

Correspondence: Alexei A. Kiselev (alexei.kiselev@kit.edu)

Received: 9 August 2023 – Discussion started: 23 August 2023

Revised: 16 January 2024 – Accepted: 18 January 2024 – Published: 7 March 2024

Abstract. Water activity (a_w) and interfacial energy or surface tension (σ) are key thermodynamic parameters to describe the hygroscopic growth of atmospheric aerosol particles and their ability to serve as cloud condensation nuclei (CCN), thus influencing the hydrological cycle and climate. Due to size effects and complex mixing states, however, these parameters are not well constrained for nanoparticles composed of organic and inorganic compounds in aqueous solution.

In this study, we determined a_w and σ by differential Köhler analysis (DKA) of hygroscopic growth measurement data for aerosol particles smaller than 100 nm composed of aqueous ammonium sulfate (AS), D-glucose (GI), and their mixtures. High-precision measurements of hygroscopic growth were performed at relative humidities (denoted RH) ranging from 2.0 % to 99.6 % with a high-humidity tandem differential mobility analyzer (HHTDMA) in three complementary modes of operation: hydration, dehydration, and restructuring. The restructuring mode (hydration followed by dehydration) enabled the transformation of initially irregular particles into compact globules and the determination of mass equivalent diameters. The HHTDMA-derived growth factors complemented by DKA allows for determination of water activity and surface tension from dilute to highly supersaturated aqueous solutions that are not accessible with other methods. Thus, for mixed AS/GI nanoparticles with mass ratios of 4 : 1 and 1 : 1, the upper limit of solute mass fraction (X_s) was 0.92 and 0.98, respectively.

For pure AS and GI, the DKA-derived a_w is in good agreement with electrodynamic balance and bulk measurement data. For AS particles, our a_w data also agree well with the Extended Aerosol Inorganics Model (E-AIM III) over the entire concentration range. In contrast, the UNIFAC model as a part of AIOMFAC (Zuend et al., 2011) was found to overestimate a_w in aqueous GI particles, which can be attributed to unaccounted intermolecular interactions.

For mixed AS and GI nanoparticles, we observed a non-monotonic concentration dependence of the surface tension that does not follow the predictions by modeling approaches constructed for mixed inorganic/organic systems. Thus, AS/GI particles with a 1 : 1 mass ratio exhibited a strong decrease of σ with increasing solute mass fraction, a minimum value of 56.5 mN m^{-1} at $X_s \approx 0.5$, and a reverse trend of increasing σ at higher concentrations. We suggest that D-glucose molecules surrounded by ammonium sulfate ions tend to associate, forming non-polar aggregates, which lowers the surface tension at the air–droplet interface.

We analyzed the uncertainty in the DKA-derived water activity and surface tension, related to the instrumental errors as well as to the morphology of the nanoparticles and their phase state. Our studies have shown that under optimal modes of operation of HHTDMA for moderate aqueous concentrations, the uncertainty in a_w and σ does not exceed 0.2 %–0.4 % and 3 %–4 %, respectively, but it increases by an order of magnitude in the case of highly concentrated nanodroplet solutions.

1 Introduction

Water uptake of aerosol particles is among the central issues of current research into atmospheric and climate processes (Pöschl, 2005; Andreae and Rosenfeld, 2008). Water activity and surface tension (interfacial energy) are key thermodynamic parameters of classical Köhler theory, which describes the hygroscopic growth of particles and their ability to serve as cloud condensation nuclei (CCN) (Swietlicki et al., 2008; Prisle et al., 2011; Forestieri et al., 2018; Ruehl et al., 2016; Ovadnevaite et al., 2017; Davies et al., 2019). Bulk methods, such as pendant drop tensiometry (Anastasiadis et al., 1987; Topping et al., 2007; Shapiro et al., 2009), electrodynamic balance (EDB), and optical tweezers operating with super-micrometric particles are mainly used to determine a_w and σ (Tang et al., 2019; Bzdek et al., 2020). The applicability of these methods to nanoparticles is limited because the water uptake, gas–particle and bulk–surface partitioning, and related phase transitions (deliquescence, efflorescence) depend on particle size (Biskos et al., 2006a, b; Prisle et al., 2010; Djikaev et al., 2001; Cheng et al., 2015). To bridge the gap between experimental and modeling results for bulk materials and nanoparticles, a new method called differential Köhler analysis (DKA) was developed recently (Cheng et al., 2015). This method allows for determination of the water activity and surface tension of supersaturated aqueous solutions based on hygroscopic growth measurement of nano-sized droplets. Recently, Lei et al. (2023) used the DKA method to estimate water activity of the levoglucosan and D-glucose aerosol nanoparticles using nano-HTDMA (hygroscopicity tandem differential mobility analyzer) in the size range of 6–100 nm with RH below 90 %. These results show a good agreement of the DKA-derived water activity with the earlier experimental data and E-AIM. However, no DKA-based surface tension data were reported in that study.

Atmospheric aerosol particles are mainly a mixture of organic and inorganic components. The interaction of organic–inorganic species in aqueous solutions affects both bulk (Raoult effect) and surface (Kelvin effect) properties of nanoparticles. In order to estimate these effects independently, model approximations for a_w or σ are introduced (Li and Lu, 2001; Tuckermann, 2007; Prisle et al., 2010; Petters and Petters, 2016; Ruehl et al., 2016; Ovadnevaite et al., 2017; Forestieri et al., 2018; Davies et al., 2019; Schmedding and Zuend, 2023; and references therein). However, it is difficult to verify the validity of these approximations with respect to the nanodroplets accounting for the bulk–surface partitioning and liquid–liquid separation effects. The advantage of the DKA method is precisely that it allows for simultaneous determination of the bulk and surface characteristics of nanodroplets in the wide concentration range, including a highly supersaturated solution. This provides a new opportunity for validation of the theoretical approaches used for both

a_w and σ in the pure and mixed organic–inorganic aerosols. In addition to pure substances, the DKA method is applied here to derive the water activity and surface tension of two-component particles comprising ammonium sulfate and D-glucose with mass ratios of 4 : 1 and 1 : 1. Both components are surface inactive and highly water soluble.

The DKA data were obtained from the measurements of hygroscopic growth factors with a high-humidity tandem differential mobility analyzer (HHTDMA) for aerosol particles with diameters in the range of 17–100 nm at relative humidity (RH) of 2.0 %–99.6 %. The high precision of growth factor measurements in the wide particle size and RH range allows for determination of the water activity and surface tension from dilute to highly supersaturated aqueous solutions under conditions that are not accessible to other methods.

We analyze and discuss our nanoparticle measurement results with respect to those obtained with bulk experimental methods and thermodynamic models. In addition, our HHTDMA measurements provide information about particle restructuring in response to water vapor adsorption at low and intermediate RH levels (Mikhailov et al., 2009, 2021), which enable the determination of mass equivalent diameters for the investigated nanoparticles as well as uncertainty analyses and refinements of the DKA-derived thermodynamic parameters at subsaturation and supersaturation conditions.

2 Material and methods

2.1 Aerosol generation

The investigated aerosols were produced by nebulizing an aqueous solution of pure ammonium sulfate (99.9 % pure, ChemCruz) and D-glucose (99.55 % pure, Fisher) at ~ 0.01 % weight concentration or their mixture with AS : Gl = 4 : 1 and 1 : 1 weight ratios.

2.2 HHTDMA setup and modes of operation

The hygroscopic properties of size-selected aerosol particles were measured in the 2.0 %–99.6 % RH range with a high-humidity tandem differential mobility analyzer (HHTDMA) (see Mikhailov and Vlasenko, 2020; Mikhailov et al., 2021). Throughout the whole relative humidity range, the absolute uncertainty of RH is less than 0.5 %, and the relative growth factor uncertainty due to RH and instrumental errors does not exceed 1 %. A detailed calculation of growth factor uncertainty is described in Mikhailov and Vlasenko (2020) (Sect. 2.5). Three operation modes are available using this HHTDMA instrument: hydration and dehydration (H&D) (also called restructuring mode), hydration, and dehydration. The H&D mode was used to determine the optimal RH range in which initial irregular particles transform into compact globules.

Inorganic and organic aerosol particles as well as their mixtures restructure upon humidification below their deliquescence (Mikhailov et al., 2020). Irregular envelope shape and porosity cause a discrepancy between the mobility-equivalent and mass-equivalent particle diameters that limits its precision of the mobility-diameter-based hygroscopicity tandem differential mobility analyzer (HTDMA) (Gysel et al., 2004; Mikhailov et al., 2020). To account for restructuring, we used the minimum mobility particle diameter, $D_{b,H\&D,min}$, obtained in the H&D HHTDMA mode as an approximation of mass-equivalent diameter of the dry solute particle, D_s (i.e., $D_s = D_{b,H\&D,min}$). The size-dependent restructuring factor was calculated as follows: $g_{b,H\&D} = D_{b,RH}/D_{b,i}$, where $D_{b,i}$ and $D_{b,RH}$ are the initial mobility particle diameter and that measured at RH of aerosol preconditioning, respectively (Mikhailov et al., 2021, Supplement S3). The description of the particle restructuring mechanism is beyond the scope of this work. It will be considered in detail in the next paper.

In this study, the H&D mode was in situ coupled with a conventional hydration or dehydration mode. In the combined modes, the initial monodisperse dry particles were first led through the preconditioning section where they underwent microstructural transformation, by means of humidification in the 2%–90% RH range (Nafion humidifier) and drying (Nafion dryer followed by a silica gel diffusion dryer). At the same time, the aerosol and sheath flow in the second differential mobility analyzer was below 3% RH (Mikhailov et al., 2020, 2021, Supplement S3). The preconditioning RH at which the particles reached their minimum diameter, $D_{b,H\&D,min}$, was then kept constant throughout the standard hydration or dehydration experiment. The mobility-equivalent particle growth factor, g_b , was calculated as the ratio of the mobility-equivalent diameter, D_b , measured after conditioning (hydration, dehydration) to the minimum mobility diameter, $D_{b,H\&D,min}$, observed in the H&D mode:

$$g_b = \frac{D_b}{D_{b,H\&D,min}}. \quad (1)$$

3 Thermodynamic models

3.1 Differential Köhler analysis (DKA)

The basis of the DKA method (Cheng et al., 2015) is the Köhler equation which describes the equilibrium relative humidity, s_w , over a spherical droplet as a function of a_w , σ , and the mass equivalent diameter D_b :

$$s_w = \frac{RH}{100} = a_w \exp\left(\frac{4\sigma v_w}{RT D_b}\right) = a_w \exp\left(\frac{4\sigma v_w}{RT D_s g_b}\right), \quad (2)$$

where v_w is the partial molar volume of water, R is the universal gas constant, and T is the temperature. D_b in Eq. (2) is often replaced by the product of D_s and g_b ; that is, $g_b =$

D_b/D_s . The logarithmic form of Eq. (2) is

$$\ln s_w = \ln a_w(g_b) + \frac{A}{D_s} \sigma(g_b), \quad (3)$$

where $A = 4v_w/RT g_b$. From Eq. (3) it follows that a_w and σ can be obtained as the fitting parameters for the $s_w(D_s)$ dependence with the same g_b . Obviously, the larger the number of experimental values of D_s that are used for the fitting, the more accurately a_w and σ can be estimated. This study utilized four sizes for ammonium sulfate and eight sizes for D-glucose and its mixture with ammonium sulfate in the D_s range 17–100 nm, where the Kelvin curvature effect has a significant impact on particle hygroscopicity. We used the Levenberg–Marquardt algorithm (implemented in Origin 9.0) to find a_w and σ in the iterative procedure with standard fitting error as a measure of their uncertainty. It should be noted that fitting of the experimental $s_w(D_s)$ dependences over the entire set of dry diameters significantly improves the reliability of the DKA-derived a_w and σ as compared to the original approach, where these quantities were calculated in pairs for available dry diameters (Cheng et al., 2015; Lei et al., 2023).

According to Eq. (3), DKA assumes that water activity and surface tension depend on g_b only (i.e., concentration). However, in the case of bulk–surface partitioning, both water activity and surface tension may also depend on D_s . To minimize this effect, we used surface inactive compounds. Previous studies on the example of NaCl and $(NH_4)_2SO_4$ nanoparticles have shown that the size effect is negligible for such compounds (Bahadur and Russell, 2008; Cheng et al., 2015, Supplement).

3.2 HHTDMA-derived activity coefficients

In a binary system at a constant temperature and pressure, the water activity, a_w , and the solute activity, a_s , are related by the Gibbs–Duhem equation (Prausnitz et al., 1999):

$$x_s d \ln a_s = -x_w d \ln a_w, \quad (4)$$

where x_w and x_s are the mole fractions of water and solute, respectively. According to Robinson and Stokes (1970), the water activity and molal osmotic coefficient of solute, Φ_s , for the real solution are related by

$$\ln a_w = -\frac{\nu_s m_s \Phi_s M_w}{1000}, \quad (5)$$

where ν_s and m_s are the stoichiometric dissociation number and molality of the solute (mol kg^{-1} of water), and M_w is the molecular mass of water (g mol^{-1}). For electrolyte solution in the molality scale,

$$a_s = a_{\pm}^{\nu_s} = (\gamma_{\pm} m_{\pm})^{\nu_s} = \gamma_{\pm}^{\nu_s} m_s (\nu_+^{\nu_+} \nu_-^{\nu_-}). \quad (6)$$

ν_+ and ν_- are the numbers of positive and negative ions produced upon dissociation of the solute ($\nu_s = \nu_+ + \nu_-$); and

m_{\pm} , γ_{\pm} , and a_{\pm} are the mean molality, activity coefficient, and activity of the solute. For ammonium sulfate with $\nu_s = 3$ and non-dissociating D-glucose with $\nu_s = 1$ from Eq. (6) it follows that

$$a_{AS} = 4^{1/3} m_{AS} \gamma_{\pm}^3 \quad \text{and} \quad a_{GI} = \gamma m_{GI}. \quad (7)$$

Combining Eqs. (4)–(6) gives (Prausnitz et al., 1999)

$$d \ln \gamma_{\pm} = d\Phi_s + \frac{1}{m_s} (\Phi_s - 1) dm_s. \quad (8)$$

Integration from $m_s = 0$ to the solution of interest yields

$$\ln \gamma_{\pm} = \Phi_s - 1 + \int_0^m \frac{\Phi_s - 1}{m_s} dm_s. \quad (9)$$

The integral in Eq. (9) was evaluated numerically by plotting values of $(\Phi_s - 1)/m_s$ against m_s .

3.3 Surface tension models

The DKA-derived σ of pure AS and GI aqueous solutions and their mixtures were compared with the model of Li and Lu (2001) and Dutcher et al. (2010). These models were moderately successful at representing the surface tensions of aqueous solutions of some soluble salts and their mixtures with organic compounds (Topping et al., 2007). The model by Dutcher et al. (2010) expresses the surface tension (σ) in terms of the surface tension of pure water (σ_w) and that of a hypothetical molten salt (σ_s):

$$\ln \sigma = x_w \ln \sigma_w + x_s \ln \sigma_s, \quad (10)$$

where x_w and x_s are the mole fractions of water (w) and salt (s). It is assumed that the surface tension of a solution at different temperatures is described by $\sigma = \sigma_w + x_s F_{ws}(T)$, with $F_{ws}(T) = a_{ws} + b_{ws}T$, where a_{ws} and b_{ws} are fitted parameters. At very high salt concentrations, a similar relationship is used: $\sigma = \sigma_s + x_w F_{sw}(T)$ with $F_{sw}(T) = a_{sw} + b_{sw}T$. The two above expressions can be applied to express the surface tension over the entire concentration range:

$$\ln \sigma(T) = x_w \ln(\sigma_w(T) + x_s F_{ws}(T)) + x_s \ln(\sigma_s(T) + x_w F_{sw}(T)). \quad (11)$$

Equation (11) was used to calculate σ of ammonium sulfate particles at $T = 298$ K with parameters $\sigma_w = 72.0$ mN m⁻¹, $\sigma_s = 185.0$ mN m⁻¹, $a_{ws} = a_{sw} = 0$, $b_{ws} = 0.366$ mN m⁻¹ K⁻¹, and $b_{sw} = -0.289$ mN m⁻¹ K⁻¹ as suggested by Dutcher et al. (2010). The Li and Lu (2001) surface tension model includes a Langmuir-type adsorption isotherm and a term dependent on the activity of dissolved solid. For an aqueous solution with one solute, this model yields

$$\sigma = \sigma_w + RT \Gamma_s^{w0} \ln \frac{1}{1 + K_s a_s}, \quad (12)$$

where Γ_s^{w0} is the saturated surface excess of solute (s), and K_s is the adsorption equilibrium constant of solute (s). For multi-component systems, Li and Lu (2001) proposed two approaches. The first approach, referred to as LiLu (1), implies that there is no interaction or competing adsorption between species at the interface:

$$\sigma = \sigma_w + RT \sum_{i=1}^k \Gamma_i^{w0} \ln \frac{1}{1 + K_i a_i}. \quad (13)$$

The second approach, that we refer to as LiLu (2), considers the interacting and competing adsorption between species in the interface for mixed aqueous systems at higher concentrations. This yields another expression for the surface tension of mixed solutions:

$$\sigma = \sigma_w + RT \sum_{i=1}^k \Gamma_i^{w0} \ln \left(1 - \frac{K_i a_i}{1 + \sum_i K_i a_i} \right). \quad (14)$$

We also compared the DKA-derived σ with simpler approximations, including those describing a supersaturated solution. Thus, for an ammonium sulfate solution, we used the parameterization by Pruppacher and Klett (1997):

$$\sigma(\text{N m}^{-1}) = 0.072 + \frac{2.34 \times 10^{-2} \times X_s}{1 - X_s}, \quad (15)$$

where X_s is the mass fraction of solute. This parameterization is expected to be valid up to $X_s \approx 0.8$. For the concentration dependence of surface tension of an aqueous D-glucose solution, the linear function given by Aumann et al. (2010) was chosen:

$$\sigma(\text{mN m}^{-1}) = \sigma_w + \frac{\Delta\sigma}{\Delta C} C, \quad (16)$$

where C is the molarity concentration (mol L⁻¹) and $\Delta\sigma/\Delta C = 1.29$ (mN m⁻¹/mol L⁻¹) with upper limit of C being 3.5 mol L⁻¹ ($m_{GI} = 6.0$ mol kg⁻¹).

4 Experimental results and discussion

4.1 Particle restructuring due to humidification

Figure 1 illustrates the restructuring of aerosol particles in the H&D mode. Both pure and mixed particles restructure with increasing RH, wherein this effect is strongest for the largest diameters. Interestingly, mixed AS/GI particles with a mass ratio of 4 : 1 undergo strong restructuring with $g_{b,H\&D}$ less than 1 (Fig. 1c), whereas AS/GI particles with a mass ratio of 1 : 1 show no structural change (Fig. 1d). For these particles, the restructuring factor $g_{b,H\&D}$ is larger than 1 and increases with RH, indicating that initial particles are compact and spherical (i.e., $D_{b,i} = D_{b,H\&D,\min} = D_s$) (Mikhailov et al., 2020). The size-dependent increase in $g_{b,H\&D}$ (Fig. 1c) is due to a residual water film on the particle surface after dehydration in the H&D mode. The higher values of

$dg_{b,H\&D}/dRH$ observed for particles with smaller diameters support this assumption. Since the H&D mode is in situ coupled with a conventional hydration or dehydration mode, the observed $D_{b,H\&D,min}$ values at specified RH were directly used to calculate the growth factor using Eq. (1). This has made it possible to eliminate the uncertainty in the growth factor caused by irregular particle morphology, which is as high as 5 % or more for particles with $D_{b,i} > 70$ nm.

4.2 Size-dependent growth factors

Figure 2 shows the size-dependent growth factors of pure and mixed particles observed upon dehydration in the HHTDMA mode, calculated using Eq. (1). The dehydration branch was used because it allows us to determine a_w and σ over a wide concentration range, including highly concentrated droplet solutions. Note that, in contrast to pure ammonium sulfate and its 4 : 1 mixture with D-glucose, in the case of pure D-glucose (Fig. 2b) and mixed AS/Gl particles with a mass ratio of 1 : 1 (Fig. 2d), the growth factors obtained in the hydration and dehydration modes are identical over the whole range of relative humidity (Fig. S1 in the Supplement). As expected, due to the Kelvin effect, for the same g_b the equilibrium values of s_w (or RH) for smaller particles shift towards higher values (Eq. 2). This is clearly visible for all particles measured at RH above 70 % (inserts in Fig. 2). For ammonium sulfate particles, this pattern is valid also for highly concentrated solutions up to efflorescence RH (ERH \sim 30 %) (Fig. 3a). However, in the cases of D-glucose (Fig. 3b) and AS/Gl (1 : 1) mixed particles (Fig. 3d), this pattern is not obvious. Moreover, in the case of AS/Gl (4 : 1) particles, the values of s_w for small particles are lower than for large particles (Fig. 3c), in contrast to the behavior of pure ammonium sulfate (Fig. 3a). The measurement data in Fig. 3b and d also demonstrate that for D-glucose and AS/Gl (1 : 1) particles in the vicinity of RH \approx 48 % the $g_b(RH)$ dependence has an inflection point (derivative dg_b/dRH has extremum; see the insert in Fig. 3b and d), indicating a distinct water sorption mechanism and hence a different phase state of the particles before and after this RH point. Most likely, at low RH, D-glucose and AS/Gl (1 : 1) mixed particles are in the semi-solid amorphous state; the increase in humidity over glassy amorphous particles leads to a moisture-induced phase transition that occurs at a “glass transition relative humidity” of $RH_g \approx 48$ % (Mikhailov et al., 2009).

Continued water uptake converts the amorphous particles into concentrated solution droplets. Due to the very low molecular diffusivity of glasses, the uptake of water vapor by glassy aerosol particles is limited to surface adsorption, whereas above RH_g the particles are in a viscous liquid state and absorb water in the particle bulk. Differences in water sorption mechanisms may explain why the growth factors of small particles at low RH are higher than for larger particles. At RH below RH_g with the same water adsorption layer, the water film contribution to g_b will be higher for small parti-

cles. For example, at a water layer thickness of 0.56 nm (i.e., two monolayers of water molecules), g_b for aerosol particles of 19 and 100 nm are 1.031 and 1.006, respectively. This pattern is clearly seen for AC/Gl (1 : 1) particles with compact initial structures (i.e., $D_{b,i} = D_{H\&D,min}$, and $g_b = g_{H\&D}$) for which the growth factor of large particles at the same RH is lower than that for small particles (Fig. 1d). Note also that moisture-induced transformation from the semi-solid state to the solute state can occur gradually within a certain RH range (Mikhailov et al., 2009). All these features in the water uptake by amorphous particles at high concentrations (low growth factor values) lead to additional uncertainties when using the DKA method.

4.3 DKA-derived a_w and a_s for single-component aqueous solution

The DKA-derived water activity for ammonium sulfate and D-glucose aerosol particles are shown in Fig. 4a and b, respectively. More information on the DKA calculation can be found in Sect. S1 in the Supplement. The estimated uncertainty of a_w for pure species is shown in Fig. S2 in the Supplement. The a_w values for ammonium sulfate (Fig. 4a) agree well with electrodynamic balance (EDB) measurements by Tang and Munkelwitz (1994), Clegg et al. (1995), and Chan et al. (1992) and are almost identical to the DKA-derived a_w values previously reported by Cheng et al. (2015) and Extended Aerosol Inorganic Model (E-AIM III) of Wexler and Clegg (2002), including highly concentrated aqueous solutions. In general, the DKA-derived water activities obtained for pure ammonium sulfate particles reproduce the literature data well.

Figure 4b shows the DKA-derived a_w values of D-glucose aerosol particles. Our data are consistent with the EDB measurements by Peng et al. (2001) up to $X_{Gl} \sim 0.8$. The observed divergence of the water activity at $X_{Gl} > 0.8$ may be due to kinetic limitations in highly viscous semi-solid amorphous particles (Mikhailov et al., 2009), which prevail in EDB experiments with micrometer-sized particles. As noted in Sect. 4.2 and shown in Fig. 3d at $RH = RH_g \sim 48$ %, particles undergo moisture-induced phase transition where semi-solid particles become concentrated solution droplets. This RH value is very close to $a_w \sim 49$ %, below which a strong deviation between DKA- and EDB-derived a_w is observed (Fig. 4b). If this is the case, the DKA-derived a_w here is closer to the real a_w .

At the same time, the DKA-derived a_w values differ from those predicted by the AIOMFAC model (Zuend et al., 2008), with a maximum deviation of 6 % at $X_{Gl} \sim 0.7$. The discrepancy between measured and AIOMFAC-predicted a_w values for carbohydrates has been noted previously and is attributed to relatively strong intramolecular interactions due to several polar groups in close proximity, not accounted for by the UNIFAC model within the AIOMFAC framework (Zuend et al., 2011).




Figure 1. RH-dependent restructuring of ammonium sulfate (a), D-glucose (b), and their mixture with mass ratio AS : GI = 4 : 1 (c) and AS : GI = 1 : 1 (d) obtained in the H&D mode as a function of particle size.

The multicomponent surface tension model of Li and Lu (2001) is based on the activity of the solute. To take advantage of this model, we calculated the activity coefficients of single solutes by numerical integration of Eq. (9) and then, using Eq. (7), their activities (see Sect. S2 in the Supplement for more details). The calculation results are shown in Fig. 5 in comparison with literature data. One can see that the ammonium sulfate activity is in good agreement with E-AIM (Fig. 5a) up to $X_{AS} = 0.75$ ($m_{AS} = 22.7 \text{ mol kg}^{-1}$). At higher X_{AS} values, there is a discrepancy between our data and the model data. This discrepancy is most likely due to the uncertainty in DKA-derived a_w in concentrated ammonium sulfate aqueous solutions (Fig. S2) and uncertainty in assessment of the integral Eq. (9) in the asymptotic region ($m_s \rightarrow 0$) (Lakhanpal and Conway, 1960).

Figure 5b shows the DKA-based activity of D-glucose. One can see that these values are in agreement with bulk measurements by Miyajima et al. (1983), obtained by the isopiestic method (blue symbols) available up to $X_{GI} = 0.52$ ($m_{GI} = 6.0 \text{ mol kg}^{-1}$). At the same time, a systematic deviation is observed between our results and the AIOMFAC model. As previously noted, the UNIFAC as a part of the AIOMFAC model does not match well with D-glucose and

other carbohydrates, owing to strong intramolecular interactions of the polar groups in close proximity.

In general, the DKA-derived water activity and activity of ammonium sulfate and D-glucose in aqueous solutions reproduce the literature data well. For future applications, we fitted DKA-derived a_w and a_s for single solutes with a polynomial function. The obtained fitting parameters are listed in Tables S1 and S2 in the Supplement.

4.4 Surface tension of the single solute solution droplet

Figure 6 shows the DKA-derived surface tension of ammonium sulfate and D-glucose aqueous solution droplets as a function of solute activity. For future use, we fit our data with the Li and Lu (2001) model (Eq. 12). The resulting values of Γ_s^{w0} and K_s together with the a_s fitting interval are listed in Table S3 in the Supplement. The negative Γ_s^{w0} values obtained for both ammonium sulfate and glucose solution droplets indicate that a negative adsorption occurs on the interface between the vapor and liquid phases, leading to surface tensions increasing with concentration, which is typical for surface-inactive compounds (Tuckermann, 2007; Aumann et al., 2010).

For comparison with the literature data, in Fig. 7 we plotted the DKA surface tension in more accessible concentra-




Figure 2. Size-dependent growth factors, g_b , of ammonium sulfate (a), D-glucose (b), and their mixture with mass ratios AS : GI = 4 : 1 (c) and AS : GI = 1 : 1 (d) obtained in the dehydration HHTDMA mode.

tion scales, i.e., in mass fraction and molality. Figure 7a shows that the DKA-derived σ for ammonium sulfate solution is in agreement, within uncertainty, with the empirical approximation of Pruppacher and Klett (1997) and also consistent with the Dutcher et al. (2010) model up to $X_{AS} \approx 0.5$ ($m_{AS} = 7.6 \text{ mol kg}^{-1}$). Above this value, the difference between DKA data and those by Dutcher et al. (2010) gradually increases with increasing concentration. This difference is due to the fact that model parameters used in Dutcher et al. (2010) (Sect. 3.3) are only valid for a concentration of ammonium sulfate not exceeding $X_{AS} = 0.43$ ($m_{AS} = 5.69 \text{ mol kg}^{-1}$), although the model structure allows for calculation of σ in the concentration range from dilute solution to molten salt (Eq. 11). Figure 7b shows the DKA-derived σ for D-glucose solution droplets as a function of mass fraction or molality together with the literature data. It can be seen that the DKA data are consistent with bulk measurements reported by Aumann et al. (2010). In the X_{GI} range of 0.24 to 0.52, the average deviation between DKA-derived and literature σ is 2%.

Note that the uncertainty in the DKA-retrieval parameters is strongly influenced by the accuracy of s_w (or RH) and g_b determination. At low g_b (highly concentrated solutions) when $\Delta g_b / \Delta s_w$ is small, the uncertainty in the DKA-derived

a_w and σ increases with decreasing g_b . In addition, phase state ambiguity and associated kinetic limitations can provide additional uncertainties. As an example, Fig. S3 in the Supplement shows experimental $s_w(D_s)$ dependences at low and high growth factors for D-glucose aerosol particles and corresponding uncertainties in the DKA-derived a_w and σ .

4.5 DKA-derived a_w and σ for mixed ammonium sulfate and D-glucose aerosol particles

In the absence of solute–solute interactions (“separate solutes”), the corresponding relationship for the water activity of the mixture is $a_w = \prod_i a_{w,i}$, where $a_{w,i}$ is the water activity of a pure aqueous solution of i at the same concentration as in the mixture (Mikhailov et al., 2004; Clegg and Seinfeld, 2006). Within this approach, referred to as “separate solute water activity” (SSWA), the water activity of mixed particles containing ammonium sulfate and D-glucose can be expressed as

$$a_w = a_{w,AS} a_{w,GI}. \quad (17)$$

Figure 8 shows the DKA-derived water activity of the mixed AS/GI particles with a mass ratio of 4 : 1 (a) and 1 : 1 (b), as well as results of the AIOMFAC and SSWA models.




Figure 3. Growth factor, g_b , of ammonium sulfate (a), D-glucose (b), and their mixture with mass ratios AS:GI = 4:1 (c) and AS:GI = 1:1 (d) at low RH values for minimum and maximum sizes. The inserts (b and d) show derivative dg_b/dRH with inflection point RH_g . The instrumental error in RH_g does not exceed the symbol size.




Figure 4. DKA-derived water activity, a_w , of ammonium sulfate (a) and D-glucose (b) aqueous solution at 298 K in comparison with literature data as a function of mass fraction and molality. The red line in panels (a) and (b) is a polynomial fit to all DKA data of a_w . The best fit coefficients are listed in Table S1.




Figure 5. Solute activity of ammonium sulfate (a) and D-glucose (b) in water at 298 K together with literature data as a function of mass fraction and molality. The red line in panels (a) and (b) are polynomial function with fitting parameters listed in Table S2.




Figure 6. Surface tension of ammonium sulfate (a) and D-glucose (b) aqueous solution droplets as a function of solute activity. The line in panels (a) and (b) is the model of Li and Lu (2001) (Eq. 12), with the best-fit parameters listed in Table S3.

One can see that the DKA-derived a_w and both models are consistent up to $X_s \sim 0.3$. A further increase in concentration is accompanied by a significant deviation of the model a_w values, both among themselves and from the DKA-derived a_w . It appears that the SSWA approximation underestimates the specific interactions between ions and molecules in the droplet volume (higher a_w), whereas the AIOMFAC model overestimates them (lower a_w), especially in concentrated solutions.

Figure 9 shows the DKA-retrieved surface tension of the mixed AS/Gl particles with mass ratios of 4 : 1 (a) and 1 : 1 (b), compared to the Li and Lu (2001) model. Both approaches, i.e., LiLu (1) and LiLu (2), do not match well with the HHTDMA–DKA-based σ . This discrepancy is especially pronounced for the mixed particles with an AS/Gl mass ratio of 1 : 1 (Fig. 9b). A slight lowering of the surface tension compared to that of water is already noticeable for a 4 : 1 mixture (Fig. 9a). The AS : Gl = 1 : 1 mixture (Fig. 9b) shows more significant surface tension depression, which is

not a monotonic function of concentration. At $X_s \approx 0.5$, the σ reaches a minimum of $56.5 \pm 3.0 \text{ mN m}^{-1}$ (22 % reduction compared to pure water) and increases again with concentration. Further increase in surface tension with concentration is the result of solidification of aerosol particles. According to Eq. (10) at $X_s \rightarrow 1$, σ tends to the surface tension of molten state, σ_s , which for AS and Gl is 185 mN m^{-1} (Dutcher et al., 2010) and 150.9 mN m^{-1} (Docoslis et al., 2000), respectively. Thus, at $X_s = 1$ the mole-fraction-weighted value of σ for AS : Gl = 4 : 1 and 1 : 1 is 179 and 170 mN m^{-1} , respectively, which reasonably agrees with the DKA-derived σ at high X_s values (Fig. 9a and b).

Note that the decrease in surface tension observed in Fig. 9b and its subsequent increase with concentration is reproduced in repeated measurements (see Fig. S4 in the Supplement). Most likely the reduction in surface tension at moderate concentrations is caused by the salting out effect (Setschenow, 1889; Kiss et al., 2005; Marcolli and Krieger, 2006; Frosch et al., 2011; Wang et al., 2014; Lin et al.,




Figure 7. DKA-derived surface tension of ammonium sulfate (a) and D-glucose (b) solution droplets as a function of mass fraction and molality in comparison with literature. The shaded area in panels (a) and (b) indicates uncertainty in the DKA retrieval σ .




Figure 8. Water activity of the aqueous solution of the mixed ammonium sulfate and D-glucose with mass ratio AS:Gl = 4 : 1 (a) and AS:Gl = 1 : 1 (b) compared to model data as a function of the mass fraction of the solute, X_s . The red line in panels (a) and (b) are polynomial function with fitting parameters listed in Table S1. SSWA denotes separate solute water activity (Eq. 17).

2020; Bzdek et al., 2020) and results from the interaction between ammonium sulfate ions and D-glucose molecules, facilitating the association of initially surface-inactive hydrophilic organic molecules into surface-active hydrophobic associates (quasi-macromolecules). Docoslis et al. (2000) analyzed the difference between polysaccharides (dextran, Ficoll) and their surface-inactive constituents (sucrose, glucose). They concluded that the contrasting results are caused by the differences in polar intermolecular reactivities of the monomeric and polymeric glucides. The singly dissolved monomeric sugars contribute strongly to the polar (Lewis acid–base) free energy of cohesion through the multiple interactions between their freely available electron donors and acceptors, through mutual interactions as well as through interactions with the surrounding water dipoles. However once covalently polymerized, these polysaccharides have lost the strong bipolarity of the monomeric sugar molecules. It can

thus be assumed that, in an aqueous AS/Gl system, NH_4^+ and SO_4^{2-} ions effectively neutralize the bipolarity of the monomeric D-glucose molecules, facilitating their association into less-polar aggregates with reduced σ values, which are more readily accommodated at the air–droplet interface. Obviously, this mechanism is more pronounced for mixed AS:Gl = 1 : 1 particles (Fig. 9b) than for their 4 : 1 mixture (Fig. 9a), due to the higher D-glucose content (by a factor of 2.5).

5 Summary and conclusions

In this study, the DKA method was applied to derive water activities and surface tension of pure ammonium sulfate and D-glucose as well as their mixtures with mass ratios of 4 : 1 and 1 : 1 based on shape-corrected hygroscopic growth factors for particles with diameter of 17–100 nm in the rel-




Figure 9. DKA-derived surface tension of the aqueous solution of the mixed ammonium sulfate and D-glucose with mass ratio AS : GI = 4 : 1 (a) and AS : GI = 1 : 1 (b) together with Li and Lu (2001) model (LiLu (1) – Eq. (13) and LiLu (2) – Eq. (14)) as a function of mass fraction of the solute, X_s . The shaded area in panels (a) and (b) indicates the uncertainty in the DKA retrieval σ , the black line is the surface tension of pure water $\sigma_w = 72.0 \text{ mN m}^{-1}$.

ative humidity range of 2.0 %–99.6 %. The obtained a_w and σ values for pure ammonium sulfate and D-glucose droplet solution are in good agreement with the bulk measurements in the available concentration range for bulk methods. The DKA method was employed for the first time to determine a_w and σ of AS/GI mixed particles. Our data show that for dilute and moderate solution concentrations ($X_s < 0.3$), the SSWA and AIOMFAC models are in agreement with the DKA-derived a_w . However, the discrepancy between the model and our data increases rapidly with increasing solution concentration, with SSWA underestimating DKA data, while AIOMFAC overestimates it. Both ammonium sulfate and D-glucose are surface-inactive compounds with a positive $\Delta\sigma/\Delta C$ slope, indicating negative solute adsorption. Interestingly, mixing them together leads to positive adsorption and reduction in surface tension at the air–surface interface (i.e., the AS/GI mixture becomes a surface-active compound due to salting out). We suggest that AS ions neutralize polar groups of D-glucose, helping them to combine into less-polar aggregates with reduced σ values at the air–surface interface.

The error analysis showed that there are some factors affecting the accuracy of a_w and σ determined by the DKA method. One of them is the irregular morphology of the initial nanoparticles, which leads to a size-dependent error in g_b . Other factors are mainly characteristic of highly concentrated nanodroplet solutions, such as the large uncertainty of g_b and ambiguity of the phase state. Both small growth factor changes at low RH and the different mechanisms of water uptake (surface adsorption vs. bulk absorption) lead to an increased uncertainty in the DKA-derived values of a_w and σ . Thus, for D-glucose aerosol particles at RH near $\text{RH}_g = 48\%$ and $g_b = 1.08$ ($X_{\text{GI}} = 0.85$; $m_{\text{GI}} = 32.4 \text{ mol kg}^{-1}$), the relative uncertainty in a_w is 3.5 % and in σ it is 41.2 %, respectively.

As mentioned above, DKA assumes that thermodynamic parameters depend only on concentration and are independent of particle size. Good agreement of DKA-derived a_w and σ with the literature data showed that, for particles in the size range of 20–100 nm, this approach is acceptable for single-component solutions of AS and GI as well as their mixtures, at least at moderate concentrations. Additional HHTDMA–DKA studies are needed to evaluate the accuracy of this approximation for nanoparticles containing surface-active molecules. We plan to conduct such studies to compare the DKA-derived a_w and σ with available experimental data and models that account for size-dependent bulk–surface partitioning.

Data availability. The raw data used in this study are archived and are available on request by contacting the corresponding author, Alexei A. Kiselev (alexei.kiselev@kit.edu).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-24-2971-2024-supplement>.

Author contributions. EFM designed the study, performed the concomitant measurements, carried out the data analysis, and wrote the manuscript with input from all coauthors. SSV and AAK contributed to the discussion of the results.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, pub-

lished maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. We would like to thank Ulrich Pöschl, Yafang Cheng, and Hang Su for their helpful recommendations that improved this paper.

Financial support. The work of EFM and SSV was supported by the Russian Science Foundation, project 22-27-00258. AAK acknowledges financial support by the Helmholtz Association under the Atmosphere and Climate program (ATMO).

The article processing charges for this open-access publication were covered by the Karlsruhe Institute of Technology (KIT).

Review statement. This paper was edited by Daniel Knopf and reviewed by two anonymous referees.

References

- Anastasiadis, S. H., Chen, J. K., Koberstein, J. T., Siegel, A. F., Sohn, J. E., and Emerson, J. A.: The determination of interfacial tension by video image processing of pendant fluid drops, *J. Colloid Interf. Sci.*, 119, 55–66, [https://doi.org/10.1016/0021-9797\(87\)90244-X](https://doi.org/10.1016/0021-9797(87)90244-X), 1987.
- Andreae, M. O. and Rosenfeld, D.: Aerosol-cloud precipitation interactions. Part 1. The nature and sources of cloud-active aerosols, *Earth-Sci. Rev.*, 89, 13–41, <https://doi.org/10.1016/j.earscirev.2008.03.001>, 2008.
- Aumann, E., Hildemann, L. M., and Tabazadeh, A.: Measuring and modeling the composition and temperature-dependence of surface tension for organic solutions, *Atmos. Environ.*, 44, 329–337, <https://doi.org/10.1016/j.atmosenv.2009.10.033>, 2010.
- Bahadur, R. and Russell, L. M.: Effect of surface tension from MD simulations on size-dependent deliquescence of NaCl nanoparticles, *Aerosol Sci. Tech.*, 42, 369–376, <https://doi.org/10.1080/02786820802104965>, 2008.
- Biskos, G., Malinowski, A., Russell, L. M., Buseck, P. R., and Martin, S. T.: Nanosize effect on the deliquescence and the efflorescence of sodium chloride particles, *Aerosol Sci. Tech.*, 40, 97–106, <https://doi.org/10.1080/02786820500484396>, 2006a.
- Biskos, G., Russell, L. M., Buseck, P. R., and Martin, S. T.: Nanosize effect on the hygroscopic growth factor of aerosol particles, *Geophys. Res. Lett.*, 33, L07801, <https://doi.org/10.1029/2005GL025199>, 2006b.
- Bzdek, B. R., Reid, J. P., Malila, J., and Prisle, N. L.: The surface tension of surfactant-containing, finite volume droplets, *P. Natl. Acad. Sci. USA*, 117, 8335–8343, <https://doi.org/10.1073/pnas.1915660117>, 2020.
- Chan, C. K., Flagan, R. C., and Seinfeld, J. H.: Water activity of $\text{NH}_4\text{NO}_3/(\text{NH}_4)_2\text{SO}_4$ solutions, *Atmos. Environ. A-Gen.*, 26, 1661–1673, [https://doi.org/10.1016/0960-1686\(92\)90065-S](https://doi.org/10.1016/0960-1686(92)90065-S), 1992.
- Cheng, Y., Su, H., Kopp, T., Mikhailov, E. F., and Pöschl, U.: Size dependence of phase transitions in aerosol nanoparticles, *Nat. Commun.*, 6, 5923, <https://doi.org/10.1038/ncomms6923>, 2015.
- Clegg, S. L. and Seinfeld, J. H.: Thermodynamic models of aqueous solutions containing inorganic electrolytes and dicarboxylic acids at 298.15 K. 1. The acids as nondissociating components, *J. Phys. Chem. A*, 110, 5692–5717, 2006.
- Clegg, S. L., Ho, S. S., Chan, C. K., and Brimblecombe, P.: Thermodynamic properties of aqueous $(\text{NH}_4)_2\text{SO}_4$ to high supersaturation as a function of temperature, *J. Chem. Eng. Data*, 40, 1079–1090, <https://doi.org/10.1021/jc00021a011>, 1995.
- Davies, J. F., Zuend, A., and Wilson, K. R.: Technical note: The role of evolving surface tension in the formation of cloud droplets, *Atmos. Chem. Phys.*, 19, 2933–2946, <https://doi.org/10.5194/acp-19-2933-2019>, 2019.
- Djikaev, Y. S., Bowles, R., Reiss, H., Hämeri, K., Laaksonen, A., and Väkevä, M.: Theory of size dependent deliquescence of nanoparticles: Relation to heterogeneous nucleation and comparison with experiments, *J. Phys. Chem. B*, 105, 7708–7722, <https://doi.org/10.1021/jp010537e>, 2001.
- Docoslis, A., Giese, R. F., and van Oss, C. J.: Influence of the water–air interface on the apparent surface tension of aqueous solutions of hydrophilic solutes, *Colloid. Surface. B*, 19, 147–162, [https://doi.org/10.1016/S0927-7765\(00\)00137-5](https://doi.org/10.1016/S0927-7765(00)00137-5), 2000.
- Dutcher, C. S., Wexler, A. S., and Clegg, S. L.: Surface tensions of inorganic multicomponent aqueous electrolyte solutions and melts, *J. Phys. Chem. A*, 114, 12216–12230, <https://doi.org/10.1021/jp105191z>, 2010.
- Forestieri, S. D., Staudt, S. M., Kuborn, T. M., Faber, K., Ruehl, C. R., Bertram, T. H., and Cappa, C. D.: Establishing the impact of model surfactants on cloud condensation nuclei activity of sea spray aerosol mimics, *Atmos. Chem. Phys.*, 18, 10985–11005, <https://doi.org/10.5194/acp-18-10985-2018>, 2018.
- Frosch, M., Prisle, N. L., Bilde, M., Varga, Z., and Kiss, G.: Joint effect of organic acids and inorganic salts on cloud droplet activation, *Atmos. Chem. Phys.*, 11, 3895–3911, <https://doi.org/10.5194/acp-11-3895-2011>, 2011.
- Gysel, M., Weingartner, E., Nyeki, S., Paulsen, D., Baltensperger, U., Galambos, I., and Kiss, G.: Hygroscopic properties of water-soluble matter and humic-like organics in atmospheric fine aerosol, *Atmos. Chem. Phys.*, 4, 35–50, <https://doi.org/10.5194/acp-4-35-2004>, 2004.
- Kiss, G., Tombácz, E., and Hansson, H.-C.: Surface tension effects of humic-like substances in aqueous extract of tropospheric fine aerosol, *J. Atmos. Chem.*, 50, 279–294, <https://doi.org/10.1007/s10874-005-5079-5>, 2005.
- Lakhanpal, M. L. and Conway, B. E.: A method of integration of the Gibbs–Duhem equation when activities of a solute are required from those of the solvent, *Can. J. Chem.*, 38, 199–203, <https://doi.org/10.1139/v60-027>, 1960.
- Lei, T., Su, H., Ma, N., Pöschl, U., Wiedensohler, A., and Cheng, Y.: Size-dependent hygroscopicity of levoglucosan and D-glucose aerosol nanoparticles, *Atmos. Chem. Phys.*, 23, 4763–4774, <https://doi.org/10.5194/acp-23-4763-2023>, 2023.
- Li, Z. B. and Lu, B. C. Y.: Surface tension of aqueous electrolyte solutions at high concentrations – representation and prediction,

- Chem. Eng. Sci., 56, 2879–2888, [https://doi.org/10.1016/S0009-2509\(00\)00525-X](https://doi.org/10.1016/S0009-2509(00)00525-X), 2001.
- Lin, J. J., Kristensen, T. B., Calderón, S. M., Malila, J., and Prisle, N. L.: Effects of surface tension time-evolution for CCN activation of a complex organic surfactant, *Environ. Sci.-Proc. Imp.*, 22, 271–284, <https://doi.org/10.1039/C9EM00426B>, 2020.
- Marcilli, C. and Krieger, U. K.: Phase changes during hygroscopic cycles of mixed organic/inorganic model systems of tropospheric aerosols, *J. Chem. Phys.*, 110, 1881–1893, <https://doi.org/10.1021/jp0556759>, 2006.
- Mikhailov, E., Vlasenko, S., Niessner, R., and Pöschl, U.: Interaction of aerosol particles composed of protein and salt with water vapor: hygroscopic growth and microstructural rearrangement, *Atmos. Chem. Phys.*, 4, 323–350, <https://doi.org/10.5194/acp-4-323-2004>, 2004.
- Mikhailov, E., Vlasenko, S., Martin, S. T., Koop, T., and Pöschl, U.: Amorphous and crystalline aerosol particles interacting with water vapor: conceptual framework and experimental evidence for restructuring, phase transitions and kinetic limitations, *Atmos. Chem. Phys.*, 9, 9491–9522, <https://doi.org/10.5194/acp-9-9491-2009>, 2009.
- Mikhailov, E. F. and Vlasenko, S. S.: High-humidity tandem differential mobility analyzer for accurate determination of aerosol hygroscopic growth, microstructure, and activity coefficients over a wide range of relative humidity, *Atmos. Meas. Tech.*, 13, 2035–2056, <https://doi.org/10.5194/amt-13-2035-2020>, 2020.
- Mikhailov, E. F., Pöhlker, M. L., Reinmuth-Selzle, K., Vlasenko, S. S., Krüger, O. O., Fröhlich-Nowoisky, J., Pöhlker, C., Ivanova, O. A., Kiselev, A. A., Krempner, L. A., and Pöschl, U.: Water uptake of subpollen aerosol particles: hygroscopic growth, cloud condensation nuclei activation, and liquid–liquid phase separation, *Atmos. Chem. Phys.*, 21, 6999–7022, <https://doi.org/10.5194/acp-21-6999-2021>, 2021.
- Miyajima, K., Sawada, M., and Nakagaki, M.: Studies on aqueous solutions of saccharides. I. Activity coefficients of monosaccharides in aqueous solutions at 25 °C, *B. Chem. Soc. Jpn.*, 56, 1620–1623, <https://doi.org/10.1246/bcsj.56.1620>, 1983.
- Ovadnevaite, J., Zuend, A., Laaksonen, A., Sanchez, K. J., Roberts, G., Ceburnis, D., Decesari, S., Rinaldi, M., Hodas, N., Facchini, M. C., Seinfeld, J. H., and O’Dowd, C.: Surface tension prevails over solute effect in organic-influenced cloud droplet activation, *Nature*, 546, 637–641, <https://doi.org/10.1038/nature22806>, 2017.
- Peng, C., Chow, A. H. L., and Chan, C. K.: Hygroscopic study of glucose, citric acid, and sorbitol using an electrodynamic balance: comparison with UNIFAC predictions, *Aerosol Sci. Tech.*, 35, 753–758, <https://doi.org/10.1080/02786820152546798>, 2001.
- Petters, S. S. and Petters, M. D.: Surfactant effect on cloud condensation nuclei for two-component internally mixed aerosols, *J. Geophys. Res.-Atmos.*, 121, 1878–1895, <https://doi.org/10.1002/2015JD024090>, 2016.
- Pöschl, U.: Atmospheric aerosols: composition, transformation, climate and health effects, *Angew. Chem. Int. Edit.*, 44, 7520–7540, <https://doi.org/10.1002/anie.200501122>, 2005.
- Prausnitz, J. M., Lichtenthaler, R. N., and de Azevedo, E. G.: *Molecular Thermodynamics of Fluid-Phase Equilibria*, Prentice Hall, Upper Saddle River, New Jersey, USA, 226, <https://doi.org/10.1002/cjce.5450780222>, 1999.
- Prisle, N. L., Raatikainen, T., Laaksonen, A., and Bilde, M.: Surfactants in cloud droplet activation: mixed organic-inorganic particles, *Atmos. Chem. Phys.*, 10, 5663–5683, <https://doi.org/10.5194/acp-10-5663-2010>, 2010.
- Prisle, N. L., Dal Maso, M., and Kokkola, H.: A simple representation of surface active organic aerosol in cloud droplet formation, *Atmos. Chem. Phys.*, 11, 4073–4083, <https://doi.org/10.5194/acp-11-4073-2011>, 2011.
- Pruppacher, H. R. and Klett, J. D.: *Microphysics of clouds and precipitation*, Kluwer Academic Publishers, <https://doi.org/10.1007/978-0-306-48100-0>, 1997.
- Robinson, R. A. and Stokes, R. H.: *Electrolyte solutions*, 2nd edn., Butterworths, London, ISBN 100408184906, 1970.
- Ruehl, C. R., Davies, J. F., and Wilson, K. R.: An interfacial mechanism for cloud droplet formation on organic aerosols, *Science*, 351, 1447–1450, <https://doi.org/10.1126/science.aad4889>, 2016.
- Schmedding, R. and Zuend, A.: A thermodynamic framework for bulk–surface partitioning in finite-volume mixed organic–inorganic aerosol particles and cloud droplets, *Atmos. Chem. Phys.*, 23, 7741–7765, <https://doi.org/10.5194/acp-23-7741-2023>, 2023.
- Setschenow, J.: Über die Konstitution der Salzlösungen auf Grund ihres Verhaltens zu Kohlensäure, *Z. Phys. Chem.*, 4, 117–125, <https://doi.org/10.1515/zpch-1889-0409>, 1889.
- Shapiro, E. L., Szprengiel, J., Sareen, N., Jen, C. N., Giordano, M. R., and McNeill, V. F.: Light-absorbing secondary organic material formed by glyoxal in aqueous aerosol mimics, *Atmos. Chem. Phys.*, 9, 2289–2300, <https://doi.org/10.5194/acp-9-2289-2009>, 2009.
- Swietlicki, E., Hansson, H.-C., Hämeri, K., Svenningsson, B., Massling, A., McFiggans, G., McMurry, P. H., Petäjä, T., Tunved, P., Gysel, M., Topping, D., Weingartner, E., Baltensperger, U., Rissler, J., Wiedensohler, A., and Kulmala, M.: Hygroscopic properties of submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various environments – a review, *Tellus B*, 60, 432–469, <https://doi.org/10.1111/j.1600-0889.2008.00350.x>, 2008.
- Tang, I. N. and Munkelwitz, H. R.: Water activities, densities, and refractive indices of aqueous sulfates and sodium nitrate droplets of atmospheric importance, *J. Geophys. Res.*, 99, 18801–18808, <https://doi.org/10.1029/94JD01345>, 1994.
- Tang, M., Chan, C. K., Li, Y. J., Su, H., Ma, Q., Wu, Z., Zhang, G., Wang, Z., Ge, M., Hu, M., He, H., and Wang, X.: A review of experimental techniques for aerosol hygroscopicity studies, *Atmos. Chem. Phys.*, 19, 12631–12686, <https://doi.org/10.5194/acp-19-12631-2019>, 2019.
- Topping, D. O., McFiggans, G. B., Kiss, G., Varga, Z., Facchini, M. C., Decesari, S., and Mircea, M.: Surface tensions of multi-component mixed inorganic/organic aqueous systems of atmospheric significance: measurements, model predictions and importance for cloud activation predictions, *Atmos. Chem. Phys.*, 7, 2371–2398, <https://doi.org/10.5194/acp-7-2371-2007>, 2007.
- Tuckermann, R.: Surface tension of aqueous solutions of water-soluble organic and inorganic compounds, *Atmos. Environ.*, 41, 6265–6275, <https://doi.org/10.1016/j.atmosenv.2007.03.051>, 2007.
- Wang, C., Lei, Y. D., Endo, S., and Wania, F.: Measuring and modeling the salting-out effect in ammonium sul-

- fate solutions, *Environ. Sci. Technol.*, 48, 22, 13238–13245, <https://doi.org/10.1021/es5035602>, 2014.
- Wexler, A. S. and Clegg, S. L.: Atmospheric Aerosol Models for Systems Including the Ions H^+ , NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- , Br^- , and H_2O , *J. Geophys. Res.*, 107, 4207, <https://doi.org/10.1029/2001JD000451>, 2002.
- Zuend, A., Marcolli, C., Luo, B. P., and Peter, T.: A thermodynamic model of mixed organic-inorganic aerosols to predict activity coefficients, *Atmos. Chem. Phys.*, 8, 4559–4593, <https://doi.org/10.5194/acp-8-4559-2008>, 2008.
- Zuend, A., Marcolli, C., Booth, A. M., Lienhard, D. M., Soonsin, V., Krieger, U. K., Topping, D. O., McFiggans, G., Peter, T., and Seinfeld, J. H.: New and extended parameterization of the thermodynamic model AIOMFAC: calculation of activity coefficients for organic-inorganic mixtures containing carboxyl, hydroxyl, carbonyl, ether, ester, alkenyl, alkyl, and aromatic functional groups, *Atmos. Chem. Phys.*, 11, 9155–9206, <https://doi.org/10.5194/acp-11-9155-2011>, 2011.