



# Role of sea spray aerosol at the air–sea interface in transporting aromatic acids to the atmosphere

Yaru Song, Jianlong Li, Narcisse Tsona Tchinda, Kun Li, and Lin Du

Qingdao Key Laboratory for Prevention and Control of Atmospheric Pollution in Coastal Cities, Environment Research Institute, Shandong University, Qingdao, 266237, China

**Correspondence:** Lin Du (lindu@sdu.edu.cn)

Received: 5 September 2023 – Discussion started: 4 October 2023

Revised: 6 April 2024 – Accepted: 11 April 2024 – Published: 23 May 2024

**Abstract.** Aromatic acids are ubiquitous in seawater (SW) and can be transported to the atmosphere via sea spray aerosol (SSA). Despite their importance in affecting the global radiative balance, the contribution of marine aromatic acids and their transport mechanisms through SSA remain unclear. Herein, the distribution of particle size and number concentration of SSA produced in SW containing nine different aromatic acids (i.e., benzoic acids, benzenedicarboxylic acids, hydroxybenzoic acids, vanillic acid, and syringic acid) was studied using a custom-made SSA simulation chamber; moreover, the enrichment of aromatic acids in SSA and their emission flux to the atmosphere were analyzed. Transmission electron microscopy (TEM) images clearly revealed that aromatic acids can be transferred to the nascent SSA. Interestingly, the morphology associated with benzenedicarboxylic-acid-coated particles showed that aromatic acids can promote the growth of other surfaces of sea salt, thus making the sea salt core spherical. Aromatic acids showed a significant enrichment behavior at the air–sea interface, which clearly indicated that SSA represents a source of aromatic acids in the atmosphere. Vanillic acid had the largest global emission flux through SSA ( $962 \text{ t yr}^{-1}$ ), even though its concentration in SW was lower. The calculated results indicated that the global annual flux of aromatic acids was affected not only by the concentration in SW, but also by their enrichment factor (EF). These data are critical for further quantifying the contribution of organic acids to the atmosphere via SSA, which may provide an estimate of the potential influence of the atmospheric feedbacks to the ocean carbon cycle.

## 1 Introduction

Aromatic acids are considered to be important environmental pollutants due to their potential toxicity and persistence (Zhao et al., 2019). They are ubiquitous in seawater (SW) and the atmosphere and have an important influence on the marine and atmospheric environment. On the one hand, aromatic acids can readily be taken up by marine organisms, leading to their enrichment within these organisms or transportation to remote areas (Fu et al., 2011; Shariati et al., 2021; Wang and Kawamura, 2006). This process poses health risks to the endocrine system of aquatic organisms and the overall marine ecosystem (Saha et al., 2006). Other than the influence on marine systems, they have also been reported to have negative impacts on human health, bioaccu-

mulation, environmental persistence, and climate change (Lu et al., 2021).

Sea spray aerosol (SSA) may contain organic acids that play a major role in Earth's climate. Previous studies suggested that these organic acids can alter the composition of SSA, subsequently influencing atmospheric processes such as cloud condensation nuclei (CCN) or ice nuclei (IN) activities (Moore et al., 2011; Zhu et al., 2019). In addition, organic acids, such as carboxylic acids and aromatic acids, also contribute significantly to ocean acidification (Kumari et al., 2022). Therefore, due to their importance in upper-ocean biogeochemistry and their environmental risk, the transport processes and accumulation potential of aromatic acids at the air–sea interface require a comprehensive understanding.

In various observations, aromatic acids have been detected in both natural and anthropogenic sources (Zhao et al., 2019; Zangrando et al., 2019; Dekiff et al., 2014). Many studies have confirmed that aromatic acids can not only be produced by biological processes but can also be considered potential sources of oil spills, marine plastics, and sunscreen product discharges (Kristensen et al., 2021; Aitken et al., 2018). An important source of marine aromatic acid is the acidic degradation of spilled oil that occurs extensively in human-impacted areas, including coastal oceans and harbors (Lu et al., 2021). Importantly, in addition to biodegradation, the identification of phthalic acid in organisms raises the possibility of its origin from the ingestion of marine plastics (Almulhim et al., 2022). This is consistent with previous studies finding that phthalic acid is primarily derived from anthropogenic sources, such as plasticizers, biomass burning, and fossil fuel combustion (Boreddy et al., 2022; Ding et al., 2021; Ren et al., 2023; Sanjuan et al., 2023; Shumilina et al., 2023; Yang et al., 2020; Zhu et al., 2022), whereas hydroxybenzoic acid has both anthropogenic and natural sources (Castillo et al., 2023; Liao et al., 2019; Lu et al., 2021, 2023; Zhao et al., 2019).

Recent laboratory studies have shown that personal-care products in SW, especially sunscreen (e.g., *o*-hydroxybenzoic acid), are reduced in levels during algal blooms (Franklin et al., 2022). This phenomenon has prompted discussions about the concept of “missing aromatic acids”. Nevertheless, the currently available data do not provide a conclusive explanation for the existence of these missing aromatic acids. One possible reason for the disappearance of these aromatic acids is their release into the atmosphere. The existing datasets obtained from remote marine areas offer evidence of the presence of these compounds in the atmosphere (Fu et al., 2010). Although recent studies outline the critical sources of aromatic acids, much less is known about their transportation via SSA (Castillo et al., 2023; Hu et al., 2022).

SSAs, generated by breaking waves and bubble bursting, are among the major sources of atmospheric particles (Andrae and Rosenfeld, 2008; Angle et al., 2021; Hasencz et al., 2020; Malfatti et al., 2019). SSAs can act as carrier agents for the vertical transport of much more than just sea salt and often include organic surfactants in the ocean, as already shown by field and laboratory studies (Cochran et al., 2016; Franklin et al., 2022; Rastelli et al., 2017).

Recent data indicate that the surface activity and octanol–water partitioning coefficients ( $K_{ow}$ ) of organic compounds may affect their transport efficiency from the aqueous phase to the atmosphere (Olson et al., 2020). Moreover, the molecular structure may induce changes in organic acids’ properties (i.e., surface tension, toxicity), which further affect their transport potential and global emission flux (Lee et al., 2017; Rastelli et al., 2017; Frossard et al., 2019; Acker et al., 2021a). While field studies have demonstrated the presence of aromatic acids in both the ocean and atmosphere

(Boreddy et al., 2017), the specific mechanisms influencing their transport at the air–sea interface require further investigation. Therefore, understanding the factors influencing the enrichment behavior of aromatic acids at the air–sea interface and determining the emission flux of aromatic acids are crucial.

In a previous study, we observed that the transfer of short-chain organic acids between the air and ocean through SSA may be influenced by seawater surface tension. This factor could, in turn, impact the enrichment behavior of organic acids (Song et al., 2022). However, extrapolation of these results to all of the organic acids may not be warranted as the mechanism for enrichment behavior may be closely related to the molecular structure.

In this study, we systematically identified the transport process of aromatic acids from SSA to the atmosphere in the natural seawater environment. To this end, we developed a sea spray aerosol simulation chamber using the plunging jet method, providing the closest proxy for natural SSA currently available in a controlled laboratory monitoring environment, and used it to probe the transport process of aromatic acids. In our SSA simulation chamber, SW and SSA samples were concurrently analyzed for their aromatic acid concentrations, allowing better characterization of any enrichment processes involved in bubble bursting. Finally, the global emission fluxes of aromatic acids via SSA were assessed by combining data on the concentration of aromatic acids in SW and experimental enrichment factor data, which provides unique insights into the enrichment behavior and atmospheric transport of aromatic-acid-containing SSA particles, particularly at the air–sea interface.

## 2 Experimental section

### 2.1 Materials

Aromatic acids were purchased from Shanghai Aladdin Bio-Chem Technology, China. Nine aromatic acids, including benzoic acid, phthalic acids (*o*-, *m*-, and *p*-isomers), hydroxybenzoic acids (*o*-, *m*-, and *p*-isomers), vanillic acid, and syringic acid, were investigated to determine the influence of functional group position and quantity on the transmission of aromatic acids at the sea–air interface (Table 1). Further details are provided in Table S1 in the Supplement, which lists sources and concentrations of these aromatic acids identified in SW and atmospheric samples over the ocean. High-performance liquid chromatography (HPLC)-grade methanol and acetonitrile were supplied by Fisher Scientific, USA. Artificial seawater (ASW) was supplied by Psaitong, China. SW collected from the coast of the Yellow Sea in Qingdao, China, was transported into the laboratory SSA simulation chamber (see Fig. S1 in the Supplement). The range of pH values measured by a pH meter (PHS-3C, Shanghai Yidian Scientific Instrument, China) in all experiments was 7.58–7.92. Ultrapure water with a resistivity of

18.2 M $\Omega$ cm was generated by a Milli-Q purification system (Merck Millipore, France), which can be used to prepare mobile phases for liquid chromatography. Each aromatic acid was first sonicated in natural seawater for 30 min at a concentration of 1 mM. For simplified consideration, aromatic acids were added separately to ASW to achieve concentrations of 1  $\mu$ M and 1 mM in order to verify the effects of background systems and concentrations on the EF of aromatic acids, which may not accurately reflect realistic conditions but provide an approximated trend of EFs instead.

## 2.2 Experimental setup

A jet-based laboratory SSA simulation chamber, made of 316 L stainless-steel material, was used to mimic SSA generation (Fig. S2). This chamber was a clamshell cuboid box (length 30 cm, width 20 cm, height 40 cm) with a viewable glass window, which has recently been adapted for air–sea transport process studies for continuously generated plunging jets to generate realistic SSA (Liu et al., 2022; Xu et al., 2023; Zhan et al., 2022). The detailed dimensions of the SSA simulation chamber are provided in Table S2. All the enrichment experiments were conducted in the SSA simulation chamber filled with approximately 9 L of SW. When used, plunging jets were cycled in SW by the pump (Shenchen V6-6, China) at a flow rate of 1 L min<sup>-1</sup> through a stainless-steel nozzle (inner diameter 4.3 mm). Then the bubble breaking on the surface of the SW was observed through the glass window attached to the SSA simulation chamber. During the experiment, it was assumed that the interaction of the bubble plume generated by the plunging jet with the wall was negligible (Fig. S3).

To avoid any contamination from indoor air, particle-free air was supplied by a compressor and zero-air generator (model 111, Thermo Scientific, USA). A mass flow controller (Beijing Sevenstar Electronics, China) was used to adjust the airflow rate entering the SSA simulation chamber, with a flow rate in the range of 3–50 L min<sup>-1</sup>. The outlet was fitted with a three-way valve to transfer the sample airflow and the excess gas. A Nafion drying tube (MD-700-06S-3, Perma Pure, USA) is used to control the relative humidity of the aerosol particles at the sampling port. The relative humidity and temperature were monitored by a two-channel thermo-hygrometer (Testo HM42, Vaisala, Finland), in the ranges of 30%–40% and 20–25 °C, respectively. Surface tension was measured with approximately 20 mL of SW samples. The tensiometer (JK99C, Powereach, China) was calibrated with 20 mL of ultrapure water at 25 °C and the sheet metal was cleaned with ethanol between each measurement. The surface tension represents collections of at least three independent measurements for each aromatic acid in order to guarantee data reproducibility.

## 2.3 Sample collection and instrumental analysis












The experiment consisted of a total of 25 sets with target compound concentrations of 10<sup>-3</sup> and 1 mM (Table S3). Each aromatic acid was added individually to SW and ASW for the experiment. Particle size distributions were measured at a relative humidity of 34.2%  $\pm$  3.9% using a scanning mobility particle sizer (SMPS, TSI, model 3936). The SMPS consisted of a differential mobility analyzer (DMA, model 3081, TSI, USA) coupled to a condensation particle counter (CPC, model 3776, TSI, USA), which measured particles with diameters between 14.1 and 710.5 nm. The SMPS measured particle number concentration and geometric mean diameter (GMD) within the SSA simulation chamber at a time resolution of 3 min, with an inlet and sheath gas flow rate of 0.3 and 3.0 L min<sup>-1</sup>, respectively. Before turning on the pump, we confirmed that the particle number concentration was < 20 cm<sup>-3</sup> during the first 30 min of each experiment to ensure no leaking and no background particles. Each group of number concentration monitoring experiments lasted approximately 1 h. After the size distribution stabilized, a total of 40 mL of SW was collected from a tap located 1.5 cm above the bottom of the chamber.

SSA samples were collected on aluminum foil (25 mm, Jowin Technology, China) with a 14-stage Dekati low-pressure impactor (DLPI+, Dekati, Finland) for 5 h and then stored at -20 °C until analysis. All samples were analyzed by attenuated total internal reflectance Fourier transform infrared (ATR-FTIR) spectroscopy (Vertex 70, Bruker, Germany) using an automated fitting algorithm and techniques described in a previous study (Xu et al., 2021). In the range of 4000–600 cm<sup>-1</sup>, the ATR-FTIR spectra of SSA particles were recorded with an average of 64 scans. An SW sample was collected as the ATR-FTIR spectra of blank aluminum foil to confirm that there was no infrared absorption of target functional groups on the aluminum foil.

In addition, laboratory SSA transmission electron microscopy (TEM) samples were collected using a single-particle sampler (DKL-2, Genstar Electronic Technology, China). TEM was performed using a TEM cryo-mount (Gatan 626) to load the samples, where the TEM grid was immersed in liquid nitrogen and then mounted on the holder by means of a cryo-transfer workstation. TEM with a high-angle annular-dark-field detector was used and then TEM images were obtained at an accelerating voltage of 200 kV. The SSA particles impacted onto copper grids films (T11023, Tianld, China) with a flow rate of 1 L min<sup>-1</sup> for 1 h. TEM images were used to characterize the morphology of coated and uncoated SSA samples.

Submicron SSA particles were impacted onto 25 mm diameter quartz-fiber filters (QFF, 1851–025, Waterman, UK), which were baked in a muffle furnace at 450 °C for 3 h. Filters collected at stages 6–10 (0.19–0.94  $\mu$ m) of the low-pressure impactor were extracted with 4 mL of ultrapure water under ultrasonication for 30 min. Thermo Scientific

**Table 1.** Summary of aromatic acids used in the experiments.

Aromatic acids				
	benzoic acid	<i>o</i> -phthalic acid	<i>m</i> -phthalic acid	<i>p</i> -phthalic acid
Position of –COOH				
Position of –OH	benzoic acid 	<i>o</i> -hydroxybenzoic acid 	<i>m</i> -hydroxybenzoic acid 	<i>p</i> -hydroxybenzoic acid 
Number of –CH <sub>3</sub> O	<i>p</i> -hydroxybenzoic acid 	vanillic acid 	syringic acid 	

series of high-performance liquid chromatography (HPLC, Vanquish, Germany) coupled with an Atlantis C18 reversed-phase column (2.1 mm × 150 mm, 3 μm particle size, 100 Å, waters) at 30 °C and a mass spectrum were performed to determine the concentration of aromatic acids in SSA and SW samples. The mobile phase consisted of 0.1 % formic acid aqueous solution (A) and acetonitrile and was kept at a flow rate of 0.2 mL min<sup>-1</sup>. An eluent gradient program was used as follows (Witkowski and Gierczak, 2017): starting with 5 % A for 5 min, increasing A from 5 % to 25 % in 2 min, keeping A at 25 % for 4 min, increasing A from 25 % to 95 % in 5 min, and holding constant for 2 min, then turning A to 5 % in 0.5 min and keeping it at 5 % for 6.5 min. ESI was performed in negative ion mode with a capillary voltage of 3.5 kV. The injection volume was set at 10 μL. The standard curves for aromatic acids were constructed over a concentration range of 0.01–1000 μM, with more than seven data points (Fig. S4).

Inorganic ions in SW and SSA samples were filtered through 0.22 μm PTFE filters and analyzed by ion chromatography (Dionex ICS-6000, Thermo Fisher Scientific, USA) coupled with conductivity detection. Cations were isolated on a Dionex IonPac CS12A (4 mm × 250 mm) column, preceded by a guard column, with 20 mM methanesulfonic

acid at a flow rate of 1 mL min<sup>-1</sup>. The sample volume was 10 μL for each injection.

## 2.4 Data analysis

SSA production was used to evaluate the particle yield at the air–sea layer. In a previous laboratory study (Christiansen et al., 2019), the production of aromatic acids in SSA was calculated as

$$\text{SSA production} = N_{\text{Total}} \times Q_{\text{Sweep}}, \quad (1)$$

where  $N_{\text{Total}}$  and  $Q_{\text{Sweep}}$  are the total number concentration of particles detected from SMPS and the airflow rate through the headspace of the chamber, respectively.

The EF of the aromatic acid was then calculated using Eq. (2), where  $[X]_{\text{SSA}}$  and  $[\text{Na}^+]_{\text{SSA}}$  are the concentrations of aromatic acid and Na<sup>+</sup> in SSA, and  $[X]_{\text{SW}}$  and  $[\text{Na}^+]_{\text{SW}}$  denote the concentrations in SW (Sha et al., 2021a).

$$\text{EF} = \frac{[X]_{\text{SSA}}/[\text{Na}^+]_{\text{SSA}}}{[X]_{\text{SW}}/[\text{Na}^+]_{\text{SW}}} \quad (2)$$

An equation to estimate organic acid global flux from SSA emission was developed by Sha et al. (2021b). Both the laboratory-derived EF and the concentration of Na<sup>+</sup> data



were obtained to quantify the global emission flux of the aromatic acid.

$$[X]_{\text{SSA}} = k_{\text{SSA}} \times [\text{Na}^+]_{\text{SSA}} \quad (3)$$

$$[X]_{\text{SSA}} = \text{EF} \times \frac{[X]_{\text{SW}}}{[\text{Na}^+]_{\text{SW}}} \times [\text{Na}^+]_{\text{SSA}} \quad (4)$$

$$\text{flux}_X = k_{\text{SSA}} \times \text{flux}_{[\text{Na}^+]} \quad (5)$$

In these equations,  $k_{\text{SSA}}$  represents the concentration of the aromatic acid transferred to the atmosphere per  $\mu\text{g}$  of  $\text{Na}^+$ . In addition,  $\text{flux}_{[\text{Na}^+]}$  means SSA annual production. Note that this quantification assumes a constant concentration of each aromatic acid reported from the literature and no atmospheric aging process of aromatic acids.

### 3 Results and discussion

#### 3.1 Surface tension of seawater containing aromatic acids

The surface tension of SW was examined using a surface tensiometer (Sigma 700, Biolin Scientific, Sweden) equipped with a Wilhelmy plate, calibrated at  $25^\circ\text{C}$  with 30 mL of ultrapure water. The variation of seawater surface tensions containing different aromatic acids at room temperature is given in Fig. 1. Surface tension measured in SW ranged from  $73.59$  to  $73.84 \text{ mN m}^{-1}$  with a median of  $73.75 \pm 0.06 \text{ mN m}^{-1}$ . However, the average surface tension value of ASW was  $74.5 \text{ mN m}^{-1}$  (Fig. S5), indicating that the presence of surfactants in SW enhances its surface activity. The surface tension of SW containing benzoic acid is  $73.67 \pm 0.03 \text{ mN m}^{-1}$  and certainly lower than that of SW, indicating that benzoic acid slightly reduces the surface tension of SW. Likewise, it is generally observed that surfactants in SW will decrease the surface tension of SW (Cravigan et al., 2020; Liu and Dutcher, 2021; Pierre et al., 2022; Keene et al., 2017; Enders et al., 2023).

The surface tension of both benzenedicarboxylic acids and hydroxybenzoic acids is also lower than that of SW, indicating that the effect of carboxyl and hydroxyl groups on the surface tension of SW is similar. However, differences in the positions of the functional groups of the compounds lead to some differences between the surface tension of SW containing the two types of aromatic acids. SW containing *p*-phthalic acid ( $73.92 \pm 0.14 \text{ mN m}^{-1}$ ) exhibited the highest surface tension among the isomers of benzenedicarboxylic acids. The seawater surface tension varied with the position of the carboxyl group in the order of *p*-phthalic acid > *m*-phthalic acid > *o*-phthalic acid. It is important to note that the benzenedicarboxylic acid used here is amphiphilic, which may be the main cause of the results observed above. The current findings on the surface tension of SW containing aromatic acids are also consistent with a recent study showing that the surface propensity of monocarboxylic acids leads in general to a reduction of surface ten-

sion compared to SW, while dicarboxylic acids give higher values (Ozgurel et al., 2022). The theory applies to hydroxybenzoic acids as the surface tension of SW containing hydroxybenzoic acid was lower than that of SW devoid of aromatic acids, with surface tension depressions ranging from  $0.01$  to  $0.17 \text{ mN m}^{-1}$ .


Unlike the order of seawater surface tension containing benzenedicarboxylic acid, the most obvious inhibition of surface tension was found with the addition of *p*-hydroxybenzoic acid ( $73.58 \pm 0.10 \text{ mN m}^{-1}$ ), while *o*-hydroxybenzoic acid had the weakest effect on seawater surface activity. Similar to the hydroxybenzoic acid, vanillic acid and syringic acid inhibited the surface tension of SW to varying degrees. The results showed that the hydrophobic group (methoxy group) enhanced the surface activity of SW in the order of syringic acid > vanillic acid > *p*-hydroxybenzoic acid. As illustrated in Fig. 1d, the seawater surface activity increased with the number of methoxy groups. And similar trends were also observed in ASW (Fig. S5). Our data demonstrated strong functional group connections to the seawater surface tension containing aromatic acids.

#### 3.2 Transfer of individual aromatic acids to submicron SSA particles

##### 3.2.1 SSA size distribution in the different seawater systems containing individual aromatic acids

The size distribution of SSA generated from the SSA simulation chamber was measured to study the influence of the molecular structure of aromatic acids on SSA production. Figure S6 shows the number size distribution of SSA generated with the SSA simulation chamber in this study, which could be fitted to a lognormal-mode distribution as observed in previous studies (Quinn et al., 2017; Saliba et al., 2019; Xu et al., 2022). In Fig. 2a, the size distribution of sea salt particles and benzoic-acid-added particles shows a peak number concentration near  $186.26 \text{ nm}$ . The number size distribution showed no significant changes in response to SSA particles after adding benzoic acid, with an SSA production rate of  $1.08 \times 10^7 \text{ particles s}^{-1}$ .

To explore the response of SSA formation to the number of carboxyl groups in aromatic acids, we measured the particle size distribution in the system after adding different benzenedicarboxylic acids. The number concentration of SSA particles with added benzenedicarboxylic acids was much higher than that with added benzoic acids. As shown in Fig. 2b, *p*-phthalic acid promoted a much higher SSA particle number concentration, which could be observed visually through SSA simulation chamber window as an increase in bubble bursting. The observed bursting phenomenon is in accordance with the conclusion that higher seawater surface tension could promote the SSA production. The most common explanation is that *p*-phthalic acid has two hydrophilic




**Figure 1.** Measured surface tension values of natural seawater and aromatic-acid-containing seawater: benzoic acids (a), benzenedicarboxylic acids (b), hydroxybenzoic acids (c), *p*-hydroxybenzoic acid, vanillic acid, and syringic acid (d). The dark spots represent the mean values of at least nine data points, the boxes represent the ranges of the 25th–50th–75th percentiles, and the whiskers represent the maximum and minimum values.

carboxyl groups at both ends, and the increase in particle number concentration cannot be attributed to the inhibition effect of surfactants on bubbles bursting. Obviously, the presence of benzenedicarboxylic acid increases the particle number concentration, as described in a recent study (Dubitsky et al., 2023). For the particle size distribution, the size distribution of particles with added benzenedicarboxylic acid was much narrower in contrast to the system with added benzoic acids. The morphology of SSA particles seems to change after the addition of benzenedicarboxylic acid, which is very similar to the SSA chamber studies described by Lv et al. (2020) and Lee et al. (2020b), further indicating that the number and position of carboxylic groups in aromatic acids could significantly affect the formation of SSA.

Further, different functional groups were investigated to determine their effect on particle generation. The distributions show a decrease in the particle number concentrations when adding hydroxybenzoic acids compared with the system with added benzoic acids, leading to the conclusion that hydroxybenzoic acids, acting as surfactants, could inhibit SSA production (Fig. 2c). Compared with the SW with added benzenedicarboxylic acids, the particle number concentration with added hydroxybenzoic acids decreased sig-

nificantly. The hydroxybenzoic acids enhanced the seawater surface activity, which would increase the bubble lifetime and then decrease the SSA production. Importantly, Fig. 2d shows that the particle number concentration decreased proportionally with the increase in the number of methoxy groups, providing further evidence that organic matter with hydrophobic functional groups has preferential atomization ability. Previous studies also showed that particle number concentration and GMD increased with the seawater surface tension (Guzmán et al., 2014; Liu et al., 2022), further indicating that the types of functional groups and the seawater surface tension both affect SSA generation. Moreover, the results showed that the effect of trends of aromatic acids on SSA production in ASW were consistent with those observed in SW (Fig. S7), eliminating the influence of organic matter.

Integrated particle size, SSA production, and mass concentration based on SMPS are shown in Fig. 3. The mode diameter of SSA particles containing *m*-hydroxybenzoic acid was the largest among all the hydroxybenzoic acid isomers, suggesting that *m*-hydroxybenzoic acid was more likely attached to sea salt particles. In all of the benzenedicarboxylic acid position isomers, *p*-phthalic-acid-containing particles gave the largest mode diameter, while SSA particles con-




**Figure 2.** Number concentration distribution of sea salt particles and SSA particles containing benzoic acids (a), benzenedicarboxylic acids (b), hydroxybenzoic acids (c), *p*-hydroxybenzoic acid, vanillic acid, and syringic acid (d). SW represents natural seawater.

taining *o*-phthalic acid expressed the smallest mode diameter. Rather, the SSA particle diameters of vanillic acid and syringic acid did not change significantly, with a deviation of less than 3.75%. For SSA production, the rate of *p*-phthalic-acid-added particles ( $1.54 \times 10^7$  particles  $s^{-1}$ ) is higher than that of *m*-phthalic-acid-added ( $1.33 \times 10^7$  particles  $s^{-1}$ ) and *o*-phthalic-acid-added particles ( $1.05 \times 10^7$  particles  $s^{-1}$ ). In addition, the SSA production of particles when adding *o*-hydroxybenzoic acid ( $1.13 \times 10^7$  particles  $s^{-1}$ ) was obviously higher than that of other isomers (*m*-hydroxybenzoic acid:  $0.90 \times 10^7$  particles  $s^{-1}$ , *p*-hydroxybenzoic acid:  $0.98 \times 10^7$  particles  $s^{-1}$ ) shown in Fig. 3 because the surface tensions of SW when adding *m*-hydroxybenzoic acid and *p*-hydroxybenzoic acid were lower than that of SW. These differences in SSA production can likely be attributed to the selective transfer of aromatic acids in the air–sea phases. As described in previous studies, the bubble formation and break-out were controlled by seawater surface tension (Lee et al., 2020a; Liu and Dutcher, 2021). Furthermore, the mass concentration was increased with SSA production, which is similar to previous findings (Sha et al., 2021a).

### 3.2.2 Morphologies of SSA particles

The particle size and number concentration of particulate matter obtained by SMPS were varied by adding different aromatic acids, which may be closely related to the interaction between aromatic acids and sea salt. To test this hypothesis, SSA particles from SW experimental systems with addition of different aromatic acids were collected and the morphologies of individual particles were characterized (Fig. 4). The TEM images visually show that the cores of the sea salt particles were coated to varying degrees, forming a typical core–shell structure. Furthermore, aluminum foil samples were also characterized by ATR-FTIR for the qualitative analysis of organic coating. The broad band in the range of  $3700\text{--}3000\text{ cm}^{-1}$  is O–H stretching in water or acids (Diniz et al., 2018). The other absorption band peak was observed at approximately  $1643\text{ cm}^{-1}$ , representing O–H stretching in acids (Jin et al., 2013). The image from Fig. S8A provides evidence that SSA produced by bubble bursting can transfer acids. The absorbances in the ranges  $1900\text{--}1670$ ,  $1600\text{--}1480$ , and  $1475\text{--}1300\text{ cm}^{-1}$  are the stretching vibration of C=O in aromatic carboxylic acid, C=C stretching in an aromatic ring, and C–H symmetrical carboxylate stretch-



**Figure 3.** SSA production, particle size, and mass concentration distribution of aromatic acids. The symbol size represents the geometric mean diameter of SSA particles, with the numbers below or above the points giving the geometric mean diameter (in nm), and the error bars are the standard deviation. The symbol color indicates the particle mass concentration, with SW representing natural seawater.

ing (Andreeva and Burkova, 2017; Diniz et al., 2018; Geng et al., 2009; Koutstaal and Ponec, 1993).

As can be seen from Fig. 4a and b, the sea salt particles had a cubic shape, and the benzoic-acid-coated sea salt particles formed a core–shell structure. With this perspective, the finding of TEM images is in agreement with previously reported morphology for SSA particles (Unger et al., 2020). Figure 4c–e clearly show the single-particle morphology characteristics of benzenedicarboxylic acids. Notably, it is evident that the core morphology of the salt particles underwent a significant change, with the cubic structure being transformed into a sphere structure. The core of SSA particles containing *p*-phthalic acid on the TEM grid became more round compared with SSA particles containing *m*-phthalic acid and *o*-phthalic acid (Fig. 4a). As previously shown by a different study, these morphologies may suggest that *o*-phthalic-acid-containing SSA forms round particles by promoting the growth of other surfaces of sea salt particles (Ballabh et al., 2006). The organic coating of sea salt is a well-known process and has been shown specifically from phytoplankton blooms to produce more organics that likely further lead to spherical structures (Ault et al., 2013). Moreover, the organic coating became thicker in the order of *p*-phthalic acid > *m*-phthalic acid > *o*-phthalic acid. This order in the organic coating clearly explained the result discussed above that the GMD of *p*-phthalic-acid-containing particles was markedly larger than those of other isomers of benzenedicarboxylic acids in SSA particles. According to the spectrum in Fig. S8B, for SSA particles containing benzenedicarboxylic acids, the very weak absorption band featuring a range of 1000–650 cm<sup>-1</sup> represents C–H out-


of-plane bending in the present study and in the literature (Chang et al., 2022; Lin et al., 2014; Świsłocka et al., 2013). The in situ ATR-FTIR identified SSA particle functional groups provide the key information on the transfer of aromatic acids from SW to the atmosphere.

Figure 4b shows the morphology images of SSA particles containing hydroxybenzoic acids, where the core maintained the crystalline phase of sea salt with its cubic structure and the coatings mainly consisted of hydroxybenzoic acids. Notably, the organic coating of *m*-hydroxybenzoic acid is the thickest among all the isomers. This is consistent with the result that the GMD of *m*-hydroxybenzoic acid is the largest measured by SMPS. The TEM images in Fig. 4c show that individual nascent SSA containing vanillic acid and syringic acid also has the cubic sea salt core, despite the coating thickness being similar. Both characteristic absorption peaks of hydroxybenzoic acid, vanillic acid, and syringic acid are observed in Fig. S8C and D, consistent with the results for benzenedicarboxylic acids discussed above. Besides, we expressed herein that sea salt particles surrounded by aromatic acid coatings would be also inferred from TEM images and ATR-FTIR spectra.

### 3.3 Enrichment of individual aromatic acids in submicron SSA particles

The enrichment factors (EFs) of different aromatic acids in SSA are characterized to visually demonstrate the influence of different functional groups on the accumulation degree of aromatic acids from SW to the atmosphere through SSA. As shown in Fig. 5, all aromatic acids exhibit varying degrees of enrichment in SSA particles. The EF ranged






**Figure 4.** Particle morphology observed using TEM of sea salt particles and benzenedicarboxylic acids (a), hydroxybenzoic acids (b), *p*-hydroxybenzoic acid, vanillic acid, and syringic-acid-coated sea salt particles (c).


from 5.97 to 24.69, with the largest and smallest being of *m*-hydroxybenzoic acid and *o*-phthalic acid, respectively. The reason for the difference in EFs may be related to the difference in surface tension of SW and the difference in octanol–water partitioning coefficients ( $\log(K_{ow})$ ) (Olson et al., 2020). In detail, the EF of benzoic acid was approximately 10, indicating that benzoic acid can be significantly transferred to the atmosphere, which is consistent with the findings of ASW (Fig. S9). In combination with the TEM images, not only is the *p*-phthalic acid coating the thickest, but its EF is also the largest among its three isomers. Interestingly, the EFs show the  $\log(K_{ow})$  dependence for the three isomers of benzenedicarboxylic acids in the order of *p*-phthalic acid > *m*-phthalic acid > *o*-phthalic acid. A negative correlation between EF and the  $\log(K_{ow})$  of surfactants has been observed (Olson et al., 2020), and the detailed

data are available on the website (<https://comptox.epa.gov/dashboard/>, last access: 17 May 2024) and summarized in Table S4. The EF order of benzenedicarboxylic acids is opposite to that of surfactants, probably due to the amphiphilicity of *o*-phthalic acid. However, the EF of *p*-phthalic acid was significantly lower in ASW, which may be due to the depletion of *p*-phthalic acid acting as an  $\cdot\text{OH}$  scavenger (Li et al., 2023). In contrast, organic matter in SW reacts preferentially with  $\cdot\text{OH}$  and therefore has less effect on *p*-phthalic acid enrichment (Anastasio and Newberg, 2007).

Notably, the correlation between the EF containing the isomers of hydroxybenzoic acid and the corresponding organic coating was also strong, among which the EF of *m*-hydroxybenzoic acid ( $\text{EF} = 24.7 \pm 8.2$ ) was the highest. Meanwhile, the GMD of SSA particles containing *m*-hydroxybenzoic acid was also the highest. One possible rea-



**Figure 5.** Enrichment factors of benzenedicarboxylic acids (a), hydroxybenzoic acids (b), *p*-hydroxybenzoic acid, vanillic acid, and syringic acid (c) from seawater to the atmosphere.




**Figure 6.** Enrichment factors of  $K^+$ ,  $Mg^{2+}$ , and  $Ca^{2+}$  in submicron SSA during the experiment.

son could be that the  $\log(K_{ow})$  for *m*-hydroxybenzoic acid (1.50) supports preferential transfer of *m*-hydroxybenzoic acid to SSA particles. The fundamental role of  $\log(K_{ow})$  in the transfer of organic matter from the aqueous phase to particulate phase has been proven (Olson et al., 2020; Mccord et al., 2018; Maagd et al., 1999). However, the enrichment behavior of *o*-hydroxybenzoic acid is weaker than those of others. Therefore,  $\log(K_{ow})$  is not the only factor affecting the EF, which should be combined with the octanol–air partitioning coefficients ( $\log(K_{oa})$ ). Just as *o*-hydroxybenzoic acid is easily transferred to the seawater surface microlayer according to  $\log(K_{ow})$ , it is not easily transferred to the atmosphere according to  $\log(K_{oa})$ . Compared with *p*-hydroxybenzoic acid, vanillic acid has one more hydropho-

bic group (methoxy) to its molecular structure. Accordingly, the EF of vanillic acid ( $13.3 \pm 2.5$ ) in SSA is slightly higher than that of *p*-hydroxybenzoic acid ( $12.1 \pm 2.5$ ). The EFs of vanillic acid and syringic acid depend on their surface activities. Syringic acid has strong surface activity, making it more easily transported to the atmosphere through SSA than vanillic acid. For most of the target aromatic acids, the enrichment trends are less affected by the compound concentration (Fig. S9). The above EF data indicate that aromatic acids have significant transport potential at the air–sea interface.

We hypothesized that the enrichment of aromatic acids is largely related to the bridging effect of  $Ca^{2+}$  as well, and as a result, the EFs of  $Ca^{2+}$  were calculated. The observed selectivity in cation transfer shown in Fig. 6 has the same trend as the cation binding affinity of aromatic acids. Moreover, another reason for the effective transfer of *m*-hydroxybenzoic acids to SSA particles is their binding to  $Ca^{2+}$ , as evidenced by the significant enrichment of  $Ca^{2+}$  (Fig. 6). It is known that  $Ca^{2+}$  has the potential to transfer organics (Shaloski et al., 2015; Hasencz et al., 2019). The fact is that aromatic acids can form complexes with cations which are then transferred to SSA by bubble bursting. The strong binding abilities of divalent cations to phenolic –OH and aromatic C=C have been reported (Jayarathne et al., 2016). Thus, the enrichment of major cations also acts as a standard for assessing the abundance of organic matter. In Fig. 6, the enrichment of  $Ca^{2+}$  is the most obvious, followed by those of  $Mg^{2+}$  and  $K^+$  among all the experiments. In SSA,  $Ca^{2+}$  always exhibited high enrichment (EF > 1), while the EFs for  $K^+$  and  $Mg^{2+}$  were around and just below 1. The above observation of the enrichment capacity of  $Ca^{2+}$  is consistent with recent studies (Salter et al., 2016; Unger et al., 2020; Lee et al., 2021; Acker et al., 2021b). Interestingly, the greater the EF for  $Ca^{2+}$ , the larger the EF for the corresponding aromatic acids. Taking *m*-hydroxybenzoic acid as an example, we calculated



**Figure 7.** Concentration range of aromatic acids in seawater (a) and the estimated range of annual global aromatic acid emission ( $\text{tyr}^{-1}$ ) via SSA (b). Yellow and blue stacked columns represent emissions based on Textor et al. (2006) and Gliß et al. (2021), respectively.

the enrichment factors of  $\text{Ca}^{2+}$  and *m*-hydroxybenzoic acid in NaCl solution, ASW and SW, and found that both follow the pattern:  $\text{EF}_{\text{NaCl}} < \text{EF}_{\text{ASW}} < \text{EF}_{\text{SW}}$  (Fig. S10). Hence, the current results further confirm that  $\text{Ca}^{2+}$  bridging is important in the formation of complexes and transport of organics.

### 3.4 Estimation of SSA contribution to global aromatic acid emissions

The  $k_{\text{SSA}}$  for aromatic acids ranged from 0.0002 to 0.623 (Table S5). These values were strongly influenced by the concentration of aromatic acid in SW, with differences of 1–4 orders of magnitude. For example, the seawater concentrations of syringic acid, *p*-hydroxybenzoic acid, and benzoic acid are 0.3, 4.58, and  $34 \text{ ng L}^{-1}$ , respectively (Zhao et al., 2019). Such large variations highlight the importance of concentration variations in SW of aromatic acids in assessing the SSA contribution. More than that, benzoic acid and *p*-hydroxybenzoic acid concentrations in the ocean were 205 and  $8.66 \text{ ng L}^{-1}$ , respectively (Zhao et al., 2019). Therefore, it is necessary to estimate their atmospheric fluxes. Notably, for syringic acid,  $k_{\text{SSA}}$  is equivalent to that of *o*-hydroxybenzoic acid, while the concentration of syringic acid in SW is obviously lower. As a result, EFs of aromatic acids also contribute to SSA emission flux. Furthermore, the enrichment process in the field is much more complicated than in controlled laboratory experiments. For example, seawater temperature affects SSA release, while wind speed at the sea surface may influence the amount and size of SSA particles emitted. As such, further research on the environmental enrichment mechanism of aromatic acids in SSA particles is required to reduce the uncertainty in the estimation of aromatic acid emissions.

On the basis of  $k_{\text{SSA}}$  and modeled annual SSA particles, the global emission fluxes of aromatic acids from SW to the atmosphere through SSA particles can be estimated as in Eq. (4). The flux of  $\text{Na}^+$  was obtained by using the annual emission of SSA calculated by Textor et al. (2006) and Gliß et al. (2021). The estimate of aromatic acid fluxes using the above  $\text{Na}^+$  global emission is included in Table S6 for reference, and the comparison between aromatic acid fluxes is also shown in Fig. 7b. For example, the laboratory-derived annual flux of benzoic acids was  $27 \text{ tyr}^{-1}$  for SSA emission of approximately  $3.65 \times 10^{12} \text{ kg yr}^{-1}$  but increased to  $71 \text{ tyr}^{-1}$  when SSA emissions were about  $9.7 \text{ kg yr}^{-1}$ . Therefore, the SSA emission used in the global annual emission of aromatic acids may be one possible reason for the observed differences. Similarly, the atmospheric fluxes measured for aromatic acids are of a similar order of magnitude as those for perfluorocarboxylic acids in a previous laboratory study based on field samples (Sha et al., 2021b). It follows that using lab-derived global annual fluxes gives results that are close to the modeling results. As illustrated in Fig. 7, vanillic acid has the highest global emission among all aromatic acids studied, although it has a lower concentration in SW than others. This demonstrates that the EF might play a very important role in the global emission fluxes of organic matter, in addition to concentration. Moreover, an in-depth quantification of the organic matter global emission flux transferred to the atmosphere through SSA particles will help estimate the CCN influence of the organic contribution.

## 4 Conclusion

Based on our experimental data, we highlight that aromatic acids can be transferred from SW to the atmosphere through

bubble bursting. The air–sea transfer efficiency of aromatic acids was evaluated by simulating SSA generation with a plunging jet and collecting the SSA particles. First and foremost, aromatic acids being transferred from SW to the SSA particles was confirmed as possible by combining infrared spectra and TEM images. TEM images intuitively show that aromatic acids are coated with sea salt particles to form a core–shell structure, which further proved that there is a good correlation between the EF of organic acids and organic coating. Our data confirm that the enrichment of aromatic acid was enhanced by the increase in hydrophobic functional groups, while the corresponding number concentration of SSA particles decreased. As a whole, the transfer capacity of aromatic acids may depend on their functional groups and on the bridging effect of cations, as well as their concentration in SW, as these factors influence the global emission flux of aromatic acids via SSA particles. Overall, our research helps to close an existing knowledge gap in studying the global annual flux of aromatic acids from SSA particles to the atmosphere. Though the current results deepen the understanding of aromatic acid transport, much detailed work is still needed to explore how different aromatic acids affect the marine carbon cycle.

**Data availability.** The data used in this study can be found online at <https://doi.org/10.5281/zenodo.10903140> (Song et al., 2024).

**Supplement.** Additional information includes six tables and 10 figures. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-24-5847-2024-supplement>.

**Author contributions.** YS: conceptualization, methodology, investigation, formal analysis, writing – original draft preparation. JL: writing – reviewing and editing. NTT: writing – reviewing and editing. KL: writing – reviewing and editing. LD: investigation, conceptualization, funding acquisition, project administration, resources, writing – reviewing and editing.

**Competing interests.** The contact author has declared that none of the authors has any competing interests.

**Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

**Acknowledgements.** We would like to thank Xiaojun Li from the State Key Laboratory of Microbial Technology of Shandong University for help and guidance in TEM.

**Financial support.** This research has been supported by the National Natural Science Foundation of China (grant nos. 22076099 and 22376121).

**Review statement.** This paper was edited by Paul Zieger and reviewed by Matthew Salter and two anonymous referees.

## References

- Acker, E. V., Huysman, S., Rijcke, M. D., Asselman, J., Schamphelaere, K. A. C. D., Vanhaecke, L., and Janssen, C. R.: Phycotoxin-enriched sea spray aerosols: Methods, mechanisms, and human exposure, *Environ. Sci. Technol.*, 55, 6184–6196, <https://doi.org/10.1021/acs.est.1c00995>, 2021a.
- Acker, E. V., Rijcke, M. D., Liu, Z., Asselman, J., Schamphelaere, K. A. C. D., Vanhaecke, L., and Janssen, C. R.: Sea spray aerosols contain the major component of human lung surfactant, *Environ. Sci. Technol.*, 55, 15989–16000, <https://doi.org/10.1021/acs.est.1c04075>, 2021b.
- Aitken, C. M., Head, I. M., Jones, D. M., Rowland, S. J., Scarlett, A. G., and West, C. E.: Comprehensive two-dimensional gas chromatography-mass spectrometry of complex mixtures of anaerobic bacterial metabolites of petroleum hydrocarbons, *J. Chromatogr. A*, 1536, 96–109, <https://doi.org/10.1016/j.chroma.2017.06.027>, 2018.
- Almulhim, F., Rossbach, S., Emwas, A.-H., Kharbatia, N. M., Jaremko, L., Jaremko, M., and Duarte, C. M.: Metabolomic study on *tridacna maxima* giant clams reveals metabolic fingerprint of environmental pollutants, *Front. Mar. Sci.*, 9, 813404, <https://doi.org/10.3389/fmars.2022.813404>, 2022.
- Anastasio, C. and Newberg, J. T.: Sources and sinks of hydroxyl radical in sea-salt particles, *J. Geophys. Res.-Atmos.*, 112, D10306, <https://doi.org/10.1029/2006jd008061>, 2007.
- Andreae, M. O. and Rosenfeld, D.: Aerosol-cloud-precipitation interactions. Part 1. The nature and sources of cloud-active aerosols, *Earth-Sci. Rev.*, 89, 13–41, <https://doi.org/10.1016/j.earscirev.2008.03.001>, 2008.
- Andreeva, O. A. and Burkova, L. A.: IR spectroscopy studies of sodium salts of some aminobenzoic acid derivatives, *Russ. J. Phys. Chem. B*, 11, 411–418, <https://doi.org/10.1134/s1990793117030149>, 2017.
- Angle, K. J., Crocker, D. R., Simpson, R. M. C., Mayer, K. J., Garofalo, L. A., Moore, A. N., Mora Garcia, S. L., Or, V. W., Srinivasan, S., Farhan, M., Sauer, J. S., Lee, C., Pothier, M. A., Farmer, D. K., Martz, T. R., Bertram, T. H., Cappa, C. D., Prather, K. A., and Grassian, V. H.: Acidity across the interface from the ocean surface to sea spray aerosol, *P. Natl. Acad. Sci. USA*, 118, e2018397118, <https://doi.org/10.1073/pnas.2018397118>, 2021.
- Ault, A. P., Moffet, R. C., Baltrusaitis, J., Collins, D. B., Ruppel, M. J., Cuadra-Rodriguez, L. A., Zhao, D., Guasco, T. L., Ebben, C. J., Geiger, F. M., Bertram, T. H., Prather, K. A., and Grassian, V. H.: Size-dependent changes in sea spray aerosol composition and



- properties with different seawater conditions, *Environ. Sci. Technol.*, 47, 5603–5612, <https://doi.org/10.1021/es400416g>, 2013.
- Ballabh, A., Trivedi, D. R., Dastidar, P., Ghosh, P. K., Pramanik, A., and Kumar, V. G.: A practical approach to produce near-spherical common salt crystals with better flow characteristics, *Cryst. Growth Des.*, 6, 1591–1594, <https://doi.org/10.1021/cg050633v>, 2006.
- Boreddy, S. K. R., Mochizuki, T., Kawamura, K., Bikkina, S., and Sarin, M. M.: Homologous series of low molecular weight (C1–C10) monocarboxylic acids, benzoic acid and hydroxyacids in fine-mode (PM<sub>2.5</sub>) aerosols over the Bay of Bengal: Influence of heterogeneity in air masses and formation pathways, *Atmos. Environ.*, 167, 170–180, <https://doi.org/10.1016/j.atmosenv.2017.08.008>, 2017.
- Boreddy, S. K. R., Hegde, P., Arun, B. S., Aswini, A. R., and Babu, S. S.: Molecular composition and light-absorbing properties of organic aerosols from west-coast of tropical India, *Sci. Total Environ.*, 845, 157163, <https://doi.org/10.1016/j.scitotenv.2022.157163>, 2022.
- Castillo, A., Celeiro, M., Lores, M., Grgić, K., Banoić, M., Jerković, I., and Jokić, S.: Bioprospecting of Targeted Phenolic Compounds of *Dictyota dichotoma*, *Gongolaria barbata*, *Ericaria amentacea*, *Sargassum hornsuschuhii* and *Ellisolandia elongata* from the Adriatic Sea Extracted by Two Green Methods, *Mar. Drugs*, 21, 97, <https://doi.org/10.3390/md21020097>, 2023.
- Chang, S., Li, M., Gao, K., Zhang, D., Duan, H., Ma, L., and Ruan, Z.: Mechanism of phthalic acid collector in flotation separation of fluorite and rare earth, *J. Rare Earths*, 40, 118–126, <https://doi.org/10.1016/j.jre.2020.11.002>, 2022.
- Christiansen, S., Salter, M. E., Gorokhova, E., Nguyen, Q. T., and Bilde, M.: Sea spray aerosol formation: Laboratory results on the role of air entrainment, water temperature, and phytoplankton biomass, *Environ. Sci. Technol.*, 53, 13107–13116, <https://doi.org/10.1021/acs.est.9b04078>, 2019.
- Cochran, R. E., Laskina, O., Jayarathne, T., Laskin, A., Laskin, J., Lin, P., Sultana, C., Lee, C., Moore, K. A., Cappa, C. D., Bertram, T. H., Prather, K. A., Grassian, V. H., and Stone, E. A.: Analysis of organic anionic surfactants in fine and coarse fractions of freshly emitted sea spray aerosol, *Environ. Sci. Technol.*, 50, 2477–2486, <https://doi.org/10.1021/acs.est.5b04053>, 2016.
- Cravigan, L. T., Mallet, M. D., Vaattovaara, P., Harvey, M. J., Law, C. S., Modini, R. L., Russell, L. M., Stelcer, E., Cohen, D. D., Olsen, G., Safi, K., Burrell, T. J., and Ristovski, Z.: Sea spray aerosol organic enrichment, water uptake and surface tension effects, *Atmos. Chem. Phys.*, 20, 7955–7977, <https://doi.org/10.5194/acp-20-7955-2020>, 2020.
- Dekiff, J. H., Remy, D., Klasmeier, J., and Fries, E.: Occurrence and spatial distribution of microplastics in sediments from Norderney, *Environ. Pollut.*, 186, 248–256, <https://doi.org/10.1016/j.envpol.2013.11.019>, 2014.
- Ding, Z., Du, W., Wu, C., Cheng, C., Meng, J., Li, D., Ho, K., Zhang, L., and Wang, G.: Summertime atmospheric dicarboxylic acids and related SOA in the background region of Yangtze River Delta, China: Implications for heterogeneous reaction of oxalic acid with sea salts, *Sci. Total Environ.*, 757, 143741, <https://doi.org/10.1016/j.scitotenv.2020.143741>, 2021.
- Diniz, L. F., Souza, M. S., Carvalho Jr., P. S., Silva, C. C. P., D’Vries, R. F., and Ellena, J.: Novel isoniazid cocrystals with aromatic carboxylic acids: Crystal engineering, spectroscopy and thermochemical investigations, *J. Mol. Struct.*, 1153, 58–68, <https://doi.org/10.1016/j.molstruc.2017.09.115>, 2018.
- Dubitsky, L., McRae, O., and Bird, J. C.: Enrichment of scavenged particles in jet drops determined by bubble size and particle position, *Phys. Rev. Lett.*, 130, 054001, <https://doi.org/10.1103/PhysRevLett.130.054001>, 2023.
- Enders, A. A., Elliott, S. M., and Allen, H. C.: Carbon on the ocean surface: Temporal and geographical investigation, *ACS Earth Space Chem.*, 7, 360–369, <https://doi.org/10.1021/acsearthspacechem.2c00248>, 2023.
- Franklin, E. B., Amiri, S., Crocker, D., Morris, C., Mayer, K., Sauer, J. S., Weber, R. J., Lee, C., Malfatti, F., Cappa, C. D., Bertram, T. H., Prather, K. A., and Goldstein, A. H.: Anthropogenic and biogenic contributions to the organic composition of coastal submicron sea spray aerosol, *Environ. Sci. Technol.*, 56, 16633–16642, <https://doi.org/10.1021/acs.est.2c04848>, 2022.
- Frossard, A. A., Gérard, V., Duplessis, P., Kinsey, J. D., Lu, X., Zhu, Y., Bisgrove, J., Maben, J. R., Long, M. S., Chang, R. Y.-W., Beaupré, S. R., Kieber, D. J., Keene, W. C., Nozière, B., and Cohen, R. C.: Properties of seawater surfactants associated with primary marine aerosol particles produced by bursting bubbles at a model air–sea interface, *Environ. Sci. Technol.*, 53, 9407–9417, <https://doi.org/10.1021/acs.est.9b02637>, 2019.
- Fu, P., Kawamura, K., and Miura, K.: Molecular characterization of marine organic aerosols collected during a round-the-world cruise, *J. Geophys. Res.*, 116, D13302, <https://doi.org/10.1029/2011jd015604>, 2011.
- Fu, P. Q., Kawamura, K., Pavuluri, C. M., Swaminathan, T., and Chen, J.: Molecular characterization of urban organic aerosol in tropical India: contributions of primary emissions and secondary photooxidation, *Atmos. Chem. Phys.*, 10, 2663–2689, <https://doi.org/10.5194/acp-10-2663-2010>, 2010.
- Geng, W., Nakajima, T., Takanashi, H., and Ohki, A.: Analysis of carboxyl group in coal and coal aromaticity by Fourier transform infrared (FT-IR) spectrometry, *Fuel*, 88, 139–144, <https://doi.org/10.1016/j.fuel.2008.07.027>, 2009.
- Gliß, J., Mortier, A., Schulz, M., Andrews, E., Balkanski, Y., Bauer, S. E., Benedictow, A. M. K., Bian, H., Checa-Garcia, R., Chin, M., Ginoux, P., Griesfeller, J. J., Heckel, A., Kipling, Z., Kirkevåg, A., Kokkola, H., Laj, P., Le Sager, P., Lund, M. T., Lund Myhre, C., Matsui, H., Myhre, G., Neubauer, D., van Noije, T., North, P., Oliví, D. J. L., Rémy, S., Sogacheva, L., Takemura, T., Tsigaridis, K., and Tsyro, S. G.: AeroCom phase III multi-model evaluation of the aerosol life cycle and optical properties using ground- and space-based remote sensing as well as surface in situ observations, *Atmos. Chem. Phys.*, 21, 87–128, <https://doi.org/10.5194/acp-21-87-2021>, 2021.
- Guzmán, E., Santini, E., Benedetti, A., Ravera, F., Ferrari, M., and Liggieri, L.: Surfactant induced complex formation and their effects on the interfacial properties of seawater, *Colloid. Surface. B*, 123, 701–709, <https://doi.org/10.1016/j.colsurfb.2014.10.010>, 2014.
- Hasencz, E. S., Kaluarachchi, C. P., Lee, H. D., Tivanski, A. V., and Stone, E. A.: Saccharide transfer to sea spray aerosol enhanced by surface activity, calcium, and protein interactions, *ACS Earth Space Chem.*, 3, 2539–2548, <https://doi.org/10.1021/acsearthspacechem.9b00197>, 2019.
- Hasencz, E. S., Jayarathne, T., Pendergraft, M. A., Santander, M. V., Mayer, K. J., Sauer, J., Lee, C., Gibson, W. S., Kruse, S.



- M., Malfatti, F., Prather, K. A., and Stone, E. A.: Marine bacteria affect saccharide enrichment in sea spray aerosol during a phytoplankton bloom, *ACS Earth Space Chem.*, 4, 1638–1649, <https://doi.org/10.1021/acsearthspacechem.0c00167>, 2020.
- Hu, J., Loh, P. S., Chang, Y.-P., and Yang, C.-W.: Multi-proxy records of paleoclimatic changes in sediment core ST2 from the southern Zhejiang-Fujian muddy coastal area since 1650 yr BP, *Cont. Shelf Res.*, 239, 104717, <https://doi.org/10.1016/j.csr.2022.104717>, 2022.
- Jayarathne, T., Sultana, C. M., Lee, C., Malfatti, F., Cox, J. L., Pendergraft, M. A., Moore, K. A., Azam, F., Tivanski, A. V., Cappa, C. D., Bertram, T. H., Grassian, V. H., Prather, K. A., and Stone, E. A.: Enrichment of saccharides and divalent cations in sea spray aerosol during two phytoplankton blooms, *Environ. Sci. Technol.*, 50, 11511–11520, <https://doi.org/10.1021/acs.est.6b02988>, 2016.
- Jin, S., Wang, D., Linhe, Q., Fu, M., Wu, S., and Ren, J.: Crystal and molecular structure of two organic acid–base salts from nicotinamide and aromatic acids, *J. Chem. Crystallogr.*, 43, 258–265, <https://doi.org/10.1007/s10870-013-0413-2>, 2013.
- Keene, W. C., Long, M. S., Kieber, D. J., Reid, J. S., Maben, J. R., Russell, L. M., Kinsey, J. D., Frossard, A. A., Quinn, P. K., and Bates, T. S.: Factors that modulate properties of primary marine aerosol generated from ambient seawater on ships at sea, *J. Geophys. Res.-Atmos.*, 122, 11961–11990, <https://doi.org/10.1002/2017jd026872>, 2017.
- Koutstaal, C. A. and Ponec, V.: FT-IR study on the adsorption of benzoic acid and its derivatives on transition-metal oxides, *Appl. Surf. Sci.*, 70, 206–210, [https://doi.org/10.1016/0169-4332\(93\)90428-E](https://doi.org/10.1016/0169-4332(93)90428-E), 1993.
- Kristensen, M., Johnsen, A. R., and Christensen, J. H.: Super-complex mixtures of aliphatic- and aromatic acids may be common degradation products after marine oil spills: A lab-study of microbial oil degradation in a warm, pre-exposed marine environment, *Environ. Pollut.*, 285, 117264, <https://doi.org/10.1016/j.envpol.2021.117264>, 2021.
- Kumari, V. R., Sarma, V. V. S. S., and Kumar, M. D.: Spatial variability in aerosol composition and its seawater acidification potential in coastal waters of the western coastal Bay of Bengal, *J. Earth Syst. Sci.*, 131, 251, <https://doi.org/10.1007/s12040-022-01996-w>, 2022.
- Lee, C., Dommer, A. C., Schiffer, J. M., Amaro, R. E., Grassian, V. H., and Prather, K. A.: Cation-driven lipopolysaccharide morphological changes impact heterogeneous reactions of nitric acid with sea spray aerosol particles, *J. Phys. Chem. Lett.*, 12, 5023–5029, <https://doi.org/10.1021/acs.jpclett.1c00810>, 2021.
- Lee, H. D., Estillore, A. D., Morris, H. S., Ray, K. K., Alejandro, A., Grassian, V. H., and Tivanski, A. V.: Direct surface tension measurements of individual sub-micrometer particles using atomic force microscopy, *J. Phys. Chem. A*, 121, 8296–8305, <https://doi.org/10.1021/acs.jpca.7b04041>, 2017.
- Lee, H. D., Morris, H. S., Laskina, O., Sultana, C. M., Lee, C., Jayarathne, T., Cox, J. L., Wang, X., Hasenecz, E. S., DeMott, P. J., Bertram, T. H., Cappa, C. D., Stone, E. A., Prather, K. A., Grassian, V. H., and Tivanski, A. V.: Organic enrichment, physical phase state, and surface tension depression of nascent core–shell sea spray aerosols during two phytoplankton blooms, *ACS Earth Space Chem.*, 4, 650–660, <https://doi.org/10.1021/acsearthspacechem.0c00032>, 2020a.
- Lee, H. D., Wigley, S., Lee, C., Or, V. W., Hasenecz, E. S., Stone, E. A., Grassian, V. H., Prather, K. A., and Tivanski, A. V.: Physicochemical mixing state of sea spray aerosols: Morphologies exhibit size dependence, *ACS Earth Space Chem.*, 4, 1604–1611, <https://doi.org/10.1021/acsearthspacechem.0c00153>, 2020b.
- Li, K., Guo, Y., Nizkorodov, S. A., Rudich, Y., Angelaki, M., Wang, X., An, T., Perrier, S., and George, C.: Spontaneous dark formation of OH radicals at the interface of aqueous atmospheric droplets, *P. Natl. Acad. Sci. USA*, 120, e2220228120, <https://doi.org/10.1073/pnas.2220228120>, 2023.
- Liao, C., Shi, J., Wang, X., Zhu, Q., and Kannan, K.: Occurrence and distribution of parabens and bisphenols in sediment from northern Chinese coastal areas, *Environ. Pollut.*, 253, 759–767, <https://doi.org/10.1016/j.envpol.2019.07.076>, 2019.
- Lin, C.-J., Xu, J.-Q., Zheng, Y.-Q., Zhu, H.-L., and Xu, W.: Synthesis, crystal structures, and properties of two new Cu(II) complexes with p-hydroxybenzoic acid, *J. Clust. Sci.*, 26, 1253–1265, <https://doi.org/10.1007/s10876-014-0810-5>, 2014.
- Liu, L., Du, L., Xu, L., Li, J., and Tsona, N. T.: Molecular size of surfactants affects their degree of enrichment in the sea spray aerosol formation, *Environ. Res.*, 206, 112555, <https://doi.org/10.1016/j.envres.2021.112555>, 2022.
- Liu, S. and Dutcher, C. S.: Measurements of static and dynamic bubble surface tension using a deformation-based microfluidic tensiometer, *J. Phys. Chem. B*, 125, 13916–13927, <https://doi.org/10.1021/acs.jpcc.1c06710>, 2021.
- Lu, S., Lin, C., Lei, K., Xin, M., Wang, B., Ouyang, W., Liu, X., and He, M.: Endocrine-disrupting chemicals in a typical urbanized bay of Yellow Sea, China: Distribution, risk assessment, and identification of priority pollutants, *Environ. Pollut.*, 287, 117588, <https://doi.org/10.1016/j.envpol.2021.117588>, 2021.
- Lu, S., Wang, J., Wang, B., Xin, M., Lin, C., Gu, X., Lian, M., and Li, Y.: Comprehensive profiling of the distribution, risks and priority of pharmaceuticals and personal care products: A large-scale study from rivers to coastal seas, *Water Res.*, 230, 119591, <https://doi.org/10.1016/j.watres.2023.119591>, 2023.
- Lv, C., Tsona, N. T., and Du, L.: Sea spray aerosol formation: Results on the role of different parameters and organic concentrations from bubble bursting experiments, *Chemosphere*, 252, 126456, <https://doi.org/10.1016/j.chemosphere.2020.126456>, 2020.
- Maagd, P. G.-J. D., Hendriks, A. J., Seinen, W., and Sijm, D. T. H. M.: pH-dependent hydrophobicity of the cyanobacteria toxin microcystin-LR, *Water Res.*, 33, 677–680, [https://doi.org/10.1016/S0043-1354\(98\)00258-9](https://doi.org/10.1016/S0043-1354(98)00258-9), 1999.
- Malfatti, F., Lee, C., Tinta, T., Pendergraft, M. A., Celussi, M., Zhou, Y., Sultana, C. M., Rotter, A., Axson, J. L., Collins, D. B., Santander, M. V., Morales, A. L. A., Aluwihare, L. I., Riemer, N., Grassian, V. H., Azam, F., and Prather, K. A.: Detection of active microbial enzymes in nascent sea spray aerosol: Implications for atmospheric chemistry and climate, *Environ. Sci. Tech. Lett.*, 6, 171–177, <https://doi.org/10.1021/acs.estlett.8b00699>, 2019.
- McCord, J., Lang, J. R., Hill, D., Strynar, M., and Chernoff, N.: pH dependent octanol-water partitioning coefficients of microcystin congeners, *J. Water Health*, 16, 340–345, <https://doi.org/10.2166/wh.2018.257>, 2018.
- Moore, M. J. K., Furutani, H., Roberts, G. C., Moffet, R. C., Gilles, M. K., Palenik, B., and Prather, K. A.: Effect of organic compounds on cloud condensation nuclei (CCN) activity of sea spray

- aerosol produced by bubble bursting, *Atmos. Environ.*, 45, 7462–7469, <https://doi.org/10.1016/j.atmosenv.2011.04.034>, 2011.
- Olson, N. E., Cooke, M. E., Shi, J. H., Birbeck, J. A., Westrick, J. A., and Ault, A. P.: Harmful algal bloom toxins in aerosol generated from inland lake water, *Environ. Sci. Technol.*, 54, 4769–4780, <https://doi.org/10.1021/acs.est.9b07727>, 2020.
- Ozgurel, O., Dufлот, D., Masella, M., Réal, F., and Toubin, C.: A molecular scale investigation of organic/inorganic ion selectivity at the air-liquid interface, *ACS Earth Space Chem.*, 6, 1698–1716, <https://doi.org/10.1021/acsearthspacechem.1c00394>, 2022.
- Pierre, J., Poujol, M., and Séon, T.: Influence of surfactant concentration on drop production by bubble bursting, *Phys. Rev. Fluids*, 7, 073602, <https://doi.org/10.1103/PhysRevFluids.7.073602>, 2022.
- Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L. M., and Bates, T. S.: Small fraction of marine cloud condensation nuclei made up of sea spray aerosol, *Nat. Geosci.*, 10, 674–679, <https://doi.org/10.1038/Ngeo3003>, 2017.
- Rastelli, E., Corinaldesi, C., Dell'Anno, A., Lo Martire, M., Greco, S., Cristina Facchini, M., Rinaldi, M., O'Dowd, C., Ceburnis, D., and Danovaro, R.: Transfer of labile organic matter and microbes from the ocean surface to the marine aerosol: an experimental approach, *Sci. Rep.-UK*, 7, 11475, <https://doi.org/10.1038/s41598-017-10563-z>, 2017.
- Ren, L., Weng, L., Chen, D., Hu, H., Jia, Y., and Zhou, J. L.: Bioremediation of PAEs-contaminated saline soil: The application of a marine bacterial strain isolated from mangrove sediment, *Mar. Pollut. Bull.*, 192, 115071, <https://doi.org/10.1016/j.marpolbul.2023.115071>, 2023.
- Saha, N. C., Bhunia, F., and Kaviraj, A.: Comparative toxicity of three organic acids to freshwater organisms and their impact on aquatic ecosystems, *Hum. Ecol. Risk Assess.*, 12, 192–202, <https://doi.org/10.1080/10807030500430625>, 2006.
- Saliba, G., Chen, C. L., Lewis, S., Russell, L. M., Rivellini, L. H., Lee, A. K. Y., Quinn, P. K., Bates, T. S., Haentjens, N., Boss, E. S., Karp-Boss, L., Baetge, N., Carlson, C. A., and Behrenfeld, M. J.: Factors driving the seasonal and hourly variability of sea-spray aerosol number in the North Atlantic, *P. Natl. Acad. Sci. USA*, 116, 20309–20314, <https://doi.org/10.1073/pnas.1907574116>, 2019.
- Salter, M. E., Hamacher-Barth, E., Leck, C., Werner, J., Johnson, C. M., Riipinen, I., Nilsson, E. D., and Zieger, P.: Calcium enrichment in sea spray aerosol particles, *Geophys. Res. Lett.*, 43, 8277–8285, <https://doi.org/10.1002/2016gl070275>, 2016.
- Sanjuan, O. N., Sait, S. T. L., Gonzalez, S. V., Tomas, J., Raga, J. A., and Asimakopoulou, A. G.: Phthalate metabolites in loggerhead marine turtles (*Caretta caretta*) from the Mediterranean Sea (East Spain region), *Environ. Toxicol. Chem.*, 5, 178–185, <https://doi.org/10.1016/j.enceco.2023.08.003>, 2023.
- Sha, B., Johansson, J. H., Benskin, J. P., Cousins, I. T., and Salter, M. E.: Influence of water concentrations of perfluoroalkyl acids (PFAAs) on their size-resolved enrichment in nascent sea spray aerosols, *Environ. Sci. Technol.*, 55, 9489–9497, <https://doi.org/10.1021/acs.est.0c03804>, 2021a.
- Sha, B., Johansson, J. H., Tunved, P., Bohlin-Nizzetto, P., Cousins, I. T., and Salter, M. E.: Sea spray aerosol (SSA) as a source of perfluoroalkyl acids (PFAAs) to the atmosphere: Field evidence from long-term air monitoring, *Environ. Sci. Technol.*, 56, 228–238, <https://doi.org/10.1021/acs.est.1c04277>, 2021b.
- Shaloski, M. A., Sobyra, T. B., and Nathanson, G. M.: DCI transport through dodecyl sulfate films on salty glycerol: Effects of seawater ions on gas entry, *J. Phys. Chem. A*, 119, 12357–12366, <https://doi.org/10.1021/acs.jpca.5b07298>, 2015.
- Shariati, S., Pourbabae, A. A., Alikhani, H. A., and Rezaei, K. A.: Anaerobic biodegradation of phthalic acid by an indigenous *Ralstonia pickettii* strain SHAn2 isolated from Anzali international wetland, *Int. J. Environ. Sci. Te.*, 19, 4827–4838, <https://doi.org/10.1007/s13762-021-03677-5>, 2021.
- Shumilina, E., Skavang, P. K., and Dikiy, A.: Application of NMR spectroscopy for the detection and quantification of phthalic acid in fish muscles: The case of Atlantic Cod from Norwegian Sea, *Mar. Environ. Res.*, 188, 105973, <https://doi.org/10.1016/j.marenvres.2023.105973>, 2023.
- Song, Y., Li, J., Tsona, N. T., Liu, L., and Du, L.: Enrichment of short-chain organic acids transferred to submicron sea spray aerosols, *Sci. Total Environ.*, 851, 158122, <https://doi.org/10.1016/j.scitotenv.2022.158122>, 2022.
- Song, Y., Li, J., Tsona Tchinda, N., Li, K., and Du, L.: Song et al. Peak areas of aromatic acids and cations in seawater and SSA filter samples & Enrichment factors for aromatic acids and cations in submicron SSA, Zenodo [data set], <https://doi.org/10.5281/zenodo.10903141>, 2024.
- Świsłocka, R., Regulska, E., Samsonowicz, M., and Lewandowski, W.: Experimental and theoretical study on benzoic acid derivatives, *J. Mol. Struct.*, 1044, 181–187, <https://doi.org/10.1016/j.molstruc.2012.12.005>, 2013.
- Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, I., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles within AeroCom, *Atmos. Chem. Phys.*, 6, 1777–1813, <https://doi.org/10.5194/acp-6-1777-2006>, 2006.
- Unger, I., Saak, C. M., Salter, M., Zieger, P., Patanen, M., and Bjorneholm, O.: Influence of organic acids on the surface composition of sea spray aerosol, *J. Phys. Chem. A*, 124, 422–429, <https://doi.org/10.1021/acs.jpca.9b09710>, 2020.
- Wang, H. and Kawamura, K.: Stable carbon isotopic composition of low-molecular-weight dicarboxylic acids and ketoacids in remote marine aerosols, *J. Geophys. Res.*, 111, D07304, <https://doi.org/10.1029/2005jd006466>, 2006.
- Witkowski, B. and Gierczak, T.: cis-Pinonic acid oxidation by hydroxyl radicals in the aqueous phase under acidic and basic conditions: Kinetics and mechanism, *Environ. Sci. Technol.*, 51, 9765–9773, <https://doi.org/10.1021/acs.est.7b02427>, 2017.
- Xu, L., Yang, Z., Tsona, N. T., Wang, X., George, C., and Du, L.: Anthropogenic-biogenic interactions at night: Enhanced formation of secondary aerosols and particulate nitrogen- and sulfur-containing organics from beta-pinene oxidation, *Environ. Sci. Technol.*, 55, 7794–7807, <https://doi.org/10.1021/acs.est.0c07879>, 2021.

- Xu, M., Tsona Tchinda, N., Li, J., and Du, L.: Insoluble lipid film mediates transfer of soluble saccharides from the sea to the atmosphere: the role of hydrogen bonding, *Atmos. Chem. Phys.*, 23, 2235–2249, <https://doi.org/10.5194/acp-23-2235-2023>, 2023.
- Xu, W., Ovadnevaite, J., Fossum, K. N., Lin, C. S., Huang, R. J., Ceburnis, D., and O’Dowd, C.: Sea spray as an obscured source for marine cloud nuclei, *Nat. Geosci.*, 15, 282–286, <https://doi.org/10.1038/s41561-022-00917-2>, 2022.
- Yang, J., Zhao, W., Wei, L., Zhang, Q., Zhao, Y., Hu, W., Wu, L., Li, X., Pavuluri, C. M., Pan, X., Sun, Y., Wang, Z., Liu, C.-Q., Kawamura, K., and Fu, P.: Molecular and spatial distributions of dicarboxylic acids, oxocarboxylic acids, and  $\alpha$ -dicarbonyls in marine aerosols from the South China Sea to the eastern Indian Ocean, *Atmos. Chem. Phys.*, 20, 6841–6860, <https://doi.org/10.5194/acp-20-6841-2020>, 2020.
- Zangrando, R., Corami, F., Barbaro, E., Grosso, A., Barbante, C., Turetta, C., Capodaglio, G., and Gambaro, A.: Free phenolic compounds in waters of the Ross Sea, *Sci. Total Environ.*, 650, 2117–2128, <https://doi.org/10.1016/j.scitotenv.2018.09.360>, 2019.
- Zhan, Y., Li, J., Tsona, N. T., Chen, B., Yan, C., George, C., and Du, L.: Seasonal variation of water-soluble brown carbon in Qingdao, China: Impacts from marine and terrestrial emissions, *Environ. Res.*, 212, 113144, <https://doi.org/10.1016/j.envres.2022.113144>, 2022.
- Zhao, X., Qiu, W., Zheng, Y., Xiong, J., Gao, C., and Hu, S.: Occurrence, distribution, bioaccumulation, and ecological risk of bisphenol analogues, parabens and their metabolites in the Pearl River Estuary, South China, *Ecotox. Environ. Safe.*, 180, 43–52, <https://doi.org/10.1016/j.ecoenv.2019.04.083>, 2019.
- Zhu, M., Jiang, B., Li, S., Yu, Q., Yu, X., Zhang, Y., Bi, X., Yu, J., George, C., Yu, Z., and Wang, X.: Organosulfur compounds formed from heterogeneous reaction between  $\text{SO}_2$  and particulate-bound unsaturated fatty acids in ambient air, *Environ. Sci. Tech. Lett.*, 6, 318–322, <https://doi.org/10.1021/acs.estlett.9b00218>, 2019.
- Zhu, Y., Tilgner, A., Hoffmann, E. H., Herrmann, H., Kawamura, K., Xue, L., Yang, L., and Wang, W.: Molecular distributions of dicarboxylic acids, oxocarboxylic acids, and  $\alpha$ -dicarbonyls in aerosols over Tuoji Island in the Bohai Sea: Effects of East Asian continental outflow, *Atmos. Res.*, 272, 106154, <https://doi.org/10.1016/j.atmosres.2022.106154>, 2022.