



Supplement of

Surface snow bromide and nitrate at Eureka, Canada, in early spring and implications for polar boundary layer chemistry

Xin Yang et al.

Correspondence to: Xin Yang (xinyang55@bas.ac.uk)

The copyright of individual parts of the supplement might differ from the article licence.

Table S1. Overview of relative standard deviations of duplicate analyses for 2019 samples (n=36), and average limits of detection (LOD)^a and limits of quantification (LOQ)^b. The units are μM .

	MSA	Cl^-	Br^-	NO_3^-	$\text{SO}_4^{=}$	Na^+	K^+	Mg^{2+}
Relative Standard deviation	-	0.034	0.023	0.037	0.011	0.109	0.263	0.391
LOD ^a (μM)	0.070	0.367	0.200 ^c	0.484 ^d	0.531	0.696	1.228	0.741
LOQ ^b (μM)	0.260	1.185	0.238 ^c	0.484 ^d	1.780	2.305	4.092	2.510

5 ^a LOD = 3x standard deviation of filter blank average peak area. ^b LOQ = 10x standard deviation of filter blank average peak area. ^c and ^d Br^- and NO_3^- did not have filter blank background and therefore LOD and LOQ were defined as 3x and 10x the minimum measurable peak area, respectively.

Table S2. Statistical results for the ionic analysis of the 2018 snow samples at the Sea ice, Onshore, PEARL, and Creek sites.

10 The units are μM .

2018									
	0-0.5cm				Column average				
	n	mean	std	median	n	mean	std	median	
Sea ice	[Br ⁻]	36	0.23	0.10	0.24	80	10.74	8.52	7.63
	[Cl ⁻]	30	12.55	6.15	12.66	80	5822.27	5166.40	3131.57
	[Ca ²⁺]	33	2.42	2.62	1.20	80	63.43	51.40	28.23
	[F ⁻]	30	0.09	0.03	0.10	73	1.16	0.40	1.21
	[K ⁺]	30	1.64	0.04	1.62	80	55.35	41.15	28.47
	[MSA]	26	0.07	0.05	0.06	2	0.32	0.04	0.32
	[Acetate]	34	15.16	9.92	16.26	24	91.49	9.74	89.94
	[Formate]	26	6.02	0.08	5.98	28	4.90	2.36	4.04
	[NH ₄ ⁺]	34	0.43	0.24	0.38	53	5.11	3.88	3.96
	[Oxalate]	34	3.69	2.54	4.40	41	8.24	7.69	2.27
	[Mg ²⁺]	32	1.86	0.77	1.84	81	338.56	259.62	172.65
	[Na ⁺]	32	3.18	2.00	3.29	75	1437.78	722.12	1120.23
	[NO ₃ ⁻]	33	3.17	1.00	3.32	87	4.95	2.75	4.66
	[SO ₄ ²⁻]	34	0.73	0.38	0.57	69	163.62	97.14	176.53
Onshore	nss-[Br ⁻]	36	0.22	0.10	0.23	82	5.69	6.79	3.71
	nss-[SO ₄ ²⁻]	32	0.49	0.26	0.43	70	87.17	115.13	66.67
	[Br ⁻]	33	0.27	0.11	0.30	60	0.33	0.23	0.25
	[Cl ⁻]	27	11.46	6.69	11.41	66	122.82	70.75	106.86
	[Ca ²⁺]	32	6.28	5.23	3.31	66	16.11	7.74	18.14
	[F ⁻]	8	0.05	0.05	0.05	63	0.13	0.06	0.11
	[K ⁺]	33	0.57	0.65	0.23	65	3.40	2.04	3.65
	[MSA]	4	0.13	0.06	0.12	34	0.01	0.01	0.01
	[Acetate]	32	7.26	0.75	7.27	56	1.21	0.31	1.18
	[Formate]	31	1.39	0.27	1.39	58	0.75	0.27	0.71
	[NH ₄ ⁺]	30	1.33	0.35	1.25	57	1.12	0.32	1.08
	[Oxalate]	33	0.19	0.05	0.20	73	0.50	0.34	0.41
	[Mg ²⁺]	26	1.40	1.19	0.96	66	19.22	12.31	18.77
	[Na ⁺]	27	7.17	4.75	7.26	65	79.70	56.94	71.62
PEARL	[NO ₃ ⁻]	34	3.22	1.45	2.83	66	3.91	0.73	3.97
	[SO ₄ ²⁻]	33	1.55	1.12	0.97	64	6.20	1.78	6.35
	nss-[Br ⁻]	33	0.23	0.09	0.24	61	0.18	0.14	0.16
	nss-[SO ₄ ²⁻]	23	0.51	0.36	0.34	65	1.11	3.12	0.97
	[Br ⁻]	8	0.08	0.06	0.05	37	0.03	0.02	0.02
	[Cl ⁻]	8	25.22	20.67	14.22	38	7.70	5.46	7.01
	[Ca ²⁺]	8	1.41	1.48	1.09	42	1.91	0.95	1.65
	[F ⁻]	6	0.06	0.03	0.06	24	0.03	0.02	0.02
	[K ⁺]	7	0.36	0.29	0.25	37	0.19	0.08	0.20
	[MSA]	6	0.01	0.00	0.01	37	0.18	0.07	0.16
	[Acetate]	7	5.90	0.16	5.84	36	1.53	1.39	1.06
	[Formate]	8	6.34	3.15	8.03	39	0.51	0.10	0.56
	[NH ₄ ⁺]	2	1.50	0.54	1.50	43	0.72	0.60	0.47
	[Oxalate]	8	0.91	0.46	1.16	37	0.08	0.03	0.09
	[Mg ²⁺]	7	1.13	1.77	0.36	38	0.99	0.62	0.79
	[Na ⁺]	7	5.92	13.78	0.65	37	6.69	4.93	5.68
Creek	[NO ₃ ⁻]	8	2.07	0.66	1.86	37	0.71	0.26	0.66
	[SO ₄ ²⁻]	8	0.94	0.40	0.87	41	1.75	0.89	1.64
	nss-[Br ⁻]	8	0.06	0.03	0.05	36	0.01	0.01	0.01
	nss-[SO ₄ ²⁻]	7	0.50	0.54	0.70	44	1.06	0.78	1.01
	[Br ⁻]	12	0.06	0.01	0.06	27	0.63	0.58	0.41
	[Cl ⁻]	12	7.53	3.22	6.82	27	0.93	104.62	162.47
	[Ca ²⁺]	13	6.26	4.26	5.43	22	9.16	6.93	8.31
	[F ⁻]	12	0.03	0.02	0.03	27	0.25	0.16	0.21
	[K ⁺]	11	0.13	0.05	0.11	22	0.56	0.40	0.47
	[MSA]	10	0.24	0.14	0.28	18	0.01	0.01	0.01
	[Acetate]	12	6.59	0.71	6.45	23	5.87	0.05	5.86
	[Formate]	13	1.34	0.54	1.22	27	5.75	3.31	8.03
	[NH ₄ ⁺]	13	1.67	0.57	1.55	20	1.10	1.30	0.14
	[Oxalate]	13	0.17	0.02	0.16	27	0.97	0.41	1.17
	[Mg ²⁺]	13	1.44	1.12	1.23	23	5.14	5.06	3.96
	[Na ⁺]	12	5.22	2.32	4.55	23	15.76	13.32	14.08
	[NO ₃ ⁻]	13	1.45	0.52	1.46	25	4.15	0.75	4.03
	[SO ₄ ²⁻]	13	1.28	0.27	1.37	26	60.48	36.41	60.95
	nss-[Br ⁻]	11	0.05	0.01	0.05	27	0.56	0.54	0.39
	nss-[SO ₄ ²⁻]	12	0.97	0.19	0.96	26	58.02	35.58	54.43

Table S3. Statistical results for the ionic analysis of the 2019 snow samples at the Sea ice, Onshore, PEARL, and 0PAL sites. Note that the airborne samples were collected by the mounted tray above the ground (~1 m at 0PAL and 11 m at PEARL). The units are μM .

2019												Column average								
	Airborne			0-0.2cm			0.2-0.5cm			0.5-1.5cm			n	mean	std	median	n	mean	std	median
Sea ice	[Br ⁻]			40	0.40	0.20	0.37	51	3.03	4.14	0.84	36	9.31	5.14	10.72	66	6.47	5.36	5.79	
	[Cl ⁻]			41	61.25	68.76	25.06	33	180.72	244.14	64.42	16	2897.67	2472.62	2718.74	33	1660.12	2080.86	510.54	
	[Mg ²⁺]			36	9.68	12.49	4.46	43	168.88	213.56	57.26	17	264.31	221.14	392.24	46	202.59	196.83	78.13	
	[Na ⁺]			38	29.52	32.25	14.98	34	149.80	197.82	43.99	12	1189.99	14170.62	320.53	19	390.39	304.56	382.34	
	[NO ₃ ⁻]			37	3.46	1.55	3.50	61	4.83	2.32	5.15	61	2.90	1.18	2.88	75	2.67	1.18	2.56	
	[SO ₄ ²⁻]			41	4.54	2.96	4.74	42	29.51	43.19	9.23	21	122.89	96.44	99.36	40	58.89	62.52	25.56	
	nss-[Br ⁻]			39	0.35	0.20	0.34	32	0.24	0.19	0.19	8	0.58	0.77	0.20	17	0.22	0.18	0.14	
	nss-[SO ₄ ²⁻]			39	2.52	1.53	2.89	28	1.85	2.73	1.24	8	3.12	3.01	2.42	19	1.54	2.67	0.69	
	[Cl ⁻]			38	0.40	0.17	0.37	58	0.38	0.22	0.31	64	0.97	0.86	0.53	104	0.99	0.93	0.49	
	[Mg ²⁺]			31	1.79	1.95	0.70	56	31.53	43.17	8.80	62	51.75	52.64	9.76	102	52.30	51.99	21.94	
Onshore	[Na ⁺]			39	17.06	15.32	10.20	46	37.60	32.62	27.57	56	278.79	343.11	52.32	93	297.66	336.14	70.87	
	[NO ₃ ⁻]			36	3.76	1.99	3.58	61	3.84	2.18	3.67	63	3.41	1.52	4.16	101	3.06	1.53	3.19	
	[SO ₄ ²⁻]			42	4.11	2.41	4.38	51	6.06	4.86	5.01	64	14.67	11.49	8.61	104	14.98	11.47	11.49	
	nss-[Br ⁻]			38	0.36	0.19	0.34	50	0.21	0.17	0.20	56	0.35	0.32	0.21	89	0.30	0.31	0.17	
	nss-[SO ₄ ²⁻]			42	2.56	2.21	2.84	46	2.40	1.90	1.90	50	-1.36	10.17	1.47	91	-4.65	13.11	1.22	
	[Br ⁻]			31	0.35	0.18	0.39	31	0.30	0.12	0.31	32	0.22	0.10	0.17	28	0.19	0.05	0.18	
	[Cl ⁻]			19	26.82	12.14	22.87	33	17.71	6.54	18.59	57	120.88	115.60	47.27	58	127.37	88.05	52.73	
	[Mg ²⁺]			19	2.06	1.92	1.11	32	1.04	0.67	0.84	56	11.23	14.65	3.84	58	13.39	11.26	5.37	
	[Na ⁺]			19	19.86	9.75	17.65	34	11.80	5.20	10.84	56	85.49	79.83	36.53	58	93.06	63.96	43.03	
	[NO ₃ ⁻]			18	2.23	1.37	2.29	23	1.24	0.50	1.13	51	1.29	0.63	1.00	58	1.42	0.63	1.15	
PEARL	[SO ₄ ²⁻]			20	5.51	3.55	5.92	35	3.82	2.16	4.04	58	7.74	4.44	7.65	57	5.74	3.17	4.41	
	nss-[Br ⁻]			21	0.28	0.20	0.23	31	0.28	0.12	0.29	33	0.00	0.13	-0.03	31	-0.05	0.08	-0.07	
	nss-[SO ₄ ²⁻]			22	4.63	4.24	4.55	36	3.19	2.20	3.46	56	2.32	2.24	2.05	56	0.01	1.53	-0.10	
	[Br ⁻]			42	0.38	0.24	0.34	5	0.15	0.01	0.15	9	0.30	0.15	0.28	5	0.35	0.07	0.39	
	[Cl ⁻]			42	57.33	55.82	35.20	6	44.90	47.59	27.38	9	86.66	53.97	108.75	5	99.09	49.44	69.82	
	[Mg ²⁺]			38	6.13	6.48	4.00	3	9.11	5.68	7.80	9	9.77	7.56	13.16	5	11.80	7.26	7.71	
	[Na ⁺]			41	36.99	38.42	23.25	6	31.33	34.37	19.07	9	62.31	39.36	80.94	5	70.64	34.89	50.67	
OPAL	[NO ₃ ⁻]			40	3.41	2.05	2.81	3	0.96	0.21	0.91	8	2.16	1.07	1.82	5	2.02	0.23	2.07	
	[SO ₄ ²⁻]			45	5.47	3.61	4.31	6	2.45	1.95	2.00	9	5.06	3.69	5.58	5	6.14	0.52	5.98	
	nss-[Br ⁻]			40	0.24	0.19	0.18	6	0.10	0.07	0.14	9	0.18	0.11	0.12	5	0.22	0.12	0.31	
	nss-[SO ₄ ²⁻]			44	2.17	2.37	2.46	4	0.56	0.10	0.59	9	1.32	1.90	0.28	5	1.90	1.75	3.08	

Table S4. Geographic heterogeneity of snow ions (Na^+ , NO_3^- , Br^- , and $\text{nss}[\text{Br}]$ in units of μM) at each sampling site with snow samples randomly collected from a small area ($2\text{m} \times 2\text{m}$) during two periods in 2019.

Period	Sites	Depth (cm)	$\text{Na}^+(\mu\text{M})$		$\text{NO}_3^-(\mu\text{M})$		$\text{Br}^-(\mu\text{M})$		$\text{nss-Br}^-(\mu\text{M})$	
			mean	std	mean	std	mean	std	mean	std
Feb 26- Mar 3, 2019	Sea ice	0.0-0.5	129.33	148.65	4.27	1.74	0.67	0.74	0.43	0.48
		0.5-1.5	336.12	316.92	2.96	0.94	7.67	5.51	0.98	0.97
	Onshore	0.0-0.5	26.13	21.04	2.27	0.71	0.30	0.13	0.25	0.13
		0.5-1.5	581.53	315.89	3.62	1.01	3.87	3.76	0.67	0.29
	PEARL	0.0-0.5	185.09	65.44	2.06	0.47	0.28	0.14	-0.05	0.07
		0.5-1.5	150.87	30.09	2.00	0.38	0.25	0.10	-0.02	0.08
Mar 4-5, 2019	Sea ice	0-0.2	-	-	1.98	0.34	0.38	0.04	-	-
		0.2-0.5	520.98	27.56	3.22	1.50	5.73	5.57	0.01	0.04
		0.5-1.5	-	-	1.77	0.19	12.91	-	-	-
	Onshore	0-0.2	32.39	0.64	-	-	0.12	0.00	0.06	0.00
		0.2-0.5	75.56	14.85	1.59	0.17	0.22	0.04	0.09	0.03
		0.5-1.5	49.31	9.41	1.92	0.58	0.23	0.02	0.14	0.03
	OPAL	0-0.2	71.22	25.51	1.19	0.00	0.14	0.02	0.01	0.03
		0.2-0.5	104.82	2.44	2.73	1.23	2.42	2.54	0.10	0.00
		0.5-1.5	-	-	-	-	-	-	-	-

20 Table S5. Regression analyses (i.e., number of observations, standard error of slope, and *p*-values) for key linear models used in this study and associated with Figures 5, 6, and 7.

Figures	linear regressions	Sites	Number of observations	Standard error of slope (μM)	<i>p</i> -value
Figure 5	(d) $[\text{Na}^+]$ vs time	sea ice	11	0.65117	0.17666
		Onshore	11	1.3248	0.018913
		PEARL	2	-	-
	(e) $[\text{NO}_3^-]$ vs time	sea ice	11	0.031625	3.37E-05
		Onshore	11	0.07233	0.047092
		PEARL	2	-	-
	(f) $[\text{Br}^-]$ vs time	sea ice	11	0.0059219	0.31536
		Onshore	11	0.0058544	0.21431
		PEARL	2	0	-
	(g) nss-[Br^-]	sea ice	11	0.0054474	0.42219
		Onshore	11	0.0050932	0.8453
		PEARL	2	-	-
Figure 6	(d) Tray $[\text{Na}^+]$ vs time	PEARL	14	1.6482	0.19785
		OPAL	24	1.1562	0.47351
	(e) Tray $[\text{NO}_3^-]$ vs time	PEARL	12	0.06437	0.35704
		OPAL	24	0.07293	0.024153
	(f) Tray $[\text{Br}^-]$ vs time	PEARL	14	0.0038572	0.00091149
		OPAL	24	0.0057058	0.0010769
	(g) Tray nss-[Br^-] vs time	PEARL	14	0.0056266	0.04216
		OPAL	24	0.0058292	0.00069418
Figure 7	(a) 0-0.2cm $[\text{NO}_3^-]$ vs time	Sea ice	21	0.10137	0.022069
		Onshore	20	0.12408	0.033649
		PEARL	13	0.033592	0.63919
	(b) 0.2-0.5cm $[\text{NO}_3^-]$ vs time	Sea ice	22	0.053913	0.00030065
		Onshore	21	0.062582	0.016509
		PEARL	18	0.013772	0.24503
	(c) 0.5-1.5cm $[\text{NO}_3^-]$ vs time	Sea ice	29	0.024566	0.027422
		Onshore	27	0.027371	0.0071189
		PEARL	26	0.0081262	7.64E-07
	(d) 0-0.2cm nss-[Br^-] vs time	Sea ice	21	0.0096136	0.019895


		Onshore	20	0.0080688	0.010627
		PEARL	17	0.0035759	0.0048613
(e) 0.2-0.5cm nss-[Br-] vs time	Sea ice		12	0.012075	0.17789
	Onshore		17	0.0088479	0.13183
	PEARL		10	0.0069285	0.035824
	Sea ice		4	0.05373	0.1602
(f) 0.5-1.5cm nss-[Br-] vs time	Onshore		23	0.0070587	0.63092
	PEARL		13	0.0048105	0.9346

25

Table S6. Nitrate and bromide analysis between morning and afternoon samples (i.e., mean (μM), one standard deviation (μM), number of observations, and p -values) for Figure 8.

		Morning				Afternoon				
		Mean	STD	N	p-value	Mean	STD	N	p-value	
[NO ₃] (μM)	Tray	PEARL	1.50	0.95	4	0.050	2.20	1.25	9	0.001
		OPAL	1.54	0.69	5	0.007	3.27	1.80	8	0.001
	0-0.2cm	Sea ice	2.83	1.27	14	1.48E-06	2.52	0.82	8	5.43E-05
		Onshore	2.48	1.34	12	5.23E-05	2.72	0.88	8	5.26E-05
	0.2-0.5cm	PEARL	1.67	0.77	4	0.023	0.63	0.10	2	0.072
		Sea ice	3.91	2.09	14	9.27E-06	3.40	2.36	10	0.001
		Onshore	3.55	1.92	14	1.05E-05	3.28	2.45	12	0.001
		PEARL	1.18	0.60	11	7.29E-05	1.11	0.58	5	0.013
	0.5-1.5cm	Sea ice	2.34	1.06	16	2.49E-07	2.79	0.97	10	7.90E-06
		Onshore	3.03	1.66	13	2.63E-05	2.26	1.72	14	2.82E-04
		PEARL	1.51	0.64	17	3.72E-08	1.84	0.74	5	0.005
[nss-Br] (μM)	Tray	PEARL	0.28	0.18	8	0.003	0.30	0.24	9	0.005
		OPAL	0.04	0.09	6	0.340	0.22	0.13	10	4.55E-04
	0-0.2cm	Sea ice	0.23	0.11	13	5.88E-06	0.24	0.25	8	0.029
		Onshore	0.25	0.12	12	1.31E-05	0.23	0.21	8	0.019
	0.2-0.5cm	PEARL	0.24	0.14	12	6.56E-05	0.15	0.03	4	0.001
		Sea ice	0.18	0.03	10	1.73E-08	0.13	0.09	6	0.022
		Onshore	0.01	0.27	14	0.852	0.06	0.22	11	0.404
		PEARL	0.01	0.13	10	0.783	0.04	0.11	5	0.400
	0.5-1.5cm	Sea ice	0.18	0.03	4	0.002	NaN	NaN	NaN	NaN
		Onshore	0.18	0.22	16	0.005	0.22	0.26	12	0.013
		PEARL	-0.08	0.08	11	0.008	-0.05	0.11	5	0.329

30



35






Figure S1. UV index at Eureka, Canada (79.989N, 85.934W, 8.7 m) during (a) February 24 - March 31, 2018 and (b) February 24 - March 31, 2019.



40

Figure S2. Snow density profiles measured in 2018 at the Sea ice, Onshore, and PEARL sampling sites (see Figure 1).



45 Figure S3. Snow salinity profiles in 2018 and 2019 samples. Top 0.5 cm snow sample mean salinities (collected in 2018) are plotted at a depth of 0.25 cm. At each site, two types of surface snow are collected: one is soft fluffy white colour snow and one is hard, light brown colour snow. The horizontal error bar represents one standard deviation value. 2019 sea ice profile includes only one column collected on Feb 27 at Sea ice site. 2018 inland profile includes two columns, one from Onshore and one from Creek site. The bottom interface snow-ice sample was collected on March 1, 2018 at Sea ice site.

50




Figure S4. Vertical profiles of 2019 ions (a) $[Ca^{2+}]$, (b) $[Mg^{2+}]$, (c) $[K^+]$, (d) $[SO_4^{2-}]$, and enrichments of (e) $[Ca^{2+}]$, (f) $[Mg^{2+}]$, and (g) $[K^+]$ (see Section 3.2 for details).

55





Figure S5. Vertical profiles of 2018 ions (a) $[\text{Na}^+]$, (b) $[\text{NO}_3^-]$, (c) $[\text{Br}^-]$, and (d) nss $[\text{Br}^-]$.



65 Figure S6. 2019 BrO column timeseries from March 5th to May 31st, in which linear regression fit and exponential fit are
inserted)




Figure S7. Scatter plot of surface snow bromide versus sodium in (a) 0–0.5 cm layer snow (2018), (b) 0–0.2 cm layer (2019), (c) 0.2–0.5 cm layer (2019), and (d) 0.5–1.5 cm layer snow. Linear regressions and corresponding correlation coefficients R

70 are given.




Figure S8. Same as Figure 7, except here nss[Br $^-$] is used in the scatter plot and relationship analysis.