



Stratospheric $\delta^{13}\text{CO}_2$ observed over Japan and its governing processes

Satoshi Sugawara¹, Shinji Morimoto², Shigeyuki Ishidoya³, Taku Umezawa^{4,2}, Shuji Aoki², Takakiyo Nakazawa², Sakae Toyoda⁵, Kentaro Ishijima⁶, Daisuke Goto⁷, and Hideyuki Honda²

¹Faculty of Education, Miyagi University of Education, Sendai 980-0845, Japan

²Center for Atmospheric and Oceanic Studies, Tohoku University, Sendai 980-8578, Japan

³National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba 305-8569, Japan

⁴National Institute for Environmental Studies, Tsukuba 305-8506, Japan

⁵School of Materials and Chemical Technology, Institute of Science Tokyo, Yokohama 226-8501, Japan

⁶Meteorological Research Institute, Japan Meteorological Agency, Tsukuba 305-0052, Japan

⁷National Institute of Polar Research, Tachikawa 190-8518, Japan

Correspondence: Satoshi Sugawara (sugawara@staff.miyakyo-u.ac.jp)

Received: 3 March 2025 – Discussion started: 4 April 2025

Revised: 16 July 2025 – Accepted: 18 July 2025 – Published: 2 October 2025

Abstract. Due to very few reports of $\delta^{13}\text{CO}_2$ (the stable carbon isotopic ratio of CO_2) observations in the stratosphere, its variations are not well understood. In order to elucidate stratospheric $\delta^{13}\text{CO}_2$ variations and their governing mechanisms, we have collected stratospheric air samples using balloon-borne cryogenic samplers over Japan since 1985 and analyzed them for $\delta^{13}\text{CO}_2$. To obtain precise $\delta^{13}\text{CO}_2$ values, we incorporated the mass-independent fractionation of ^{17}O and ^{18}O in the $\delta^{13}\text{CO}_2$ calculation. $\delta^{13}\text{CO}_2$ has decreased through time in the mid-stratosphere with an average rate of change of $-0.026 \pm 0.001 \text{‰ yr}^{-1}$ for the period 1985–2020, consistent with that in the troposphere. However, mid-stratospheric $\delta^{13}\text{CO}_2$ values did not show a time delay compared to the tropical tropospheric values. This could be explained by the production of CO_2 by CH_4 oxidation and the gravitational separation of $^{13}\text{CO}_2$ and $^{12}\text{CO}_2$. To confirm this hypothesis, we used a two-dimensional model to simulate the stratospheric $\delta^{13}\text{CO}_2$ values while accounting for these processes. The results indicate that these two effects strongly impact the vertical distribution of $\delta^{13}\text{CO}_2$. We newly defined “stratospheric potential $\delta^{13}\text{C}^*$ ” ($\delta^{13}\text{C}_\text{p}$) as a quasi-conservative parameter incorporating the kinetic isotope effect of CH_4 oxidation and gravitational separation, and we found that $\delta^{13}\text{C}_\text{p}$ in the mid-latitude mid-stratosphere decreases over time with about a 5 yr lag relative to the tropical upper troposphere. This fact strongly supports that stratospheric $\delta^{13}\text{CO}_2$ variations are governed by the airborne production of ^{13}C -depleted CO_2 by CH_4 oxidation, the gravitational separation, and the propagation of the decreasing tropospheric $\delta^{13}\text{CO}_2$ trend into the stratosphere.

1 Introduction

The isotopic ratio of carbon in CO_2 (here, we use “ $\delta^{13}\text{CO}_2$ ” to distinguish it from that in CH_4 , “ $\delta^{13}\text{CH}_4$ ”) is thought to provide information on the carbon cycle, and observations have been carried out mainly using ground stations, ships, and aircraft in the troposphere (e.g., Keeling et al., 1995; Francey et al., 1995; Nakazawa et al., 1993; Morimoto et al., 2000). However, there have been very few reports of $\delta^{13}\text{CO}_2$

observations in the mid-stratosphere, mainly because of the difficulty in collecting air samples at such altitudes; the only available reports are balloon observations over Japan and Scandinavia (Gamo et al., 1995; Aoki et al., 2003). It is well known that tropospheric $\delta^{13}\text{CO}_2$ values have decreased with time due to anthropogenic emissions with low $\delta^{13}\text{CO}_2$ values resulting from fossil fuel consumption. Gamo et al. (1995) reported that $\delta^{13}\text{CO}_2$ values observed in the mid-stratosphere were higher than those in the troposphere over the same pe-

riod, and that the average stratospheric $\delta^{13}\text{CO}_2$ value during 1986–1992 was about -7.5‰ and decreasing with time. Aoki et al. (2003) reported that the mid-stratosphere (25–32 km altitude) $\delta^{13}\text{CO}_2$ value over Japan in 1997 was approximately -7.7‰ ; this value is consistent with the expected value extrapolated from the results of Gamo et al. (1995), suggesting that their estimated secular decreasing trend is plausible.

Anomalous ^{18}O enrichment in stratospheric CO_2 was first reported in balloon observations (Gamo et al., 1989), which spurred extended studies of the mass-independent isotopic effect (MIE) on the triple oxygen isotope system (e.g., Thiemens, 1999; Kawagucci et al., 2008). The isotopic fractionation is usually caused by the effect of mass differences between isotopologues (i.e., the mass-dependent isotopic effect). In this case, the fractionation between ^{18}O and ^{16}O is almost twice as large as that between ^{17}O and ^{16}O . However, the isotopic fractionation that does not follow this relation occurs mainly in photochemical processes. These MIE studies have clarified that the relationship between ^{17}O and ^{18}O enrichment in stratospheric CO_2 is quite different from that in the troposphere. This fact is very important for stratospheric $\delta^{13}\text{CO}_2$ measurements because mass spectrometry measures the ratios $^{45}\text{CO}_2/^{44}\text{CO}_2$ and $^{46}\text{CO}_2/^{44}\text{CO}_2$, and an assumed relationship between ^{17}O and ^{18}O enrichment is needed to calculate $\delta^{13}\text{CO}_2$. The influence of the MIE on oxygen in stratospheric $\delta^{13}\text{CO}_2$ has not been taken into account in previous studies. Although Aoki et al. (2003) found that $\delta^{13}\text{CO}_2$ values increase with increasing altitude in the stratosphere over Japan and suggested that the MIE on ^{17}O and ^{18}O should be considered when interpreting such a vertical distribution, no quantitative investigation of these factors has been conducted.

Stratospheric methane (CH_4) is destroyed by reactions with hydroxyl (OH), excited singlet oxygen ($\text{O}(^1\text{D})$), and chlorine (Cl) radicals, and thus plays an important role in stratospheric chemistry. However, the quantitative contributions of these chemical processes to CH_4 loss and their temporal variations remain poorly understood. It is thought that $\delta^{13}\text{CH}_4$ provides useful information about CH_4 destruction in the stratosphere because characteristic isotopic fractionations occur in each reaction (e.g., Saueressig et al., 1995; Sugawara et al., 1997; Rice et al., 2003; Röckmann et al., 2011). However, only a few measurements have been made so far in the mid-stratosphere. Oxidation of CH_4 produces CO , which is eventually oxidized to CO_2 . In general, the isotopic fractionations in the above reactions cause the faster destruction of $^{12}\text{CH}_4$ relative to $^{13}\text{CH}_4$, making the chemical products depleted in ^{13}C (decreased $\delta^{13}\text{C}$) while the remaining CH_4 becomes enriched in ^{13}C (increased $\delta^{13}\text{CH}_4$). Indeed, Brenninkmeijer et al. (1996) found that CO in the lower Antarctic stratosphere is extremely depleted in ^{13}C (decreased $\delta^{13}\text{CO}$) mainly because of the local production of CO via the reaction of CH_4 with Cl . Recently, Gromov et al. (2018) and Röckmann et al. (2024) have discussed $\delta^{13}\text{CO}$

variations associated with the $\text{CH}_4 + \text{Cl}$ reaction. From these studies, it is expected that the $\text{CH}_4\text{--CO--CO}_2$ chain reaction could be an airborne source of CO_2 with extremely low $\delta^{13}\text{C}$ values, depressing stratospheric $\delta^{13}\text{CO}_2$ values. However, no study has yet quantitatively assessed this effect.

The age of stratospheric air is the transit time of air from around the tropical tropopause to a certain location in the stratosphere, which is a powerful tool for diagnosing stratospheric transport (see Garny et al., 2024 for a review of the latest developments in air age studies). It is expected that possible changes to the Brewer–Dobson circulation induced by climate change are detectable as long-term changes in the mean age of air. For this purpose, age tracers, such as mole fractions of CO_2 and SF_6 , have been measured in air samples collected by scientific balloons and satellites (e.g., Engel et al., 2009, 2017; Stiller et al., 2012). CO_2 mole fractions observed in the mid-stratosphere over Japan have been combined with SF_6 data to provide the longest record of the mean age of air, as reported by Engel et al. (2009, 2017) and Ray et al. (2014). Engel et al. (2017) reported that the mean age of air in the mid-stratosphere does not indicate a significant trend over the period 1975–2016, with only a slightly positive increase of $+0.15 \pm 0.18 \text{ yr decade}^{-1}$. However, recent models predict accelerating circulation throughout the stratosphere (e.g., Butchart, 2014), which will decrease the mean age of air. Such a decreasing trend is consistent with observational results in the lower stratosphere (Ray et al., 2014). Despite many efforts to elucidate the trend of age of air, a discrepancy remains between mid-stratosphere age trends in models and observations. One reason for this discrepancy is the large uncertainties on the observational trends. To resolve this problem, it is necessary to increase the frequency of high-altitude balloon observations and observe additional age tracers beyond CO_2 and SF_6 . Perfluorocarbons and hydrofluorocarbons with extremely long lifetimes could be alternative age tracers (Leedham Elvidge et al., 2018; Laube et al., 2025; Umezawa et al., 2025), and observations of such multiple age tracers should reduce the uncertainties on mean age of air estimates.

The gravitational separation (hereafter, “GS”) of major atmospheric constituents in the stratosphere was first reported from balloon observations (Ishidoya et al., 2006, 2008a, b, 2013). They showed that vertical gradients of the isotopic and elemental ratios of major atmospheric components (N_2 , O_2 , and Ar) in the stratosphere are caused by molecular diffusion and depend on molecular mass. Stratospheric GS has now been observed in polar and equatorial regions (Ishidoya et al., 2018; Sugawara et al., 2018) and reproduced in numerical models (Belikov et al., 2019; Birner et al., 2020). Alongside age of air, stratospheric GS can be used to diagnose stratospheric transport processes (Ishidoya et al., 2013; Sugawara et al., 2018; Birner et al., 2020). These findings also suggest that GS could affect the mole fractions and isotopic ratios of all stratospheric constituents, although no studies have addressed the impact of GS on stratospheric $\delta^{13}\text{CO}_2$ values.

The main objective of this study is to clarify the mechanisms governing $\delta^{13}\text{CO}_2$ variations in the stratosphere and understand the long-term variations. A comprehensive understanding of the variations of stratospheric $\delta^{13}\text{CO}_2$ requires that the MIE on oxygen, CH_4 oxidation and associated isotopic fractionation, mean age of air, and GS be considered. Balloon observations provide a unique opportunity to evaluate these effects by studying a variety of atmospheric constituents using high-precision measurements. In this study, we constructed a new $\delta^{13}\text{CO}_2$ record for the stratosphere over Japan spanning the 35 yr from 1985 to 2020, primarily by newly analyzing archived CO_2 samples and in part by including previously published data (Gamo et al., 1995; Aoki et al., 2003). We then examined vertical and temporal variations in the record. Furthermore, we investigated in detail the various mechanisms affecting the vertical distribution of stratospheric $\delta^{13}\text{CO}_2$ and propose herein a new concept called “stratospheric potential $\delta^{13}\text{C}$.” We also investigated the possibility of estimating the age of stratospheric air from the $\delta^{13}\text{CO}_2$ record.

2 Experimental procedures

2.1 Air sample collection

We have collected stratospheric air samples over Japan since 1985 using balloon-borne cryogenic samplers (e.g., Nakazawa et al., 1995). A large scientific balloon equipped with a sampler was launched nearly once a year during 1985–2002 from the Sanriku Balloon Center (SBC; $39^\circ 10' \text{N}$, $141^\circ 50' \text{E}$). The frequency of observations has decreased since 2004, with the longest interval between observations being 5 yr. During this period, the balloon launch site was relocated from SBC to the Taiki Aerospace Research Field in Hokkaido (TARF; $42^\circ 30' \text{N}$ $143^\circ 26' \text{E}$) in 2008. Balloon observations conducted so far over Japan are summarized in Table S1 of the Supplement. The cryogenic air sampler consists mainly of a liquid helium dewar, stainless steel bottles, motor-driven valves, and a control unit (Honda et al., 1996). Liquid helium was used as a refrigerant for collecting stratospheric air cryogenically under low atmospheric pressures, and 20–30 L STP of stratospheric air was collected in each bottle, depending on the altitude. Air sampling was performed typically at 11 different altitudes, spaced about 2 km apart, between 14 and 35 km altitude. Occasionally, anomalously low CO_2 mole fractions were observed, probably due to sample deterioration in the bottle. In such cases, both CO_2 mole fraction and $\delta^{13}\text{CO}_2$ data were excluded from the data record. In 2020, the CO_2 mole fraction and $\delta^{13}\text{CO}_2$ value at 28.8 and 30.9 km altitude were not available due to water contamination of the air samples.

2.2 Analytical methods for mole fractions and isotopic ratios

Collected air samples were distributed to several institutes in Japan for analysis of the mole fractions and isotopic compositions of various gases. The CO_2 mole fraction was measured using a nondispersive infrared gas analyzer at Tohoku University (TU) with an analytical precision of $<0.02 \mu\text{mol mol}^{-1}$; the CO_2 measurement protocol has been described in detail in previous studies (Nakazawa et al., 1995; Aoki et al., 2003; Sugawara et al., 2018). N_2O and CH_4 mole fractions were measured using gas chromatographs equipped with an electron capture detector and flame ionization detector, respectively, at TU. The N_2O mole fraction was used both for mass spectrometry analyses to determine $\delta^{13}\text{CO}_2$ and to investigate the MIE between ^{17}O and ^{18}O in CO_2 , as described below. The analytical precisions for the N_2O and CH_4 mole fractions were 0.3 and 3 nmol mol^{-1} , respectively (Nakazawa et al., 2002; Aoki et al., 2003; Ishijima et al., 2007).

The analytical procedures for $\delta^{13}\text{CO}_2$ in air samples collected after 1988 have already been described in previous studies (Nakazawa et al., 1993, 1997a; Morimoto et al., 2000), and is briefly summarized here. An aliquot of each air sample (about 550 mL STP) was introduced into a CO_2 extraction system, and the pure extracted CO_2 was divided and sealed in small glass tubes; some tubes were archived for future study. The carbon and oxygen isotopic ratios ($\delta^{13}\text{C}$ and $\delta^{18}\text{O}$) of the sample CO_2 were measured using a mass spectrometer (Finnigan MAT- δS) installed at TU against an in-house reference CO_2 calibrated to Vienna Pee Dee Belemnite (VPDB). The analytical precision of $\delta^{13}\text{CO}_2$ was $\pm 0.02 \text{‰}$ (Morimoto et al., 2000). Air samples collected in 1985, 1986, 1988, and 1989 were analyzed for $\delta^{13}\text{CO}_2$ using a mass spectrometer (Finnigan MAT250) with a reproducibility of less than $\pm 0.02 \text{‰}$ at the University of Tokyo (Gamo et al., 1989, 1995).

In this study, we modified the calculation of isotopic ratios to take into account the stratospheric MIE and GS. Hereafter, the isotopic ratios of ^{13}C to ^{12}C , ^{17}O and ^{18}O to ^{16}O , and $^{45}\text{CO}_2$ and $^{46}\text{CO}_2$ to $^{44}\text{CO}_2$ are expressed as $^{13}R = n(^{13}\text{C})/n(^{12}\text{C})$, $^{17}R = n(^{17}\text{O})/n(^{16}\text{O})$, $^{18}R = n(^{18}\text{O})/n(^{16}\text{O})$, $^{45}R = n(^{45}\text{CO}_2)/n(^{44}\text{CO}_2)$, and $^{46}R = n(^{46}\text{CO}_2)/n(^{44}\text{CO}_2)$, respectively, where n is the amount of each isotopologue. $\delta^{13}\text{CO}_2$ is defined as

$$\delta^{13}\text{CO}_2 = \left(\frac{^{13}R}{^{13}R_{\text{VPDB}}} - 1 \right) \times 10^3 (\text{‰}), \quad (1)$$

where $^{13}R_{\text{VPDB}}$ is the known ^{13}R value of VPDB (Coplen and Shrestha, 2016). The isotopic ratios ^{45}R and ^{46}R are expressed as (Santrock et al., 1985)

$$^{45}R = ^{13}R + 2 \cdot ^{17}R, \quad (2a)$$

$$^{46}R = 2 \cdot ^{18}R + 2 \cdot ^{13}R \cdot ^{17}R + (^{17}R)^2 \quad (2b)$$

To calculate $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values from ^{45}R and ^{46}R , the relationship between ^{17}O and ^{18}O should be known, because $^{45}\text{CO}_2$ contains $^{12}\text{C}^{17}\text{O}^{16}\text{O}$, whereas $^{46}\text{CO}_2$ contains both $^{13}\text{C}^{17}\text{O}^{16}\text{O}$ and $^{12}\text{C}^{17}\text{O}_2$. The relationship between ^{17}R and ^{18}R can be expressed as (e.g., Meijer and Li, 1998)

$$\frac{^{17}R}{^{17}R_{\text{VSMOW}}} = \left(\frac{^{18}R}{^{18}R_{\text{VSMOW}}} \right)^\beta, \quad (2c)$$

where $^{17}R_{\text{VSMOW}}$ and $^{18}R_{\text{VSMOW}}$ are the known ratios of Vienna standard mean ocean water (VSMOW; Coplen and Shrestha, 2016). In measurements of tropospheric $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, the mass-dependent relationship between ^{17}O and ^{18}O can be assumed, i.e., $\beta = 0.528$ (Meijer and Li, 1998; Asonov and Brenninkmeijer, 2003). However, the MIE between ^{17}O and ^{18}O is non-negligible in the stratosphere (Gamo et al., 1995; Thiemens, 1999). Therefore, the $\delta^{17}\text{O}$ value is essential for accurately determining the $\delta^{13}\text{C}$ value in the stratosphere. Our air samples were also measured for triple oxygen isotopes, as reported by Kawagucci et al. (2008). However, $\delta^{17}\text{O}$ measurements were not performed for all of our stratospheric samples. In such cases, the following method was applied to obtain $\delta^{17}\text{O}$ values. The values of ^{17}R and ^{18}R calculated from $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ observed by Kawagucci et al. (2008) were fitted to Eq. (2c), and the average value of β was calculated to be 1.7 ± 0.2 using the least squares method. For samples for which $\delta^{17}\text{O}$ was not analyzed, we used Eq. (2c) with this β value to calculate $\delta^{13}\text{C}$. By substituting Eqs. (2a) and (2b) into (2c), we obtained the following equation:

$$3(^{17}R)^2 - 2 \cdot ^{45}R \cdot ^{17}R - 2\gamma (^{17}R)^{\frac{1}{\beta}} + ^{46}R = 0, \quad (3a)$$

where γ is defined as

$$\gamma = ^{18}R_{\text{VSMOW}} \left(^{17}R_{\text{VSMOW}} \right)^{-\frac{1}{\beta}}. \quad (3b)$$

Equation (3a) was solved for ^{17}R numerically using the bisection method and the measured ^{45}R and ^{46}R values, and then ^{13}R and $\delta^{13}\text{C}$ were calculated by Eqs. (2a) and (1), respectively. We note that GS occurs in CO_2 molecules with different mass numbers (44, 45, 46). Therefore, it is not accurate to correct $\delta^{13}\text{C}$ by assuming a mass number difference of uniquely $\Delta m = 1$. To obtain the correct $\delta^{13}\text{C}$ of stratospheric CO_2 , it is necessary to first apply the GS correction to the measured values of ^{45}R and ^{46}R , and then to calculate $\delta^{13}\text{C}$ using the method described above. The method used to correct ^{45}R and ^{46}R for GS is discussed in Sect. 3.3.

The MIE calculation described above was not considered in Gamo et al. (1995) and Aoki et al. (2003), and it is therefore likely that their $\delta^{13}\text{C}$ values are not correct, especially

at higher altitudes. $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ in CO_2 increase rapidly with increasing altitude in the stratosphere (Gamo et al., 1995; Kawagucci et al., 2008), and the deviations from mass-dependent values also increase with altitude, meaning that the overestimation of $\delta^{13}\text{C}$ also increases with altitude. The influence of the oxygen MIE on $\delta^{13}\text{C}$ is almost zero near the tropopause, but at an altitude of about 35 km over Japan, it is approximately 0.6‰. Therefore, this effect significantly impacts the vertical distributions of $\delta^{13}\text{C}$. We reanalyzed some of the CO_2 samples used by Gamo et al. (1995) and Aoki et al. (2003) to determine the differences between their results (not accounting for the oxygen MIE) and ours to allow us to correct their data and add them to our long-term record. The MIE on $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ occurs through photochemical reactions in O_2 , O_3 , and CO_2 (e.g., Thiemens, 1999), which are relatively tightly correlated with the mole fraction of N_2O , which is also destroyed through photochemical processes. Figure 1 shows the relationships between the N_2O mole fraction and $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ in CO_2 in the stratosphere over Japan. These results show that $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ values calculated using $\beta = 1.7$ are consistent with the values observed by Kawagucci et al. (2008).

While cryogenically extracting CO_2 from the air sample, N_2O was simultaneously trapped with the CO_2 . Because N_2O and CO_2 have the same mass number, we must correct for N_2O to obtain the true $\delta^{13}\text{C}$ value. The correction method was described in detail by Nakazawa et al. (1993), and we summarize it here. The ionization efficiency of N_2O was measured in advance, and the measured $\delta^{13}\text{C}$ value was corrected using the $\text{N}_2\text{O}/\text{CO}_2$ mole fraction ratio of the air sample. In the troposphere, the $\text{N}_2\text{O}/\text{CO}_2$ mole fraction ratio does not change significantly, and the N_2O correction is almost constant (approximately +0.2‰; Nakazawa et al., 1997a). However, because the N_2O mole fraction decreases rapidly with increasing altitude in the stratosphere (e.g., Toyoda et al., 2001), the $\text{N}_2\text{O}/\text{CO}_2$ mole fraction ratio decreases with altitude, and the magnitude of the N_2O correction also varies vertically. Therefore, the N_2O correction also strongly influences the observed vertical distribution of $\delta^{13}\text{C}$ in the stratosphere.

Stratospheric air samples were also measured for $\delta^{13}\text{CH}_4$ using a gas chromatograph-combustion-isotope ratio spectrometer developed at the National Institute of Polar Research (NIPR), Japan. The method has been described in detail by Morimoto et al. (2006) and is briefly summarized here. An air sample was flushed by pure helium and passed through a trap containing HayeSep D at a temperature of -120°C to concentrate CH_4 in the sample. Then, the CH_4 was released and transferred into a PoraBOND Q cryofocusing trap at -197°C . The CH_4 released from the trap when brought to room temperature was separated from the residual components using a PoraPLOT Q capillary column and combusted into CO_2 in a furnace with a Pt-Ni-Cu catalyst at 940°C . The converted CO_2 was analyzed using a continuous-flow mass spectrometer (Finnigan MAT252).




Figure 1. The relationship between the N_2O mole fraction and (a) $\delta^{18}\text{O}$ in CO_2 and (b) $\delta^{17}\text{O}$ in CO_2 . $\delta^{18}\text{O}$ and $\delta^{17}\text{O}$ values measured by Kawagucci et al. (2008) are shown by open circles. $\delta^{18}\text{O}$ and $\delta^{17}\text{O}$ values calculated assuming an MIE factor of 1.7 (see text) in this study are shown by closed circles.

The precision was reported to be 0.06‰ (Morimoto et al., 2006). Before 1996, $\delta^{13}\text{CH}_4$ measurements were performed using off-line oxidation equipment (Sugawara et al., 1997), in which CH_4 in the sample was converted to CO_2 using a catalyst (0.5‰ Pt supported on alumina pellets) at 750 °C , and the pure CO_2 sample thus prepared was analyzed using the mass spectrometer installed at TU in the same way as for $\delta^{13}\text{CO}_2$, except using a micro-volume inlet system. The overall analytical precision was 0.07‰ (Sugawara et al., 1997). To produce a high-quality data record, we needed to verify that these two methods produced the same results. Therefore, several sets of stratospheric air samples were analyzed using both methods, and the $\delta^{13}\text{CH}_4$ results differed by about 0.85‰ . To correct for this difference, the off-line measurement results were adjusted to match the on-line measurements. In this study, the $\delta^{13}\text{CH}_4$ value is reported relative to VPDB. The international comparison of $\delta^{13}\text{CH}_4$ measurements described in detail by Umezawa et al. (2018) revealed a difference of 0.2‰ between NIPR and the Institute of Arctic and Alpine Research (INSTAAR) results. Therefore, all $\delta^{13}\text{CH}_4$ values observed by the National Oceanic and Atmospheric Administration Global Monitoring Laboratory (NOAA/GML) and INSTAAR were shifted to match the NIPR scale.

The method used to analyze GS is described in Appendix A.

3 Results and discussion

3.1 Vertical profiles of CO_2 mole fraction and $\delta^{13}\text{CO}_2$

Figure 2 shows vertical profiles of the CO_2 mole fraction and $\delta^{13}\text{CO}_2$ over Japan throughout the study period. CO_2 mole fractions and $\delta^{13}\text{CO}_2$ values have increased and decreased, respectively, from 1985 to 2020. Vertical $\delta^{13}\text{CO}_2$ profiles observed by Gamo et al. (1995) in 1985 and 1986 are included

in this figure after MIE correction. Previous studies reported that the CO_2 mole fraction decreases with increasing altitude from the tropopause to around 20–25 km, and then it becomes almost constant at higher altitudes (Nakazawa et al., 1995, 2002). Our CO_2 mole fraction data show a similar vertical distribution during the observation period. Because our $\delta^{13}\text{CO}_2$ data are more scattered with respect to altitude than the CO_2 mole fraction, we calculated detrended vertical $\delta^{13}\text{CO}_2$ profiles as follows. First, the long-term trend of the tropospheric $\delta^{13}\text{CO}_2$ at Mauna Loa, Hawaii (hereafter, MLO) (Michel et al., 2025) was calculated as a representative tropospheric trend, and all our stratospheric $\delta^{13}\text{CO}_2$ observation data were shifted up or down based on that trend to produce values corresponding to August 2016 for comparison with the model results described below. Then, the $\delta^{13}\text{CO}_2$ values thus obtained for the entire study period were grouped into 11 2 km vertical bins, and the average value for each bin was calculated (Fig. 2c). The results showed that $\delta^{13}\text{CO}_2$ values increased with increasing altitude from near the tropopause to around 25 km and then decreased slightly above that altitude. Gamo et al. (1989) originally reported that $\delta^{13}\text{CO}_2$ values increased by approximately 0.4‰ from the tropopause to 25 km altitude in September 1985, whereas our recalculated results showed an increase of about 0.2‰ (Fig. 2b). The average increase shown in Fig. 2c was even smaller, only about 0.1‰ . This difference between the vertical profiles originally reported by Gamo et al. (1989) and our present results is due to our incorporation of the MIE on ^{17}O and ^{18}O (Sect. 2.2).

Below an altitude of ~ 17 km, $\delta^{13}\text{CO}_2$ values are quite variable, often being irregularly high or low depending on the observation year, which is exhibited by the large error bars in Fig. 2c. Based on detailed aircraft observations of $\delta^{13}\text{CO}_2$ variations in the troposphere over Japan, Nakazawa et al. (1993) reported that the seasonal cycle of $\delta^{13}\text{CO}_2$ between 8 km altitude and the tropopause show a minimum




Figure 2. Vertical profiles of (a) CO₂ mole fraction and (b) $\delta^{13}\text{CO}_2$ over Japan during the period 1985–2020. (c) $\delta^{13}\text{CO}_2$ values averaged over the study period and grouped in 11 2 km vertical bins (open squares). Curves show 2-D modeling results without the tropospheric trend, seasonal cycle, gravitational separation (GS), or airborne sources (thick dotted line), including only the tropospheric trend (dashed line), including the tropospheric trend, seasonal cycle, and GS (dotted-dashed line), and including the tropospheric trend, seasonal cycle, GS, and airborne sources (solid line).

around May and a maximum around September, with a peak-to-peak amplitude of 0.36‰. Indeed, our balloon observations were conducted in either spring (late May to early June) or summer (late July to early September), and the observed $\delta^{13}\text{CO}_2$ values at lower altitudes tended to be lower in spring and higher in summer. Accordingly, we attribute the large $\delta^{13}\text{CO}_2$ variations below an altitude of 17 km to the upper tropospheric air itself, or stratospheric air directly influenced by the troposphere.

Our data clearly reveal a secular decrease in mid-stratospheric $\delta^{13}\text{CO}_2$ values. To compare the temporal variations of $\delta^{13}\text{CO}_2$ in the mid-stratosphere with those in the troposphere, the $\delta^{13}\text{CO}_2$ values obtained by averaging mid-stratospheric data above 24 km altitude (30 hPa) from each year and monthly average MLO data are shown in Fig. 3. When comparing our data with the NOAA/GML and INSTAAR data at MLO (Michel et al., 2025), any possible offset of $\delta^{13}\text{CO}_2$ measurements should be taken into account. The inter-comparison results of the WMO/IAEA Round Robin Comparison Experiment (<https://gml.noaa.gov/ccgg/wmorrr/>, last access: 20 February 2025) show that our values are approximately 0.07‰ lower than INSTAAR values. In this study, we directly compared $\delta^{13}\text{CO}_2$ data at Ny-Ålesund, Svalbard observed by Goto et al. (2017) with the INSTAAR data at the Zeppelin Observatory (ZEP) in Ny-Ålesund because we used the same $\delta^{13}\text{CO}_2$ measurement system and standard scale as Goto et al. (2017). Monthly average $\delta^{13}\text{CO}_2$ values at Ny-Ålesund from both studies were calculated by applying a curve-fitting procedure (Nakazawa et al., 1997b) over the years compared, 1996–2014. As a result, we estimated the difference between this study and INSTAAR data to be insignificant (-0.005 ± 0.067 ‰) and did not apply any further correction to our $\delta^{13}\text{CO}_2$ data. The average rate of change of $\delta^{13}\text{CO}_2$ in the mid-stratosphere was calculated to be -0.026 ± 0.001 ‰ yr⁻¹ for the period 1985–2020, consistent with the average rate of change at MLO from 1990 to 2024 (-0.026 ± 0.001 ‰ yr⁻¹) despite the slightly different time periods. Tropospheric $\delta^{13}\text{CO}_2$ values are known to decrease over time due to the emission of CO₂ with low $\delta^{13}\text{CO}_2$ by anthropogenic fossil fuel consumption, and our results show that a similar decreasing trend is observable in the stratosphere. It is also well known that mid-stratospheric air over the mid-latitudes is “aged,” mainly because of the slow transport of air from the tropical tropopause to the mid-latitude mid-stratosphere. Indeed, the age of mid-stratospheric air estimated from CO₂ mole fractions was approximately 4.4 yr older on average than that of tropical tropospheric air (see Sect. 3.5). Nevertheless, the $\delta^{13}\text{CO}_2$ values observed in the mid-stratosphere over Japan did not seem to show any significant time delay compared with the observed values in the tropical troposphere. Although systematic observations of $\delta^{13}\text{CO}_2$ have not yet been performed in the tropical upper troposphere where stratospheric air originates, Assonov et al. (2010) found from CARIBIC aircraft observations that $\delta^{13}\text{CO}_2$ variations in the tropical upper tro-




Figure 3. Average mid-stratospheric (>24 km altitude) $\delta^{13}\text{CO}_2$ values over Japan (closed squares). The linear function fitted to the average values using the least-squares method is shown by the dashed line. Values reported by Gamo et al. (1995) are shown by open squares. Monthly average $\delta^{13}\text{CO}_2$ observed at Mauna Loa by NOAA/GML and INSTAAR (Michel et al., 2025) is also shown by the dotted line.

posphere are close to those at MLO. Therefore, if we assume no difference between the $\delta^{13}\text{CO}_2$ values at MLO and those in the tropical upper troposphere, the lack of a delay between the values in the mid-stratosphere over Japan and those at MLO clearly contradicts the concept of the age of air. Simply following the logic of the age of air, we would expect the decrease of stratospheric $\delta^{13}\text{CO}_2$ to lag behind the tropospheric trend, such that the $\delta^{13}\text{CO}_2$ value at any given time should be higher in the mid-stratosphere than in the troposphere. This discrepancy suggests that $\delta^{13}\text{CO}_2$ is not a conserved quantity in the stratosphere and is instead modified not only by air transport but also by other mechanisms. In the following sections we discuss possible mechanisms in detail.

3.2 Airborne CO_2 source from CH_4 oxidation

In the stratosphere, CH_4 is destroyed by reactions with OH, $\text{O}(^1\text{D})$, and Cl, which produce intermediate products such as CH_3O_2 and CH_2O . These products are immediately converted to CO because of their short lifetimes. However, CO is also destroyed by reaction with OH. Therefore, the chemical budget of CO in the stratosphere is expressed as (Minschwaner et al., 2010)

$$\begin{aligned} \frac{d[\text{CO}]}{dt} = & [\text{CH}_4] (k_{\text{CH}_4+\text{OH}} [\text{OH}] \\ & + k_{\text{CH}_4+\text{Cl}} [\text{Cl}] + k_{\text{CH}_4+\text{O}} [\text{O}(^1\text{D})]) \\ & + J_{\text{CO}_2} [\text{CO}_2] - k_{\text{CO}+\text{OH}} [\text{CO}] [\text{OH}], \end{aligned} \quad (4)$$

where $k_{\text{CH}_4+\text{OH}}$, $k_{\text{CH}_4+\text{Cl}}$, $k_{\text{CH}_4+\text{O}}$, and $k_{\text{CO}+\text{OH}}$ are the reaction rates for the respective chemical reactions, and J_{CO_2} is the rate of photodissociation of CO_2 . The last term on the right-hand side of this equation represents the oxidation of

CO to CO_2 . Although the amount of CO_2 produced this way is small compared to the stratospheric abundance, it cannot be ignored as an airborne source, especially in terms of isotopic signatures. The timescales for the photochemical loss and production of CO in the mid-latitude mid-stratosphere (~ 35 km altitude) are about 30 d (Minschwaner et al., 2010), quite short compared to the timescale of the stratospheric mean meridional circulation (more than several years). If we assume that the chemical budget of CO is in a steady state and that CO_2 photodissociation is negligibly small below the mid-stratosphere, the rate of CH_4 loss should equal the rate of CO_2 production. Indeed, to estimate the age of stratospheric air from the CO_2 mole fraction, a small amount of CO_2 produced by the chemical destruction of CH_4 was taken into account (e.g., Engel et al., 2002; Sugawara et al., 2018). To correct for this effect, each observed CO_2 mole fraction was adjusted using the CH_4 mole fraction measured from the corresponding air sample prior to age calculation (see Appendix C). The adjustment increases with increasing altitude and becomes larger than $1 \mu\text{mol mol}^{-1}$ at the highest altitudes. As with the CO_2 mole fraction, ^{13}C can be corrected by assuming a steady-state ^{13}CO budget in which ^{13}C is added to $^{13}\text{CO}_2$ from $^{13}\text{CH}_4$ through a series of oxidation reactions from CH_4 to CO_2 . Indeed, it is well known that isotopic fractionation occurs during CH_4 oxidation (Saueressig et al., 1995; Sugawara et al., 1997; Rice et al., 2003; Röckmann et al., 2011).

Figure 4a, b shows the vertical CH_4 mole fraction and $\delta^{13}\text{CH}_4$ profiles observed over Japan, respectively. A common feature of both profiles is their rapid change with altitude, although $\delta^{13}\text{CH}_4$ values increase, whereas the mole fractions decrease with increasing altitude. Such vertical distributions have been attributed to large kinetic isotopic effects (KIEs) associated with the chemical destruction of CH_4 in the stratosphere (Sugawara et al., 1997; Rice et al., 2003; Röckmann et al., 2011). KIEs on CH_4 have also been studied using numerical models (Saueressig et al., 2001; Wang et al., 2002; McCarthy et al., 2003). In previous studies, the apparent fractionation factor was derived from the Rayleigh distillation model by assuming that the chemical destruction of CH_4 occurs in a closed system (Sugawara et al., 1997; Rice et al., 2003). In general, the Rayleigh distillation model for the CH_4 mole fraction and $\delta^{13}\text{CH}_4$ is expressed as

$$\delta^{13}\text{CH}_4 = \left(\delta^{13}\text{CH}_{4,0} + 1 \right) \mu^{\alpha-1} - 1, \quad (5)$$

where $\delta^{13}\text{CH}_{4,0}$, μ , and α are the isotopic ratio before CH_4 is destroyed, the ratio of mole fractions ($\mu = [\text{CH}_4]/[\text{CH}_4]_0$), and the apparent fractionation factor, respectively. The apparent fractionation factor can be estimated by applying Eq. (5) to observed data. We note that $[\text{CH}_4]_0$ and $\delta^{13}\text{CH}_{4,0}$ change with time and should be taken here as the values in the tropical upper troposphere at the time when air entered the stratosphere through the tropical tropopause layer (TTL). Therefore, the tropical upper tropospheric records of

CH_4 and $\delta^{13}\text{CH}_4$ were used to give $[\text{CH}_4]_0$ and $\delta^{13}\text{CH}_{4,0}$ values going back in time based on the mean age estimated from CO_2 mole fractions (see Appendix C). Details of the tropical upper tropospheric records of CH_4 and $\delta^{13}\text{CH}_4$ are described in Sect. 3.5. Figure 4c shows that the relationship between $\ln\{(\delta^{13}\text{CH}_4 + 1)/(\delta^{13}\text{CH}_{4,0} + 1)\}$ and $\ln([\text{CH}_4]/[\text{CH}_4]_0)$ can be approximated by linear functions if we divided the data between the lower (<24 km) and mid-stratosphere (>24 km), with the upper layer showing larger isotopic fractionation effects. The apparent fractionation factors calculated using the least-squares method were 0.9889 ± 0.0003 and 0.9866 ± 0.0007 for the lower and mid-stratosphere, respectively. We attributed these differences in apparent fractionation factors to differences in the relative contributions of the CH_4 reactions with OH, $\text{O}(^1\text{D})$, or Cl over the air transport pathway, and/or the effect of air mixing (Röckmann et al., 2011).

Equation (5) expresses the relationship between the mole fraction and isotopic ratio of the remaining CH_4 . Similarly, focusing on the CH_4 lost via the $\text{CH}_4\text{--CO--CO}_2$ chain reaction, the relationship between the CH_4 mole fraction and isotopic ratio of lost CH_4 (hereafter, $\delta^{13}\text{CH}_{4,\text{L}}$) is derived from Eq. (5) as

$$\delta^{13}\text{CH}_{4,\text{L}} = \left(\delta^{13}\text{CH}_{4,0} + 1 \right) \frac{1 - \mu^\alpha}{1 - \mu} - 1. \quad (6)$$

Here, the removed CH_4 includes the total amount of CH_4 loss integrated over the air transport pathway. Based on this equation, we calculated $\delta^{13}\text{CH}_{4,\text{L}}$ for each year's data from the observed CH_4 mole fractions and apparent fractionation factors by grouping them into altitudes below and above 24 km. Typical $\delta^{13}\text{CH}_{4,\text{L}}$ values were about -57‰ , -56‰ , and -55‰ around altitudes of 20, 30, and 35 km, respectively. The stratospheric $\delta^{13}\text{CO}_2$ values were corrected for airborne sources simply by assuming that carbon with the isotopic ratio of $\delta^{13}\text{CH}_{4,\text{L}}$ was added to ambient CO_2 . We note that the amount of CO_2 added by CH_4 oxidation was estimated from the difference between the CH_4 mole fraction observed in the mid-latitude stratosphere and that estimated at the tropical upper troposphere; this does not represent chemical production at the observed location, but apparent production, because CH_4 oxidation occurs along the transport pathway of the air mass and is accompanied by air mixing. Consequently, the depression of CH_4 integrated over the air transport pathway is observed at a certain location in the stratosphere. Because we assumed the simple Rayleigh distillation model, the $\delta^{13}\text{CH}_{4,\text{L}}$ estimated here is also an apparent value in a similar sense as described above. Accordingly, our assumption described above was a first-order approximation to correct $\delta^{13}\text{CO}_2$ for airborne CO_2 sources.

As an example, we corrected the vertical distributions of the CO_2 mole fraction and $\delta^{13}\text{CO}_2$ (Fig. 5a) and compared with the CH_4 mole fraction and $\delta^{13}\text{CH}_4$ (Fig. 5b) observed on 22 August 2010. The results indicated that the $\delta^{13}\text{CO}_2$ value decreased by 0.14‰ and the CO_2 mole fraction in-

creased by approximately $1.1 \mu\text{mol mol}^{-1}$ at the highest altitude (~ 34 km) when corrected for airborne CO_2 sources. Therefore, such a correction for the airborne CO_2 source with low $\delta^{13}\text{CO}_2$ values produced by CH_4 oxidation is essential for understanding CO_2 mole fraction and $\delta^{13}\text{CO}_2$ variations in the mid-stratosphere. This effect is particularly strong on $\delta^{13}\text{CO}_2$ and significantly impacted the shape of the vertical distribution.

We note that the transport of CO from the troposphere into the stratosphere and its oxidation in the lower stratosphere are not considered in the above discussion. Because the photochemical timescales of CO are longer in the lower stratosphere than in the mid-stratosphere, tropospheric CO could be a stratospheric CO_2 source associated with troposphere–stratosphere exchange and may be important in the lower stratospheric CO budget. Observational results on the stable carbon isotopic ratio of CO, $\delta^{13}\text{CO}$, have been reported not only for the troposphere (Brenninkmeijer, 1993; Röckmann and Brenninkmeijer, 1997; Röckmann et al., 1999, 2002; Kato et al., 2000) but also for the lower stratosphere (Brenninkmeijer et al., 1996; Röckmann et al., 2024). In particular, Brenninkmeijer et al. (1996) found that extremely low $\delta^{13}\text{CO}$ values in the southern high-latitude lower stratosphere are accompanied by increasing $\delta^{13}\text{CH}_4$ values and suggested that the large KIE of CH_4 destruction by Cl plays an important role. The local production of CO by CH_4 -destruction-depleted ^{13}C in stratospheric CO is consistent with the above discussion if CO is rapidly oxidized to CO_2 depleted in ^{13}C . However, tropospheric CO could be an additional airborne CO_2 source if it is transported into the lower stratosphere and oxidized to CO_2 . Previous studies have reported that $\delta^{13}\text{CO}$ values in the free troposphere are around -30‰ to -25‰ (Brenninkmeijer, 1993; Röckmann and Brenninkmeijer, 1997; Röckmann et al., 1999, 2002; Kato et al., 2000), and the KIE on carbon in the reaction $\text{CO} + \text{OH}$ depends on atmospheric pressure (Stevens and Wagner, 1989; Bergamaschi et al., 2000). Its fractionation factor is 0.994 (expressed as the ratio of rate constants, $^{13}k/^{12}k$) at 1013 hPa, but increases to about 1.000 at 400 hPa and further to 1.003–1.005 above the tropopause. Therefore, CO_2 produced from the reaction $\text{CO} + \text{OH}$ should be slightly enriched in ^{13}C in the stratosphere, contrary to CO_2 produced in the lower troposphere. Accordingly, tropospheric CO could be an airborne CO_2 source with $\delta^{13}\text{C}$ values lower than around -25‰ to -20‰ in the lower stratosphere. Assuming that tropospheric air containing $250 \text{ nmol mol}^{-1}$ CO with $\delta^{13}\text{CO} = -30\text{‰}$ is transported to the stratosphere where $200 \text{ nmol mol}^{-1}$ CO is lost by reaction with OH, $\delta^{13}\text{CO}_2$ values should decrease by only about 0.01‰ when $0.2 \mu\text{mol mol}^{-1}$ CO_2 with $\delta^{13}\text{CO}_2 = -28\text{‰}$ is added to $390 \mu\text{mol mol}^{-1}$ CO_2 with $\delta^{13}\text{CO}_2 = -8.5\text{‰}$. Therefore, this effect would be negligibly small compared with the influence of CH_4 oxidation. Unfortunately, we could not further evaluate the effect of tropospheric CO from our




Figure 4. Vertical profiles of (a) CH₄ mole fraction and (b) $\delta^{13}\text{CH}_4$ over Japan, and (c) the relationship between $\ln\left(\frac{(\delta^{13}\text{CH}_4 + 1)}{(\delta^{13}\text{CH}_4)_0 + 1}\right)$ and $\ln\left(\frac{[\text{CH}_4]}{[\text{CH}_4]_0}\right)$. In (c), the data are divided into the lower (<24 km, closed squares) and mid-stratosphere (>24 km, open circles), and the linear functions fitted to each dataset using the least-squares method are shown by lines.

observations, and further study including measurements of $\delta^{13}\text{CO}$ is needed.

3.3 Gravitational separation of $\delta^{13}\text{CO}_2$

The standardized GS is defined as $\langle\delta_G\rangle$, which is normalized by the mass number differences of corresponding molecules (see Appendix A). With respect to $\delta^{13}\text{CO}_2$, the upward decrease due to GS is equal to $\langle\delta_G\rangle$ for $^{13}\text{C}^{16}\text{O}_2$ and $^{12}\text{C}^{17}\text{O}^{16}\text{O}$ ($\Delta m = 1$), and is twice as large for $^{12}\text{C}^{18}\text{O}^{16}\text{O}$, $^{13}\text{C}^{17}\text{O}^{16}\text{O}$, and $^{12}\text{C}^{17}\text{O}_2$ ($\Delta m = 2$). Therefore, GS equal to and twice the magnitude of $\langle\delta_G\rangle$ can be considered to act on $\delta^{45}\text{CO}_2$ and $\delta^{46}\text{CO}_2$, respectively:

$$\Delta_G(\delta^{45}\text{CO}_2) = \langle\delta_G\rangle - \langle\delta_G\rangle_0, \quad (7a)$$

$$\Delta_G(\delta^{46}\text{CO}_2) = 2(\langle\delta_G\rangle - \langle\delta_G\rangle_0). \quad (7b)$$

Here, $\Delta_G(x)$ and $\langle\delta_G\rangle_0$ represent the change in isotopic value x due to GS and the value of $\langle\delta_G\rangle$ before GS oc-

curs (i.e., corresponding to the relevant value in the upper troposphere), respectively. Because $\langle\delta_G\rangle - \langle\delta_G\rangle_0$ is negative in the stratosphere (see Appendix A), $\Delta_G(\delta^{45}\text{CO}_2)$ and $\Delta_G(\delta^{46}\text{CO}_2)$ are also negative. When $\delta^{13}\text{CO}_2$ was corrected for GS, Eqs. (7a) and (7b) were first applied to $\delta^{45}\text{CO}_2$ and $\delta^{46}\text{CO}_2$, respectively, and then $\delta^{13}\text{CO}_2$ was determined using the MIE algorithm described in Sect. 2.2. As shown in Fig. 5c, $|\Delta_G(\delta^{13}\text{CO}_2)|$ was slightly smaller than (about 82 % of) $|\Delta_G(\delta^{45}\text{CO}_2)|$ in the mid-stratosphere, mainly because of $^{12}\text{C}^{17}\text{O}^{16}\text{O}$ enrichment due to the MIE in the stratosphere.

The same method can be applied to the mole fraction of a specific molecule (Ishidoya et al., 2006):

$$\Delta_G(C) = C_0 \times (m - m_{\text{air}}) \times (\langle\delta_G\rangle - \langle\delta_G\rangle_0). \quad (8)$$

Here, C_0 and C denote the mole fractions of the molecule before and after GS, respectively, and m and m_{air} are the respective mass numbers of the molecule and air. The maximum depression of the CO₂ mole fraction amounts to about $0.6 \mu\text{mol mol}^{-1}$ at the highest altitude, assuming $C_0 = 400 \mu\text{mol mol}^{-1}$ and $\langle\delta_G\rangle - \langle\delta_G\rangle_0 = -100$ per meg (10^{-6}). As an example, the vertical profile of the CO₂ mole frac-




Figure 5. (a) Vertical profiles of the CO_2 mole fraction (closed circles) and $\delta^{13}\text{CO}_2$ (open circles) observed over Japan on 22 August 2010. Values corrected for airborne CO_2 sources (squares) and additionally for gravitational separation (GS; triangles) are also shown. (b) Vertical profiles of the CH_4 mole fraction (closed circles) and $\delta^{13}\text{CH}_4$ (open circles). (c) Vertical profiles of the magnitude of GS corrections, $|\Delta_G|$, for $\delta^{45}\text{CO}_2$ (circles), $\delta^{46}\text{CO}_2$ (triangles), $\delta^{13}\text{CO}_2$ (squares), $\delta^{18}\text{O}$ (diamonds), and $\delta^{17}\text{O}$ (crosses) in CO_2 . (d) Observed $\delta^{13}\text{CO}_2$ plotted against CO_2 mole fraction (closed circles) and those corrected for airborne CO_2 sources and GS (triangles). Lines are linear functions fitted to each trend by the least-squares method.

tion obtained by correcting the 22 August 2010 observations for GS is shown in Fig. 5a. Compared to the profiles corrected for airborne CO_2 sources, GS correction increases the CO_2 mole fraction by $0.4 \mu\text{mol mol}^{-1}$ and the $\delta^{13}\text{CO}_2$ by approximately 0.06‰ at 34 km altitude. Although the effect of GS on the CO_2 mole fraction is small, that on $\delta^{13}\text{CO}_2$ is non-negligible, significantly influencing the vertical profile along with CH_4 oxidation. In this regard, the mean age estimated from the CO_2 mole fraction without GS correction has an older age bias that increases with increasing altitude. The age bias is negligibly small (< 1 month) in the lower stratosphere but not negligible in the mid-stratosphere (> 2 months at 35 km altitude). Therefore, we applied the GS correction to

the CO_2 mole fraction to estimate the CO_2 age, as described in Appendix C.

Correcting for both CH_4 oxidation and GS, $\delta^{13}\text{CO}_2$ values increase with altitude and become nearly constant above ~ 25 km; they are clearly anticorrelated with the CO_2 mole fraction. Assuming a linear relationship between the CO_2 mole fraction and $\delta^{13}\text{CO}_2$ (Fig. 5d), the slope calculated by least squares is $-0.005 \pm 0.003 \text{‰} (\mu\text{mol mol}^{-1})^{-1}$ for the raw observational data and $-0.023 \pm 0.002 \text{‰} (\mu\text{mol mol}^{-1})^{-1}$ for the corrected data. This rate of change for the corrected data is less negative than approximately $-0.05 \text{‰} (\mu\text{mol mol}^{-1})^{-1}$ for the seasonal cycles in the troposphere (Nakazawa et al., 1993) and is near the rate of change for their secular trends (Morimoto et al., 2000; Goto




Figure 6. Schematic representation of the mechanisms influencing the vertical profile of stratospheric $\delta^{13}\text{CO}_2$ (see text).

et al., 2017). Generally, because of the Brewer–Dobson circulation, air at higher altitudes is older. Therefore, we consider that this rate of change for the corrected data basically reflects different ages of air.

3.4 Two-dimensional model of stratospheric $\delta^{13}\text{CO}_2$

The mechanisms governing the vertical $\delta^{13}\text{CO}_2$ profile in the stratosphere are shown schematically in Fig. 6, including the influence of the age of air as well as the effects of CH_4 oxidation and GS discussed in Sects. 3.2 and 3.3. Tropospheric $\delta^{13}\text{CO}_2$ values have been decreasing through time due to the combustion of fossil fuels. If it is assumed that $^{12}\text{CO}_2$ and $^{13}\text{CO}_2$ have no sinks or sources in the stratosphere, the air that intrudes from the tropical upper troposphere is slowly transported to the mid-latitude stratosphere by the Brewer–Dobson circulation, conserving the $\delta^{13}\text{CO}_2$ value. Therefore, based on the concept of age of air, it is expected that $\delta^{13}\text{CO}_2$ values should also decrease with time in the stratosphere after a certain lag time. Accordingly, $\delta^{13}\text{CO}_2$ values should be higher in the mid-stratosphere than in the troposphere (dotted line in Fig. 6). In contrast, GS decreases $\delta^{13}\text{CO}_2$ values at higher altitudes because $^{13}\text{C}^{16}\text{O}_2$ is heavier than $^{12}\text{C}^{16}\text{O}_2$ (dashed line in Fig. 6). In addition, ^{13}C -depleted CO_2 produced by CH_4 oxidation further decreases $\delta^{13}\text{CO}_2$ values at high altitudes (solid line in Fig. 6). The observed stratospheric vertical $\delta^{13}\text{CO}_2$ profile should therefore be formed by a combination of these effects.

To verify this hypothesis, we performed numerical simulations using a two-dimensional model of the middle atmosphere (SOCRATES) developed by the National Center for Atmospheric Research (NCAR; Huang et al., 1998; Park et al., 1999; Khosravi et al., 2002). Details of the model calculations are described in Appendix B. The monthly average

meridional $\delta^{13}\text{CO}_2$ distribution calculated by SOCRATES for August 2016, which includes the effects of CH_4 oxidation and GS, is shown in Fig. 7. The effect of the CO_2 production associated with the KIE on CH_4 destruction is included in a simplified manner (see Appendix B). The $\delta^{13}\text{CO}_2$ values decrease with increasing altitude up to about 14 km and then increase up to about 20 km in the mid-latitudes of the northern hemisphere during summer. The distribution of $\delta^{13}\text{CO}_2$ from the troposphere to the lower stratosphere should therefore be closely related to the propagation of its seasonal cycle in the troposphere. $\delta^{13}\text{CO}_2$ values reach a maximum at around 20 km and then gradually decrease. At high latitudes in the southern (winter) hemisphere, the vertical decrease of $\delta^{13}\text{CO}_2$ is significant, suggesting that the effects of CH_4 oxidation and GS are enhanced by the descending upper atmosphere.

To investigate the vertical $\delta^{13}\text{CO}_2$ profile in detail, we compared this model with our observations in Fig. 2c. The modeled $\delta^{13}\text{CO}_2$ profile calculated without accounting for CH_4 oxidation and GS increased with increasing altitude, and it was 0.14‰ higher at altitudes around 35 km than near the tropopause. Because the average rate of change of the long-term $\delta^{13}\text{CO}_2$ record at MLO is -0.026‰ yr^{-1} for the period 1990–2022, the lag time at 35 km (0.14‰ divided by 0.026‰ yr^{-1}) should be approximately 5.4 yr, nearly consistent with the calculated mean age of air (see Appendix C). If the effect of GS is further taken into account, the $\delta^{13}\text{CO}_2$ decreased by approximately 0.07‰ at 35 km. The most significant impact on the modeled vertical profile was the effect of CH_4 oxidation, which imparted a decrease of 0.26‰ at around 35 km. The vertical profile calculated including all effects exhibited a maximum at an altitude of about 20 km and then decreased monotonically at greater altitudes, mainly because of the effect of CH_4 oxidation. However, the observed $\delta^{13}\text{CO}_2$ profile increased to about 25 km altitude, indicating that the model results clearly overestimated the ^{13}C depletion at higher altitudes. We note, though, that this model calculation included the seasonal cycle of tropospheric $\delta^{13}\text{CO}_2$ and nearly reproduced the vertical distribution observed at 14–20 km altitude, which is mainly due to the upward propagation of the seasonal cycle from the upper troposphere to the lower stratosphere. Therefore, the difference between the observed and modeled results above 20 km was likely due mainly to an overestimation of the effect of CH_4 oxidation.

One possible reason for this difference was an overestimation of CH_4 loss at high altitudes. The observed CH_4 mole fraction at ~ 35 km altitude ranged between about 500 and 900 nmol mol^{-1} (Fig. 4), but the model results indicated a monthly average of about 600 nmol mol^{-1} for August, which was lower than the observed average value (see Fig. B1c in Appendix B). This discrepancy may be related to the fact that we weakened the mass stream function to reproduce a realistic age of air (see Appendix B). Another reason may be that the model simulation set the KIE on CH_4 destruction in a simplified manner, with the apparent fractionation




Figure 7. Monthly average meridional distribution of $\delta^{13}\text{CO}_2$ for August 2016 calculated using the SOCRATES model. Values lower than -9.0‰ are colored gray.

factors set uniformly at 0.9889 and 0.9866 for the lower and mid-stratosphere, respectively. In reality, the apparent fractionation factor for CH_4 oxidation varies with altitude and latitude, and it is determined not only by a complex process that depends on the paths of air masses, but also by the chemical reactions that occur along the pathway (e.g., Röckmann et al., 2011). In this study, the KIEs on the CH_4 – CO – CO_2 reaction processes were not calculated explicitly, but rather in a simplified manner, which may have resulted in an overestimation of the effect on $\delta^{13}\text{CO}_2$. However, the present model simulation was able to prove our hypothesis that three factors – CH_4 oxidation, GS, and air age – are essential to determining vertical $\delta^{13}\text{CO}_2$ profiles.

3.5 Stratospheric potential $\delta^{13}\text{C}$

From the above discussion, we considered that the total amount of ^{13}C in both CO_2 and CH_4 , $\delta^{13}\text{C}_\text{T}$, was conserved. We defined $\delta^{13}\text{C}_\text{T}$ as

$$\delta^{13}\text{C}_\text{T} = \frac{n(\text{CO}_2) \times \delta^{13}\text{CO}_2 + n(\text{CH}_4) \times \delta^{13}\text{CH}_4}{n(\text{CO}_2) + n(\text{CH}_4)}, \quad (9)$$

which is quasi-conservative with respect to the KIEs associated with CH_4 oxidation and consequent CO_2 production in the stratosphere. Furthermore, as mentioned above, the effect of GS cannot be ignored in the stratosphere. Taking this into account, we defined the stratospheric potential $\delta^{13}\text{C}$, $\delta^{13}\text{C}_\text{P}$, by correcting $\delta^{13}\text{C}_\text{T}$ for GS:

$$\delta^{13}\text{C}_\text{P} = \delta^{13}\text{C}_\text{T} - \Delta_\text{G}(\delta^{13}\text{CO}_2), \quad (10)$$

where $\Delta_\text{G}(\delta^{13}\text{CO}_2)$ denotes the GS correction for $\delta^{13}\text{CO}_2$. Normally, $\Delta_\text{G}(\delta^{13}\text{CO}_2)$ gives a negative value, which means that it has a positive effect on $\delta^{13}\text{C}_\text{P}$. This equation does not include the effect of GS on $\delta^{13}\text{CH}_4$, but that contribution to $\delta^{13}\text{C}_\text{P}$ is negligibly small (about 10^{-4}‰). Equations (9) and (10) were applied to the observed results to calculate stratospheric $\delta^{13}\text{C}_\text{T}$ and $\delta^{13}\text{C}_\text{P}$ profiles. For some air samples without $\delta^{13}\text{CH}_4$ measurements, values were estimated from the CH_4 mole fraction using a linear function fitted to the log–log plot of the observed data shown in Fig. 4. The $\langle\delta_\text{G}\rangle$ values were also partially complemented by applying the average vertical profile. The average vertical profile of $\delta^{13}\text{C}_\text{P}$ thus calculated is shown in Fig. 8, alongside the CO_2 mole fraction. The data plotted in this figure were detrended and normalized to August 2016. The CO_2 mole fractions were also corrected for CH_4 oxidation and GS, although the magnitude of these corrections was relatively small. $\delta^{13}\text{C}_\text{P}$ increased with increasing altitude from the tropopause to about 25 km, then was almost constant with increasing altitude. Unlike $\delta^{13}\text{CO}_2$ (Fig. 2c), $\delta^{13}\text{C}_\text{P}$ was inversely correlated with the CO_2 mole fraction. Figure 9 shows the temporal variations of the averaged $\delta^{13}\text{CO}_2$, $\delta^{13}\text{C}_\text{T}$, and $\delta^{13}\text{C}_\text{P}$ values above 24 km altitude. $\delta^{13}\text{C}_\text{T}$ and $\delta^{13}\text{C}_\text{P}$ changed almost in parallel with $\delta^{13}\text{CO}_2$, showing a similar decreasing trend through time. The average difference from $\delta^{13}\text{CO}_2$ was $-0.079 \pm 0.025\text{‰}$ for $\delta^{13}\text{C}_\text{T}$ and $-0.034 \pm 0.027\text{‰}$ for $\delta^{13}\text{C}_\text{P}$.

If $\delta^{13}\text{C}_\text{P}$ is truly a conserved quantity in the stratosphere, it should decrease over time with a time lag relative to the variation in the tropical upper troposphere. However, there are no long-term data records of $\delta^{13}\text{C}$ values in CO_2 and




Figure 8. Vertical profiles of CO_2 mole fraction (closed squares) and stratospheric potential $\delta^{13}\text{C}$ ($\delta^{13}\text{C}_p$, open squares) based on their averages in 11 vertical bins over the period 1985–2020. All values were detrended and normalized to values in August 2016.

CH_4 in the tropical upper troposphere. Therefore, we hypothetically synthesized a $\delta^{13}\text{C}_p$ record in the tropical upper troposphere based on surface data observed at MLO and in the equatorial Pacific region by INSTAAR (Michel et al., 2025, 2023). In our previous study, we employed CO_2 mole fraction data obtained in the tropical upper troposphere using the Automatic Air Sampling Equipment in the Comprehensive Observation Network for TRace gases by AirLiner (CONTRAIL) program (Machida et al., 2008; Sawa et al., 2008; Matsueda et al., 2015) to estimate the mean age of stratospheric air (Sugawara et al., 2018). A similar method was used in this study. We first estimated the average seasonal cycle of $\delta^{13}\text{CO}_2$ in the tropical upper troposphere from the observed seasonal cycle of the CO_2 mole fraction, assuming that the correlation between the seasonal cycles of the CO_2 mole fraction and $\delta^{13}\text{CO}_2$ in the upper troposphere was approximately $-0.05\text{‰}(\mu\text{mol mol}^{-1})^{-1}$ (Nakazawa et al., 1993). The long-term $\delta^{13}\text{CO}_2$ trend was calculated from the observation data at MLO and then adjusted to match the observations in the equatorial Pacific region. The average seasonal cycles were then added to the long-term $\delta^{13}\text{CO}_2$ trend estimated in the equatorial Pacific region. The difference between $\delta^{13}\text{CO}_2$ values at the surface and those in the upper troposphere in the equatorial region is currently unknown. However, Sugawara et al. (2018) reported that the CO_2 and SF_6 ages in the TTL are 0.5–0.6 yr, consistent with the SF_6 age of 0–1.5 yr in the TTL based on Michelson Interferometer for Passive Atmospheric Sounding (Stiller et al.,




Figure 9. Average mid-stratospheric (>24 km altitude) values of $\delta^{13}\text{CO}_2$ (closed circles), $\delta^{13}\text{C}_T$ (closed triangles), and $\delta^{13}\text{C}_p$ (closed squares) over Japan. $\delta^{13}\text{CO}_2$ and $\delta^{13}\text{C}_p$ in the tropical upper troposphere are shown by dashed and solid lines, respectively. Red squares show the mid-stratospheric $\delta^{13}\text{C}_p$ shifted by -5.0 yr.

2008). Taking this into account, we assumed the difference between surface and upper troposphere $\delta^{13}\text{CO}_2$ values in the equatorial region to be equivalent to 0.5 yr. Because systematic measurements of CH_4 mole fraction and $\delta^{13}\text{CH}_4$ in the upper troposphere were performed in the CONTRAIL program (Umezawa et al., 2012), the MLO data obtained by INSTAAR were adjusted to fit the CONTRAIL data in the equatorial upper troposphere to produce long-term records of CH_4 mole fraction and $\delta^{13}\text{CH}_4$ in the equatorial upper troposphere. Because the seasonal cycle of $\delta^{13}\text{CH}_4$ in the equatorial upper troposphere is not clear (Umezawa et al., 2012), the seasonal cycle was ignored in the calculation of the upper tropospheric $\delta^{13}\text{C}_p$. Since GS, i.e., $\langle\delta_G\rangle - \langle\delta_G\rangle_0$, should be zero in the equatorial upper troposphere, tropospheric $\delta^{13}\text{C}_p = \delta^{13}\text{C}_T$.

As seen in Fig. 9, $\delta^{13}\text{CO}_2$ values did not differ significantly between the troposphere and the mid-stratosphere. However, the $\delta^{13}\text{C}_p$ values calculated from observed values in the mid-stratosphere were, on average, about $0.14 \pm 0.03\text{‰}$ higher than those calculated for the tropical troposphere. This means that $\delta^{13}\text{C}_p$ is conservative in the stratosphere and that the decreasing tropospheric $\delta^{13}\text{C}_p$ trend through time propagates into the stratosphere with a time delay. Roughly calculated, this lag time is about 5 yr, consistent with the age of air estimated from CO_2 mole fractions (Appendix C) and the model calculation. If the mid-stratospheric $\delta^{13}\text{C}_p$ values were shifted by -5 yr, they were in good agreement with the tropospheric $\delta^{13}\text{C}_p$ trend (Fig. 9).

It is thus possible to estimate the mean age of air using $\delta^{13}\text{C}_p$, which can serve as an additional air age tracer. Therefore, we estimated the mean age of air not only from the CO_2 mole fraction but also from $\delta^{13}\text{C}_p$ using a convolution method. Details of the age estimation method are described




Figure 10. Temporal variations of averaged $\delta^{13}\text{C}_\text{p}$ ages (closed squares) and CO_2 ages (open squares) in the mid-stratosphere (>24 km altitude) over Japan. The dashed line is a linear least-squares fit to the CO_2 ages. The linear fit to the $\delta^{13}\text{C}_\text{p}$ age is not shown because the trend is not significant.

in Appendix C. The total uncertainty on individual $\delta^{13}\text{C}_\text{p}$ ages was estimated to be ± 1.9 yr. This rather large uncertainty means that $\delta^{13}\text{C}_\text{p}$, with its smaller tropospheric secular change compared to other age tracers, was somewhat disadvantageous for age estimations. Because some of the $\delta^{13}\text{C}_\text{p}$ values in the lower stratosphere showed high variability, with values lower than the tropospheric reference record, the convolution method could not provide reasonable age estimations for those data. We attributed these irregularities and low values to the large seasonal variations in the upper troposphere and the stratosphere–troposphere air exchange in the lower stratosphere. Therefore, only mid-stratospheric (>24 km altitude) data were taken into account for age estimations, and the average values of the mean age were calculated.

The results are shown in Fig. 10, along with the mean age calculated from the CO_2 mole fraction using the same method. The average $\delta^{13}\text{C}_\text{p}$ and CO_2 ages were 5.5 ± 1.6 and 4.4 ± 0.6 yr, respectively. Because the uncertainty of the $\delta^{13}\text{C}_\text{p}$ age was large, no significant difference was apparent between the two values. To examine the long-term trend of mean age, a linear function was fitted to both the CO_2 and $\delta^{13}\text{C}_\text{p}$ age data using the least squares method. A slightly positive trend (0.04 ± 0.01 yr yr^{-1}) was observed for the CO_2 age, but no specific trend was observed for the $\delta^{13}\text{C}_\text{p}$ age. It is important to note that our data were unevenly distributed over time, with sparse observations in recent years and more frequent observations before 2007. This made it difficult to clarify the exact 35 yr trend from our data alone. At present, the $\delta^{13}\text{C}_\text{p}$ age has a relatively large uncertainty, making it unsuitable for use as an additional constraint for the age estimation. However, considering that the average $\delta^{13}\text{C}_\text{p}$ age was estimated to be 5.5 ± 1.6 yr, the concept of stratospheric $\delta^{13}\text{C}_\text{p}$ itself would be valid.

4 Conclusions

Previous studies on $\delta^{13}\text{CO}_2$ have mainly focused on observations in the troposphere and on elucidating the global CO_2 budget based on monitoring programs. However, little research has been performed on $\delta^{13}\text{CO}_2$ in the stratosphere, and the mechanisms of its distribution and variations are not fully understood. This study presented a comprehensive analysis of stratospheric $\delta^{13}\text{CO}_2$ obtained from balloon observations over Japan during the period 1985–2020. The following four factors were found to be important for understanding the variations of $\delta^{13}\text{CO}_2$ in the stratosphere. (1) Appropriate corrections for the MIE on ^{17}O and ^{18}O in isotopic analyses are essential. However, even after applying this correction, we found that the average values of $\delta^{13}\text{CO}_2$ in the northern mid-latitude mid-stratosphere and the tropical troposphere were almost identical, despite the secular decreasing $\delta^{13}\text{CO}_2$ trend and the expected time lag (age of air). We attributed this strange agreement to the other three factors: (2) airborne production of ^{13}C -depleted CO_2 by the oxidation of CH_4 in the stratosphere, (3) GS of $^{12}\text{CO}_2$ and $^{13}\text{CO}_2$ in the stratosphere, and (4) propagation of the decreasing tropospheric $\delta^{13}\text{CO}_2$ trend into the stratosphere. Based on these considerations, we introduced “stratospheric potential $\delta^{13}\text{C}$ ” ($\delta^{13}\text{C}_\text{p}$) as a new concept and inspected $\delta^{13}\text{C}_\text{p}$ in terms of the age of air. The average $\delta^{13}\text{C}_\text{p}$ age was calculated to be 5.5 ± 1.6 yr for the period of 1985–2020. Because the $\delta^{13}\text{C}_\text{p}$ age has larger uncertainties than the CO_2 age at present, it is difficult to refine the mean age estimation. However, $\delta^{13}\text{C}_\text{p}$ in the mid-latitude mid-stratosphere decreased over time with a time delay and it was found to be quasi-conservative in the stratosphere.

The observation of numerous stratospheric air constituents, as in this study, requires the collection of a large amount of stratospheric air with a high-quality method and multi-component gas analyses of each air sample. GS of atmospheric constituents in the stratosphere was first revealed by our previous balloon observations, and this knowledge is indispensable for a better understanding of the variations of stratospheric $\delta^{13}\text{CO}_2$. The quality of the air samples is particularly important for observing GS in the mid-stratosphere, because even slight separations due to molecular diffusion during air sampling could prevent GS observations. At present, the only method capable of such sampling is our cryogenic air sampler.

The 2-D model reproduced the basic structure of stratospheric $\delta^{13}\text{CO}_2$ and elucidated the mechanisms governing $\delta^{13}\text{CO}_2$ behavior. However, our numerical model simulations remained insufficient because the mean age of stratospheric air could not be reproduced adequately without arbitrarily tuning the model. Accordingly, our model overestimated the effect of airborne CO_2 production via CH_4 oxidation because the reaction process was treated in a very coarse manner. To improve upon this model, it will be necessary to incorporate KIEs into all reactions related to stratospheric carbon.

Although a two-dimensional model was used in this proof-of-concept study, a three-dimensional model is desirable for more realistic simulations in the future.

Appendix A: Gravitational separation of major atmospheric components

The gravitational separation (GS) of atmospheric components in the stratosphere was first reported from our balloon observations in Japan (Ishidoya et al., 2006, 2008a, 2013), in which the isotopic and elemental ratios of major atmospheric compositions, such as $\delta^{15}\text{N}$ in N_2 , $\delta^{18}\text{O}$ in O_2 , and $\delta(\text{Ar}/\text{N}_2)$, were measured by mass spectrometry at the National Institute of Advanced Industrial Science and Technology (AIST; e.g., Ishidoya et al., 2006, 2008a; Sugawara et al., 2018). Technical aspects of our mass spectrometry analyses are described in detail in Ishidoya and Murayama (2014). In this study, we used $\delta^{15}\text{N}$, $\delta^{18}\text{O}$, and $\delta(\text{Ar}/\text{N}_2)$ to evaluate GS, which are respectively defined as

$$\delta^{15}\text{N} = \frac{\left[\frac{n(^{15}\text{N}^{14}\text{N})}{n(^{14}\text{N}^{14}\text{N})} \right]_{\text{sp}}}{\left[\frac{n(^{15}\text{N}^{14}\text{N})}{n(^{14}\text{N}^{14}\text{N})} \right]_{\text{st}}} - 1, \quad (\text{A1a})$$

$$\delta^{18}\text{O} = \frac{\left[\frac{n(^{18}\text{O}^{16}\text{O})}{n(^{16}\text{O}^{16}\text{O})} \right]_{\text{sp}}}{\left[\frac{n(^{18}\text{O}^{16}\text{O})}{n(^{16}\text{O}^{16}\text{O})} \right]_{\text{st}}} - 1, \quad (\text{A1b})$$

and

$$\delta(\text{Ar}/\text{N}_2) = \frac{\left[\frac{n(^{40}\text{Ar})}{n(^{14}\text{N}^{14}\text{N})} \right]_{\text{sp}}}{\left[\frac{n(^{40}\text{Ar})}{n(^{14}\text{N}^{14}\text{N})} \right]_{\text{st}}} - 1, \quad (\text{A1c})$$

where “sp” and “st” denote the sample and standard gases, respectively. The 1σ reproducibilities of our $\delta^{15}\text{N}$, $\delta^{18}\text{O}$, and $\delta(\text{Ar}/\text{N}_2)$ measurements were ± 2 , ± 5 , and ± 7 per meg, respectively, which was sufficient to detect GS in the stratosphere. Ishidoya et al. (2013) concluded that vertical variations of the isotopic ratios of major atmospheric components are dominated by differences in molecular mass, based on mass-dependent relationships among related molecules such as $\delta^{15}\text{N}$ in N_2 , $\delta^{18}\text{O}$ in O_2 , and the Ar/N_2 ratio. Such GS in the stratosphere is known to be proportional to the mass difference, as would be expected for pure molecular diffusion, even though eddy diffusion obviously far exceeds molecular diffusion in the stratosphere. Therefore, GS can be expressed as

$$\delta - \delta_0 = \Delta m \times (\langle \delta_G \rangle - \langle \delta_G \rangle_0), \quad (\text{A2})$$

where Δm is the mass number difference, the suffix “0” refers to the value before GS occurs (i.e., corresponding to the relevant value in the upper troposphere), and $\langle \delta_G \rangle$ is the average of δ values normalized to $\Delta m = 1$ as reported by Ishidoya et al. (2013) and Sugawara et al. (2018). $\langle \delta_G \rangle$ is defined by

$$\langle \delta_G \rangle = \frac{1}{3} \left[\delta^{15}\text{N} + \delta^{18}\text{O}/2 + \delta(\text{Ar}/\text{N}_2)/12 \right]. \quad (\text{A3})$$




Figure A1. Vertical profiles of GS, $\langle \delta_G \rangle$, observed over Japan (Ishidoya et al., 2013).

Figure A1 shows vertical profiles of $\langle \delta_G \rangle$ obtained from balloon observations over Japan (Ishidoya et al., 2013). The value of $\langle \delta_G \rangle$ decreases gradually with increasing altitude. By fitting a linear function to the vertical profiles, the average difference of $\langle \delta_G \rangle$ between the lowermost and uppermost altitudes was found to be -66 ± 17 per meg. Here, it is important to mention the relationship between the GS of $\delta^{18}\text{O}$ and photochemical processes. As mentioned in Sect. 2.2., in the stratosphere, O_3 and CO_2 are significantly enriched in ^{17}O and ^{18}O by MIEs associated with photochemical reaction processes. Because stratospheric O_2 molecules are the source of the ^{17}O and ^{18}O enrichment, $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ values of O_2 are thought to be lower in the stratosphere than in the troposphere due to photochemical processes. This assumption is crucial not only for the Dole–Morita effect, but also for the ^{17}O budget of tropospheric O_2 (Bender et al., 1994; Luz et al., 1999; Luz and Barkan, 2011; Ishidoya et al., 2025). However, $\delta^{18}\text{O}$ values of stratospheric O_2 measured in our balloon observations show a very large decrease, i.e., by approximately -130 per meg, at around 35 km that is almost entirely dominated by GS. This decrease is probably also due in part to photochemical effects, but their contribution should be negligible, amounting to only a few per meg (Ishidoya et al., 2025).

Appendix B: Two-dimensional model

We used a two-dimensional model of the middle atmosphere (SOCRATES) developed by the National Center for Atmospheric Research (NCAR; Huang et al., 1998; Park et al., 1999; Khosravi et al., 2002). The advantage of using this model is that we have already calculated $^{13}\text{C}^{16}\text{O}_2$ and $^{12}\text{C}^{16}\text{O}_2$ and examined stratospheric GS in our previous studies (Ishidoya et al., 2013; Sugawara et al., 2018). Therefore, we give only a brief description of the model calculation here. To reproduce GS, the molecular diffusion flux must be calculated. The vertical component of the molecular diffusion flux ($F_{m_i,z}$) is given by

$$F_{m_i,z} = -D_{m_i} \left\{ \frac{\partial n_i}{\partial z} + \frac{m_i g}{RT} n_i + (1 + \alpha_{T_i}) \frac{\partial (\ln T)}{\partial z} n_i \right\}, \quad (\text{B1})$$

where n_i , D_{m_i} , m_i , and α_{T_i} are the number density, molecular diffusion coefficient, molecular mass, and thermal diffusion factor of atmospheric constituent i , respectively, and g , R , and T are the gravitational acceleration, gas constant, and temperature, respectively (Banks and Kockarts, 1973). However, because the thermal diffusion flux is not important in this case, we ignored it in the model calculations. Because it is cumbersome to include all CO_2 isotopologues in the model calculations, only $^{12}\text{C}^{16}\text{O}_2$ and $^{13}\text{C}^{16}\text{O}_2$ were calculated independently in the model, and $\delta^{13}\text{CO}_2$ was approximated using the respective mole fractions of $n(^{13}\text{C}^{16}\text{O}_2)$ and $n(^{12}\text{C}^{16}\text{O}_2)$ as

$$\delta^{13}\text{CO}_2 = \left(\frac{1}{^{13}\text{R}_{\text{VPDB}}} \frac{n(^{13}\text{C}^{16}\text{O}_2)}{n(^{12}\text{C}^{16}\text{O}_2)} - 1 \right). \quad (\text{B2})$$

Because $\delta^{13}\text{CO}_2$ has a large seasonal cycle in the troposphere, we used NOAA/GML and INSTAAR data to give latitude-dependent seasonal cycles as boundary conditions at the model surface. In addition, a secular decrease of $\delta^{13}\text{CO}_2$ in the troposphere was given by the MLO data (Michel et al., 2025). SOCRATES originally included a series of carbon reactions starting with CH_4 oxidation. Model calculations were repeated with and without GS, the secular trend and seasonal cycle of tropospheric $\delta^{13}\text{CO}_2$, and airborne CO_2 production, and comparisons were made between them (Fig. 2c). The chemical budget of CO_2 in SOCRATES is calculated as

$$\begin{aligned} \frac{d[\text{CO}_2]}{dt} = & k_{\text{CO}+\text{OH}} [\text{CO}][\text{OH}] \\ & + k_{\text{CO}+\text{O}(^3\text{P})} [M][\text{CO}][\text{O}(^3\text{P})] \\ & - J_{\text{CO}_2} [\text{CO}_2], \end{aligned} \quad (\text{B3})$$

where $k_{\text{CO}+\text{OH}}$ and $k_{\text{CO}+\text{O}(^3\text{P})}$ are the reaction rates of their respective chemical reactions, and M is the third body which carries off excess energy in a termolecular reaction. Because CO_2 photodissociation is important only above the mesosphere (e.g., Garcia et al., 2014), it was ignored in this study.

KIEs in the CH_4 – CO – CO_2 chain reaction were not included explicitly in these simulations but were rather treated in a coarse manner as follows. Simulations were performed with and without the chemical production of CO_2 . Taking the CO_2 mole fractions simulated with (“wP”) and without chemical production (“nP”) to be $n(\text{CO}_2)_{\text{wP}}$ and $n(\text{CO}_2)_{\text{nP}}$, respectively, the CO_2 mole fraction difference between the two simulations, $n(\text{CO}_2)_{\text{wP}} - n(\text{CO}_2)_{\text{nP}}$, was calculated as the apparent CO_2 production in the stratosphere. The apparent CO_2 production is distinct from the local production calculated in Eq. (B3) because it is an apparent value influenced by the accumulated chemical processes along the pathway of the air mass, including transport and mixing. A typical value of the apparent CO_2 production was simulated to be about $1.5 \mu\text{mol mol}^{-1}$ at 35 km altitude in the mid-latitudes. Similarly, the $\delta^{13}\text{CO}_2$ value simulated excluding chemical CO_2 production was expressed by $\delta^{13}\text{CO}_{2,\text{nP}}$. After both these simulations, the $\delta^{13}\text{CO}_2$ value including airborne CO_2 sources from the CH_4 destruction, $\delta^{13}\text{CO}_{2,\text{wP}}$, was calculated as

$$\delta^{13}\text{CO}_{2,\text{wP}} = \frac{\delta^{13}\text{CO}_{2,\text{nP}} \times n(\text{CO}_2)_{\text{nP}} + \delta^{13}\text{CH}_{4,\text{L}} \times [n(\text{CO}_2)_{\text{wP}} - n(\text{CO}_2)_{\text{nP}}]}{n(\text{CO}_2)_{\text{wP}}}. \quad (\text{B4})$$

In this equation, $\delta^{13}\text{CH}_{4,\text{L}}$ was calculated using Eq. (6) assuming apparent fractionation factors of 0.9889 and 0.9866 at lower (<24 km) and upper (>24 km) altitudes, respectively (see Sect. 3.2). Note that $\delta^{13}\text{CH}_{4,\text{L}}$ values were estimated from the apparent fractionation factors by assuming Rayleigh distillation relationships between vertical profiles of CH_4 mole fraction and $\delta^{13}\text{CH}_4$. Therefore, these apparent fractionation factors differed from fractionation factors *sensu stricto* and were influenced by the accumulated KIEs along the pathway of the air mass, as with the apparent CO_2 production described above. Equation (B4) means that $\delta^{13}\text{CO}_2$ was simulated by combining the apparent fractionation factors estimated from observations with the apparent CO_2 production simulated by the model.

A virtual clock tracer was used to calculate the mean age of air. However, previous research has shown that the speed of the Brewer–Dobson circulation in SOCRATES is fast, leading to an underestimation of age of air. Therefore, similar to Sugawara et al. (2018), we reduced the mass stream function to approximate a realistic mean age, which also improved the reproduction of GS. The model results of GS reproduced well the almost linear decrease with increasing altitude above the tropopause in the observational results (Fig. A1). The model calculations were carried out over the 25 yr from 1995 to 2020. The monthly average meridional distribution of the mean air age, GS, and CH_4 mole fraction calculated by SOCRATES for August 2016 is shown in Fig. B1.




Figure B1. (a) Monthly average meridional distributions of the mean age of air for August 2016 calculated using the SOCRATES model. (b) As in (a), but for gravitational separation, $\langle \delta_G \rangle$. Values lower than -150 per meg are shown in gray. (c) As in (a), but for the CH_4 mole fraction.

Appendix C: Age calculation

We estimated the mean age of air from $\delta^{13}\text{C}_\text{P}$ using the same method that was previously applied to the CO_2 age estima-

tion (Ray et al., 2017; Sugawara et al., 2018). The calculation of CO_2 age based on our balloon observations has been partly reported in previous studies (Engel et al., 2002; Ishidoya et al., 2013; Umezawa et al., 2025). The CO_2 ages estimated using air samples collected from 1995 to 2010 were published as supplementary data in Ishidoya et al. (2013). However, the method of age estimation was subsequently updated in Sugawara et al. (2018). In this study, we further improved the estimation method by correcting the $\delta^{13}\text{C}_\text{P}$ and CO_2 mole fraction data for GS, creating a new 35 yr record. We estimated the mean age of air using the convolution method (Ray et al., 2017; Leedham Elvidge et al., 2018; Sugawara et al., 2018). This method has been described in detail by Sugawara et al. (2018), so only a brief explanation is given here. Hypothetical age spectra were used in the convolution method to estimate the mean age (Waugh and Hall, 2002). Expected temporal variations in the age tracer in the stratosphere, $x(\Gamma, t)$, were calculated by convolution of the tropospheric reference curve, $x_0(t)$, and the age spectrum, $G(\Gamma, t)$:

$$x(\Gamma, t) = \int_{t-T_B}^t x_0(t') G(\Gamma, t-t') dt', \quad (\text{C1})$$

where T_B is the integration time interval. T_B should theoretically be ∞ , but we truncated it to 20 yr for the calculations. Actual age spectra, $G(\Gamma, t)$, are usually unknown. Therefore, we used the inverse Gaussian distribution (Waugh and Hall, 2002):

$$G(\Gamma, t) = \left(\frac{\Gamma^3}{4\pi \Delta^2 t^3} \right)^{1/2} \exp \left[\frac{-\Gamma(t-\Gamma)^2}{4\Delta^2 t} \right]. \quad (\text{C2})$$

Here, Δ denotes the width of the age spectrum, and it was parameterized using the mean age (Γ), i.e., the ratio of moments, Δ^2/Γ . This value has usually been assumed to be 0.7 yr, as suggested by Hall and Plumb (1994) based on results from a general circulation model calculation, and it was used to estimate SF_6 - and CO_2 -derived mean ages in the northern mid- and high-latitude stratosphere (Engel et al., 2002). Recently, however, Fritsch et al. (2020) reported that a value of 1.25 yr is better for estimating ages from the SF_6 mole fraction. Therefore, we used 1.25 yr for the ratio of moments in this study. After calculating the convolutions, the mean age was determined by substituting the observed values, x_{obs} , into the inverse function, $\Gamma(x, t)$. The convolutions are shown in Fig. C1 along with the calculated $\delta^{13}\text{C}_\text{P}$ values.

It should be noted that uncertainties of individual mole fraction or isotopic ratio measurements, as well as of the tropospheric reference record, lead to uncertainties of individual age estimates (e.g., Leedham Elvidge et al., 2018). Umezawa et al. (2025) estimated a total uncertainty of 0.72 yr for our CO_2 ages. Similarly, we estimated the uncertainty for $\delta^{13}\text{C}_\text{P}$ ages by adding normal pseudo-random numbers to all measured values and then calculating $\delta^{13}\text{C}_\text{P}$ values using Eqs. (9)




Figure C1. $\delta^{13}\text{C}_p$ observed over Japan (circles) and convolutions of $\delta^{13}\text{C}_p$ (colored lines) calculated from the tropospheric reference record and age spectrum. Observation altitudes of $\delta^{13}\text{C}_p$ are indicated by the gray scale of the symbol colors.

and (10). The uncertainty of the individual $\delta^{13}\text{C}_p$ values was nearly identical to that of $\delta^{13}\text{C}$, $\pm 0.02\text{‰}$. On the other hand, the uncertainty of the tropospheric reference record for $\delta^{13}\text{C}_p$ was calculated to be $\pm 0.03\text{‰}$. This value is in close agreement with the standard deviation obtained by applying curve fitting to the MLO $\delta^{13}\text{C}$ data used as the tropospheric reference record. Taking into account all these uncertainties and the tropospheric trend of -0.026‰ yr^{-1} , the total uncertainty of the individual $\delta^{13}\text{C}_p$ ages was estimated to be $\pm 1.9\text{ yr}$. The ratio of moments and its influence on age estimates have been examined in previous studies (e.g., Hauck et al., 2019; Nguyen et al., 2021). In this study, the mean age calculation was repeated as a sensitivity test for different ratios of moments ranging from 0.05 to 2.00 yr. As a result, the mean ages derived from $\delta^{13}\text{C}_p$ were found to be within the observational uncertainties of the $\delta^{13}\text{C}_p$ age. The $\delta^{13}\text{C}_p$ age is larger than the CO_2 age ($4.4 \pm 0.6\text{ yr}$) by about 1.1 yr on average. In this regard, $\delta^{13}\text{C}_T$ was calculated from the observed CH_4 mole fraction and its $\delta^{13}\text{C}$, assuming a closed system (Eq. (9)). However, actual chemical processes do not occur in a closed system, and atmospheric mixing processes always result in apparent fractionation being smaller than true fractionation (Rahn et al., 1998; Kaiser et al., 2006; Toyoda et al., 2018). In this study, the isotopic effect of CH_4 oxidation was calculated based on the apparent fractionation factor. This would result in an underestimation of the CH_4 oxidation effect in Eq. (9) and an overestimation of the $\delta^{13}\text{C}_p$ age. To solve this problem, it is necessary to explicitly incorporate the isotope effect of CH_4 into the model, which will be a future challenge.

Corresponding to the secular decrease of $\delta^{13}\text{CO}_2$, an increase in the CO_2 mole fraction was also clearly observed in the mid-stratosphere. To compare the temporal variations of the CO_2 mole fraction in the mid-stratosphere with those




Figure C2. Average values of mid-stratospheric CO_2 mole fraction ($>24\text{ km}$ altitude) over Japan (closed squares). Monthly average CO_2 mole fractions at Mauna Loa observed by NOAA/GML are shown by the dotted line. Quadratic functions fitted to the mid-stratospheric data (dashed line) and annual average MLO data from 1980 to 2015 (dashed-dotted line) are also shown.

in the troposphere, the average values above 24 km altitude were calculated for each year, shown in Fig. C2 alongside the monthly average data obtained by NOAA/GML at MLO. The results of the sixth WMO/IAEA Round Robin Comparison Experiment showed that our CO_2 mole fraction values were approximately 0.2 µmol mol^{-1} higher than the NOAA/GML values on the WMO- CO_2 -X2007 scale; accordingly, our CO_2 mole fraction values are plotted in Fig. C2 only after subtracting 0.2 µmol mol^{-1} . The CO_2 mole fraction in the mid-stratosphere over Japan increased monotonically and reached about 400 µmol mol^{-1} in 2020, lagging behind that at MLO by about 5 yr. The average rate of change of the CO_2 mole fraction in the mid-stratosphere was calculated to be $1.68 \pm 0.04\text{ µmol mol}^{-1}\text{ yr}^{-1}$ by applying the least-squares method to the observed values. However, this increasing trend was not linear, gradually steepening over the last 40 yr. Therefore, a quadratic function of the form $n(\text{CO}_2) = K_0 + (K_1 \times t) + (K_2 \times t^2)$ would better represent the observed variations. Here, t (years) is the elapsed time since 1980. The coefficients of the CO_2 mole fraction in the mid-stratosphere, K_0 , K_1 , and K_2 , were calculated to be 333.7, 1.0448, and 0.014755, respectively. The standard deviation of residuals was 0.8 µmol mol^{-1} . The same procedure was applied to the annual mean CO_2 mole fractions at MLO for the period 1980–2015, considering a time lag of 5 yr between the mid-stratosphere and the troposphere. The coefficients K_0 , K_1 , and K_2 for the MLO data were calculated to be 338.85, 1.2307, and 0.014577, respectively. As shown in Fig. C2, the two quadratic functions thus obtained agreed well to within $\pm 0.7\text{ µmol mol}^{-1}$. This result suggests that even temporal changes in the rate of increase in the tro-




Figure C3. CO_2 mole fractions observed over Japan (circles) and convolutions (colored lines) calculated from the tropospheric reference record and age spectrum. Observation altitudes of the CO_2 mole fraction are indicated by the gray scale of the symbol colors. Note that the CO_2 mole fractions plotted have been corrected for CH_4 oxidation and GS.

posphere propagate to the mid-stratosphere with a certain time delay.

As described in the main text, the influences on CO_2 production of CH_4 oxidation and GS should be considered to precisely estimate the CO_2 age. Therefore, the CO_2 mole fraction was corrected as follows. At first, the tropical upper tropospheric CH_4 data, $x_{\text{CH}_4\text{-tp}}(t)$, were created by adjusting the annual mean data at MLO to fit the CONTRAIL data in the tropical upper troposphere (Umezawa et al., 2012). Then, we corrected x_{obs} as

$$x_{\text{cor}}(t) = x_{\text{obs}}(t) - [x_{\text{CH}_4\text{-tp}}(t - \Gamma) - x_{\text{CH}_4\text{-obs}}(t)] - \Delta_{\text{G}}(x_{\text{obs}}(t)), \quad (\text{C3})$$

where x_{cor} and $x_{\text{CH}_4\text{-obs}}$ denote the corrected CO_2 mole fraction and observed CH_4 mole fraction, respectively. Δ_{G} is the correction for GS (see Sect. 3.3.). The CO_2 mole fractions corrected for CH_4 oxidation and GS are shown with the convolutions in Fig. C3. Finally, the CO_2 age was determined as $\Gamma_{\text{CO}_2} = \Gamma(x_{\text{cor}}, t)$. We note that Eq. (C3) contains the mean age of air, Γ , because the CH_4 mole fraction in the troposphere has increased with time and CH_4 is destroyed over the air transport pathway from the tropical upper troposphere to the observation altitude in the stratosphere. Therefore, Γ_{CO_2} was solved iteratively by starting from $\Gamma_{\text{CO}_2} = 0$. This iteration converged sufficiently after the second time of Γ_{CO_2} calculation. The vertical profiles of Γ_{CO_2} , corresponding to CO_2 age, are shown in Fig. C4.




Figure C4. Vertical profiles of the CO_2 age calculated from CO_2 mole fractions from 1985 to 2020. The monthly average vertical profile of the mean age of air for August 2016 calculated using the SOCRATES model is shown by the dashed line.

Data availability. The observational data obtained by our balloon measurements are included as a Supplement to the paper.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/acp-25-11895-2025-supplement>.

Author contributions. SS designed the study, conducted the measurements of mole fractions of greenhouse gases, and drafted the manuscript. SM, TN, SA, and SS conducted the measurements of carbon isotopic ratios of CO_2 . SM, SS, and TU conducted the measurements of carbon isotopic ratios of CH_4 . SI conducted the measurements of isotopic and elemental ratios of atmospheric major compositions. KI conducted the measurements of N_2O mole fraction. HH, TN, SA, SM, and SS conducted the balloon experiments. SI, ST, and DG participated in balloon experiments. All authors approved the final paper.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes ev-

ery effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. We deeply thank the Scientific Ballooning (DAIKIKYU) Research and Operation Group of the Institute of Space and Astronautical Science (ISAS), JAXA, Japan. We also deeply thank Sylvia E. Michel, Kerstin Braun, Justin Reid Clark, and John Ortega at INSTAAR, University of Colorado, and John Mund at NOAA GML for providing us their isotopic data.

Financial support. This study was supported by Japan Society for the Promotion of Science KAKENHI grants (grant nos. 22H05006, 23H00513, and 23K11396).

Review statement. This paper was edited by Gabriele Stiller and reviewed by two anonymous referees.

References

- Aoki, S., Nakazawa, T., Machida, T., Sugawara, S., Morimoto, S., Hashida, G., Yamanouchi, T., Kawamura, K., and Honda, H.: Carbon dioxide variations in the stratosphere over Japan, Scandinavia and Antarctica, *Tellus B*, 55, 178–186, <https://doi.org/10.1034/j.1600-0889.2003.00059.x>, 2003.
- Assonov, S. S. and Brenninkmeijer, C. A. M.: A redetermination of absolute values for 17RVPDB- CO_2 and 17RVSMOW, *Rapid Commun Mass Spectrom.*, 17, 1017–1029, <https://doi.org/10.1002/rcm.10111>, 2003.
- Assonov, S. S., Brenninkmeijer, C. A. M., Schuck, T. J., and Taylor, P.: Analysis of ^{13}C and ^{18}O isotope data of CO_2 in CARIBIC aircraft samples as tracers of upper troposphere/lower stratosphere mixing and the global carbon cycle, *Atmos. Chem. Phys.*, 10, 8575–8599, <https://doi.org/10.5194/acp-10-8575-2010>, 2010.
- Belikov, D., Sugawara, S., Ishidoya, S., Hasebe, F., Maksyutov, S., Aoki, S., Morimoto, S., and Nakazawa, T.: Three-dimensional simulation of stratospheric gravitational separation using the NIES global atmospheric tracer transport model, *Atmos. Chem. Phys.*, 19, 5349–5361, <https://doi.org/10.5194/acp-19-5349-2019>, 2019.
- Banks, P. M. and Kockarts, G.: *Aeronomy, Parts A and B*, Academic Press, Inc. New York, 1973.
- Bender, M., Sowers, T., and Labeyrie, L.: The Dole effect and its variations during the last 130,000 years as measured in the Vostok ice core, *Global Biogeochem. Cy.*, 8, 363–376, 1994.
- Bergamaschi, P., Hein, R., Brenninkmeijer, C. A. M., and Crutzen P. J.: Inverse modeling of the global CO cycle: 2. Inversion of $^{13}\text{C}/^{12}\text{C}$ and $^{18}\text{O}/^{16}\text{O}$ isotope ratios, *J. Geophys. Res.*, 105, 1929–1945, <https://doi.org/10.1029/1999JD900819>, 2000.
- Birner, B., Chipperfield, M. P., Morgan, E. J., Stephens, B. B., Linz, M., Feng, W., Wilson, C., Bent, J. D., Wofsy, S. C., Severinghaus, J., and Keeling, R. F.: Gravitational separation of Ar/N_2 and age of air in the lowermost stratosphere in airborne observations and a chemical transport model, *Atmos. Chem. Phys.*, 20, 12391–12408, <https://doi.org/10.5194/acp-20-12391-2020>, 2020.
- Brenninkmeijer, C. A. M.: Measurement of the abundance of ^{14}CO in the atmosphere and the $^{13}\text{C}/^{12}\text{C}$ and $^{18}\text{O}/^{16}\text{O}$ ratio of atmospheric CO , with application in New-Zealand and Antarctica, *J. Geophys. Res.*, 98, 10595–10614, 1993.
- Brenninkmeijer, C. A. M., Müller, R., Crutzen, P. J., Lowe, D. C., Manning, M. R., Sparks, R. J., and Velthoven, P. J. V.: A large ^{13}C deficit in the lower Antarctic stratosphere due to “ozone hole” chemistry: part 1, observations, *Geophys. Res. Lett.*, 23, 2125–2128, <https://doi.org/10.1029/96GL01471>, 1996.
- Butchart, N.: The Brewer-Dobson circulation, *Rev. Geophys.*, 52, 157–184, <https://doi.org/10.1002/2013RG000448>, 2014.
- Coplen, T. B. and Shrestha, Y.: Isotope-abundance variations and atomic weights of selected elements: 2016 (IUPAC Technical Report), *Pure Appl. Chem.*, 88, 1203–1224, <https://doi.org/10.1515/pac-2016-0302>, 2016.
- Engel, A., Strunk, M., Müller, M., Haase, H.-P., Poss, C., Levin, I., and Schmidt, U.: Temporal development of total chlorine in the high-latitude stratosphere based on reference distributions of mean age derived from CO_2 and SF_6 , *J. Geophys. Res.*, 107, <https://doi.org/10.1029/2001JD000584>, 2002.
- Engel, A., Möbius, T., Bönisch, H., Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S., Nakazawa, T., Sugawara, S., Moore, F., Hurst, D., Elkins, J., Schauffler, S., Andrews, A., and Boering, K.: Age of stratospheric air unchanged within uncertainties over the past 30 years, *Nat. Geosci.*, 2, 28–31, <https://doi.org/10.1038/Ngeo388>, 2009.
- Engel, A., Bönisch, H., Ullrich, M., Sitals, R., Membrive, O., Danis, F., and Crevoisier, C.: Mean age of stratospheric air derived from AirCore observations, *Atmos. Chem. Phys.*, 17, 6825–6838, <https://doi.org/10.5194/acp-17-6825-2017>, 2017.
- Francey, R. J., Tans, P. P., Allison, C. E., Enting, I. G., White, J. W. C., and Trolier, M.: Changes in oceanic and terrestrial carbon uptake since 1982, *Nature*, 373, 326–330, <https://doi.org/10.1038/373326a0>, 1995.
- Fritsch, F., Garny, H., Engel, A., Bönisch, H., and Eichinger, R.: Sensitivity of age of air trends to the derivation method for non-linear increasing inert SF_6 , *Atmos. Chem. Phys.*, 20, 8709–8725, <https://doi.org/10.5194/acp-20-8709-2020>, 2020.
- Gamo, T., Tsutsumi, M., Sakai, H., Nakazawa, T., Tanaka, M., Honda, H., Kubo, H., and Itoh, T.: Carbon and oxygen isotopic ratios of carbon dioxide of a stratospheric profile over Japan, *Tellus B*, 41, 127–133, <https://doi.org/10.1111/j.1600-0889.1989.tb00130.x>, 1989.
- Gamo, T., Tsutsumi, M., Sakai, H., Nakazawa, T., Machida, T., Honda, H., and Itoh, T.: Long-term monitoring of carbon and oxygen isotope ratios of stratospheric CO_2 over Japan, *Geophys. Res. Lett.*, 22, 397–400, 1995.
- García, R. R., López-Puertas, M., Funke, B., Marsh, D. R., Kinnison, D. E., Smith, A. K., and González-Galindo, F.: On the distribution of CO_2 and CO in the mesosphere and lower thermosphere, *J. Geophys. Res.-Atmos.*, 119, 5700–5718, <https://doi.org/10.1002/2013JD021208>, 2014.
- Garny, H., Ploeger, F., Abalos, M., Bönisch, H., von Clarmann, T., Diallo, M., Engel, A., Laube, J. C., Linz, M., Neu, J. L., Podglajen, A., Ray, E., Rivoire, L., Saunders, L. N., Stiller, G., Voet, F., Wagenhäuser, T., and Walker, K. A.: Age of stratospheric air: Progress on processes, observations, and long-term trends, *Rev. Geophys.*, 62, e2023RG000832, <https://doi.org/10.1029/2023RG000832>, 2024.

- Goto, D., Morimoto, S., Ishidoya, S., Aoki, S., and Nakazawa, T.: Terrestrial biospheric and oceanic CO_2 uptakes estimated from long-term measurements of atmospheric CO_2 mole fraction, $\delta^{13}\text{C}$, and $\delta(\text{O}_2/\text{N}_2)$ at Ny-Ålesund, Svalbard, *J. Geophys. Res.-Biogeosci.*, 122, 1192–1202, <https://doi.org/10.1002/2017JG003845>, 2017.
- Gromov, S., Brenninkmeijer, C. A. M., and Jöckel, P.: A very limited role of tropospheric chlorine as a sink of the greenhouse gas methane, *Atmos. Chem. Phys.*, 18, 9831–9843, <https://doi.org/10.5194/acp-18-9831-2018>, 2018.
- Hall, T. M. and Plumb, R. A.: Age as a diagnostic of stratospheric transport, *J. Geophys. Res.*, 99, 1059–1070, 1994.
- Hauck, M., Fritsch, F., Garny, H., and Engel, A.: Deriving stratospheric age of air spectra using an idealized set of chemically active trace gases, *Atmos. Chem. Phys.*, 19, 5269–5291, <https://doi.org/10.5194/acp-19-5269-2019>, 2019.
- Honda, H., Aoki, S., Nakazawa, T., Morimoto, S., and Yajima, N.: Cryogenic air sampling system for measurements of the concentrations of stratospheric trace gases and their isotopic ratios over Antarctica, *J. Geomagn. Geoelectr.*, 48, 1145–1155, 1996.
- Huang, T., Walters, S., Brasseur, G., Hauglustaine, D., and Wu, W.: Description of SOCRATES – A chemical dynamical radiative two-dimensional model, NCAR/TN-440+EDD NCAR TECHNICAL NOTE, 1998.
- Ishidoya, S. and Murayama, S.: Development of high precision continuous measuring system of the atmospheric O_2/N_2 and Ar/N_2 ratios and its application to the observation in Tsukuba, Japan, *Tellus*, 66B, 22574, <https://doi.org/10.3402/tellusb.v66.22574>, 2014.
- Ishidoya, S., Sugawara, S., Hashida, G., Morimoto, S., Aoki, S., Nakazawa, T., and Yamanouchi, T.: Vertical profiles of the O_2/N_2 ratio in the stratosphere over Japan and Antarctica, *Geophys. Res. Lett.*, 33, L13701, <https://doi.org/10.1029/2006GL025886>, 2006.
- Ishidoya, S., Sugawara, S., Morimoto, S., Aoki, S., and Nakazawa, T.: Gravitational separation of major atmospheric components of nitrogen and oxygen in the stratosphere, *Geophys. Res. Lett.*, 35, L03811, <https://doi.org/10.1029/2007GL030456>, 2008a.
- Ishidoya, S., Morimoto, S., Sugawara, S., Watai, T., Machida, T., Aoki, S., Nakazawa, T., and Yamanouchi, T.: Gravitational separation suggested by O_2/N_2 , $\delta^{15}\text{N}$ of N_2 , $\delta^{18}\text{O}$ of O_2 , Ar/N_2 observed in the lowermost part of the stratosphere at northern middle and high latitudes in the early spring of 2002, *Geophys. Res. Lett.*, 35, L03812, <https://doi.org/10.1029/2007GL031526>, 2008b.
- Ishidoya, S., Sugawara, S., Morimoto, S., Aoki, S., Nakazawa, T., Honda, H., and Murayama, S.: Gravitational separation in the stratosphere – a new indicator of atmospheric circulation, *Atmos. Chem. Phys.*, 13, 8787–8796, <https://doi.org/10.5194/acp-13-8787-2013>, 2013.
- Ishidoya, S., Sugawara, S., Inai, Y., Morimoto, S., Honda, H., Ikeda, C., Hashida, G., Machida, T., Tomikawa, Y., Toyoda, S., Goto, D., Aoki, S., and Nakazawa, T.: Gravitational separation of the stratospheric air over Syowa, Antarctica and its connection with meteorological fields, *Atmos. Sci. Lett.*, 19, e857, <https://doi.org/10.1002/asl.857>, 2018.
- Ishidoya, S., Sugawara, S., and Okazaki, A.: Diurnal, seasonal, and interannual variations in $\delta(^{18}\text{O})$ of atmospheric O_2 and its application to evaluate natural and anthropogenic changes in oxygen, carbon, and water cycles, *Atmos. Chem. Phys.*, 25, 1965–1987, <https://doi.org/10.5194/acp-25-1965-2025>, 2025.
- Ishijima, K., Sugawara, S., Kawamura, K., Hashida, G., Morimoto, S., Murayama, S., Aoki, S., and Nakazawa, T.: Temporal variations of the atmospheric nitrous oxide concentration and its $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ for the latter half of the 20th century reconstructed from firn air analyses, *J. Geophys. Res.*, 112, D03305, <https://doi.org/10.1029/2006JD007208>, 2007.
- Kaiser, J., Engel, A., Borchers, R., and Röckmann, T.: Probing stratospheric transport and chemistry with new balloon and aircraft observations of the meridional and vertical N_2O isotope distribution, *Atmos. Chem. Phys.*, 6, 3535–3556, <https://doi.org/10.5194/acp-6-3535-2006>, 2006.
- Kato, S., Kajii, Y., Akimoto, H., Bräunlich, M., Röckmann, T., and Brenninkmeijer, C. A. M.: Observed and modeled seasonal variation of ^{13}C , ^{18}O , and ^{14}C of atmospheric CO at Happo, a remote site in Japan, and a comparison with other records, *J. Geophys. Res.*, 105, 8891–8900, <https://doi.org/10.1029/1999JD901144>, 2000.
- Kawagucci, S., Tsunogai, U., Kudo, S., Nakagawa, F., Honda, H., Aoki, S., Nakazawa, T., Tsutsumi, M., and Gamo, T.: Long-term observation of mass-independent oxygen isotope anomaly in stratospheric CO_2 , *Atmos. Chem. Phys.*, 8, 6189–6197, <https://doi.org/10.5194/acp-8-6189-2008>, 2008.
- Keeling, C. D., Wholf, T. P., Wahlen, M., and van der Plicht, J.: Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980, *Nature*, 375, 666–670, <https://doi.org/10.1038/375666a0>, 1995.
- Khosravi, R., Brasseur, G., Smith, A., Rusch, D., Walters, S., Chabrilat, S., and Kockarts, G.: Response of the mesosphere to human-induced perturbations and solar variability calculated by a 2-D model, *J. Geophys. Res.*, 107, 4358, <https://doi.org/10.1029/2001JD001235>, 2002.
- Laube, J. C., Schuck, T. J., Chen, H., Geldenhuys, M., van Heuven, S., Keber, T., Popa, M. E., Tuffnell, E., Vogel, B., Wagenhäuser, T., Zanchetta, A., and Engel, A.: Vertical distribution of halogenated trace gases in the summer Arctic stratosphere determined by two independent in situ methods, *EGUSphere* [preprint], <https://doi.org/10.5194/egusphere-2024-4034>, 2025.
- Leedham Elvidge, E. C., Bönisch, H., Brenninkmeijer, C. A. M., Engel, A., Fraser, P. J., Gallacher, E., Langenfelds, R., Mühle, J., Oram, D. E., Ray, E. A., Ridley, A. R., Röckmann, T., Sturges, W. T., Weiss, R. F., and Laube, J. C.: Evaluation of stratospheric age of air from CF_4 , C_2F_6 , C_3F_8 , CHF_3 , HFC-125 , HFC-227ea and SF_6 ; implications for the calculations of halocarbon lifetimes, fractional release factors and ozone depletion potentials, *Atmos. Chem. Phys.*, 18, 3369–3385, <https://doi.org/10.5194/acp-18-3369-2018>, 2018.
- Luz, B. and Barkan, E.: The isotopic composition of atmospheric oxygen, *Global Biogeochem. Cy.*, 25, GB3001, <https://doi.org/10.1029/2010GB003883>, 2011.
- Luz, B., Barkan, E., Bender, M. L., Thieme, M. H., and Boering, K. A.: Triple-isotope composition of atmospheric oxygen as a tracer of biosphere productivity, *Nature*, 400, 547–550, <https://doi.org/10.1038/22987>, 1999.
- Machida, T., Matsueda, H., Sawa, Y., Nakagawa, Y., Hirokuni, K., Kondo, N., Goto, K., Nakazawa, N., Ishikawa, K., and Ogawa, T.: Worldwide measurements of atmospheric CO_2 and other trace gas species using commercial airlines, *J. Atmos. Ocean. Tech.*,

- 25, 1744–1754, <https://doi.org/10.1175/2008JTECHA1082.1>, 2008.
- Matsueda, H., Machida, T., Sawa, Y., and Niwa, Y.: Long-term change of CO_2 latitudinal distribution in the upper troposphere, *Geophys. Res. Lett.*, 42, <https://doi.org/10.1002/2014GL062768>, 2015.
- McCarthy, M. C., Boering, K. A., Rice, A. L., Tyler, S. C., Connell P., and Atlas, E.: Carbon and hydrogen isotopic compositions of stratospheric methane: 2. Two-dimensional model results and implications for kinetic isotope effects, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2002JD003183>, 2003.
- Meijer, H. A. J. and Li, W. J.: The use of electrolysis for accurate $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ isotope measurements in water, *Isotopes in Environmental and Health Studies*, 34, 349–369, <https://doi.org/10.1080/10256019808234072>, 1998.
- Michel, S., Braun, K., Ortega, J., Tans, P., Li, J., Baugh, K., Crotwell, M., Madronich, M., Moglia, E., Mund, J., NOAA Global Monitoring Laboratory, and University Of Colorado Institute Of Arctic And Alpine Research (INSTAAR): Monthly means of Stable Isotopic Composition of Atmospheric Carbon Dioxide ($\text{d}13\text{C}-\text{CO}_2$) from the NOAA GML Carbon Cycle Cooperative Global Air Sampling Network, 1990–2024, NOAA Global Monitoring Laboratory [data set], <https://doi.org/10.15138/6E9X-QA28>, 2025.
- Michel, S. E., Clark, J. R., Vaughn, B. H., Crotwell, M., Madronich, M., Moglia, E., Neff, D., and Mund, J., University of Colorado, Institute of Arctic and Alpine Research (INSTAAR), Stable Isotopic Composition of Atmospheric Methane (^{13}C) from the NOAA GML Carbon Cycle Cooperative Global Air Sampling Network, 1998–2022 Version: 2023-09-21 [data set], <https://doi.org/10.15138/9p89-1x02>, 2023.
- Minschwaner, K., Manney, G. L., Livesey, N. J., Pumphrey, H. C., Pickett, H. M., Froidevaux, L., Lambert, A., Schwartz, M. J., Bernath, P. F., Walker, K. A., and Boone, C. D.: The photochemistry of carbon monoxide in the stratosphere and mesosphere evaluated from observations by the Microwave Limb Sounder on the Aura satellite, *J. Geophys. Res.*, 115, D13303, <https://doi.org/10.1029/2009JD012654>, 2010.
- Morimoto, S., Nakazawa, T., Higuchi, K., and Aoki, S.: Latitudinal distribution of atmospheric CO_2 sources and sinks inferred by $\delta^{13}\text{C}$ measurements from 1985 to 1991, *J. Geophys. Res.*, 105, 24315–24326, <https://doi.org/10.1029/2000JD900386>, 2000.
- Morimoto, S., Aoki, S., Nakazawa T., and Yamanouchi, T.: Temporal Variations of the Carbon Isotopic Ratio of Atmospheric Methane Observed at Ny Ålesund, Svalbard from 1996 to 2004, *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2005GL024648>, 2006.
- Nakazawa, T., Morimoto, S., Aoki, S., and Tanaka, M.: Time and space variations of the carbon isotopic ratio of tropospheric carbon dioxide over Japan, *Tellus B*, 45, 258–274, <https://doi.org/10.3402/tellusb.v45i3.15728>, 1993.
- Nakazawa, T., Machida, T., Sugawara, S., Murayama, S., Morimoto, S., Hashida, G., Honda, H., and Itoh, T.: Measurements of the stratospheric carbon dioxide concentration over Japan using a balloon-borne cryogenic sampler, *Geophys. Res. Lett.*, 22, 1229–1232, <https://doi.org/10.1029/95GL01188>, 1995.
- Nakazawa, T., Morimoto, S., Aoki, S., and Tanaka, M.: Temporal and spatial variations of the carbon isotopic ratio of atmospheric carbon dioxide in the western Pacific region, *J. Geophys. Res.*, 102, 1271–1285, 1997a.
- Nakazawa, T., Ishizawa, M., Higuchi, K., and Trivett, N. B. A.: Two curve fitting methods applied to CO_2 flask data, *Environmetrics*, 8, 197–218, [https://doi.org/10.1002/\(SICI\)1099-095X\(199705\)8:3<197::AID-ENV248>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1099-095X(199705)8:3<197::AID-ENV248>3.0.CO;2-C), 1997b.
- Nakazawa, T., Aoki, S., Kawamura, K., Saeki, T., Sugawara, S., Honda, H., Hashida, G., Morimoto, S., Yoshida, N., Toyoda, S., Makide, Y., and Shirai, T.: Variations of stratospheric trace gases measured using a balloon-borne cryogenic sampler, *Adv. Space Res.*, 30, 1349–1357, [https://doi.org/10.1016/S0273-1177\(02\)00551-3](https://doi.org/10.1016/S0273-1177(02)00551-3), 2002.
- Nguyen, T. H., Ishijima, K., Sugawara, S., and Hasebe, F.: Application of a nudged general circulation model to the interpretation of the mean age of air derived from stratospheric samples in the tropics, *J. Meteorol. Soc. Jpn. II*, 99, 1149–1167, <https://doi.org/10.2151/jmsj.2021-056>, 2021.
- Park, J. H., Ko, M. K. W., Jackman, C. H., Plumb, R. A., Kaye, J. A., and Sage, K. H.: Models and Measurements Intercomparison II, NASA/TM-1999-209554, available at: <http://www.cs.odu.edu/~mln/ltrs-pdfs/NASA-99-tm209554.pdf> (last access: 19 July 2025), 1999.
- Rahn, T., Zhang, H., Wahlen, M., and Blake, G. A.: Stable isotope fractionation during ultraviolet photolysis of N_2O , *Geophys. Res. Lett.*, 25, 4489–4492, <https://doi.org/10.1029/1998GL900186>, 1998.
- Ray, E. A., Moore, F. L., Rosenlof, K. H., Davis, S. M., Sweeney, C., Tans, P., Wang, T., Elkins, J. W., Bönnisch, H., Engel, A., Sugawara, S., Nakazawa, T., and Aoki, S.: Improving stratospheric transport trend analysis based on SF_6 and CO_2 measurements, *J. Geophys. Res.-Atmos.*, 119, 14110–14128, <https://doi.org/10.1002/2014JD021802>, 2014.
- Ray, E. A., Moore, F. L., Elkins, J. W., Rosenlof, K. H., Laube, J. C., Röckmann, T., Marsh, D. R., and Andrews, A. E.: Quantification of the SF_6 lifetime based on mesospheric loss measured in the stratospheric polar vortex, *J. Geophys. Res.-Atmos.*, 122, 4626–4638, <https://doi.org/10.1002/2016JD026198>, 2017.
- Rice, A. L., Tyler, S. C., McCarthy, M. C., Boering K. A., and Atlas, E.: Carbon and hydrogen isotopic compositions of stratospheric methane: 1. High-precision observations from the NASA ER-2 aircraft, *J. Geophys. Res.*, 108, 4460, <https://doi.org/10.1029/2002JD003042>, 2003.
- Röckmann, T. and Brenninkmeijer, C. A. M.: CO and CO_2 isotopic composition in Spitsbergen during the 1995 ARCTOC campaign, *Tellus*, 49B, 455–465, 1997.
- Röckmann, T., Brenninkmeijer, C. A. M., Crutzen, P. J., and Platt, U.: Short-term variations in the $^{13}\text{C}/^{12}\text{C}$ ratio of CO as a measure of Cl activation during tropospheric ozone depletion events in the Arctic, *J. Geophys. Res.*, 104, 1691–1697, <https://doi.org/10.1029/1998JD100020>, 1999.
- Röckmann, T., Jöckel, P., Gros, V., Bränlich, M., Possnert, G., and Brenninkmeijer, C. A. M.: Using ^{14}C , ^{13}C , ^{18}O and ^{17}O isotopic variations to provide insights into the high northern latitude surface CO inventory, *Atmos. Chem. Phys.*, 2, 147–159, <https://doi.org/10.5194/acp-2-147-2002>, 2002.
- Röckmann, T., Brass, M., Borchers, R., and Engel, A.: The isotopic composition of methane in the stratosphere: high-altitude balloon sample measurements, *Atmos. Chem. Phys.*, 11, 13287–13304, <https://doi.org/10.5194/acp-11-13287-2011>, 2011.

- Röckmann T., van Herpen, M., Brashear, C., van der Veen, C., Grovov, S., Li, Q., Saiz-Lopez, A., Meidan, D., Barreto, A., Prats, N., Mármol, I., Ramos, R., Baños, I., Arrieta, J. M., Zahnle, S., Jordan, A., Moossen, H., Timas, H., Young, D., Sperlich, P., Moss, R., and Johnson, M. S.: The use of $\delta^{13}\text{C}$ in CO to determine removal of CH_4 by Cl radicals in the atmosphere, *Environ. Res. Lett.*, 19, 064054, <https://doi.org/10.1088/1748-9326/ad4375>, 2024.
- Santrock, J., Studley, S. A., and Hayes, J. M.: Isotopic analyses based on the mass spectra of carbon dioxide, *Anal. Chem.*, 57, 1444–1448, <https://doi.org/10.1021/ac00284a060>, 1985.
- Saueressig, G., Bergamaschi, P., Crowley, J. N., and Fischer, H.: Carbon KIE in the reaction of CH_4 with Cl atoms, *Geophys. Res. Lett.*, 22, 1225–1228, 1995.
- Saueressig, G., Crowley, J. N., Bergamaschi, P., Brühl, C., Brenninkmeijer, C. A. M., and Fischer, H.: Carbon-13 and D kinetic isotope effects in the reactions of CH_4 with $\text{O}(^1\text{D})$ and OH: New laboratory measurements and their implications for the isotopic composition of stratospheric methane, *J. Geophys. Res.*, 106, 23127–23138, 2001.
- Sawa, Y., Machida, T., and Matsueda, H.: Seasonal variations of CO_2 near the tropopause observation by commercial aircraft, *J. Geophys. Res.*, 113, D23301, <https://doi.org/10.1029/2008JD010568>, 2008.
- Stevens, C. M. and Wagner, A. F.: The Role of isotope fractionation effects in atmospheric chemistry, *Zeitschrift für Naturforschung A*, 44, 376–384, <https://doi.org/10.1515/zna-1989-0505>, 1989.
- Stiller, G. P., von Clarmann, T., Höpfner, M., Glatthor, N., Grabowski, U., Kellmann, S., Kleinert, A., Linden, A., Milz, M., Reddmann, T., Steck, T., Fischer, H., Funke, B., López-Puertas, M., and Engel, A.: Global distribution of mean age of stratospheric air from MIPAS SF_6 measurements, *Atmos. Chem. Phys.*, 8, 677–695, <https://doi.org/10.5194/acp-8-677-2008>, 2008.
- Stiller, G. P., von Clarmann, T., Haenel, F., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., and López-Puertas, M.: Observed temporal evolution of global mean age of stratospheric air for the 2002 to 2010 period, *Atmos. Chem. Phys.*, 12, 3311–3331, <https://doi.org/10.5194/acp-12-3311-2012>, 2012.
- Sugawara, S., Nakazawa, T., Shirakawa, Y., Kawamura, K., Aoki, S., Machida, T., and Honda, H.: Vertical profile of the carbon isotopic ratio of stratospheric methane over Japan, *Geophys. Res. Lett.*, 24, 2989–2992, 1997.
- Sugawara, S., Ishidoya, S., Aoki, S., Morimoto, S., Nakazawa, T., Toyoda, S., Inai, Y., Hasebe, F., Ikeda, C., Honda, H., Goto, D., and Putri, F. A.: Age and gravitational separation of the stratospheric air over Indonesia, *Atmos. Chem. Phys.*, 18, 1819–1833, <https://doi.org/10.5194/acp-18-1819-2018>, 2018.
- Thiemens, M. H.: Mass-independent isotope effects in planetary atmospheres and the early solar system, *Science*, 283, 341–345, <https://doi.org/10.1126/science.283.5400.341>, 1999.
- Toyoda, S., Yoshida, N., Urabe, T., Aoki, S., Nakazawa, T., Sugawara, S., and Honda, H.: Fractionation of N_2O isotopomers in the stratosphere, *J. Geophys. Res.*, 106, 7515–7522, <https://doi.org/10.1029/2000JD900680>, 2001.
- Toyoda, S., Yoshida, N., Morimoto, S., Aoki, S., Nakazawa, T., Sugawara, S., Ishidoya, S., Uematsu, M., Inai, Y., Hasebe, F., Ikeda, C., Honda, H., and Ishijima, K.: Vertical distributions of N_2O isotopocules in the equatorial stratosphere, *Atmos. Chem. Phys.*, 18, 833–844, <https://doi.org/10.5194/acp-18-833-2018>, 2018.
- Umezawa, T., Machida, T., Ishijima, K., Matsueda, H., Sawa, Y., Patra, P. K., Aoki, S., and Nakazawa, T.: Carbon and hydrogen isotopic ratios of atmospheric methane in the upper troposphere over the Western Pacific, *Atmos. Chem. Phys.*, 12, 8095–8113, <https://doi.org/10.5194/acp-12-8095-2012>, 2012.
- Umezawa, T., Brenninkmeijer, C. A. M., Röckmann, T., van der Veen, C., Tyler, S. C., Fujita, R., Morimoto, S., Aoki, S., Sowers, T., Schmitt, J., Bock, M., Beck, J., Fischer, H., Michel, S. E., Vaughn, B. H., Miller, J. B., White, J. W. C., Brailsford, G., Schaefer, H., Sperlich, P., Brand, W. A., Rothe, M., Blunier, T., Lowry, D., Fisher, R. E., Nisbet, E. G., Rice, A. L., Bergamaschi, P., Veidt, C., and Levin, I.: Interlaboratory comparison of $\delta^{13}\text{C}$ and $\delta^{13}\text{C}$ and δD measurements of atmospheric CH_4 for combined use of data sets from different laboratories, *Atmos. Meas. Tech.*, 11, 1207–1231, <https://doi.org/10.5194/amt-11-1207-2018>, 2018.
- Umezawa, T., Sugawara, S., Hikichi, S., Morimoto, S., Saito, T., Krummel, P. B., Fraser, P. J., and Weiss, R. F.: Evaluation of stratospheric age of air estimated from halocarbon measurements of air samples collected by a balloon-borne cryogenic air sampler over Japan, *SOLA*, 21, 237–243, <https://doi.org/10.2151/sola.2025-029>, 2025.
- Wang, J. S., McElroy, M. B., Spivakovsky, C. M., and Jones, D. B. A.: On the contribution of anthropogenic Cl to the increase in $\delta^{13}\text{C}$ of atmospheric methane, *Global Biogeochem. Cy.*, 16, 1047, <https://doi.org/10.1029/2001GB001572>, 2002.
- Waugh, D. W. and Hall, T. M.: Age of stratospheric air: Theory, observations, and models, *Rev. Geophys.*, 40, 1010, <https://doi.org/10.1029/2000RG000101>, 2002.