



The importance of stratocumulus clouds for projected warming patterns and circulation changes

Philipp Breul¹, Paulo Ceppi¹, and Peer Nowack^{2,3}

¹Department of Physics, Imperial College London, London, United Kingdom

²Institute of Theoretical Informatics, Karlsruhe Institute of Technology, 76131 Karlsruhe, Germany

³Institute of Meteorology and Climate Research (IMK-ASF), Karlsruhe Institute of Technology,
76131 Karlsruhe, Germany

Correspondence: Philipp Breul (pyb18@ic.ac.uk)

Received: 17 January 2025 – Discussion started: 3 February 2025

Revised: 17 July 2025 – Accepted: 17 July 2025 – Published: 2 October 2025

Abstract. Stratocumulus clouds are thought to exert a strong positive radiative feedback on climate change, but recent analyses suggest that this feedback is widely under-represented in global climate models. To assess the broader implications of this model error for the simulated climate change responses, we investigate the impact of Pacific stratocumulus cloud feedback on projected warming patterns, equilibrium climate sensitivity and tropical atmospheric circulation under increased CO₂ concentrations. Using the Community Earth System Model, with modifications to enhance low-cloud-cover sensitivity to sea surface temperature (SST) anomalies in Pacific stratocumulus regions, we find increased tropical SST variability and persistence, a higher equilibrium climate sensitivity, an enhanced east–west warming contrast across the tropical Pacific, and a stronger slowdown of the Walker circulation under 4×CO₂ conditions. Our findings are supported by inter-model relationships across CMIP6 4×CO₂ simulations. These results underscore the importance of accurately representing cloud feedback in climate models to predict future climate change impacts not only globally but also on a regional scale, such as warming patterns or circulation change.

1 Introduction

Clouds play a major role in shaping both our current climate and future climate change through their impact on incoming shortwave radiation as well as outgoing terrestrial longwave radiation. Any external forcing on the climate system can potentially change cloud properties, therefore feeding back to the global climate.

Cloud feedback has been studied extensively in terms of its implications for *global mean* surface temperature change, particularly the equilibrium climate sensitivity (ECS) (e.g. Zelinka et al., 2020; Zhu and Poulsen, 2020; Myers et al., 2021; Ceppi and Nowack, 2021). However, cloud feedback is potentially also key for the *pattern* of surface warming, influencing the local radiation budget and atmospheric circulation (e.g. Voigt and Shaw, 2015; Ceppi and Hartmann, 2016). This is all the more important as general circulation models

(GCMs) feature a wide spread of sensitivity of low clouds to sea surface temperatures (SSTs) in stratocumulus regions, with most GCMs underestimating this sensitivity relative to observations (Myers et al., 2021; Ceppi et al., 2024).

Previous studies have addressed the impact of clouds on the internal variability of climate modes and SST, e.g. the amplitude and variability of El Niño–Southern Oscillation (ENSO) (Bellomo et al., 2014; Rädcl et al., 2016; Middlemas et al., 2019; Yang et al., 2022) and decadal-scale ocean variability (Brown et al., 2016; Burgman et al., 2017; Hsiao et al., 2022). Clouds can influence SSTs through feedbacks locally as well as remotely through teleconnections – for example through the wind–evaporation–SST (WES) and Bjerknes feedbacks – which communicate subtropical cloud-induced SST variability into the deeper tropics (e.g. Hsiao et al., 2022; Kim et al., 2022). The impact of low clouds on the future CO₂-forced warming pattern and circulation change

has received less attention, with some studies finding a feedback contribution to warming in the East Pacific (Chalmers et al., 2022; Fu and Fedorov, 2023).

In this study, we test the climate impact of increasing stratocumulus cloud sensitivity to SST in a coupled climate model in the Northeast and Southeast Pacific subsidence regions. These are of special interest as they couple to major climate modes in the Pacific (Bellomo et al., 2014; Rädel et al., 2016; Yuan et al., 2018; Myers and Mechoso, 2020). Our focus is on understanding the impact of enhanced low-cloud sensitivity on projected warming patterns and resulting circulation change. Previous studies that analysed cloud impacts on abrupt CO₂ responses mostly used model experiments, with all clouds decoupled from the meteorology (often described as “cloud locking”); by contrast, we couple the Pacific stratocumulus clouds more strongly to the underlying SSTs, which is arguably more physical. The results provide insight into the potential global implications of model bias in stratocumulus cloud feedback.

2 Data and methods

2.1 Experimental setup

We use the Community Earth System Model version 2.1.3 (CESM2.1.3), with the atmosphere model CAM4 (Neale et al., 2010) and the ocean model POP2 (Danabasoglu et al., 2012). We use a T31 (3.8°) horizontal grid with 26 vertical levels for CAM4 and a 3° horizontal grid with 60 depth levels for POP2. Our choice of an already widely used model configuration with a relatively coarse spatial grid is aimed at computational efficiency when testing the effects of large-scale changes in cloud sensitivities to SSTs. Our primary goal is to qualitatively assess the importance of the role of clouds on coupled climate change, rather than achieving exact alignment with observations through our modifications to cloud sensitivities (see below).

Cloud sensitivities are calculated following the cloud-controlling-factor analysis framework of Ceppi and Nowack (2021) and Ceppi et al. (2024). In this framework, cloud sensitivities to controlling factors (shown in Fig. A2) are calculated as the coefficients of regularised ridge regression. Contrary to a classical multiple regression analysis, the regularisation of ridge regression allows us to include more (correlated) predictors, in this case from neighbouring grid cells, leading to improved sensitivity estimates. We direct the reader to Ceppi and Nowack (2021) for further detail.

In this CESM-CAM4 configuration, relative to observations, the sensitivity of low-cloud cover (LCC; a model output calculated using the maximum–random overlap assumption for all clouds at or lower than 700 hPa) to SST anomalies in Pacific subsidence regions is too low (Fig. A1). In parts of the subsidence regions, sensitivities even have the opposite sign compared to observations, causing the regional average to be close to zero (Fig. A2a). We thus modify the

low-cloud sensitivity to local SST by adding a perturbation at every radiation time step, proportional to the instantaneous SST anomaly to all clouds at or below 700 hPa. Importantly, this instantaneous perturbation is only applied to the cloud amount “seen” by the radiative transfer code, ensuring that any direct effect is only on the *radiative* properties of the cloud, and the perturbation is not carried over to the next time step. Instantaneous SST anomalies are calculated relative to the 450-year monthly climatology of the control simulation with unperturbed clouds, termed $1 \times \text{Orig}$ (Table 1). This setup is similar to that of Bellomo et al. (2014) and, to a lesser extent, Erfani and Burls (2019). Although CESM-CAM4 is biased in terms of its sensitivities to factors other than SST (Fig. A2a), the feedback is dominated by the SST contribution, so we target that.

The cloud amount perturbation magnitude is set to -3 percentage points of local cloud amount anomaly per degree of local SST anomaly. Note that this modification does not necessarily translate to a 3-percentage-point sensitivity decrease of LCC, due to the random overlap statistics. While the model previously significantly underestimated the LCC–SST sensitivity averaged over the subsidence regions compared to observations, the modifications lead to an overshoot in sensitivity (Fig. A2a). Although this large modification exaggerates the effect of model bias, it also provides a larger signal of the impact of enhanced cloud sensitivity. Sensitivities to other controlling factors are changed as well, although only by comparatively small amounts (Fig. A2a). The modifications are restricted to the Pacific subsidence regions as calculated from the ECMWF Reanalysis version 5 (ERA5; Hersbach et al., 2020), following Scott et al. (2020). The resulting regions are shown in Fig. A2b.

We use four different experimental setups, which are combinations of modified or unmodified cloud sensitivity and control or quadrupled CO₂ concentrations (Table 1). Comparing $1 \times \text{Orig}$ and $1 \times \text{Mod}$ allows us to analyse changes to internal variability. Comparing the $4 \times \text{CO}_2$ responses with unperturbed versus perturbed clouds enables us to estimate the impact of enhanced cloud sensitivities under greenhouse gas (GHG) forcings. We spun up the model for 1250 years to reach equilibrium, then branched off and spun up $1 \times \text{Orig}$ and $1 \times \text{Mod}$ for an additional 50 years. The $4 \times \text{CO}_2$ experiments ($4 \times \text{Orig}$ and $4 \times \text{Mod}$) include three 150-year ensemble members, branched off from their respective $1 \times \text{CO}_2$ simulations at 150-year intervals to ensure approximate independence.

2.2 CMIP6 data

In addition to our CESM-CAM4 experiments, we analyse monthly mean output from 22 models from the Coupled Model Intercomparison Project Phase 6 (CMIP6; Eyring et al., 2016), using the piControl and $4 \times \text{CO}_2$ experiments. The models were selected based on the availability of necessary data for calculating the cloud radiative sensitivity to SST anomalies, following the cloud-controlling-factor anal-

Table 1. Summary of the climate model experiments conducted.

Name	Cloud modifications	CO ₂	Length (years)	Ensemble members
1×Orig	No	1×CO ₂	450	1
1×Mod	Yes	1×CO ₂	450	1
4×Orig	No	4×CO ₂	150	3
4×Mod	Yes	4×CO ₂	150	3

ysis method of Ceppi and Nowack (2021). The models used are ACCESS-CM2, ACCESS-ESM1-5, BCC-CSM2-MR, BCC-ESM1, CESM2, CNRM-CM6-1, CNRM-ESM2-1, EC-Earth3-Veg, FGOALS-f3-L, GFDL-CM4, GISS-E2-1-G, HadGEM3-GC31-LL, INM-CM4-8, INM-CM5-0, IPSL-CM6A-LR, MIROC6, MIROC-ES2L, MPI-ESM1-2-HR, MPI-ESM1-2-LR, MRI-ESM2-0, NESM3 and UKESM1-0-LL.

2.3 Definitions of indices and warming patterns

We calculate the commonly used Niño3.4 index as a measure of El Niño–Southern Oscillation (ENSO) strength, defined as the deseasonalised SST anomaly box-averaged over (5° S, 5° N) latitude and (190° E, 240° E) longitude (e.g. Trenberth and Stepaniak, 2001). To assess the Walker circulation strength, we follow Vecchi et al. (2006), calculating the difference in surface pressure between two boxes: (5° S, 5° N) and (200, 280° E) in the equatorial East Pacific minus (5° S, 5° N) and (80, 160° E) over the Indo-Pacific warm pool. For the Walker circulation index in the 4×CO₂ experiments, we use anomalies relative to the corresponding 1×CO₂ experiments (noting that 1×Mod and 1×Orig have near-identical climatologies), as this allows us to compare between experiments and to evaluate the evolution in the 4×CO₂ experiments.

We calculate warming maps from the 4×CO₂ experiments by regressing local surface temperature onto global mean temperature. We will distinguish between fast, slow and total responses, calculated over the years 1–20, 21–150 and 1–150 since CO₂ quadrupling (as in e.g. Andrews et al., 2015; Rugenstein et al., 2016; Lin et al., 2019).

3 Results

We first discuss the results from the two 1×CO₂ experiments 1×Orig and 1×Mod, which allow us to analyse changes in internal variability stemming from the cloud sensitivity modifications. We will then turn our attention to the 4×CO₂ experiments 4×Orig and 4×Mod. Finally, we will analyse the extent to which the results from these experiments hold across CMIP6 models.

3.1 Mean state and internal variability

Although changes to the mean state and internal variability are not the primary focus of this paper, results from the 1×CO₂ experiments enable us to test whether the cloud modifications result in behaviour in line with both our physical understanding and findings from previous studies.

The climatological SSTs are almost unaffected between the 1×Orig and 1×Mod experiments (Fig. A3). While a few regions do show differences in SST, the corresponding anomalies are very small (~0.1 K) and are located in neither the subsidence regions nor the tropical Pacific. Changes in climatology are therefore unlikely to have a significant impact on the observed changes in internal variability discussed below.

The cloud modifications cause enhanced SST variance in the tropical Pacific and subtropical East Pacific (Fig. 1a). This behaviour is in line with our physical understanding of positive feedback between LCC and SSTs, which previous work established is not confined to local SST changes (e.g. Bellomo et al., 2014; Zhou et al., 2017; Erfani and Burls, 2019). Here, enhanced positive feedback between SST anomalies and LCC (locally higher SSTs lead to lower LCC, which increases the SSTs) leads to locally more variable SSTs. In the equatorial Pacific, this enhanced variability is advected westward by the trade winds along the cold tongue region and possibly enhanced further by the Bjerknes feedback (Kim et al., 2022; Fu and Fedorov, 2023).

Due to the enhanced LCC–SST feedback, we find a higher SST variability in the Niño3.4 region and also an increase in extreme ENSO events, which is reflected in heavier tails of the estimated probability density of the Niño3.4 index (Fig. 1b). Additionally, ENSO becomes more persistent and larger in amplitude, which is reflected in an amplitude increase and a shift to lower frequencies of the power spectrum (Fig. 1c). Increased power is also found at periods of around a decade, suggesting possible changes in the amplitude of decadal Pacific variability. The interdecadal Pacific oscillation power spectrum (not shown), on the other hand, shows reduced amplitude in response to enhanced low-cloud feedback, which points to a more complicated interaction on long timescales. We therefore leave a detailed investigation of such changes for future work. Overall, however, the findings shown in Fig. 1 are in agreement with the physical understanding of the enhanced feedback mechanisms detailed above.

The influence of clouds on ENSO variability has been documented in previous literature, with studies generally finding that clouds amplify ENSO variability and persistence (e.g. Rädel et al., 2016). The importance of low clouds specifically has been studied as well, with some studies highlighting the importance of Northeast Pacific low clouds affecting local SST variability, in turn modulating ENSO through the wind–evaporation–SST (WES) feedback mechanism (Yang et al., 2022); in contrast, other studies found low clouds of




Figure 1. Changes to surface temperature variability between the $1\times\text{Mod}$ and $1\times\text{Orig}$ experiments. **(a)** Differences in surface temperature standard deviations. The grey contours denote the subsidence regions where the LCC sensitivity to SST anomalies was enhanced in the Mod experiments. The black box shows the Niño3.4 region. **(b)** Estimated probability densities of the Niño3.4 index. **(c)** Power spectrum of the Niño3.4 index, estimated using the Welch method (Welch, 1967) with 44 20-year-long half-overlapping windows. The shaded areas show 95 % confidence intervals based on a χ^2 test.

the Southeast Pacific to be especially important (Bellomo et al., 2014). Similar cloud influences extend to tropical Atlantic SST variability (Bellomo et al., 2015; Brown et al., 2016). By contrast, Middlemas et al. (2019) found, through cloud-locking experiments, that clouds can decrease ENSO persistence. This finding might be related to our findings of reduced decadal persistence.

In summary, increasing East Pacific low-cloud sensitivities to SSTs increases SST variability and persistence both locally and remotely; this is reflected in a more persistent and variable ENSO. These results are in line with our physical understanding as well as with most of the previous literature.

3.2 Response to $4\times\text{CO}_2$ forcing

We now analyse the impact of increased cloud sensitivity on the coupled climate response to $4\times\text{CO}_2$ forcing. As a first step, we estimate the effective radiative forcing (ERF) and equilibrium climate sensitivity (ECS) in the two experiments ($4\times\text{Orig}$ and $4\times\text{Mod}$) by extrapolating the relationship between the global mean radiative imbalance and the global mean surface air temperature during the fast and slow response in each experiment (Fig. 2a; Gregory et al., 2004). We find that $\text{ERF}_{\text{Orig}} = 7.35 \text{ W m}^{-2}$ and $\text{ERF}_{\text{Mod}} = 7.26 \text{ W m}^{-2}$, $\text{ECS}_{\text{Orig}} = 6.25 \text{ K}$ and $\text{ECS}_{\text{Mod}} = 6.63 \text{ K}$. The low-cloud modification therefore leaves ERF essentially unchanged (as expected, given that our modification should

only affect the SST-mediated cloud response; the small decrease is most likely sampling bias, as the ERF ranges between $4\times\text{Orig}$ and $4\times\text{Mod}$ completely overlap, Fig. 2a), while enhancing ECS by approximately 6.2 %. This is in line with across-model relationships between ECS and cloud feedback (although not necessarily from Pacific stratocumulus clouds) that have been reported in previous studies (e.g. Zelinka et al., 2020; Ceppi and Nowack, 2021; Myers et al., 2021).

With the ERF unchanged, the change in ECS must come from climate feedback, which we calculate here as the slopes in Fig. 2a. Our expectation is for more positive (or less negative) cloud feedback and thus less negative net feedback. We find a 7 % decrease in net feedback magnitude in the fast $4\times\text{CO}_2$ response, in line with a higher ECS. The slow response only shows minimal change, however, suggesting that the fast response is the main driver of global mean warming differences. To assess the sources of the feedback differences, we perform a radiative kernel decomposition (Soden et al., 2008) of the differences in fast, slow and total feedback in the $4\times\text{Mod}$ and $4\times\text{Orig}$ experiments (Fig. 3). A caveat of the analysis is that the kernel method can only explain about half of the total feedback increase in the fast response (Fig. 3a). As expected, we find more positive cloud feedback in the fast, slow and total response. The enhanced cloud feedback is the largest contributor to the net feedback increase; it stays quantitatively the same in the fast, slow and




Figure 2. Ensemble mean $4\times\text{CO}_2$ responses as a function of global mean temperature anomaly. We show yearly averages for the first 20 years as dots and decadal averages as triangles over the complete 150 years, together with linear fits to both the fast and slow periods. We additionally plot slope m for the fast and slow components. (a) Gregory plot of the top-of-the-atmosphere radiative imbalance plotted against global mean temperature anomaly. The crosses show the mean projected equilibrium temperature on the x axis and the approximated effective radiative forcing on the y axis, with the vertical lines indicating the ensemble spread. (b) Anomalous Walker circulation strength index plotted against global mean temperature.

total response. Changes in lapse rate and albedo feedback between the fast and slow response lead to a much smaller net feedback in the slow response compared to the fast response. Analysing maps of the cloud feedback difference (Fig. 4a–c), we find a clear fingerprint of the contribution from the perturbed Pacific subsidence regions, which is also reflected in a very similar pattern in the LCC response difference (Fig. 4d–f).

In our experiments with modified stratocumulus sensitivity to SST, the global climate feedbacks respond not only to the imposed change in low-cloud sensitivity but also to any resulting changes in the SSTs themselves, both locally and potentially through a global “pattern effect” (Andrews and Webb, 2018; Dong et al., 2019; Myers et al., 2023), where warming patterns remotely alter tropospheric stability via circulation changes. To characterise the SST pattern response, we show the warming pattern maps of $4\times\text{Orig}$ for the fast, slow and total responses in Fig. 5a–c, respectively, and the difference in warming maps between $4\times\text{Mod}$ and $4\times\text{Orig}$ in Fig. 5d–f.

First considering the overall warming response in $4\times\text{Orig}$ (Fig. 5a–c), our experiments show an enhanced East Pacific warming compared to other tropical regions, amplified Arctic warming in the fast response and Antarctic amplification in the slow response. Differences between the fast and slow responses are likely attributable to ocean dynamical changes that are slow to manifest. Similar fast and slow responses to CO_2 increase have been observed in previous studies (e.g. Andrews et al., 2015; Ceppi et al., 2018).

Increasing the low-cloud sensitivities to SST results in a further enhancement of East Pacific warming (Fig. 5d–f).

These results are also in line with previous work: in an abrupt CO_2 -doubling experiment but with shortwave fluxes taken from a control simulation, Fu and Fedorov (2023) found a weakening of the east–west tropical Pacific warming contrast. This is in line with our physical expectation of an enhanced LCC–SST feedback causing a larger weakening of the east–west temperature difference in the modified experiments. Given that the impact of enhanced cloud sensitivity is spatially similar for the SST warming patterns compared to internal variability (Figs. 5d–f and 1a), we hypothesise that similar mechanisms are at play; i.e. local feedbacks between clouds and SSTs are communicated via WES and Bjerknes feedbacks to remote areas outside of the perturbed cloud regions.

A weakening of the Walker circulation in response to CO_2 forcing has been documented in prior literature (Vecchi et al., 2006; He and Soden, 2015; Nowack et al., 2017; Malik et al., 2020; Heede et al., 2020). Enhanced East Pacific warming should weaken the circulation even further since the resulting impact on convection counteracts the Walker circulation (e.g. Tokinaga et al., 2012). This is indeed what we find in Fig. 2b: the Walker circulation weakens under CO_2 forcing ($\sim -20(\pm 11) \text{ Pa K}^{-1}$ in the fast response and $\sim -9(\pm 6) \text{ Pa K}^{-1}$ in the slow response), but $4\times\text{Mod}$ shows a greater weakening compared to $4\times\text{Orig}$ ($\sim -34(\pm 16) \text{ Pa K}^{-1}$ and $\sim -14(\pm 5) \text{ Pa K}^{-1}$), with the brackets giving a 95 % confidence interval based on Newey–West standard error estimates. This corresponds to ~ 70 % additional reduction in the fast response and ~ 49 % additional reduction in the slow response, although the numbers should be interpreted with care, given the relatively small ensemble size and large vari-




Figure 3. Global mean feedback differences, decomposed following Soden et al. (2008) with relative humidity as a state variable (Held and Shell, 2012) and using seven different kernels, calculated using ClimKern (Janoski et al., 2025). Shown are the ensemble mean differences of the experiments $4\times\text{Mod}$ and $4\times\text{Orig}$ over the (a) fast, (b) slow and (c) total period. Calculated are the Planck, lapse rate (LR), relative humidity (RH), albedo and cloud feedbacks as well as their sum (net). The red cross shows the actual net feedback calculated from the global mean TOA radiative flux shown in Fig. 2a.




Figure 4. (a–c) Maps of the cloud feedback difference, calculated with the same method as Fig. 3. Shown are (a) fast, (b) slow and (c) total differences of cloud feedback. (d–f) Difference in ensemble mean low-cloud cover responses (regressed against global mean temperature) between the $4\times\text{Mod}$ and $4\times\text{Orig}$ experiments for the (d) fast, (e) slow and (f) total low-cover cloud response. Stippling shows where all combinations of ensemble member differences agree on the sign.

ability in the data. While the changes in the warming pattern are a plausible explanation for the Walker circulation weakening, additional contributions from other mechanisms, e.g. a reduction in cloud-top longwave radiative cooling due to the LCC decrease, are possible. We checked that differences in the global mean lapse rate change, which might cause Walker circulation changes (Knutson and Manabe, 1995), are small (not shown) and are therefore unlikely to be the reason for the additional reduction.

3.3 Direct and pattern-mediated low-cloud responses

We now interpret the low-cloud feedback changes resulting from the LCC sensitivity perturbation. Because our GCM experiments are fully coupled, changes in cloud feedback between $4\times\text{Orig}$ and $4\times\text{Mod}$ result not only from the modified LCC sensitivities but also from the coupled climate response to cloud feedback: modifying clouds change the SST warming pattern, which in turn modifies the clouds further. Thus, our applied LCC sensitivity perturbation can affect low-cloud feedback in two ways:




Figure 5. Warming patterns for (a) fast, (b) slow and (c) total response of the $4\times\text{Orig}$ experiments with 1 K K^{-1} (in units of local surface warming per global surface warming) subtracted. (d)–(f) As for (a)–(c), respectively, but for the difference in warming pattern between $4\times\text{Mod}$ and $4\times\text{Orig}$. The stippling shows the regions where the signs of all nine possible differences between the two ensembles agree on the sign of change. (g) Fast, (h) slow and (i) total CMIP6 across-model regression maps of warming patterns onto the CRE–SST Pacific cloud sensitivity index. The stippling shows a statistical significance of $p < 0.05$, based on a Student t test. Note that adjusting for multiple hypothesis testing following Wilks (2016) would remove any statistical significance.

- Through the imposed change in LCC sensitivity to SST;
- Through the change in the SST warming pattern and its impact on cloud controlling factors, henceforth the “pattern-mediated cloud response” or simply “pattern effect”.

To explain this further, we describe the total change in LCC as an expanded total derivative, following the cloud-controlling-factor analysis framework (e.g. Stevens and Brenguier, 2009; Myers et al., 2021):

$$\frac{dC}{dT} = \sum \frac{\partial C}{\partial Y_i} \frac{dY_i}{dT}, \quad (1)$$

with C being the low-cloud cover in either of the $4\times\text{CO}_2$ experiments, T being the corresponding global mean surface temperature and Y_i being the relevant controlling factors; all variables are defined locally (except for T), but we drop the location index for conciseness. We also neglect $\frac{\partial C}{\partial T}$ as usual, since T is a global mean quantity and therefore has no direct physical connection to local cloud cover.

We approximate the total derivatives in Eq. (1) by finite differences and obtain an approximate expression of LCC change,

$$\Delta C = \sum \frac{\partial C}{\partial Y_i} \Delta Y_i = \sum \theta_i \Delta Y_i, \quad (2)$$

with $\theta_i \equiv \frac{\partial C}{\partial Y_i}$ being the cloud sensitivities to controlling factors. For the experiments $4\times\text{Orig}$ and $4\times\text{Mod}$, we get, respec-

tively,

$$\Delta C_{\text{Orig}} = \sum \theta_{i,\text{Orig}} \Delta Y_{i,\text{Orig}}, \quad (3)$$

$$\Delta C_{\text{Mod}} = \sum \theta_{i,\text{Mod}} \Delta Y_{i,\text{Mod}} = \sum (\theta_{i,\text{Orig}} + \delta_i) \Delta Y_{i,\text{Mod}}, \quad (4)$$

with $\delta_i \equiv \theta_{i,\text{Mod}} - \theta_{i,\text{Orig}}$ being the difference in cloud sensitivities between the two experiments, by definition. Taking the difference of ΔC_{Mod} and ΔC_{Orig} , we obtain

$$\Delta C_{\text{Mod}} - \Delta C_{\text{Orig}} = \sum [(\theta_{i,\text{Orig}} + \delta_i) \Delta Y_{i,\text{Mod}} - \theta_{i,\text{Orig}} \Delta Y_{i,\text{Orig}}], \quad (5)$$

$$= \sum [\theta_{i,\text{Orig}} (\Delta Y_{i,\text{Mod}} - Y_{i,\text{Orig}}) + \delta_i \Delta Y_{i,\text{Orig}} + \delta_i (\Delta Y_{i,\text{Mod}} - \Delta Y_{i,\text{Orig}})]. \quad (6)$$

The first term on the right-hand side of Eq. (6) is the contribution from changes in the cloud-controlling factors while holding the sensitivities fixed, the second is from changes in the cloud sensitivities while holding the cloud-controlling factors fixed and the third is a cross term. Note that the sensitivity contribution (second term on the right-hand side of Eq. 6) will not necessarily be equal to the imposed change of 3 percentage points per K at each model level: cloud overlap statistics imply that the vertically integrated LCC sensitivity change should be slightly greater than this (not shown); here, the subsidence-averaged LCC sensitivity increases from $-0.2\% \text{ K}^{-1}$ to $-4.0\% \text{ K}^{-1}$.

We plot the simulated cloud response difference $\Delta C_{\text{Mod}} - \Delta C_{\text{Orig}}$ in Fig. 6, along with the reconstructed difference and

the different contributions derived from Eq. (6). We find the reconstruction to work very well, which allows us to quantify the relative contributions of the different terms. To this end, we adjust the derivation of Eq. (6) to consider quantities normalised by global mean temperature change, to yield

$$\frac{\Delta C_{\text{Mod}}}{\Delta T_{\text{Mod}}} - \frac{\Delta C_{\text{Orig}}}{\Delta T_{\text{Orig}}} = \sum \left[\theta_{i,\text{Orig}} \left(\frac{\Delta Y_{i,\text{Mod}}}{\Delta T_{\text{Mod}}} - \frac{Y_{i,\text{Orig}}}{\Delta T_{\text{Orig}}} \right) + \delta_i \frac{\Delta Y_{i,\text{Orig}}}{\Delta T_{\text{Orig}}} + \delta_i \left(\frac{\Delta Y_{i,\text{Mod}}}{\Delta T_{\text{Mod}}} - \frac{\Delta Y_{i,\text{Orig}}}{\Delta T_{\text{Orig}}} \right) \right], \quad (7)$$

with the first term again describing the pattern contribution, the second one describing the sensitivity contribution and the third term describing the cross term.

Estimating the relative contributions of these terms using linear regression for the fast and slow components, we find that the SST sensitivity change dominates the overall Pacific subsidence low-cloud responses (65 % and 79 % for the fast and slow responses, respectively). Meanwhile, the pattern and the cross term make secondary contributions (11 % and 24 % in the fast response, 4 % and 17 % in the slow response). The pattern contribution is mostly driven by estimated inversion strength (EIS) changes (Fig. A4).

In summary, our experiments show that, in CESM-CAM4, a higher and more realistic low-cloud sensitivity to SST leads to a greater decrease in the low-cloud amount under abrupt $4\times\text{CO}_2$ increase. This occurs both directly, due to the imposed increased low-cloud sensitivity to SST, and indirectly, through additional changes in cloud sensitivities and changes in the warming pattern and associated controlling-factor changes. This pattern effect makes only a minor additional contribution to the Pacific subsidence regions and is mostly mediated by changes in EIS.

3.4 CMIP6

We now analyse whether our experimental findings in the previous sections can also be traced in a CMIP6 model ensemble. As a first step, we plot the ECS, derived from $4\times\text{CO}_2$ experiments, against the index of Pacific subsidence-region cloud radiative effect (CRE) SST sensitivity – where the SST sensitivity in each model is quantified through a cloud-controlling-factor analysis, as in Ceppi and Nowack (2021). We use CRE sensitivity in this section rather than LCC sensitivity owing to limited data availability of LCC output in CMIP6 models. In agreement with our findings in Fig. 2a, we find that a higher CRE sensitivity correlates with an increased ECS ($r = 0.67$), with an increase of 0.7 K ($p = 0.001$) per unit CRE sensitivity increase (Fig. 7a), in line with previous studies (e.g. Zelinka et al., 2020; Ceppi and Nowack, 2021; Myers et al., 2021). For comparison, we show the values calculated from the Orig and Mod experiments as black and red crosses, respectively. While the relationship between the CMIP models qualitatively fits our experiments, the former have a stronger dependency on the CRE sensitivity index. This is most likely due to CRE sensitivity in the Pacific

subsidence regions correlating with CRE sensitivity in other regions in the CMIP6 models – whereas in our experiments, the CRE sensitivity difference is restricted to the subsidence regions. As a side note, we find that our $1\times\text{Orig}$ setup does not underestimate CRE sensitivity to SST relative to observations (Fig. 7a) as much as was the case for the LCC sensitivity (Fig. A2a). This difference may reflect a greater contribution of mid- and upper-level clouds on CRE in CESM-CAM4 compared to observations, as well as possible differences in the low-cloud optical depth contribution.

Next, we determine the impact of the CRE–SST sensitivity on the warming patterns across models. To this end, we regress the models' warming patterns against their CRE sensitivity, averaged over the Pacific subsidence regions. Note that the maps in Fig. 5 have different units: for CMIP models (panels g–i), we regress onto a cloud radiative sensitivity index in $\text{W m}^{-2} \text{K}^{-1}$, whereas for our model experiments (panels d–f), we are simply considering warming pattern differences. Nevertheless, the maps should be physically comparable, since in both cases we are considering the effect of an increase in cloud–SST sensitivity in the Pacific subsidence regions.

The fast response shows excellent agreement between our experiments (Fig. 5d) and the CMIP6 analysis (Fig. 5g), with similar patterns of enhanced warming in the East Pacific and reduced warming in the North and Southwest Pacific (although the latter feature is not as pronounced in our experiments). While the slow patterns (Fig. 5e, h) do not match as closely as the fast patterns, we still observe enhanced Southeast Pacific warming both in our experiments and in the CMIP6 analysis. The reduced warming along the cold tongue region is in contrast to our experiments; however, this seems to be mostly driven by three outlier models (GISS-E2-1-G, MIROC6 and MIROC-ES2L), with exceptionally strong warming in this region. Excluding these three models reduces the effect significantly (Fig. A5). The total warming patterns (Fig. 5f, i) also show a qualitative agreement of enhanced East Pacific warming. In CMIP6, both the slow and total warming patterns include a hemispherically asymmetric component with enhanced warming in the Southern Hemisphere – likely related to changes in the Atlantic Meridional Overturning Circulation (Lin et al., 2019) – which is absent in our CESM-CAM4 simulations.

Given the increase in relative Southeast Pacific warming due to cloud feedback across CMIP6 models, we now turn to analysing the relationship with Walker circulation change. Since a greater CRE–SST sensitivity is associated with an enhanced reduction in the east–west SST contrast across the tropical Pacific under CO_2 forcing (Fig. 5i), we expect an enhanced weakening of the Walker circulation. Figure 7b shows that this is indeed the case, where we plot the slope of decadal averages of the Walker circulation anomaly per degree of global mean temperature change in the $4\times\text{CO}_2$ experiments against the Pacific subsidence-region cloud sensitivity index. This gives a moderate Pearson correlation coefficient




Figure 6. Difference in low-cloud-cover change between the ensemble mean $4\times_{\text{Mod}}$ and $4\times_{\text{Orig}}$ experiments averaged over the Pacific subsidence regions. Shown are the simulation results (black), the reconstruction using a cloud-controlling-factor analysis (red) and the different contributions derived in Eq. (6), summed over all cloud-controlling factors. These are the changes in cloud sensitivity (blue), changes in cloud-controlling factors (orange) and a cross term (green).

of $r = -0.48$, $p = 0.03$ and a decline of -4.5 Pa K^{-1} per unit of CRE sensitivity increase. We note, however, that two outlier models (namely MPI-ESM1-2-HR and MPI-ESM1-2-LR) were excluded from the analysis. The across-model relationship should therefore be considered with caution until the drivers of the large Walker circulation responses in the two MPI models are understood. Our experimental results (black and red crosses in Fig. 7b) agree with the CMIP6 inter-model regression, providing confidence that the relationship is causal. The higher CRE–SST sensitivity in observations compared to most models therefore implies an enhanced weakening of the Walker circulation in the future compared to model projections.

4 Conclusions

This study investigated the global climate impacts of increasing stratocumulus cloud sensitivity to sea surface temperature (SST) anomalies in the Pacific subsidence regions, targeting a common climate model bias (Myers et al., 2021; Ceppi et al., 2024). Our main focus was on warming patterns and the atmospheric circulation response under abrupt CO_2 forcing, but we also considered changes in unforced SST variability. We conducted simulations with modified low-cloud sensitivity under both $1\times_{\text{Mod}}$ and $4\times_{\text{Mod}}$, and compared them to control simulations with default sensitivity, labelled $1\times_{\text{Orig}}$ and $4\times_{\text{Orig}}$.

The key impacts of stronger Pacific stratocumulus cloud feedback identified from our experiments are as follows:

- *Increased SST variability.* In the control climate, enhanced cloud sensitivity to SST leads to more variable, persistent and extreme SSTs in the tropical and subtropical East Pacific as well as the Niño3.4 region, consistent with positive feedback between low-cloud amount and SST and in line with similar previous findings (e.g. Bellomo et al., 2014; Rädel et al., 2016; Yang et al., 2022).
- *Higher equilibrium climate sensitivity (ECS).* ECS increases by approximately 6 % in $4\times_{\text{Mod}}$ compared to $4\times_{\text{Orig}}$, driven by a change in the climate feedback.
- *Change in SST warming pattern.* In addition to the ECS change, enhancing cloud sensitivity results in relatively higher warming in the East Pacific and thus a reduced east–west contrast across the tropical Pacific under CO_2 forcing. This SST “pattern effect” acts to moderately amplify the low-cloud reduction in the Pacific subsidence regions, contributing about 11 % in the fast and 4 % in the slow LCC response.
- *Walker circulation weakening.* While a slowdown of the Walker circulation is expected under CO_2 forcing, we observe an additional slowdown in Walker circulation in response to the change in the warming pattern.

To evaluate the broader implications of these findings, we compared our results with an across-model analysis of




Figure 7. CMIP6 relationships between CRE–SST Pacific cloud sensitivity index and $4\times\text{CO}_2$ (a) equilibrium climate sensitivity (ECS) and (b) Walker circulation change with global warming. The plot titles show the Pearson correlation coefficient r as well as its p value, calculated using a Student t test. The empty circles in (b) denote outliers that were excluded from the regression analysis. The dashed line shows the CRE sensitivity index calculated from the CERES data. Black and red crosses show the values of the Orig and Mod experiments, respectively.

22 CMIP6 models. We found a connection between ECS and Pacific subsidence cloud–SST sensitivity across models ($r = 0.67$, $p = 0.001$), qualitatively in line with previous results (e.g. Zelinka et al., 2020; Ceppi and Nowack, 2021; Myers et al., 2021). Furthermore, through an inter-model regression analysis, we found excellent alignment in terms of the fast warming pattern between our experiments and CMIP6 models. Agreement was more limited for the slow response, with discrepancies likely arising from model-dependent slower oceanic processes, such as changes in the Atlantic meridional overturning. This analysis indicates that our CESM-CAM4 findings have significant implications for climate projections, especially as models typically predict intensified warming in the East Pacific in the slow response to CO_2 forcing (Rugenstein et al., 2023), which may even further amplify the global impacts of stratocumulus feedback.

Additionally, we identified a potential across-model relationship in CMIP6 between cloud sensitivity and Walker circulation weakening ($r = -0.48$, $p = 0.03$, excluding two outlier models). This relationship agrees well with both our physical understanding and our experimental results, suggesting that stratocumulus cloud sensitivity has substantial impacts on projected regional climate changes. This finding is especially important given that climate models still commonly exhibit biases in cloud feedback representation (Zelinka et al., 2022; Ceppi et al., 2024). Future work could address the importance of biases in other cloud types and their sensitivities to environmental factors, in particular high clouds and their effects on atmospheric circulation (Wilson Kemsley et al., 2024).

Appendix A: Figures




Figure A1. Sensitivities of low-cloud cover (LCC) to surface temperature for (a) MODIS observational data, (b) $1\times\text{Orig}$ experiment and (c) $1\times\text{Mod}$ experiment.




Figure A2. (a) Sensitivities of LCC to controlling factors, expressed as a percentage of LCC anomaly per ERA5 standard deviation of each factor, averaged over Pacific subsidence regions. These sensitivities are presented for MODIS observations and for the $1 \times \text{Orig}$ and $1 \times \text{Mod}$ experiments. The controlling factors used are surface temperature (T_s), estimated inversion strength (EIS), near-surface horizontal temperature advection (T_{adv}), near-surface wind speed (WS), 700 hPa vertical velocity (OMEGA700) and 700 hPa relative humidity (RH700). (b) Location of the Pacific subsidence regions, calculated from ERA5 following Scott et al. (2020).




Figure A3. Climatological surface temperatures for (a) $1 \times \text{Orig}$, (b) $1 \times \text{Mod}$ and (c) their difference. The stippling in (c) marks significant differences at the $p < 0.05$ level using a two-sample t test for autocorrelated data following Wilks (2019) on the annually averaged surface temperature.




Figure A4. Contributions to differences in low-cloud-cover change between the ensemble mean $4\times_{\text{Mod}}$ and $4\times_{\text{Orig}}$ experiments averaged over the Pacific subsidence regions, derived in Eq. (6) and resolved by the following cloud-controlling factors: surface temperature (T_s), estimated inversion strength (EIS), advected surface temperature (T_{adv}), total surface wind speed (WS), relative humidity at 700 hPa (RH700) and vertical wind at 700 hPa (OMEGA700). The subfigures show the different terms in Eq. (6): (a) the contribution from changes in the cloud-controlling factors, (b) the contribution from changes in the cloud sensitivity, (c) the contribution of the cross term of co-varying sensitivity and controlling-factor change, and (d) the total sum of (a)–(c).




Figure A5. Same as for Fig. 5g–i but excluding GISS-E2-1-G, MIROC6 and MIROC-ES2L from the analysis.

Code and data availability. The code written for the presented analysis was archived under <https://doi.org/10.5281/zenodo.15977831> (Breul, 2025); the accompanying data were archived under <https://doi.org/10.5281/zenodo.16025001> (Breul and Ceppi, 2025).

Author contributions. PB designed and performed the model experiments and analysis and wrote the paper. PC contributed to the design of the experiments, the interpretation of the results and the writing of the paper. PN contributed to the interpretation of the results and the writing of the paper.

Competing interests. At least one of the (co-)authors is a member of the editorial board of *Atmospheric Chemistry and Physics*. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. We are grateful to two anonymous reviewers for their constructive comments on the manuscript. This work used JASMIN, the UK collaborative data analysis facility. This work also used the ARCHER2 UK National Supercomputing Service (<https://www.archer2.ac.uk>, last access: 18 September 2025). We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modelling groups for producing and making available their model output. We also thank the Earth System Grid Federation (ESGF) for archiving the model output and providing access, and we thank the multiple funding agencies who support CMIP and ESGF.

Financial support. This research has been supported by the Natural Environment Research Council (grant nos. NE/V012045/1 and NE/T006250/1) and the UK Research and Innovation organisation (grant no. EP/Y036123/1).

Review statement. This paper was edited by Raphaela Vogel and reviewed by two anonymous referees.

References

- Andrews, T. and Webb, M. J.: The Dependence of Global Cloud and Lapse Rate Feedbacks on the Spatial Structure of Tropical Pacific Warming, *J. Climate*, 31, 641–654, <https://doi.org/10.1175/JCLI-D-17-0087.1>, 2018.
- Andrews, T., Gregory, J. M., and Webb, M. J.: The Dependence of Radiative Forcing and Feedback on Evolving Patterns of Surface Temperature Change in Climate Models, *J. Climate*, 28, 1630–1648, <https://doi.org/10.1175/JCLI-D-14-00545.1>, 2015.
- Bellomo, K., Clement, A., Mauritsen, T., Rädel, G., and Stevens, B.: Simulating the Role of Subtropical Stratocumulus Clouds in Driving Pacific Climate Variability, *J. Climate*, 27, 5119–5131, <https://doi.org/10.1175/JCLI-D-13-00548.1>, 2014.
- Bellomo, K., Clement, A. C., Mauritsen, T., Rädel, G., and Stevens, B.: The Influence of Cloud Feedbacks on Equatorial Atlantic Variability, *J. Climate*, 28, 2725–2744, <https://doi.org/10.1175/JCLI-D-14-00495.1>, 2015.
- Breul, P.: Code accompanying the publication “egusphere-2025-221”, Zenodo [code], <https://doi.org/10.5281/zenodo.15977831>, 2025.
- Breul, P. and Ceppi, P.: Dataset accompanying the publication “egusphere-2025-221”, Zenodo [data set], <https://doi.org/10.5281/zenodo.16025001>, 2025.
- Brown, P. T., Lozier, M. S., Zhang, R., and Li, W.: The necessity of cloud feedback for a basin-scale Atlantic Multidecadal Oscillation, *Geophys. Res. Lett.*, 43, 3955–3963, <https://doi.org/10.1002/2016GL068303>, 2016.
- Burgman, R. J., Kirtman, B. P., Clement, A. C., and Vazquez, H.: Model evidence for low-level cloud feedback driving persistent changes in atmospheric circulation and regional hydroclimate, *Geophys. Res. Lett.*, 44, 428–437, <https://doi.org/10.1002/2016GL071978>, 2017.
- Ceppi, P. and Hartmann, D. L.: Clouds and the Atmospheric Circulation Response to Warming, *J. Climate*, 29, 783–799, <https://doi.org/10.1175/JCLI-D-15-0394.1>, 2016.
- Ceppi, P. and Nowack, P.: Observational evidence that cloud feedback amplifies global warming, *P. Natl. Acad. Sci. USA*, 118, e2026290118, <https://doi.org/10.1073/pnas.2026290118>, 2021.
- Ceppi, P., Zappa, G., Shepherd, T. G., and Gregory, J. M.: Fast and Slow Components of the Extratropical Atmospheric Circulation Response to CO₂ Forcing, *J. Climate*, 31, 1091–1105, <https://doi.org/10.1175/JCLI-D-17-0323.1>, 2018.
- Ceppi, P., Myers, T. A., Nowack, P., Wall, C. J., and Zelinka, M. D.: Implications of a Pervasive Climate Model Bias for Low-Cloud Feedback, *Geophys. Res. Lett.*, 51, e2024GL110525, <https://doi.org/10.1029/2024GL110525>, 2024.
- Chalmers, J., Kay, J. E., Middlemas, E. A., Maroon, E. A., and DiNezio, P.: Does Disabling Cloud Radiative Feedbacks Change Spatial Patterns of Surface Greenhouse Warming and Cooling?, *J. Climate*, 35, 1787–1807, <https://doi.org/10.1175/JCLI-D-21-0391.1>, 2022.
- Danabasoglu, G., Bates, S. C., Briegleb, B. P., Jayne, S. R., Jochum, M., Large, W. G., Peacock, S., and Yeager, S. G.: The CCSM4 Ocean Component, *J. Climate*, 25, 1361–1389, <https://doi.org/10.1175/JCLI-D-11-00091.1>, 2012.
- Dong, Y., Proistosescu, C., Armour, K. C., and Battisti, D. S.: Attributing Historical and Future Evolution of Radiative Feedbacks to Regional Warming Patterns using a Green's Function Ap-

- proach: The Preeminence of the Western Pacific, *J. Climate*, 32, 5471–5491, <https://doi.org/10.1175/JCLI-D-18-0843.1>, 2019.
- Erfani, E. and Burls, N. J.: The Strength of Low-Cloud Feedbacks and Tropical Climate: A CESM Sensitivity Study, *J. Climate*, 32, 2497–2516, <https://doi.org/10.1175/JCLI-D-18-0551.1>, 2019.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Fu, M. and Fedorov, A.: The Role of Bjerknes and Short-wave Feedbacks in the Tropical Pacific SST Response to Global Warming, *Geophys. Res. Lett.*, 50, e2023GL105061, <https://doi.org/10.1029/2023GL105061>, 2023.
- Gregory, J. M., Ingram, W. J., Palmer, M. A., Jones, G. S., Stott, P. A., Thorpe, R. B., Lowe, J. A., Johns, T. C., and Williams, K. D.: A new method for diagnosing radiative forcing and climate sensitivity, *Geophys. Res. Lett.*, 31, <https://doi.org/10.1029/2003GL018747>, 2004.
- He, J. and Soden, B. J.: Anthropogenic Weakening of the Tropical Circulation: The Relative Roles of Direct CO₂ Forcing and Sea Surface Temperature Change, *J. Climate*, 28, 8728–8742, <https://doi.org/10.1175/JCLI-D-15-0205.1>, 2015.
- Heede, U. K., Fedorov, A. V., and Burls, N. J.: Time Scales and Mechanisms for the Tropical Pacific Response to Global Warming: A Tug of War between the Ocean Thermostat and Weaker Walker, *J. Climate*, 33, 6101–6118, <https://doi.org/10.1175/JCLI-D-19-0690.1>, 2020.
- Held, I. M. and Shell, K. M.: Using Relative Humidity as a State Variable in Climate Feedback Analysis, *J. Climate*, 25, 2578–2582, <https://doi.org/10.1175/JCLI-D-11-00721.1>, 2012.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G. D., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P. d., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hsiao, W.-T., Hwang, Y.-T., Chen, Y.-J., and Kang, S. M.: The Role of Clouds in Shaping Tropical Pacific Response Pattern to Extratropical Thermal Forcing, *Geophys. Res. Lett.*, 49, e2022GL098023, <https://doi.org/10.1029/2022GL098023>, 2022.
- Janoski, T. P., Mitevski, I., Kramer, R. J., Previdi, M., and Polvani, L. M.: ClimKern v1.2: a new Python package and kernel repository for calculating radiative feedbacks, *Geosci. Model Dev.*, 18, 3065–3079, <https://doi.org/10.5194/gmd-18-3065-2025>, 2025.
- Kim, H., Kang, S. M., Kay, J. E., and Xie, S.-P.: Subtropical clouds key to Southern Ocean teleconnections to the tropical Pacific, *P. Natl. Acad. Sci. USA*, 119, e2200514119, <https://doi.org/10.1073/pnas.2200514119>, 2022.
- Knutson, T. R. and Manabe, S.: Time-Mean Response over the Tropical Pacific to Increased CO₂ in a Coupled Ocean-Atmosphere Model, *J. Climate*, 8, 2181–2199, [https://doi.org/10.1175/1520-0442\(1995\)008<2181:TMROTT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1995)008<2181:TMROTT>2.0.CO;2), 1995.
- Lin, Y.-J., Hwang, Y.-T., Ceppi, P., and Gregory, J. M.: Uncertainty in the Evolution of Climate Feedback Traced to the Strength of the Atlantic Meridional Overturning Circulation, *Geophys. Res. Lett.*, 46, 12331–12339, <https://doi.org/10.1029/2019GL083084>, 2019.
- Malik, A., Nowack, P. J., Haigh, J. D., Cao, L., Atique, L., and Plancherel, Y.: Tropical Pacific climate variability under solar geoengineering: impacts on ENSO extremes, *Atmos. Chem. Phys.*, 20, 15461–15485, <https://doi.org/10.5194/acp-20-15461-2020>, 2020.
- Middlemas, E. A., Clement, A. C., Medeiros, B., and Kirtman, B.: Cloud Radiative Feedbacks and El Niño–Southern Oscillation, *J. Climate*, 32, 4661–4680, <https://doi.org/10.1175/JCLI-D-18-0842.1>, 2019.
- Myers, T. A. and Mechoso, C. R.: Relative Contributions of Atmospheric, Oceanic, and Coupled Processes to North Pacific and North Atlantic Variability, *Geophys. Res. Lett.*, 47, e2019GL086321, <https://doi.org/10.1029/2019GL086321>, 2020.
- Myers, T. A., Scott, R. C., Zelinka, M. D., Klein, S. A., Norris, J. R., and Caldwell, P. M.: Observational constraints on low cloud feedback reduce uncertainty of climate sensitivity, *Nat. Clim. Change*, 11, 501–507, <https://doi.org/10.1038/s41558-021-01039-0>, 2021.
- Myers, T. A., Zelinka, M. D., and Klein, S. A.: Observational Constraints on the Cloud Feedback Pattern Effect, *J. Climate*, 36, 6533–6545, <https://doi.org/10.1175/JCLI-D-22-0862.1>, 2023.
- Neale, R. B., Richter, J. H., Conley, A. J., Park, S., Lauritzen, P. H., Gettelman, A., Williamson, D. L., Rasch, P. J., Vavrus, S. J., Taylor, M. A., Collins, W. D., Zhang, M., and Lin, S.-J.: Description of the NCAR Community Atmosphere Model (CAM 4.0), NCAR Tech. Note NCAR/TN-485+STR, 120 pp., 2010.
- Nowack, P. J., Braesicke, P., Luke Abraham, N., and Pyle, J. A.: On the role of ozone feedback in the ENSO amplitude response under global warming, *Geophys. Res. Lett.*, 44, 3858–3866, <https://doi.org/10.1002/2016GL072418>, 2017.
- Rädel, G., Mauritsen, T., Stevens, B., Dommenges, D., Matei, D., Bellomo, K., and Clement, A.: Amplification of El Niño by cloud longwave coupling to atmospheric circulation, *Nat. Geosci.*, 9, 106–110, <https://doi.org/10.1038/ngeo2630>, 2016.
- Rugenstein, M., Zelinka, M., Karanaukas, K., Ceppi, P., and Andrews, T.: Patterns of Surface Warming Matter for Climate Sensitivity, *Eos*, 104, <https://doi.org/10.1029/2023EO230411>, 2023.
- Rugenstein, M. A. A., Caldeira, K., and Knutti, R.: Dependence of global radiative feedbacks on evolving patterns of surface heat fluxes, *Geophys. Res. Lett.*, 43, 9877–9885, <https://doi.org/10.1002/2016GL070907>, 2016.
- Scott, R. C., Myers, T. A., Norris, J. R., Zelinka, M. D., Klein, S. A., Sun, M., and Doelling, D. R.: Observed Sensitivity of Low-Cloud Radiative Effects to Meteorological Perturbations over the Global Oceans, *J. Climate*, 33, 7717–7734, <https://doi.org/10.1175/JCLI-D-19-1028.1>, 2020.
- Soden, B. J., Held, I. M., Colman, R., Shell, K. M., Kiehl, J. T., and Shields, C. A.: Quantifying Climate Feedbacks Using Radiative Kernels, *J. Climate*, 21, 3504–3520, <https://doi.org/10.1175/2007JCLI2110.1>, 2008.
- Stevens, B. and Brenguier, J.-L.: Cloud-controlling Factors: Low Clouds, in: *Clouds in the Perturbed Climate System*, edited by: Heintzenberg, J. and Charlson, R. J.,

- The MIT Press, 173–196, ISBN 978-0-262-25544-8, <https://doi.org/10.7551/mitpress/8300.003.0010>, 2009.
- Tokinaga, H., Xie, S.-P., Deser, C., Kosaka, Y., and Okumura, Y. M.: Slowdown of the Walker circulation driven by tropical Indo-Pacific warming, *Nature*, 491, 439–443, <https://doi.org/10.1038/nature11576>, 2012.
- Trenberth, K. E. and Stepaniak, D. P.: Indices of El Niño Evolution, *J. Climate*, 14, 1697–1701, [https://doi.org/10.1175/1520-0442\(2001\)014<1697:LIOENO>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<1697:LIOENO>2.0.CO;2), 2001.
- Vecchi, G. A., Soden, B. J., Wittenberg, A. T., Held, I. M., Leetmaa, A., and Harrison, M. J.: Weakening of tropical Pacific atmospheric circulation due to anthropogenic forcing, *Nature*, 441, 73–76, <https://doi.org/10.1038/nature04744>, 2006.
- Voigt, A. and Shaw, T. A.: Circulation response to warming shaped by radiative changes of clouds and water vapour, *Nat. Geosci.*, 8, 102–106, <https://doi.org/10.1038/ngeo2345>, 2015.
- Welch, P.: The use of fast Fourier transform for the estimation of power spectra: A method based on time averaging over short, modified periodograms, *IEEE T. Acoust. Speech*, 15, 70–73, <https://doi.org/10.1109/TAU.1967.1161901>, 1967.
- Wilks, D.: Statistical methods in the atmospheric sciences, Elsevier Science, ISBN 978-0-12-816527-0, <https://doi.org/10.1016/C2017-0-03921-6>, 2019.
- Wilks, D. S.: “The Stippling Shows Statistically Significant Grid Points”: How Research Results are Routinely Overstated and Overinterpreted, and What to Do about It, *B. Am. Meteorol. Soc.*, 97, 2263–2273, <https://doi.org/10.1175/BAMS-D-15-00267.1>, 2016.
- Wilson Kemsley, S., Ceppi, P., Andersen, H., Cermak, J., Stier, P., and Nowack, P.: A systematic evaluation of high-cloud controlling factors, *Atmos. Chem. Phys.*, 24, 8295–8316, <https://doi.org/10.5194/acp-24-8295-2024>, 2024.
- Yang, L., Xie, S.-P., Shen, S. S. P., Liu, J.-W., and Hwang, Y.-T.: Low Cloud–SST Feedback over the Subtropical Northeast Pacific and the Remote Effect on ENSO Variability, *J. Climate*, 36, 441–452, <https://doi.org/10.1175/JCLI-D-21-0902.1>, 2022.
- Yuan, T., Oreopoulos, L., Platnick, S. E., and Meyer, K.: Observations of Local Positive Low Cloud Feedback Patterns and Their Role in Internal Variability and Climate Sensitivity, *Geophys. Res. Lett.*, 45, 4438–4445, <https://doi.org/10.1029/2018GL077904>, 2018.
- Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., Klein, S. A., and Taylor, K. E.: Causes of Higher Climate Sensitivity in CMIP6 Models, *Geophys. Res. Lett.*, 47, e2019GL085782, <https://doi.org/10.1029/2019GL085782>, 2020.
- Zelinka, M. D., Klein, S. A., Qin, Y., and Myers, T. A.: Evaluating Climate Models’ Cloud Feedbacks Against Expert Judgment, *J. Geophys. Res.-Atmos.*, 127, e2021JD035198, <https://doi.org/10.1029/2021JD035198>, 2022.
- Zhou, C., Zelinka, M. D., and Klein, S. A.: Analyzing the dependence of global cloud feedback on the spatial pattern of sea surface temperature change with a Green’s function approach, *J. Adv. Model. Earth Sy.*, 9, 2174–2189, <https://doi.org/10.1002/2017MS001096>, 2017.
- Zhu, J. and Poulsen, C. J.: On the Increase of Climate Sensitivity and Cloud Feedback With Warming in the Community Atmosphere Models, *Geophys. Res. Lett.*, 47, e2020GL089143, <https://doi.org/10.1029/2020GL089143>, 2020.