



Impacts of shipping emissions on ozone pollution in China

Zhenyu Luo^{1,★}, Li Peng^{1,★}, Zhaofeng Lv¹, Junchao Zhao², Tingkun He¹, Wen Yi¹, Yongyue Wang¹,
Kebin He¹, and Huan Liu¹

¹State Key Laboratory of Regional Environment and Sustainability, School of Environment,
Tsinghua University, Beijing 100084, China

²Key Laboratory of Vehicle Emission Control and Simulation of Ministry of Ecology and Environment,
Vehicle Emission Control Center, Chinese Research Academy of environmental Sciences, Beijing, China

★These authors contributed equally to this work.

Correspondence: Huan Liu (liu_env@tsinghua.edu.cn)

Received: 29 April 2025 – Discussion started: 4 June 2025

Revised: 13 August 2025 – Accepted: 10 September 2025 – Published: 24 October 2025

Abstract. With the Two Phases of Clean Air Actions in China, the shipping sector has emerged as a significant source with substantial emission reduction potential compared to land-based anthropogenic sectors. Therefore, understanding the contribution of shipping emissions to ozone (O₃) pollution is therefore essential for advancing China's air pollution control efforts. In this study, a coupled framework including a chemical transport model with machine learning techniques was developed to systematically investigate the interannual and seasonal impacts of shipping emissions on O₃ concentrations across China during the period from 2016 to 2020, and explore mechanisms by which shipping emissions influence O₃ formation. Results indicate that shipping emissions increase O₃ concentrations by a five-year average of 3.5 ppb nationwide, exhibiting significant spatial and temporal heterogeneity across different regions and seasons. Although significant differences exist between the emissions of ocean vessels and inland vessels, their contributions to O₃ formation are becoming increasingly comparable. Solely controlling shipping emissions may have limited impact on O₃ mitigation. Instead, coordinated reductions targeting both shipping and land-based anthropogenic sources, along with region-specific and targeted emission control strategies, are critical for achieving substantial improvements in O₃ pollution mitigation.

1 Introduction

Over the past decades, China's rapid industrialization and urbanization have boosted the economy but also exacerbated air pollution (Zhang et al., 2019). To address air pollution and the health burden from anthropogenic emissions, China introduced the Air Pollution Prevention and Control Action Plan in 2013 (Zheng et al., 2018). Although China has implemented synergistic control of VOC and NO_x emissions, the warm-season mean maximum daily 8 h average ozone (MDA8 O₃) increased by 2.6 μg m⁻³ yr⁻¹ in China between 2013–2020, especially in urban areas where declining PM_{2.5} levels offset gains in O₃ mitigation (Liu et al., 2023). Numerous epidemiological studies have shown that ground-level O₃

pollution leads to a range of adverse health effects, including increased incidence and mortality of respiratory diseases (Ito et al., 2005; Jerrett et al., 2009; Tao et al., 2012). Therefore, the health benefits achieved by reducing PM_{2.5} pollution are partially offset by the increase in O₃ pollution (Wang et al., 2021a; Xie et al., 2019), indicating a need to explore strategies to mitigate O₃ pollution in China.

During the promotion of China's emission control actions, emissions from the industry and power sectors declined substantially, with NO_x reductions exceeding 50 %, while the transportation sector still retains significant potential for further cuts (Liu et al., 2023). Ships emit both gaseous and particulate pollutants, including sulfur dioxide (SO₂), nitrogen oxides (NO_x), particulate matter, and volatile organic

compounds (VOC). As the largest maritime trading nation, China has a higher share of shipping emissions among anthropogenic sources compared to international levels (Fu et al., 2017; Yi et al., 2025). From 2016 to 2019, shipping emission controls in China focused on reducing SO₂ and PM emissions through the adoption of low-sulfur fuels, while NO_x and VOC emissions from shipping continued to rise by approximately 13 % due to increasing trade volumes (Wang et al., 2021b). Additionally, the use of low-sulfur fuel may further increase VOC emissions (Wu et al., 2020). Therefore, clarifying the historical and current contribution of shipping emissions to the formation of O₃ is critically important for further pollution control in China.

Previous studies have quantified the impacts of shipping emissions on O₃ pollution in China. In the southern coastal region, shipping emissions contributed approximately 0.9 µg m⁻³ to annual O₃ pollution (Cheng et al., 2023), with a peak winter contribution of up to 10 % (Feng et al., 2023). In the eastern coastal region during summer, the shipping-related O₃ concentration ranged from −15 to 15 ppb (Fu et al., 2023; Wang et al., 2019a). In the Bohai Rim Area (BRA), shipping emissions showed a maximum annual negative contribution of 0.5 µg m⁻³ (Wan et al., 2023), while summer O₃ concentration in Shandong Province increased by up to 10 ppb due to shipping (Wang et al., 2022). However, these studies predominantly focused on coastal areas, such as Yangtze River Delta (YRD), Pearl River Delta (PRD), and BRA, while the potential inland impacts are not well studied. However, recent studies have indicated that the air quality effects of inland shipping should not be neglected (Huang et al., 2022; Luo et al., 2024). The formation of O₃ exhibits strong spatial heterogeneity and is influenced by multiple factors, such as meteorological conditions and the emission intensities from both shipping and land-based sources. Currently, there is a lack of comprehensive studies employing a unified methodology to assess the nationwide impact of shipping emissions on O₃. Furthermore, previous studies were limited to restricted periods, resulting in a deficiency in comprehensive assessments of multi-year scenarios that encompass coordinated variations in both land and shipping emissions.

O₃ is generated by photochemical reactions between NO_x and VOC under solar radiation, thus, the impacts of shipping emissions on O₃ concentrations are attributed by the nonlinear response of O₃ to changes in NO_x and VOC emissions (Wang et al., 2019a, 2017). Additionally, the titration of O₃ by NO from shipping emissions, particularly within a few kilometers of ship tracks, can further complicate the simulation and interpretation of O₃ concentrations at the local scale (Merico et al., 2016). However, previous studies commonly used the zero-out method to assess ship's impacts by comparing scenario differences simulated by chemical transport models, which does not fully involve the nonlinear response of O₃ to its precursors and would result in considerable uncertainty in the evaluations (Cheng et al., 2023; Feng et al.,

2023; Fu et al., 2023; Wan et al., 2023; Wang et al., 2022, 2019a). Furthermore, although model-based assessments can generate large amounts of simulation data to investigate the impacts of shipping emissions, the number of scenarios that can be simulated by chemical transport models remains limited due to computational constraints. As a result, current analyses struggle to struggle the mechanism of how shipping emissions contribute to O₃ formation from these discrete scenarios. Recently, the advancement of machine learning techniques, with strong capabilities in capturing nonlinear relationships, provides a valuable approach for uncovering underlying patterns in such datasets (Luo et al., 2025).

In this study, we conducted source-oriented chemical transport model with a spatial resolution of 36 km × 36 km, to investigate the annual and seasonal impacts of shipping emissions on O₃ concentrations in China, especially for key coastal and inland regions from 2016 to 2020. We also apportion the contribution of shipping emissions from ocean-going vessels (OGVs), coastal vessels (CVs), and river vessels (RVs) to O₃ pollution to identify the influences of regionally differentiated shipping emission control policies. Furthermore, an explainable machine learning model was applied to explore investigate the potential source-receptor relationships between shipping emissions and the O₃ formation based on five-years simulated data. Our study provides a nationwide and long-term analysis of the impacts shipping emissions on China's O₃ pollution, and provide new insights for shipping control measures in the future.

2 Methods

2.1 Shipping emissions

The Shipping Emission Inventory Model (SEIM v2.0) is a disaggregate dynamic method (Wang et al., 2021b) driven by (a) the high-frequency ship Automatic Identification System (AIS) data, including signal time, coordinate location, navigational speed, and operating status, and (b) the integrated Ship Technical Specifications Database (STSD) (updated to 2020), which describes ship static properties, including vessel type, maximum designed speed, DWT and engine power. First, the originally collected raw AIS data and ship profile data from multiple sources are combined to form a ship activity database and STSD. Second, a route restoration module is applied for cross-land trajectory with a long distance in the AIS data, in which the 10 min linear interpolation will be applied on the shorted paths instead. Third, the instantaneous emission along with the movement of the ship's trajectory will be calculated based on the ship's static technical parameters, dynamic load changes, and extra parameters and factors. In the SEIM, shipping emissions for both air pollutants (e.g. SO₂, PM, NO_x, CO, and VOC) and greenhouse gases (e.g. CO₂, CH₄, and N₂O) from the main engines, auxiliary engines and boilers were calculated, detailed information of SEIM is described in our previous study (Wang et al., 2021b).

Here, emissions beyond 200 nautical miles from the Chinese mainland's territorial sea baseline were excluded from the domain by applying GIS-based spatial processing to the global shipping emission inventory, and only the annual shipping emissions from 2016 to 2020 within 200 nautical miles were used in the simulation.

In this study, vessels were classified as ocean-going vessels (OGVs), coastal vessels (CVs) and river vessels (RVs) for emission estimation according the following rules: (a) OGVs were identified by both valid International Maritime Organization (IMO) numbers and the Maritime Mobile Service Identity (MMSI) numbers, since they are mostly engaged in international trade following the management of the IMO; (b) RVs were identified by frequency distribution method based on the navigation trajectories for each vessel. Vessels with more than 50 % of the AIS signals throughout the entire year occurring on inland rivers (14–43° N, 104–130° E) were considered as RVs; and (c) vessels that are not identified as OGVs or RVs are regarded as CVs. Figure 1 shows the interannual variation of shipping NO_x and VOC emissions from 2016 to 2020. Overall, the growing trade demands and total cargo throughput of Chinese ports contributed to increased shipping activities, which in turn resulted in a steady rise in shipping NO_x and VOC emissions, especially for OGVs. Additionally, following the implementation of the global sulfur cap (IMO, 2018), the shift to low-sulfur fuels, which are typically richer in short-chain hydrocarbons (Wu et al., 2020), has contributed to a rise in shipping VOC emissions. It is worth noting that changes in vessel operating conditions, such as idling time and engine load, also influenced emissions. Although the COVID-19 pandemic had a temporary effect on maritime activity, its impact on annual shipping emissions was relatively minor due to the rapid rebound in trade during the second half of the year (Yi et al., 2024).

2.2 Air quality model

The Weather Research and Forecasting (WRF, version 3.8.1, using meteorological fields from 2018, as detailed later) – Community Multiscale Air Quality (CMAQ, version 5.4) model was applied to simulate the air quality in China from 2016 to 2020. Considering the relatively stable monthly anthropogenic emissions, this study simulated the O_3 concentrations during January, April, July, and November to represent winter, spring, summer, and fall, respectively, for the calculation of annual and seasonal mean values. As shown in Fig. S2 in the Supplement, the modeling domain covered all of China and some parts of East Asia with a horizontal resolution of $36\text{ km} \times 36\text{ km}$. Here, we have defined four key regions: BRA, YRD, PRD and inland river areas (Moirangthem, 2016), in which we focus on shipping-related O_3 pollution.

Here, we primarily focused on examining the impact of anthropogenic emission changes on shipping-related O_3 from

a historical perspective. To eliminate the impact of interannual meteorological variability, we used meteorological field of 2018 (Zhao et al., 2022), which simulated by WRF and identified as a typical meteorological year due to its relatively stable climate conditions, to drive the CMAQ simulations for the period 2016–2020. The first guess field and boundary conditions for WRF were generated from the 6 h NCEP FNL Operational Model Global Tropospheric Analyses dataset. The four-dimensional data assimilation (FDDA) was enabled using the NCEP ADP global surface and upper air observational weather data (<http://rda.ucar.edu>, last access: 25 March 2023). WRF and CMAQ used 32 vertical layers up to 100 hPa, and the lowest layer had a thickness of approximately 37 m. The major physical options in WRF included a Morrison two-moment microphysics scheme (Morrison et al., 2009), a Kain–Fritsch cumulus cloud parameterization (Kain, 2004), the Rapid Radiative Transfer Model (RRTM) longwave and shortwave radiation scheme (Iacono et al., 2008), the Pleim–Xiu Land Surface Model (Xiu and Pleim, 2001), and the Asymmetric Convective Model version 3.0 for the PBL parameterization (Pleim, 2007). The distribution of meteorological stations for validation and WRF performance is shown in Fig. S3 and Table S1 in the Supplement, respectively.

Atmospheric gas-phase chemistry in the CMAQ was simulated with the SAPRC07ic chemical mechanism, and aerosols were predicted using the AERO7. The chemical boundary conditions of CMAQ inputs, corresponding to each simulation period, were collected from the Community Atmosphere Model with Chemistry (CAM-chem) simulation output of global tropospheric and stratospheric compositions (Buchholz et al., 2019). Each run included a 3 d spin-up period. In this study, the Integrated Source Apportionment Method (ISAM) was applied to determine the source contribution to the ambient O_3 concentrations. Here, ISAM-OP3 was applied to attribute all secondary products to sources emitting NO_x or reactive VOC species and radicals when present in the parent reactants, and otherwise assign them based on stoichiometric reaction rates (Shu et al., 2023). We divided the emissions into five groups to trace them separately in the ISAM, including the land-based anthropogenic emission (the mobiles, industry, power, domestic, and agriculture) from the MEIC and the open burning emissions from Cai's study (Cai et al., 2017), the RVs' emission, the CVs' emission, the OGVs' emission, and the other emission (the nature sources emission from the MEGANv3 and the anthropogenic emission from other countries within the modeling domain from the MIX (Li et al., 2017), details of emissions are shown in Table S2.

We evaluated the simulated MDA8 O_3 concentrations for the of 2018 against 1455 available ground-based observations (Fig. S3) for model validation. As shown in Table S3, the simulated MDA8 O_3 agreed well with observations, with the overall model performance within the performance criteria suggested by Boylan and Rus-




Figure 1. The interannual variation of shipping (a) NO_x and (b) VOC emissions from 2016 to 2020.

sell (Boylan and Russell, 2006) (mean fractional bias (MFB) $\leq \pm 60\%$ and mean fractional error (MFE) $\leq \pm 75\%$), while the model overestimated O₃ a little, mainly due to uncertainties in emission inventory and unavoidable deficiencies during meteorological and air quality simulation. Meteorological performance for simulated periods was described in our previous study (Zhao et al., 2022).

2.3 Explainable machine learning model

Although CMAQ-ISAM can generate large amounts of simulation data to investigate the impacts of shipping emissions, the number of scenarios remains limited due to computational constraints. As a result, current analyses struggle to elucidate the mechanisms by which shipping emissions contribute to O₃ formation from these discrete scenarios. In particular, capturing nonlinear interactions between emission sources, meteorological conditions, and chemical processes is challenging when only a limited number of emission perturbations are available. Recently, the advancement of machine learning techniques, especially explainable models, has provided a promising complementary approach (Liu et al., 2025; Yao et al., 2024a, b). These models can learn from existing models to approximate the source-receptor relationships embedded in the simulation results. By identifying key emission drivers, quantifying their nonlinear contributions to O₃, and revealing latent patterns across spatiotemporal scales.

Here, based on the simulated data from 2016 to 2020 using WRF-CMAQ-ISAM, the RF model was used to simulate the monthly average O₃ concentration. In the RF model, the input predictor variables included relative humidity, temperature, wind speed, wind direction, solar radiation, land anthropogenic NO_x emissions, land anthropogenic VOC emissions, shipping NO_x emission and shipping VOC emission. Notably, the emissions selected were the sum of emissions from each grid and its eight neighboring grids. We trained four RF models specifically for the BRA, YRD, PRD, and IRA regions, respectively. The simulation data from 2016 to 2019 were used as the training samples, while the simulation

data from 2020, under the scenario of the most significant changes in shipping emissions, used as the test samples to validate the generalization capability of the RF models. By comparing the root mean squared error (RMSE) for testing datasets across models with candidate parameter combinations, we set *mtry* and *NumTrees* as 6 and 200 in RF, respectively. Additionally, the 10-fold cross-validation repeated 10 times was considered to evaluate the prediction performance of our models. The total dataset was randomly divided into 10 subsets, where 9 subsets was used to train the model and another was applied for validation. As shown in Fig. 2, averages of RMSE and correlation coefficient (R^2) in the CV of the RF models were 2.12–2.47 ppb, and 0.90–0.98, respectively, indicating an acceptable performance.

In order to identify the sensitivity and response relationship between prediction variables and results in the RF models, the SHapley Additive exPlanations (SHAP) technique, a game-theoretic framework introduced by Lundberg et al. (Lundberg et al., 2020; Lundberg and Lee, 2017), was employed to interpret the pattern learned from the 2016–2019 simulation data by the RF model using the Python scikit-learn library. The SHAP approach enables the quantification of both global and local influences of input variables on the model's predictions, thereby improving the interpretability of factors contributing to air pollution. Additionally, the study examined feature interactions, which can affect the model's predictive accuracy, to gain a more comprehensive understanding of the intricate relationships among the variables.

2.4 Limitations

In this study, the spatial resolution of 36 km $\times$ 36 km may not fully capture the fine-scale spatial heterogeneity of O₃ concentrations, particularly in coastal urban areas where emissions and photochemical reactions exhibit strong spatial variability. This resolution is relatively coarse for accurately representing O₃ exceedances and local photochemical processes, which often occur at much finer spatial scales. Consequently, localized O₃ peaks and gradients may be underes-




Figure 2. Performances of RF models for (a) BRA, (b) YRD, (c) PRD, and (d) IRA. Each point represents the monthly average O₃ concentration at each CMAQ grid cell.

timated or smoothed in the model outputs. Despite this limitation, the selected resolution represents a practical compromise that enables multi-year simulations across the national domain.

Only four representative months were simulated each year to reflect annual and seasonal patterns. While this captures broad seasonal variability, it may overlook intra-seasonal fluctuations and short-term anomalies. Using these months to estimate annual and seasonal means introduces uncertainty, especially for sources with stronger monthly variation. Although monthly changes in anthropogenic and shipping emissions are generally modest (except in 2020), future work could benefit from higher temporal resolution to improve accuracy.

Anthropogenic emissions from other countries within the modeling domain were held fixed at 2010 levels, and open burning emissions were fixed at 2015 levels throughout the simulation period (2016–2020). Although this assumption simplifies the modeling framework and is unlikely to significantly alter the relative changes in shipping-related O₃ assessed in this work, it may still introduce some degree of uncertainty, particularly in regions where long-range trans-

port or fire-related emissions could have contributed more dynamically during specific years. Future studies could benefit from incorporating temporally varying background emissions to further reduce potential uncertainties and improve the representation of external influences.

Explainable machine learning model relies on the structure and quality of the input dataset and cannot account for unmeasured or omitted variables, such as hemispheric background O₃ concentrations. As a result, the derived feature importance reflects statistical associations rather than causal relationships. It should be noted that if one seeks to determine whether a given variable promotes or suppresses O₃ pollution using machine learning methods, additional field observations, experimental data, and corresponding simulation results may be required as supporting evidence. Considering the interactions among variables, even if individual contributions are small, the SHAP estimates for each explanatory variable are unlikely to perfectly reflect their actual contributions in the underlying physical processes. Furthermore, in the presence of strong collinearity or complex nonlinear interactions, SHAP values may not fully disentangle overlapping influences among features.

In this study, monthly and annual mean O_3 concentrations were derived from hourly model outputs, rather than the widely used MDA8 O_3 . While this approach is consistent with the study's focus on long-term trends and average responses, it may introduce bias due to the well-known overestimation of nighttime ozone in chemical transport models. A sensitivity test comparing shipping-related O_3 contributions based on hourly averages and MDA8 revealed that over oceanic areas, the difference may reach 2–5 ppb, while over land, it remains within 2 ppb. Given that the relative contribution of shipping emissions to total O_3 is generally low, the impact of this bias is expected to be limited.

3 Results and discussions

3.1 Annual O_3 impact from shipping emissions

Figure 3a shows the five-year average of shipping-related O_3 calculated based on hourly values, which is defined as the sum of O_3 concentration caused by emissions of OGVs, CVs, and RVs traced by CMAQ-ISAM. Overall, the shipping emissions increases O_3 concentrations by 3.5 ppb nationwide (Table S4), showing a decrease trend from the southeast coast toward inland areas. This result is greater than the findings in other countries such as 1.97 ppb in Europe, and 2.08 ppb in the US under ambient temperature and pressure (Sun et al., 2024). Due to the high coastal shipping emissions, the shipping-related O_3 could exceed 15.0 ppb in southeast coastal regions, especially in YRD and PRD, where maximum values reach 25.4 and 26.3 ppb, respectively. For the regions with low shipping emissions, the shipping-related O_3 is relatively low, not exceeding 5 ppb. For example, in IRA and BRA, the shipping-related O_3 is 3.9 and 4.5 ppb respectively, but is slightly higher than the national average. Meteorological factors are just as important as anthropogenic emission influences in O_3 production (Liu et al., 2023; Zhang et al., 2024). For example, the PRD region is characterized by a persistently warm and humid climate, which provides favorable conditions for ozone formation. In contrast, although the BRA is also a coastal region, it experiences lower temperatures and weaker solar radiation (Fig. S4), which reduces the formation of hydroxyl radicals, weakening the atmospheric photochemical oxidizing capacity and ultimately limiting the O_3 production.

Figure 3b shows the five-year average of the relative contribution of shipping emissions to O_3 . Nationwide, the shipping emissions accounts for a 8.6 % increase in O_3 , showing a similar decreasing trend of from the southeast coast to inland regions. This result is higher than 3.7 % reported for the Mediterranean region in 2015 (Fink et al., 2023), but lower than the 12 %–21 % reported in another European study for 2010 (Lupascu and Butler, 2019). Notably, some coastal cities exhibit particularly high values. For example, in PRD region, such as in Shenzhen, Guangdong Province, the relative shipping-related O_3 exceeds 30.4 %, due to the high

intensity of shipping emissions combined with relatively low emissions from land-based anthropogenic sources such as industry and power generation. Similarly, the IRA is relatively low at 10.4 %, mainly because of the higher background O_3 concentrations from land-based anthropogenic sources.

Figure 4 illustrates the interannual trend in shipping-related O_3 in key regions from 2016 to 2020. Nationwide, the shipping-related O_3 shows a slight upward trend, with an average annual growth rate of 1.7 %, primarily observed in coastal regions. This trend aligns with the changes in shipping NO_x and VOC emissions, especially in 2020 when a 0.2–0.3 ppb rise in shipping-related O_3 was observed, partly attributable to the notable increase in VOC emissions following the implementation of the global sulfur cap. However, this increase in O_3 concentrations is substantially lower than the growth in precursor emissions, highlighting the complex and nonlinear response of O_3 formation to changes in shipping emissions. This is because the formation of O_3 depends on photochemical reactions involving NO_x and VOC under solar radiation, and is influenced not only by the level of shipping emissions but also by land-based anthropogenic emissions, meteorological conditions, and long-range transport (Ye et al., 2023). Therefore, changes in shipping-related O_3 do not scale linearly with the changes in shipping NO_x and VOC emissions. The Chinese government's two phases of clean air actions (Phase I, 2013–2017; Phase II, 2018–2020) resulted in increasing trend of O_3 nationwide (Liu et al., 2023), and the relative contribution of shipping emissions to O_3 also rose slightly during the same period. It is worth noting that, despite continuous increases in shipping NO_x and VOC emissions, their relative contributions to O_3 decreased in 2018 and 2020. This pattern may result from simultaneous land-based emission reductions, which can affect atmospheric oxidizing capacity (Lv et al., 2020).

3.2 Contribution of different types of vessels

We further investigated the relationship between O_3 pollution and shipping emissions from sub-ship sectors. Figure 5 and Table S4 shows the five-year average contribution of emissions from different ship types to the shipping-related O_3 and the total O_3 , respectively. Nationwide, OGVs, CVs, and RVs contributed 2.6 %, 2.6 %, and 3.3 % to the total O_3 , respectively. In coastal regions, OGVs contribute more than 50 % to the shipping-related O_3 , and 9.7 % to the total O_3 . CVs are the second-largest contributor to O_3 pollution with an average contribution of 20 %–30 %, and contribute up to 40 % in the southern coastal regions near Zhejiang and Fujian Provinces. Notably, RVs can still contribute over 30 % to shipping-related O_3 pollution in some coastal regions, particularly in the YRD and PRD. This is primarily due to the presence of major inland waterways such as the lower Yangtze River and the Pearl River, respectively, as well as the influence of regional pollutant transport that enhances the impact of RVs' emissions in these areas. In inland regions, RVs re-




Figure 3. Contributions of shipping emissions to the five-year average O_3 pollution, including (a) absolute contributions and (b) relative contributions. Maps created with MeteoinfoMap (<http://www.meteothink.org>, last access: 16 October 2025).




Figure 4. The interannual trend in shipping-related O_3 , including (a) absolute contributions and (b) relative contributions, in key regions from 2016 to 2020.

main the main source of shipping-related O_3 , with contributions in the middle reaches of the Yangtze River reaching 50 %.

The interannual variation in the contribution of the different types of ship to shipping-related O_3 between 2016–2020 for different regions is illustrated in Fig. 6. Nationwide, the contribution of OGVs and RVs increased by 2.7 % and 0.6 %, respectively, while the contribution of CVs decreased by 3.3 %. This pattern was observed in all coastal regions and IRA. The interannual variation in the contribution of the different types of ship to shipping-related O_3 follows a similar pattern to that of shipping-related NO_x and VOC (Fig. 1), particularly NO_x emissions, which shows an upward trend for OGVs and RVs but a downward trend for CVs. As a result, the difference in the contribution of different types of ships to air quality is gradually narrowing. Notably, although RVs emissions significantly less than those of OGVs and CVs, its contribution to O_3 is comparable to that of other ship types, even exceeds that of CVs in some coastal regions. In addition, although China has required certain categories of ships to install AIS equipment since 2010, a large part of

small RVs in China have not been equipped with AIS (Zhang et al., 2017), which is not considered in this study. This result suggests the importance of paying greater attention to RVs in future emission control strategies.

3.3 Seasonal O_3 impact from shipping emissions

The five-years-average seasonal variations in the contribution of shipping emissions to O_3 concentrations across different regions are shown in Fig. 7 and Tabel 1, with January, April, July, and November representing winter, spring, summer, and fall, respectively. For cold seasons, including winter and fall, due to weaker solar radiation and lower temperatures that limit O_3 formation (Fig. S4), the shipping-related O_3 remains relatively lower than warm seasons (spring and summer), with national average and relative contribution of 1.53 ppb (5.6 %) and 2.41 ppb (7.9 %), respectively (Fig. S5). However, in the south of PRD, especially Guangdong and Hainan Provinces (Fig. 7a and d, and Table 1), the average and maximum of seasonal shipping-related O_3 exceeds 5 and 21 ppb, respectively. Notably, fall pollution even severer than that in summer. This is mainly because the PRD remains




Figure 5. Contributions of emissions from (a) OGVs, (b) CVs, and (c) RVs to the five-year average shipping-related O_3 pollution. Maps created with MeteoInfoMap (<http://www.meteothink.org>, last access: 16 October 2025).




Figure 6. The interannual trend in Contributions of emissions from OGVs, CVs, and RVs to shipping-related O_3 in key regions from 2016 to 2020.

warm and humid in fall, and prevailing monsoon winds are more likely to transport ship-borne pollutants from the sea to inland areas (Figs. S4, S6, and S7). Another distinct pattern is observed in BRA, where shipping-related O_3 formation tends to be more localized during the cold seasons, as indicated by a larger difference between the median and average values (Table 1). During this period, mainland China is under the influence of the Mongolian High Pressure System, and continental winds generally suppress the inland transport of ship-related O_3 (Cheng et al., 2023; Zhao et al., 2023). Therefore, significant shipping-related O_3 pollution only appears in major port cities with intensive maritime activity.

In spring, shipping-related O_3 reached its peak in YRD and PRD, with the maximum value exceeding 30 ppb (Fig. 7b and Table 1), consistent with the results of previous studies (Cheng et al., 2023; Schwarzkopf et al., 2022). Although spring is generally less favorable for O_3 formation compared

to summer in terms of temperature and humidity, strong onshore winds may play an important role in reduce the influence of shipping emissions (Cheng et al., 2023; Ma et al., 2022) (Fig. S4, S6, and S7). In addition, more complex physicochemical interactions may drive springtime O_3 (Cao et al., 2024; Zhang et al., 2024), which needs further investigation. In summer, shipping emissions significantly increased O_3 concentrations nationwide by 4.77 ppb and responsible for 13.7 % of national O_3 pollution (Figs. 7c and S5). Notably, even in IRA, where shipping emissions are much lower than in coastal regions, shipping-related O_3 were comparable to those along the coast. This is primarily because central China lies in a perennial monsoon region, where summer monsoons can carry shipping-related air pollutants inland from coastal cities (Zheng et al., 2024).

These findings indicate that seasonal variations for shipping-related O_3 are driven by meteorological factors,




Figure 7. Contributions of shipping emissions to the seasonal mean O_3 concentrations for (a) winter (JAN), (b) spring (APR), (c) summer (JUL), and (d) fall (NOV). Maps created with MeteoInfoMap (<http://www.meteothink.org>, last access: 16 October 2025).

particularly changes in prevailing wind direction, which are crucial for the diffusion and long-range transport of shipping emissions. Researchers suggest that the mixing emissions between shipping and local anthropogenic emissions can amplify complicated O_3 chemical formation in coastal cities (Wang et al., 2019b). Therefore, seasonal mitigation strategies and a better understanding of regional monsoon dynamics and their interaction with local anthropogenic emissions are crucial for effectively reducing shipping-related O_3 pollution.

3.4 Effects of shipping emission on O_3 formation

Although the features in the RF model include both meteorological factors and emissions, this section focuses on impacts of anthropogenic emissions, especially shipping emissions, on O_3 pollution. Figure 8 presents the SHAP summary plots for selected features in the RF model for BRA, YRD, PRD, and IRA, showing the magnitude, prevalence, and direction of each feature's impact on the model output (O_3 concentration). In the summary plot, the further a fea-

ture's SHAP value is from zero, the greater its influence; positive SHAP values indicate a positive contribution, while negative values indicate a negative effect. For example, in Fig. 8a, land-based NO_x and VOC emissions both have a significant impact on O_3 formation in the BRA, with contributions of 16.8 % and 11.0 %, respectively, suggesting that the atmospheric chemistry in this area is significantly affected by land-based anthropogenic emissions. Moreover, lower NO_x and VOC emissions leads to higher SHAP values, indicating a negative correlation between land-based anthropogenic emissions and O_3 pollution. In the coastal areas of BRA, shipping emissions are much smaller than land-based emissions, therefore, contribute only approximately 2.9 % to O_3 formation, and exhibit a similar negative correlation as land-based sources.

As the share of shipping emissions increases within total anthropogenic emissions in the coastal areas of YRD and PRD, the difference between the contributions of shipping and land-based emissions to O_3 pollution regions decreases. Especially in the PRD, shipping NO_x contributes up to 9.7 %, exceeding the land-based VOC contribution of 5.3 %. In the




Figure 8. Feature importance results of the random forest regression model for (a) BRA, (b) YRD, (c) PRD, and (d) IRA. The x axis shows SHAP values representing the impact of each feature on O_3 predictions (positive: increasing O_3 ; negative: decreasing O_3). Each dot is a grid-month sample, with color indicating the feature value. Instances with identical x values are stacked, and the stack height signifies the density.

IRA, emissions from inland vessels are much lower than those from ocean-going and coastal ships, thus, shipping NO and VOC contributions only 1.5 % and 0.8 % to O_3 pollution, respectively.

For meteorological factors, the contributions of solar radiation (25.2 %), temperature (20.0 %), wind speed (11.5 %), and wind direction (4.9 %), relative humidity (4.8 %) to O_3 formation exhibit clear regional heterogeneity. Overall, meteorological influences are greater than those of shipping emissions. This may be attributed to the highly complex physical and chemical processes involved, including cloud–radiation interactions, air mass transport, and water-related reactions.

The dependence plot (Fig. 9) further quantifies how changes in shipping emissions affect O_3 concentrations. Notably, due to the uneven distribution of shipping emissions, the x axis in Fig. 9 is set to non-uniform intervals to better illustrate their impact. Overall, as previously mentioned, changes in shipping NO_x and VOC emissions have a relatively minor effect on regional O_3 levels, approximately be-

tween −2.5 to 8 ppb, although there is clear regional heterogeneity. In the BRA region, shipping emissions are negatively correlated with O_3 , with increases in shipping NO_x and VOC emissions leading to a reduction of about 1–2 ppb in O_3 . Differently, in the YRD and PRD, increases in shipping NO_x emissions slightly promote O_3 formation—especially in the PRD region, where O_3 may increase by up to 3 ppb, which is opposite to the impact of land-based NO_x emissions (Fig. S7). This difference may be because, in the YRD and PRD, land-based NO_x emissions do not dominate the overall anthropogenic emissions as they do in the BRA, allowing shipping NO_x to also influence atmospheric chemistry. Furthermore, changes in shipping VOC emissions have almost no impact, consistent with the effect of changes in land-based VOC emissions (Fig. S7). Only when shipping VOC emissions increase dramatically in the YRD region is O_3 formation suppressed – though such cases are rare, as reflected by the partially negative SHAP values for shipping VOC emissions in Fig. 8b. In the IRA, similar to the BRA region, shipping emissions are very small compared to land-

Table 1. Seasonal ranges of shipping-related O₃ across BRA, YRD, PRD, and IRA (unit: ppb).

Region	Metric	Winter	Spring	Summer	Fall
BRA	Minimum	0.01	0.83	1.25	0.06
	25 % Quartile	0.05	3.09	7.13	0.39
	Median	0.11	4.02	9.32	0.78
	75 % Quartile	0.91	6.55	11.90	1.63
	Maximum	6.59	23.22	32.40	14.39
	Mean	0.71	5.36	10.13	1.74
YRD	Minimum	0.20	1.92	1.45	0.55
	25 % Quartile	1.01	5.02	6.11	1.91
	Median	1.79	6.64	7.29	3.23
	75 % Quartile	2.93	8.35	9.03	5.14
	Maximum	16.32	31.47	25.86	24.59
	Mean	2.47	7.39	7.92	4.41
PRD	Minimum	1.11	3.91	0.11	1.91
	25 % Quartile	2.79	7.94	3.19	4.85
	Median	5.19	10.07	5.65	7.97
	75 % Quartile	7.68	15.17	7.77	11.25
	Maximum	21.98	33.46	26.19	28.78
	Mean	5.96	11.91	5.77	8.79
IRA	Minimum	0.17	0.91	2.32	0.10
	25 % Quartile	1.19	2.88	5.65	1.44
	Median	1.61	4.85	9.07	2.85
	75 % Quartile	2.29	5.73	11.31	4.16
	Maximum	5.53	11.91	26.19	7.52
	Mean	1.76	4.58	9.03	2.87

based sources, so that changes in shipping NO_x and VOC emissions have a similar negative effect on O₃ as those from land-based sources.

4 Conclusions

In this study, we conducted multi-year CMAQ-ISAM simulations to investigate the how shipping emissions impacted O₃ across China, with a focus on three coastal regions and an inland region. From 2016 to 2020, shipping emissions increased national average O₃ concentrations by 3.5 ppb, accounting for 8.6 % of total O₃, with a spatial gradient decreasing from coastal to inland regions. Despite the increasing intensity of shipping activity and the implementation of the global sulfur cap, shipping NO_x and VOCs emissions rose significantly during this period. However, the national average shipping-related O₃ increased by only 0.23 ppb, while the relative contribution of shipping emissions to O₃ pollution rose by approximately 0.5 %. Notably, this relative contribution did not increase continuously; instead, a decline was observed in 2018 and 2020. This non-linear response, under conditions of simultaneous changes in multiple pollutants from different sectors, highlights the complexity and need for further investigation of attribution of O₃ pollution. For the four focus regions, the contribution of shipping to O₃

levels exceeded the national average, with more pronounced interannual increases.

We further disaggregate ship types to OGVs, CVs, and RVs. The result revealed that OGVs were the dominant contributors to shipping-related O₃ in coastal areas, followed by CVs, whereas RVs were the main source in inland river areas. Although OGVs, CVs, and RVs differ significantly in their emission magnitudes, the difference in their contributions to O₃ pollution is gradually narrowing. This trend suggests that the influence of RVs on regional O₃ levels should no longer be overlooked and that emission control efforts for RVs deserve renewed attention. However, from the perspective of sulfur emission control, RVs in China had already reached the final stage of sulfur regulation by 2018 under the implementation of domestic emission control policies. In contrast, NO_x control for inland vessels remains largely unaddressed. Globally, there is limited precedent or experience in regulating NO_x emissions from inland waterways, leaving China without a clear reference framework for RVs NO_x mitigation. Future control of shipping NO_x emissions needs to take into account both inland waterways and coastal areas.

The impacts of shipping emissions on O₃ also exhibited significant seasonal and regional characteristics. While shipping-related O₃ levels were generally lower in colder seasons, fall pollution in southern coastal regions exceeded that of summer due to favorable land–sea monsoon transport. Peak shipping-related O₃ levels occurred in spring over YRD and PRD, and in summer over inland areas. These patterns highlight the importance of implementing seasonal and region-specific control strategies to mitigate shipping-related O₃ pollution effectively. In particular, quality-oriented management policies such as seasonal routing adjustments, port operation scheduling, or dynamic emission monitoring, may play a more immediate role than emission control policies, which are typically less adaptable to seasonal variability and require long-term infrastructure or regulatory changes. Therefore, combining flexible operational measures with long-term emission reduction plans could enhance the overall effectiveness of O₃ mitigation.

Interpretable machine learning analysis further revealed significant spatial differences in the contribution of shipping emissions to O₃. In BRA and IRA, O₃ formation was primarily driven by land-based NO_x and VOC emissions, with shipping emissions playing a minor role and even showing a suppressive effect on O₃ formation. In contrast, in coastal regions such as YRD and PRD, the increasing share of shipping emissions in the total anthropogenic emissions enhanced their contribution to O₃, with shipping NO_x emissions showing a slight promoting effect on O₃ formation. This regional difference suggests that solely controlling shipping emissions may lead to unexpected atmospheric chemical responses and, under certain conditions, could even cause an increase in O₃ concentrations. Therefore, effective O₃ pollution control requires a coordinated reduction of both land-




Figure 9. SHAP dependence plot for (a) BRA, (b) YRD, (c) PRD, and (d) IRA.

based and shipping emissions, based on regional emission structures and atmospheric oxidation characteristics.

Code availability. Codes used during the current study are available from the corresponding author upon reasonable request.

Data availability. Data used during the current study are available from the corresponding author upon reasonable request.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/acp-25-13635-2025-supplement>.

Author contributions. Conceptualization: ZLuo, LP, HL; Methodology: ZLuo, ZLv, JZ; Investigation: WY, TH, YW; Visualization: ZLuo, LP; Supervision: HL, KH; Writing – original draft: ZLuo, LP; Writing – review and editing: ZLuo, HL.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Financial support. This research has been supported by the National Natural Science Foundation of China (grant no. 42325505 to H. Liu), the China National Postdoctoral Program for Innovative Talents (No. BX20250327 to Z. Luo), the Shuimu Tsinghua Scholar 25 Program (no. 2024SM028 to Z. Luo).

Review statement. This paper was edited by Xavier Querol and reviewed by two anonymous referees.

References

- Boylan, J. W. and Russell, A. G.: PM and light extinction model performance metrics, goals, and criteria for three-dimensional air quality models, *Atmos. Environ.*, 40, 4946–4959, <https://doi.org/10.1016/j.atmosenv.2005.09.087>, 2006.
- Buchholz, R. R., Emmons, L. K., Tilmes, S., and The CESM2 Development Team: CESM2.1/CAM-chem instantaneous output for boundary conditions, UCAR/NCAR – Atmospheric Chemistry Observations and Modeling Laboratory [data set], <https://doi.org/10.5065/NMP7-EP60>, 2019
- Cai, S., Wang, Y., Zhao, B., Wang, S., Chang, X., and Hao, J.: The impact of the “air pollution prevention and control action plan” on PM_{2.5} concentrations in Jing-Jin-Ji region during 2012–2020, *Sci. Total Environ.*, 580, 197–209, <https://doi.org/10.1016/j.scitotenv.2016.11.188>, 2017.
- Cao, T., Wang, H., Chen, X., Li, L., Lu, X., Lu, K., and Fan, S.: Rapid increase in spring ozone in the Pearl River Delta, China during 2013–2022, *npj Clim. Atmos. Sci.*, 7, 309, <https://doi.org/10.1038/s41612-024-00847-3>, 2024.
- Cheng, Q., Wang, X., Chen, D., Ma, Y., Zhao, Y., Hao, J., Guo, X., Lang, J., and Zhou, Y.: Impact of ship emissions on air quality in the Guangdong-Hong Kong-Macao Greater Bay Area (GBA):

- with a particular focus on the role of onshore wind, *Sustainability*, 15, 8820, <https://doi.org/10.3390/su15118820>, 2023.
- Feng, X., Ma, Y., Lin, H., Fu, T.-M., Zhang, Y., Wang, X., Zhang, A., Yuan, Y., Han, Z., Mao, J., Wang, D., Zhu, L., Wu, Y., Li, Y., and Yang, X.: Impacts of ship emissions on air quality in Southern China: opportunistic insights from the abrupt emission changes in early 2020, *Environ. Sci. Technol.*, 57, 16999–17010, <https://doi.org/10.1021/acs.est.3c04155>, 2023.
- Fink, L., Karl, M., Matthias, V., Oppo, S., Kranenburg, R., Kuenen, J., Moldanova, J., Jutterström, S., Jalkanen, J.-P., and Majamäki, E.: Potential impact of shipping on air pollution in the Mediterranean region – a multimodel evaluation: comparison of photooxidants NO₂ and O₃, *Atmos. Chem. Phys.*, 23, 1825–1862, <https://doi.org/10.5194/acp-23-1825-2023>, 2023.
- Fu, M., Liu, H., Jin, X., and He, K.: National- to port-level inventories of shipping emissions in China, *Environ. Res. Lett.*, 12, 114024, <https://doi.org/10.1088/1748-9326/aa897a>, 2017.
- Fu, X., Chen, D., Wang, X., Li, Y., Lang, J., Zhou, Y., and Guo, X.: The impacts of ship emissions on ozone in eastern China, *Sci. Total Environ.*, 903, 166252, <https://doi.org/10.1016/j.scitotenv.2023.166252>, 2023.
- Huang, H., Zhou, C., Huang, L., Xiao, C., Wen, Y., Li, J., and Lu, Z.: Inland ship emission inventory and its impact on air quality over the middle Yangtze River, China, *Sci. Total Environ.*, 843, 156770, <https://doi.org/10.1016/j.scitotenv.2022.156770>, 2022.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: calculations with the AER radiative transfer models, *J. Geophys. Res.-Atmos.*, 113, <https://doi.org/10.1029/2008JD009944>, 2008.
- International Maritime Organization (IMO): Resolution MEPC.305(73), Amendments to the Annex of the Protocol of 1997 to Amend the International Convention for the Prevention of Pollution from Ships, [https://wwwcdn.imo.org/localresources/en/KnowledgeCentre/IndexofIMOResolutions/MEPCDocuments/MEPC.305\(73\).pdf](https://wwwcdn.imo.org/localresources/en/KnowledgeCentre/IndexofIMOResolutions/MEPCDocuments/MEPC.305(73).pdf) (last access: 23 October 2025), 2018.
- Ito, K., De Leon, S. F., and Lippmann, M.: Associations between ozone and daily mortality: analysis and meta-analysis, *Epidemiology*, 16, 446–457, <https://doi.org/10.1097/01.ede.0000165821.90114.7f>, 2005.
- Jerrett, M., Burnett, R. T., Pope, C. A., Ito, K., Thurston, G., Krewski, D., Shi, Y., Calle, E., and Thun, M.: Long-term ozone exposure and mortality, *N. Engl. J. Med.*, 360, 1085–1095, <https://doi.org/10.1056/NEJMoa0803894>, 2009.
- Kain, J. S.: The Kain–Fritsch Convective Parameterization: An Update, [https://doi.org/10.1175/1520-0450\(2004\)043<0170:TKCPAU>2.0.CO;2](https://doi.org/10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2), 2004.
- Li, M., Zhang, Q., Kurokawa, J.-I., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935–963, <https://doi.org/10.5194/acp-17-935-2017>, 2017.
- Liu, J., Chen, M., Chu, B., Chen, T., Ma, Q., Wang, Y., Zhang, P., Li, H., Zhao, B., Xie, R., Huang, Q., Wang, S., and He, H.: Assessing the significance of regional transport in ozone pollution through machine learning: a case study of Hainan island, *ACS ES&T Air*, 2, 416–425, <https://doi.org/10.1021/acsestair.4c00297>, 2025.
- Liu, Y., Geng, G., Cheng, J., Liu, Y., Xiao, Q., Liu, L., Shi, Q., Tong, D., He, K., and Zhang, Q.: Drivers of increasing ozone during the two phases of clean air actions in China 2013–2020, *Environ. Sci. Technol.*, 57, 8954–8964, <https://doi.org/10.1021/acs.est.3c00054>, 2023.
- Lundberg, S. M. and Lee, S.-I.: A unified approach to interpreting model predictions, in: *Proceedings of the 31st International Conference on Neural Information Processing Systems*, Red Hook, NY, USA, 4768–4777, <https://dl.acm.org/doi/10.5555/3295222.3295230> (last access: 23 October 2025), 2017.
- Lundberg, S. M., Erion, G., Chen, H., DeGrave, A., Prutkin, J. M., Nair, B., Katz, R., Himmelfarb, J., Bansal, N., and Lee, S.-I.: From local explanations to global understanding with explainable AI for trees, *Nat. Mach. Intell.*, 2, 56–67, <https://doi.org/10.1038/s42256-019-0138-9>, 2020.
- Luo, Z., Lv, Z., Zhao, J., Sun, H., He, T., Yi, W., Zhang, Z., He, K., and Liu, H.: Shipping-related pollution decreased but mortality increased in Chinese port cities, *Nat Cities*, 1, 295–304, <https://doi.org/10.1038/s44284-024-00050-8>, 2024.
- Luo, Z., He, T., Lv, Z., Zhao, J., Zhang, Z., Wang, Y., Yi, W., Lu, S., He, K., and Liu, H.: Insights into transportation CO₂ emissions with big data and artificial intelligence, *Patterns*, 101186, <https://doi.org/10.1016/j.patter.2025.101186>, 2025.
- Lupaşcu, A. and Butler, T.: Source attribution of European surface O₃ using a tagged O₃ mechanism, *Atmos. Chem. Phys.*, 19, 14535–14558, <https://doi.org/10.5194/acp-19-14535-2019>, 2019.
- Lv, Z., Wang, X., Deng, F., Ying, Q., Archibald, A. T., Jones, R. L., Ding, Y., Cheng, Y., Fu, M., Liu, Y., Man, H., Xue, Z., He, K., Hao, J., and Liu, H.: Source–receptor relationship revealed by the halted traffic and aggravated haze in Beijing during the COVID-19 lockdown, *Environ. Sci. Technol.*, 54, 15660–15670, <https://doi.org/10.1021/acs.est.0c04941>, 2020.
- Ma, Y., Chen, D., Fu, X., Shang, F., Guo, X., Lang, J., and Zhou, Y.: Impact of sea breeze on the transport of ship emissions: a comprehensive study in the Bohai rim region, China, *Atmosphere-Basel*, 13, 1094, <https://doi.org/10.3390/atmos13071094>, 2022.
- Merico, E., Donato, A., Gambaro, A., Cesari, D., Gregoris, E., Barbaro, E., Dinoi, A., Giovanelli, G., Masieri, S., and Contini, D.: Influence of in-port ships emissions to gaseous atmospheric pollutants and to particulate matter of different sizes in a Mediterranean harbour in Italy, *Atmos. Environ.*, 139, 1–10, <https://doi.org/10.1016/j.atmosenv.2016.05.024>, 2016.
- Moirangthem, K.: Alternative Fuels for Marine and Inland Waterways: An Exploratory Study, <https://doi.org/10.2790/227559>, 2016.
- Morrison, H., Thompson, G., and Tatarskii, V.: Impact of cloud microphysics on the development of trailing stratiform precipitation in a simulated squall line: comparison of one- and two-moment schemes, *Mon. Wea. Rev.*, 137, 991–1007, <https://doi.org/10.1175/2008MWR2556.1>, 2009.
- Pleim, J. E.: A combined local and nonlocal closure model for the atmospheric boundary layer. Part I: model description and testing, *J. Appl. Meteor. Climatol.*, 46, 1383–1395, <https://doi.org/10.1175/JAM2539.1>, 2007.
- Schwarzkopf, D. A., Petrik, R., Matthias, V., Quante, M., Yu, G., and Zhang, Y.: Comparison of the impact of ship emissions in

- northern europe and eastern China, *Atmosphere-Basel*, 13, 894, <https://doi.org/10.3390/atmos13060894>, 2022.
- Shu, Q., Napelenok, S. L., Hutzell, W. T., Baker, K. R., Henderson, B. H., Murphy, B. N., and Hogrefe, C.: Comparison of ozone formation attribution techniques in the north-eastern United States, *Geosci. Model Dev.*, 16, 2303–2322, <https://doi.org/10.5194/gmd-16-2303-2023>, 2023.
- Sun, W., Jiang, W., and Li, R.: Global health benefits of shipping emission reduction in early 2020, *Atmos. Environ.*, 333, 120648, <https://doi.org/10.1016/j.atmosenv.2024.120648>, 2024.
- Tao, Y., Huang, W., Huang, X., Zhong, L., Lu, S.-E., Li, Y., Dai, L., Zhang, Y., and Zhu, T.: Estimated acute effects of ambient ozone and nitrogen dioxide on mortality in the Pearl River Delta of Southern China, *Environ. Health Persp.*, 120, 393–398, <https://doi.org/10.1289/ehp.1103715>, 2012.
- Wan, Z., Cai, Z., Zhao, R., Zhang, Q., Chen, J., and Wang, Z.: Quantifying the air quality impact of ship emissions in China's Bohai Bay, *Mar. Pollut. Bull.*, 193, 115169, <https://doi.org/10.1016/j.marpolbul.2023.115169>, 2023.
- Wang, F., Qiu, X., Cao, J., Peng, L., Zhang, N., Yan, Y., and Li, R.: Policy-driven changes in the health risk of PM_{2.5} and O₃ exposure in China during 2013–2018, *Sci. Total Environ.*, 757, 143775, <https://doi.org/10.1016/j.scitotenv.2020.143775>, 2021a.
- Wang, H., Gao, Y., Sheng, L., Wang, Y., Zeng, X., Kou, W., Ma, M., and Cheng, W.: The impact of meteorology and emissions on surface ozone in Shandong province, China, during summer 2014–2019, *International Journal of Environmental Research and Public Health*, 19, 6758, <https://doi.org/10.3390/ijerph19116758>, 2022.
- Wang, P., Chen, Y., Hu, J., Zhang, H., and Ying, Q.: Attribution of tropospheric ozone to NO_x and VOC emissions: considering ozone formation in the transition regime, *Environ. Sci. Technol.*, 53, 1404–1412, <https://doi.org/10.1021/acs.est.8b05981>, 2019a.
- Wang, R., Tie, X., Li, G., Zhao, S., Long, X., Johansson, L., and An, Z.: Effect of ship emissions on O₃ in the Yangtze River delta region of China: analysis of WRF-chem modeling, *Sci. Total Environ.*, 683, 360–370, <https://doi.org/10.1016/j.scitotenv.2019.04.240>, 2019b.
- Wang, T., Xue, L., Brimblecombe, P., Lam, Y. F., Li, L., and Zhang, L.: Ozone pollution in China: a review of concentrations, meteorological influences, chemical precursors, and effects, *Sci. Total Environ.*, 575, 1582–1596, <https://doi.org/10.1016/j.scitotenv.2016.10.081>, 2017.
- Wang, X., Yi, W., Lv, Z., Deng, F., Zheng, S., Xu, H., Zhao, J., Liu, H., and He, K.: Ship emissions around China under gradually promoted control policies from 2016 to 2019, *Atmos. Chem. Phys.*, 21, 13835–13853, <https://doi.org/10.5194/acp-21-13835-2021>, 2021b.
- Wu, Z., Zhang, Y., He, J., Chen, H., Huang, X., Wang, Y., Yu, X., Yang, W., Zhang, R., Zhu, M., Li, S., Fang, H., Zhang, Z., and Wang, X.: Dramatic increase in reactive volatile organic compound (VOC) emissions from ships at berth after implementing the fuel switch policy in the Pearl River Delta Emission Control Area, *Atmos. Chem. Phys.*, 20, 1887–1900, <https://doi.org/10.5194/acp-20-1887-2020>, 2020.
- Xie, Y., Dai, H., Zhang, Y., Wu, Y., Hanaoka, T., and Matsui, T.: Comparison of health and economic impacts of PM_{2.5} and ozone pollution in China, *Environ. Int.*, 130, 104881, <https://doi.org/10.1016/j.envint.2019.05.075>, 2019.
- Xiu, A. and Pleim, J. E.: Development of a Land Surface Model. Part I: Application in a Mesoscale Meteorological Model, *J. Appl. Meteor. Climatol.*, 40, 192–209, [https://doi.org/10.1175/1520-0450\(2001\)040<0192:DOALSM>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<0192:DOALSM>2.0.CO;2), 2001.
- Yao, L., Han, Y., Qi, X., Huang, D., Che, H., Long, X., Du, Y., Meng, L., Yao, X., Zhang, L., and Chen, Y.: Determination of major drive of ozone formation and improvement of O₃ prediction in typical north China plain based on interpretable random forest model, *Sci. Total Environ.*, 934, 173193, <https://doi.org/10.1016/j.scitotenv.2024.173193>, 2024a.
- Yao, T., Lu, S., Wang, Y., Li, X., Ye, H., Duan, Y., Fu, Q., and Li, J.: Revealing the drivers of surface ozone pollution by explainable machine learning and satellite observations in Hangzhou Bay, China, *J. Cleaner Prod.*, 440, 140938, <https://doi.org/10.1016/j.jclepro.2024.140938>, 2024b.
- Ye, C., Lu, K., Ma, X., Qiu, W., Li, S., Yang, X., Xue, C., Zhai, T., Liu, Y., Li, X., Li, Y., Wang, H., Tan, Z., Chen, X., Dong, H., Zeng, L., Hu, M., and Zhang, Y.: HONO chemistry at a suburban site during the EXPLORE-YRD campaign in 2018: formation mechanisms and impacts on O₃ production, *Atmos. Chem. Phys.*, 23, 15455–15472, <https://doi.org/10.5194/acp-23-15455-2023>, 2023.
- Yi, W., He, T., Wang, X., Soo, Y. H., Luo, Z., Xie, Y., Peng, X., Zhang, W., Wang, Y., Lv, Z., He, K., and Liu, H.: Ship emission variations during the COVID-19 from global and continental perspectives, *Sci. Total Environ.*, 954, 176633, <https://doi.org/10.1016/j.scitotenv.2024.176633>, 2024.
- Yi, W., Wang, X., He, T., Liu, H., Luo, Z., Lv, Z., and He, K.: The high-resolution global shipping emission inventory by the Shipping Emission Inventory Model (SEIM), *Earth Syst. Sci. Data*, 17, 277–292, <https://doi.org/10.5194/essd-17-277-2025>, 2025.
- Zhang, L., Wang, L., Ji, D., Xia, Z., Nan, P., Zhang, J., Li, K., Qi, B., Du, R., Sun, Y., Wang, Y., and Hu, B.: Explainable ensemble machine learning revealing the effect of meteorology and sources on ozone formation in megacity hangzhou, China, *Sci. Total Environ.*, 922, 171295, <https://doi.org/10.1016/j.scitotenv.2024.171295>, 2024.
- Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu, W., Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L., Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu, B., Liu, Z., Yang, F., He, K., and Hao, J.: Drivers of improved PM_{2.5} air quality in China from 2013 to 2017, *P. Natl. Acad. Sci. USA*, 116, 24463–24469, <https://doi.org/10.1073/pnas.1907956116>, 2019.
- Zhang, Y., Yang, X., Brown, R., Yang, L., Morawska, L., Ristovski, Z., Fu, Q., and Huang, C.: Shipping emissions and their impacts on air quality in China, *Sci. Total Environ.*, 581–582, 186–198, <https://doi.org/10.1016/j.scitotenv.2016.12.098>, 2017.
- Zhao, C.-B., Li, Q.-Q., Nie, Y., Wang, F., Xie, B., Dong, L.-L., and Wu, J.: The reversal of surface air temperature anomalies in China between early and late winter 2021/2022: observations and predictions, *Advances in Climate Change Research*, 14, 660–670, <https://doi.org/10.1016/j.accre.2023.09.004>, 2023.
- Zhao, J., Lv, Z., Qi, L., Zhao, B., Deng, F., Chang, X., Wang, X., Luo, Z., Zhang, Z., Xu, H., Ying, Q., Wang, S., He, K., and Liu, H.: Comprehensive assessment for the impacts of S/IVOC emissions from mobile sources on SOA for-

- mation in China, *Environ. Sci. Technol.*, 56, 16695–16706, <https://doi.org/10.1021/acs.est.2c07265>, 2022.
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095–14111, <https://doi.org/10.5194/acp-18-14095-2018>, 2018.
- Zheng, S., Jiang, F., Feng, S., Liu, H., Wang, X., Tian, X., Ying, C., Jia, M., Shen, Y., Lyu, X., Guo, H., and Cai, Z.: Impact of marine shipping emissions on ozone pollution during the warm seasons in China, *J. Geophys. Res.-Atmos.*, 129, e2024JD040864, <https://doi.org/10.1029/2024JD040864>, 2024.