



Clear-sky and cloudy-sky differences in NO₂ concentrations over the United States: implications for satellite measurement applications

Daniel L. Goldberg¹, M. Omar Nawaz¹, Congmeng Lyu^{2,3}, Jian He^{2,3}, Annmarie G. Carlton⁴, Shobha Kondragunta⁵, and Susan C. Anenberg¹

¹Department of Environmental and Occupational Health, Milken Institute School of Public Health, George Washington University, Washington, DC, USA

²Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

³NOAA Chemical Sciences Laboratory, Boulder, CO, USA

⁴Department of Chemistry, University of California – Irvine, Irvine, CA, USA

⁵NOAA NESDIS Center for Satellite Applications and Research, College Park, MD, USA

Correspondence: Daniel L. Goldberg (dgoldberg@gwu.edu)

Received: 21 March 2025 – Discussion started: 15 April 2025

Revised: 3 October 2025 – Accepted: 10 October 2025 – Published: 20 November 2025

Abstract. Satellite measurements of tropospheric trace gases are often only used when there are few clouds, which screens out 20 %–70 % of the data, depending on geographic region. Although clouds cause satellite data gaps, in situ surface measurements and model simulations can provide insight on NO₂ during cloudy conditions. Here, we intercompare surface observations, meteorological reanalysis (ERA5), satellite measurements (TROPOMI and TEMPO), and a model (WRF-Chem) during 2019 over the contiguous US to quantify how NO₂ concentrations differ under clear and cloudy skies. We find that in situ surface NO₂ measurements are, on average, +17 % larger on all days compared to clear-sky days and +36 % larger during cloudy days versus clear-sky days, with a wide distribution based on geographic region and roadway proximity: largest in the Northeast US and smallest in the Southwest US and near major roadways. WRF-Chem simulated surface NO₂ between cloudy and clear conditions is larger than the observed differences: +59 % on cloudy days vs. clear-sky days for the model. We additionally find modeled *j*NO₂ values are reasonable, suggesting this WRF-Chem NO₂ bias could be arising from overly rapid OH removal of NO₂ in sunlight, too slow NO_x regeneration of NO₂ in sunlight, or differing boundary layer depth biases under cloudy versus clear skies. Finally, using in situ NO₂ matched to provisional TEMPO data, we find the NO₂ differences between cloudy and clear conditions to be larger in the afternoon than morning. This study quantifies some of the biases in satellite measurements introduced by using only clear-sky data.

1 Introduction

Nitrogen dioxide (NO₂) is an air pollutant that adversely affects the human respiratory system (Health Effects Institute, 2022; Khreis et al., 2017) and can lead to premature mortality (Burnett et al., 2004; He et al., 2020). NO₂ is also an important precursor for ozone (O₃) and fine particulates (PM_{2.5}),

which also have serious health impacts. NO_x (= NO + NO₂; most NO_x is emitted as NO which rapidly cycles to NO₂) is released into the atmosphere by biogenic microbial activity in soils, high-temperature lightning, wildfires, and thermal fossil fuel combustion; in urban areas, the majority of ambient NO₂ originates from the latter (Crippa et al., 2021). Although end-of-pipe controls (Busca et al., 1998; Koltsakis

and Stamatelos, 1997) can reduce the amount of NO_x emitted from fossil-fuel engines and boilers, these technologies do not recover 100 % of the NO_x generation during combustion. As a consequence, NO₂ accumulates in our atmosphere and many urban areas have NO₂ concentrations that exceed the World Health Organization guideline of 5.3 ppb for an annual average (Anenberg et al., 2022).

Observing local air pollution is typically done by in situ surface monitors, which are spaced throughout a region with a higher density of monitors typically in areas of high population density and known pollution sources. In the United States, there are over 2200 in situ monitoring sites measuring some combination of O₃, PM_{2.5}, NO₂, volatile organic compounds (VOCs), and CO (U.S. EPA, 2023). While the US monitoring network is more comprehensive than most other countries (Martin et al., 2019), 79 % of US counties lack a single monitor and an additional 10 % of counties have only a single monitor, leaving only 11 % of US counties with more than 1 monitor (Sullivan and Krupnick, 2018). Although a robust and accurate ground-monitoring network is needed, the high operating cost of these instruments can be an important barrier (Kelly et al., 2017). Spatial gaps remain in-between the regulatory monitors, and sometimes these monitors are inadequate for understanding the true ambient air pollution exposure of most US residents, especially those that live and/or work several kilometers away from a regulatory monitor. Satellite data provide a way to fill in the gaps of the in situ monitoring network. Methodologies to obtain robust surface air pollutant measurement data from satellite instruments have improved dramatically in the past ten years (Bechle et al., 2015; Cao, 2023; Ghahremanloo et al., 2021, 2023; Larkin et al., 2023; Nawaz et al., 2025; Shetty et al., 2024; Sun et al., 2024).

NO₂ can be observed by remote sensing instruments due to its unique spectroscopic features (Vandaele et al., 1998). The Tropospheric Monitoring Instrument (TROPOMI) (Veefkind et al., 2012) has been measuring column densities of NO₂ pollution up to $7 \times 3.5 \text{ km}^2$ before 6 August 2019 and up to $5.5 \times 3.5 \text{ km}^2$ spatial resolution (van Geffen et al., 2024) since 6 August 2019. Because of TROPOMI's higher spatial resolution over predecessor instruments, such as the Ozone Monitoring Instrument (OMI) ($24 \times 13 \text{ km}^2$ at nadir) (Levelt et al., 2018), TROPOMI has ~ 50 daily satellite pixel measurements within a typical city ($\sim 1000 \text{ km}^2$) during clear skies, while OMI may have only 1–3 daily measurements within the borders of each city. This increased measurement capacity within a city allows us to discern spatial variability undetectable by previous instruments (de Foy et al., 2009; Sun et al., 2018).

Level 2 satellite NO₂ measurements – which are retrieved from observed NO₂ spectra using geophysical and model-based assumptions – are of the tropospheric column. In many cases, NO₂ column measurements are strongly correlated with the spatial patterns of surface NO₂ concentrations (Acker et al., 2025; Harkey and Holloway, 2024; Kim

et al., 2024) and surface NO_x emissions (Goldberg et al., 2024). For TROPOMI, studies have shown a strong correlation between tropospheric column measurements and collocated surface NO₂ for both the 13:30 average ($r^2 = 0.67$) and the 24 h average ($r^2 = 0.68$) (Goldberg et al., 2021; Kerr et al., 2023). However, there are rare instances in which NO_x emissions and NO₂ enhancements stay aloft and do not affect the surface; these are often situations associated with lightning NO_x (Nault et al., 2017), wildfire NO_x (Jin et al., 2021; Lin et al., 2024), and aircraft NO_x (Maruhashi et al., 2024). In these instances, it can be difficult to determine if the column NO₂ enhancements are also leading to surface NO₂ enhancements. These misinterpretations are more likely to occur over rural regions and/or individual days, as upper-tropospheric NO₂ enhancements near urban regions often dwarf NO₂ enhancements within the boundary layer especially over monthly or longer timescales (Goldberg et al., 2022).

Satellite measurements of trace gases are typically only used when there are few or no clouds; this is often referred to as the clear-sky bias of satellite data. In the US, this results in 20 %–70 % of the satellite data being filtered out depending on geographic region. The clear-sky bias affects NO₂ more so than other trace gases (such as CO and CH₄) because NO₂ is very photochemically active in the presence of strong sunlight; its effective lifetime during summer daytime is 2–7 h (Liu et al., 2016) and conversely can be up to 30 h during winter daytime (Kenagy et al., 2018). The speed at which it transforms into other chemical species is determined by the irradiation, ambient temperature, and oxidative capacity (Laughner and Cohen, 2019; Shah et al., 2020). More specifically, strong irradiation creates the OH radical which can react with NO₂ to create HNO₃ – a major terminal sink of NO₂ – and also accelerates the photolysis of NO₂ into NO and O(³P) leading to an accumulation of O₃ in the presence of VOCs; without VOCs, NO₂ cycles more rapidly to NO. Warm temperatures increase biogenic VOC emissions and VOC can react with NO₂ directly to create organic nitrates (e.g., peroxyacetyl nitrates and alkyl nitrates) (Zare et al., 2018) which act as a temporary sink of NO₂. Another daytime terminal sink for NO₂ is dry deposition; while this removal mechanism is often secondary to photochemical loss in urban environments and is not directly affected by sunlight, it is indirectly affected as cloudy conditions are often associated with increased relative humidity and shallower boundary layer depths, both of which increase dry deposition fluxes. Therefore, increased NO₂ dry deposition fluxes in cloudy conditions could offset some of the decreased NO₂ photochemical loss rates. The net result is that NO₂ concentrations are typically larger during cloudy conditions (Geddes et al., 2012).

However, outside of the Geddes et al. (2012) study, little has been done to observationally quantify the bias of NO₂ being larger during cloudy conditions particularly because there are no column measurements to validate the satellite

during cloudy conditions. With that said, there are surface in situ measurements during cloudy conditions that can give us an idea of how the clear-sky bias may affect the estimate of surface concentrations. In this project, we intercompare surface observations, meteorological reanalysis (ERA5), satellite measurements (TROPOMI and TEMPO), and a model (WRF-Chem) under clear and cloudy skies to better quantify the amount of surface and column bias of NO₂ concentrations that is being introduced when clouds are screened from the satellite data. Our analysis is focused on the United States during 2019 due the high density of in situ monitors and availability of high-resolution regional chemical transport models. The motivation of this project is to determine what the scientific community may be missing when excluding clouds from satellite-based NO₂ analyses.

2 Methods

2.1 EPA AQS Data

Hourly in situ NO₂ measurements were obtained from the pre-generated EPA Air Quality System (AQS) database: https://aq5.epa.gov/aq5web/airdata/download_files.html (last access: 9 July 2025). These routine measurements are operated and maintained by various state and federal agencies. 91 % of the “NO₂” measurements in 2019 were acquired through a chemiluminescence technique which converts NO₂ and unintentionally some NO_x species – such as alkyl nitrates, peroxy nitrates (PAN), and nitric acid (HNO₃) – to NO using a heated molybdenum converter; the NO is measured by quantifying the luminescence of NO when reacted in excess O₃ (Dickerson et al., 2019). Lamsal et al. (2008) suggested a correction factor for converting midday chemiluminescence NO₂^{*} (= NO₂ + unintended NO_x) measurements to NO₂ using modelled information of PAN, alkyl nitrates, and HNO₃. In Eq. (1), we show the Lamsal et al. (2008) correction factor with a modification to exclude alkyl nitrates which are not explicitly included in our WRF-Chem simulation.

$$[\text{NO}_2]^* = F_{\text{int}} \times [\text{NO}_2]$$
$$\text{where } F_{\text{int}} = \frac{[\text{NO}_2] + 0.95[\text{PAN}] + 0.35[\text{HNO}_3]}{[\text{NO}_2]} \quad (1)$$

Typically, correction factors are in the range of ~ 1.0 for fresh urban plumes and can be as large as ~ 3.0 for rural areas during summer, with averages typically in the 1–1.5 range for moderate and very polluted regimes, and are important to use for model vs. monitor intercomparisons (Kuhn et al., 2024; Lamsal et al., 2008; Poraicu et al., 2023). Other methods to measure in situ NO₂ include Cavity Attenuated Phase Shift (Kebabian et al., 2008) and Laser Induced Fluorescence (Thornton et al., 2000), but these methods are less common (9 % of all NO₂ monitors in 2019).

Annual and seasonal averages at 13:30 local standard time (between 13:00–14:00) of the in situ data were considered

valid and used if more than 75 % of the days of the year/season had valid data. There were 449 monitoring locations in 2019 in the US that achieved these criteria for an annual average, which equates to 1 monitor per $\sim 730\,000$ US residents. For the baseline analysis, we further remove data from the 75 monitoring locations (17 % of the locations) that are classified as “near-road” by the EPA, which means that they are installed within 20 m from major interstates since these in situ measurements are not representative of a $\sim 20\text{ km}^2$ satellite pixel measurement; we include the “near-road” NO₂ monitoring data in sensitivity analyses. NO₂ measurements between cloudy and clear-sky days are intercompared using the normalized mean change (NMC) as described in Eq. (2), where $\bar{x}$ and $\bar{y}$ are means of the two datasets being analyzed.

$$\text{NMC}(\%) = 100 \times \left(\frac{\bar{y} - \bar{x}}{\bar{x}} \right) \quad (2)$$

2.2 Satellite NO₂ Instruments

NO₂ slant column densities are derived from radiance measurements in the 405–465 nm spectral window of the UV-VIS-NIR spectrometer (van Geffen et al., 2022; Nowlan et al., 2025). Satellite instruments observe NO₂ by comparing observed spectra with a reference spectrum to derive the amount of NO₂ in the atmosphere between the instrument and the surface; this technique is called differential optical absorption spectroscopy (DOAS) (Platt, 1994). Tropospheric vertical column density data, which represent the vertically integrated NO₂ concentrations between the surface and the tropopause, are then calculated by subtracting the stratospheric portion and then converting the tropospheric slant column to a vertical column using an air mass factor (AMF) (Boersma et al., 2011). The AMF is a unitless quantity used to convert the slant column into a vertical column and is a function of the satellite viewing angles, solar angles, the effective cloud radiance fraction and pressure, the vertical profile shape of NO₂ provided by a chemical transport model simulation, and the surface reflectivity (Lorente et al., 2017; Palmer et al., 2001).

2.2.1 TROPOMI

TROPOMI was launched by the European Space Agency (ESA) on 13 October 2017, and data from the instrument became available on 30 April 2018, after an approximately 6-month calibration period. The satellite follows a sun-synchronous, low-earth (825 km) orbit with an equator overpass time of approximately 13:30 local solar time. TROPOMI measures total column densities of several trace gases: NO₂, HCHO, O₃, CO, CH₄, among others. At nadir, pixel sizes are $3.5 \times 7\text{ km}^2$ (modified to $3.5 \times 5.5\text{ km}^2$ on 6 August 2019) with the edges having slightly larger pixel sizes ($\sim 14\text{ km}$ wide) across a 2600 km swath, equating to 450 rows (van Geffen et al., 2020).

For our analysis we use the TROPOMI NO₂ version 2.4 (V2.4) re-processed algorithm during 1 January 2019–31 December 2019. We also conducted a sensitivity study using the version 2.3.1 (V2.3.1) algorithm. The TROPOMI NO₂ V2.4 product has a documented median low bias of -34.8% in moderately polluted locations (when NO₂ measurements are between $3-14 \times 10^{15}$ molec cm⁻²) when compared to a MAX-DOAS network (Lambert et al., 2023). Some of this low bias is due to the operational AMF which uses a $1^\circ \times 1^\circ$ model to assume vertical shape profiles; when vertical shape profiles from a regional model are instead used, the bias decreases to between -1% and -23% (Nawaz et al., 2024; Judd et al., 2020; Tack et al., 2021). Prior work has demonstrated a strong correlation between TROPOMI NO₂ column measurements and NO₂ surface concentrations in urban areas (Demetillo et al., 2020; Dressel et al., 2022; Goldberg et al., 2021; Nawaz et al., 2025). For our baseline, we screened TROPOMI pixels for quality assurance flag values greater than 0.75, and conduct a sensitivity analysis of filtering only with a cloud radiative fraction filter of 0.5. The cloud radiative fraction is calculated from the O₂ A-band using the FRESCO-S algorithm. Due to differences in wavelength between the O₂ A-band and the NO₂ retrieval window, the cloud fraction retrieved in the O₂ A-band is not exactly representative for the cloud fraction in the NO₂ window, but it is similar.

The filtered data were re-gridded to a $0.01^\circ \times 0.01^\circ$ resolution, to create a custom “Level-3” data product (Goldberg et al., 2021) during cloud-free and cloudy conditions. Single pixel TROPOMI tropospheric vertical column NO₂ uncertainties have been quantified to be between 25% – 50% under clear skies and this uncertainty is dominated by uncertainty in the tropospheric air mass factor (Glissenaar et al., 2025; Liu et al., 2021; Rijdsdijk et al., 2025); uncertainties of measurements with cloud fractions > 0.5 are larger. Oversampled NO₂ measurements over monthly and annual timeframes (10s–100s of measurements) have a smaller amount of uncertainty, approximately 10% – 20% depending on location and season (Glissenaar et al., 2025).

2.2.2 TEMPO

TEMPO was launched by SpaceX on 7 April 2023 and is hosted on Maxar Intelsat 40e. Data from the instrument became available on 2 August 2023, after an approximately 4-month dry-out, cool-down, and calibration period. The satellite is in geostationary orbit centered over the United States with north-south coverage extending from Mexico City ($\sim 17^\circ$ N) to the Canadian Oil Sands ($\sim 58^\circ$ N) and east-west coverage from Puerto Rico to the Pacific coast. TEMPO operationally measures total column densities of NO₂, HCHO, and O₃ with additional products forthcoming. At nadir, pixel sizes are 4.75×2 km² with the North-east and North-west edges having slightly larger pixels sizes. The

instrument observes the full east-west swath approximately once every hour.

For our analysis we use the TEMPO NO₂ version 3 algorithm during 2 August 2023–31 August 2024. The data was filtered to include pixels only where the effective cloud fractions are less than 0.15 and the main data quality flags are equal to 0. The filtered data was re-gridded to a $0.01^\circ \times 0.01^\circ$ resolution, to create a custom “Level-3” data product (Goldberg et al., 2021) during cloud-free and cloudy conditions. Single pixel TEMPO tropospheric vertical column NO₂ uncertainties can be assumed to be similar to the uncertainty of TROPOMI measurements (Glissenaar et al., 2025), which are between 25% – 50% under clear skies for individual pixels, and 10% – 20% for oversampled averages; future work will better quantify the uncertainties of TEMPO NO₂ measurements.

2.3 ERA5 Re-analysis

We intercompare the cloud radiative fractions from TROPOMI to the ERA5 re-analysis (Hersbach et al., 2020) of total cloud fractions in the early afternoon (18:00 Z for Eastern Time, 19:00 Z for Central Time, 20:00 Z for Mountain Time, 21:00 Z for Pacific Time), which approximates the overpass time of TROPOMI over the contiguous United States. The ERA5 total cloud fraction is a unitless quantity representing how much of a grid cell is covered by a cloud (e.g., condensed water vapor) at any vertical level of the atmosphere and does not differentiate between the optical properties of those clouds. The ERA5 re-analysis data are reported at a $0.25^\circ \times 0.25^\circ$ spatial resolution and the cloud fractions are interpolated, using bilinear interpolation, to the $0.01^\circ \times 0.01^\circ$ oversampled TROPOMI NO₂ grid.

2.4 WRF-Chem

The Weather Research and Forecasting with Chemistry (WRF-Chem) model was run at $12 \text{ km} \times 12 \text{ km}$ over the Continental US for all days of 2019: 1 January–31 December 2019 as described in He et al. (2024). For anthropogenic emissions, the Fuel-based Inventory of Vehicle Emissions (FIVE) was used to provide on-road and off-road mobile emissions, the Fuel-based Oil and Gas (FOG) inventory was used for emissions associated with oil and natural gas production, power plant emissions were provided by Continuous Emissions Monitoring Systems (CEMS), and all other anthropogenic emissions were obtained from the 2014 or 2017 National Emissions Inventory (NEI). Biogenic emissions were estimated using Biogenic Emissions Inventory System (BEIS) version 3.13. Gas-phase chemistry was from the RACM_ESRL_VCP scheme. Boundary conditions were provided from the Realtime Air Quality Modeling System (RAQMS, <http://raqms-ops.ssec.wisc.edu/>, last access: 12 November 2025) developed by the University of Wisconsin-

Madison. The cloud fractions used in this project are from the total cloud fraction “CLDFRA” variable.

3 Results

3.1 CONUS Cloud Patterns

We first conduct an analysis of cloud patterns across the contiguous United States, and inter-compare clear-sky days estimated by TROPOMI, the ERA5 re-analysis, and the WRF-Chem model (Fig. 1). For TROPOMI, we define clear skies as the percentage of days with $qa_value > 0.75$, which almost exclusively filters based on cloud fractions < 0.5 ; cloud-free snow-covered scenes typically have a $qa_value > 0.75$ (Es-kes et al., 2022). For ERA5 and WRF-Chem, we define clear skies as the percentage of days with the total cloud fractions < 0.5 . ERA5 and WRF-Chem have similar clear-sky spatial patterns as TROPOMI but show systematically lower amounts of clear-skies by 8 %. The small systematic difference between TROPOMI and ERA5 when filtering for cloud fractions at 13:30 is likely driven by how optically thin cirrus-like clouds are handled; for TROPOMI these are being observed based on optical properties and therefore optically thin clouds are not assumed to be a cloud, whereas in weather models (ERA5 and WRF-Chem) these are being computed as vertical layers in the atmosphere with condensed water vapor. Overall, there is very strong agreement between the three datasets in the estimation of clouds giving us confidence that TROPOMI, ERA5, and WRF-Chem are all good estimators of daily clear-sky amounts.

For the remainder of this project, we define “clear sky” based on the TROPOMI NO₂ retrieval and use days with observations exceeding a qa_value of 0.75. According to TROPOMI – which is the only true observational dataset – the Southwest US has the most amount of clear-sky days per year (~ 80 % of days at 13:30 local time), while the interior Northeast US and coastal Northwest has the fewest (~ 30 % of days at 13:30 local time). The major US city with the most clear-sky days is Phoenix (79 % of days), while the major US city with the least clear-sky days is Seattle (29 % of days).

Annualized spatial cloud patterns are similar throughout the daylight hours with marginally more clear skies in the morning hours especially in the eastern US (Fig. S1 in the Supplement). Despite this, clouds are often transient, and there are opportunities to observe a clear sky measurement at a different hour of the day if the 13:30 observation is obstructed by clouds. In Fig. 2, we demonstrate that between 68 %–93 % of days have a clear sky measurement during any hour of the daytime as compared to the 33 %–69 % range at 13:30.

3.2 Surface NO₂: Clouds vs. No Clouds

We then link whether TROPOMI is observing a clear sky or not (i.e., $qa_value > 0.75$) to the daily in situ ground-

level NO₂ observations to determine how clouds are affecting surface NO₂ concentrations (hereafter referred to as surface NO₂). In Fig. 3, we show that surface NO₂ at 13:30 local time is +12.9 % larger (NMC = normalized mean change) [−3.8 % (10th percentile), +32.1 % (90th percentile)] on days with clouds at 13:30 compared to the annualized 13:30 average when all days of data are included. We also note the very strong correlation between the NO₂ on cloudy days and all days, which suggests that the presence of clouds drives a systematic change from the mean rather than a random change. We next show that the NO₂ during the average of all days is +17.2 % larger [−1.8 %, +38.7 %] than on days with only clear skies. We further show that surface NO₂ at 13:30 is +36.0 % larger [−6.1 %, +72.9 %] on days with clouds compared to days with clear skies.

The difference in surface NO₂ between cloudy and clear sky days can vary dramatically based on geographic region and proximity to a major roadway (Table 1). For the purposes of the sensitivity study, we focus on the cloudy versus cloud free days, while the directional changes of “cloudy versus all days” and “all days versus no clouds” values are similar (Tables S1 and S2 in the Supplement).

First, we find that NO₂ during cloudy days is larger in the northern US (+41.5 %) than the southern US (+28.5 %) and largest in the Northeast US (+61.7 %) (Fig. 4); for this analysis, 37° N is the dividing latitude between North and South, 100° W is the dividing longitude between East and West. Although the calculated cloudy versus no cloud change is independent of the number of days of clear-skies, areas of perpetually cloudy skies also have cooler temperatures, decreased photolysis, and shallower boundary layers which could cause much larger NO₂ on cloudy days. Interestingly, the Phoenix and Salt Lake City areas – two areas with large number of days with clear skies – also have a relatively large difference between cloudy and clear sky days demonstrating that the bias is independent of the number of days with clear skies. However, the *annualized* difference between cloudy and clear sky days in the Southwest US is modest (+4.8 %) (Table S1) because there are fewer individual days affected by clouds. Approximately 13 % of monitoring sites, mostly concentrated in the Los Angeles and San Diego areas, have lower NO₂ on cloudy days, and this may be driven by enhanced westerly winds on cloudy days bringing in cleaner marine air more than offsetting the photochemically driven larger NO₂ on cloudy days. Overall, while there are a few locations with lower NO₂ on cloudy days, 87 % of locations exhibit larger NO₂ on cloudy days and this is driven by the slower photochemistry on these days.

Proximity to roadways and large sources of NO_x is another driver of whether a location will experience a small – but generally still positive – difference in NO₂ on cloudy and clear sky days. For areas in close proximity to roadways (i.e., the near-road sites) ($n = 76$), the difference in NO₂ between cloudy and clear sky days is weaker: a smaller positive change (+15.9 %) and only 77 % of sites displaying a posi-




Figure 1. Percentage of clear-sky days over the contiguous US during 2019 from the TROPOMI NO₂ V2.4 product, ERA5 re-analysis, and WRF-Chem. (Top left) Normalized frequency diagram of the binned percentage of clear sky days for the three products. (Top right) Percentage of days in which the qa_value of the TROPOMI NO₂ V2.4 measurement was greater than 0.75. (Bottom left) Percentage of days in which the total cloud cover (tcc) from the ERA5 was less than 0.5. (Bottom right) Percentage of days in each grid cell in which the total cloud fraction from the WRF-Chem was less than 0.5.

tive mean change, which is less than the difference at all other NO₂ monitoring locations (+36.0 %).

We find that seasonal effects on the differences in NO₂ between cloudy and clear days are modest. The NO₂ on cloudy days in the Spring is largest and marginally smaller in other seasons. Other factors that were not associated with strong changes to the differences in NO₂ between cloudy and clear days bias are: the version of the TROPOMI NO₂ algorithm, whether the site was using a chemiluminescence or Cavity Attenuated Phase Shift measurement technique, and population/roadway density within a 0.5° radius.

3.3 TROPOMI NO₂: Clouds versus No Clouds

We then compare TROPOMI NO₂ measurements under varying sky conditions to understand how the retrieved NO₂ columns differ under cloudy and clear-sky conditions, but not to answer how they actually differ. For this exercise, we fil-

ter the TROPOMI NO₂ data strictly based on cloud radiative fraction (crf). Although it is recommended for most applications to use data when the crf < 0.5, sometimes measurements are usable in the presence of optically thick clouds (i.e., crf > 0.5). In Fig. 5, we average TROPOMI NO₂ measurements below and above a crf = 0.5 threshold to gain an understanding of how TROPOMI column NO₂ measurements intercompare in the presence and lack of optically thick clouds. In the figure we show the tropospheric vertical columns on the top row, and tropospheric slant columns in the middle row, which have been interconverted using the tropospheric air mass factor shown on the bottom row. As discussed in Sect. 2.2.1, the tropospheric air mass factor can be a large source of uncertainty when calculating tropospheric vertical columns from slant columns (Glissenaar et al., 2025; Liu et al., 2021; Rijdsdijk et al., 2025).

In Fig. 5, we demonstrate that the vertical column NO₂ spatial patterns in the presence of clouds are much different




Figure 2. Percentage of days over the contiguous US during 2019 with cloud fractions less than 0.5 as simulated by WRF-Chem at various local times: (Left) 13:30, (Right) any time between 07:00–19:00.




Figure 3. Scatterplots intercomparing annualized surface NO₂ from the EPA AQS at 13:30 local time during all days, cloudy days, and cloud free days. (Left) Annualized surface NO₂ during cloudy days compared to annualized surface NO₂ during all days. (Center) Annualized surface NO₂ during all days compared to annualized surface NO₂ during cloud free days. (Right) Annualized surface NO₂ during cloudy days compared to annualized surface NO₂ during no cloud. Each scatter point corresponds to each of the 374 measurement stations. A “cloudy” vs “no cloud” day is determined via the qa_{value} of 0.75 from the TROPOMI NO₂ V2.4 product.




Figure 4. (Left) Ratio of the annualized surface NO₂ during cloudy and cloud free days at the EPA AQS sites not classified as “near-road”. (Right) Same image but with an inverse distance weighting underlaid to infer geographic distribution of the ratio.

Table 1. Slope, r^2 , Normalized Mean Change (NMC), and number of sites of the “cloudy vs. no clouds” bias by further filtering out AQS data using various additional sensitivity analyses. Tables S1 and S2 show the sensitivity analyses for the “cloudy vs. all days” bias, and “all days vs. no clouds” bias respectively.

	Slope	r^2	Normalized Mean Change (%)	# sites of monitoring sites used
Baseline (V2.4)	1.20	0.85	+36.0 %	374
V2.3.1	1.18	0.86	+40.4 %	374
V2.4 crf < 0.5	1.25	0.83	+80.8 %	373
V2.4 all sites	1.05	0.90	+32.7 %	449
V2.4 near road only	0.89	0.84	+15.9 %	76
V2.4 no chemiluminescence	1.20	0.87	+53.1 %	26
V2.4 Summer only	1.17	0.86	+23.8 %	366
V2.4 Winter only	1.14	0.82	+27.8 %	373
V2.4 Spring only	1.28	0.88	+31.9 %	364
V2.4 Fall only	1.07	0.77	+30.9 %	359
V2.4 North only	1.31	0.89	+41.5 %	217
V2.4 South only	0.98	0.82	+28.5 %	157
V2.4 NorthEast only	1.36	0.93	+61.7 %	106
V2.4 SouthEast only	1.27	0.94	+33.8 %	73
V2.4 NorthWest only	1.12	0.88	+22.2 %	111
V2.4 SouthWest only	0.91	0.79	+23.9 %	84
V2.4 lowPopDensity only	1.34	0.86	+36.3 %	216
V2.4 highPopDensity only	1.13	0.76	+37.5 %	167
V2.4 lowRoadDensity only	1.19	0.82	+33.6 %	216
V2.4 highRoadDensity only	1.18	0.80	+40.8 %	165

in magnitude than the slant column NO₂ whereas the vertical column NO₂ spatial patterns in the absence of clouds are similar to the slant column NO₂. This is primarily driven by the assumed scattering weights in the retrieval. During cloudy scenes, scattering and reflection by clouds reduce the satellite’s sensitivity to near-surface NO₂, leading to smaller air mass factors as shown. Also, during measurements when the crf > 0.5, the uncertainty of the TROPOMI vertical column measurements rises, and this is driven by the difficulty in calculating the air mass factor in the presence of clouds; in addition to needing to know the vertical NO₂ profile for its calculation, we also need to know the pressure level and thickness of the clouds. Such errors can generate nonlinear responses. This analysis confirms that the assumed air mass factor is the driving factor causing the retrieved differences in the tropospheric vertical column NO₂ between clear and cloudy sky days, as the slant tropospheric column NO₂ is smaller during cloudy skies due to a lack of instrument sensitivity to the surface during cloudy conditions. Therefore, special care should be used when interpreting retrieved tropospheric satellite measurements in the presence of clouds.

Qualitatively, the ratio of the column NO₂ with and without clouds is spatially similar to the ratio from the AQS analysis – with the largest ratios occurring in the Northeast US and smallest ratios occurring in the Southwest US. However, quantitatively, the column ratio observed by TROPOMI is much larger in magnitude in the eastern US than the surface

ratio observed at the AQS surface sites. It is difficult to determine whether the quantitative magnitude is correct because there are no ground-based instruments to accurately measure column NO₂ in the presence of clouds.

3.4 WRF-Chem NO₂: Clouds vs. No Clouds

We then compare the differences in NO₂ between cloudy and clear days observed by the EPA AQS surface network to the differences in NO₂ between cloudy and clear days of surface NO₂ simulated by WRF-Chem. The 13:30 local time differences in NO₂ between cloudy and clear days of surface NO₂ in WRF-Chem (+58.7 %) is substantially larger than from the AQS observations (+36.0 %) during collocations. This directional change is consistent among all geographic regions suggesting that NO₂ concentrations are too responsive to sunlight in WRF-Chem.

There could be several reasons for this discrepancy. First, 91 % of monitors in the EPA monitoring network measure using the chemiluminescence method, NO₂^{*}, which quantifies NO₂ in addition to some fraction of PAN, alkyl nitrates, and HNO₃. The latter is problematic because the NO₂ + OH → HNO₃ reaction is a photochemically-driven pathway for NO₂ during daytime and if HNO₃ is additionally being measured then this would appear to buffer photolytically driven changes. We further conducted a sensitivity test in WRF-Chem and found that the NMC is only +42.1 % down from +58.7 % when a chemiluminescence correction factor from




Figure 5. (Left column) Annual 2019 TROPOMI NO₂ filtered using only a cloud radiative fraction (crf) filter less than 0.5. (Center column) Annual 2019 TROPOMI NO₂ filtered using only a crf filter greater than 0.5. (Right column) Ratio between the two annual averages. (Top row) Vertical tropospheric column NO₂ data. (Center row) Slant tropospheric column NO₂ data. (Bottom row) Tropospheric air mass factors.

Eq. (1) is used (Fig. 6c), indicating that some of the perceived differences between WRF-Chem and EPA monitors could be due to monitor interferences from PAN and HNO₃. Second, it is possible that radical concentrations (OH, HO₂, and/or RO₂) in WRF-Chem are fluctuating improperly in the presence of and lack of clouds (Duncan et al., 2024) causing NO₂ to either be removed too rapidly in the model or regenerated too slowly. Third, there might be insufficient photolysis of organic nitrates and/or particulate nitrates in the model which could buffer NO₂ photolysis-related changes; recent work has suggested that particulate nitrate can meaningfully photolyze back to NO₂ (Sarwar et al., 2024; Shah et al., 2024). Fourth, WRF-Chem may not simulate PBL depth properly and may have different biases during cloudy and clear sky conditions (Hegarty et al., 2018; Kuhn et al., 2024; Liu et al., 2023). For example, if the predicted PBL is too shallow during cloudy conditions, this could be a contributing factor to the simulated surface NO₂ bias. Errors in surface j NO₂ do not appear to be a primary driver of the cloudy versus clear sky disagreements as the j NO₂ values from WRF-Chem

seem reasonable as compared to UV-B measurements from the NOAA Surface Radiation Budget (SURFRAD) monitoring network (Fig. S4) and is consistent with other work showing small biases in j NO₂ in WRF-Chem (Ryu et al., 2018). Follow-up work will address some of these shortcomings by adding particulate nitrate photolysis into the chemical mechanism and evaluating PBL depths during cloudy conditions using ceilometers.

We can then use WRF-Chem as a transfer standard to suggest how column NO₂ may change in relation to the surface NO₂, and we find that the relative change in column NO₂ and surface NO₂ in response to clouds are very similar (Fig. 7). This makes intuitive sense because most NO₂ over the contiguous US is located within the boundary layer, and typically clouds (if they exist) are located at the top of the boundary layer. Any sunlight obstructed by clouds will also obstruct the NO₂ both at the surface and in the full boundary layer.




Figure 6. Scatterplots intercomparing annualized surface NO₂ at 13:30 local time during cloudy days vs. cloud free days. (Left) EPA AQS data which is a repeat of Fig. 3c. (Center) WRF-Chem collocated with the AQS monitoring sites, and using the WRF-Chem cloud filter in lieu of the TROPOMI cloud filter. (Right) WRF-Chem collocated with the AQS monitoring sites, comparing NO₂* instead of NO₂.




Figure 7. Ratio of the annualized surface NO₂ at 13:30 local time from WRF-Chem during cloudy and cloud free days. (Left) Surface NO₂ (Right) Tropospheric column NO₂.

3.5 Impacts of clouds on geostationary observations

Finally, we use provisional TEMPO NO₂ data, TROPOMI NO₂ data, and AQS NO₂ data from 2 August 2023 through 30 June 2024 to understand how the changes of NO₂ during clear and cloudy conditions may be altered at different hours of the day (Fig. 8). In this analysis, the threshold between high quality and lower quality data for both satellite products is a cloud radiative fraction = 0.15. Any TEMPO NO₂ or TROPOMI NO₂ measurement with $\text{crf} < 0.15$ was assumed to be “clear sky”, while all other measurements are assumed to be cloudy. Hours with low solar zenith angles (before 08:00 and after 16:00) have been excluded from this analysis. We find that the difference in surface NO₂ between clear and cloudy days is small in the early morning hours and increases throughout the day.

Surface AQS NO₂ at 08:30 local time is +8.3 % larger on cloudy days than clear sky days, while at 15:30 it is +52.2 % larger. The calculated 13:30 difference in surface NO₂ between cloudy and clear sky days using TEMPO (+25.1 %) is

similar to the analogous value from TROPOMI (+35.4 %). Differences between TEMPO and TROPOMI are expected because the cloud algorithms and instrument characteristics are different, even though the timeframe and cloud filter threshold used for this analysis are the same.

4 Discussion

In this project we quantify how NO₂ satellite data could be biased in estimating annualized surface NO₂ concentrations due to having high quality measurements only in the absence of clouds. We find that surface in situ NO₂ measurements are on average +17 % larger on all days compared to restricting to clear sky days and +36 % larger during cloudy days vs. clear sky days, with a wide distribution based on geographic region and proximity to roadway. Using the United States as a case study, we find the clear-sky bias to be largest in the Northeast US; conversely, the clear-sky bias is smallest in the Southwest US and near major roadways. In some areas of the urban Western US, Los Angeles and San Diego, we




Figure 8. Normalized mean percentage change in the surface NO₂ during days with cloudy skies as opposed to days with clear skies. Red dot shows the mean percentage change using TROPOMI clouds as shown in Fig. 2c. Black line uses the same procedure for August 2023–June 2024 data and TEMPO cloud data.

find that NO₂ is lower on cloudy days, but these instances are rare (13 % of monitoring sites) and are driven by unique transport patterns on cloudy days. Transport patterns are a significant driver of the regional clear vs. cloudy sky differences of surface NO₂ concentrations. Although the analysis was computed for both TROPOMI and TEMPO data, it should be re-emphasized that the cloud algorithms used by both instruments are different. However, the qualitative finding that surface NO₂ differences between cloudy and clear conditions tend to be larger in the afternoon than morning is consistent with a hypothesis that active photochemistry during periods of stronger afternoon sunlight would cause this change.

This work also highlights how NO₂ concentrations are different on days when satellite instruments are not acquiring a valid measurement. Our initial hypothesis of NO₂ being consistently larger on cloudy days was only partially proven true. In many cases, surface NO₂ concentrations and column NO₂ are larger, but this is not always the case. This project demonstrates the balancing act of the reduced NO₂+OH sink and local climatological patterns (wind speed/direction, PBL depth, etc.) driving surface NO₂ during cloudy conditions. Although one of the original goals of this study was to better gap-fill satellite tropospheric vertical column NO₂ measurements in the presence of clouds, ultimately, we were not comfortable doing this yet. Reliance on a model as a transfer standard to convert surface concentrations into column concentrations exhibited too many biases under cloudy

conditions. WRF-Chem model simulations of surface NO₂ suggest that the clear-sky bias in WRF-Chem is on average much larger than the observed clear-sky bias: +59 % on cloudy days vs. clear days for the model, and +36 % for the AQS data. We hypothesized that errors in OH chemistry, NO₂ recycling speeds, and PBL mixing depths could all be contributing to this high bias. Future work should target these three research topics. Future work could also use a machine-learning approach to account for some of these model biases.

Another consideration with the interpretation of satellite measurements is the impact of lightning NO_x, wildfire NO_x, and aircraft NO_x emissions, mostly staying aloft, which could be misinterpreted as surface NO₂ enhancements. While lightning NO_x and wildfire NO_x emissions are often screened out when applying a cloud filter because they occur in optically thick clouds/smoke, it is possible for the NO₂ to remain aloft for several days after the initial thunderstorm/fire and be observed during clear skies. An algorithm to detect and screen out downwind NO₂ attributed to upwind lightning NO_x and wildfire NO_x emissions could be especially helpful. At minimum, care should be taken during timeframes and regions where there are large pulses of these types of emissions, such as our findings during summer.

In some ways, the chosen year 2019 was an ideal year to conduct the analysis because it preceded the 2020 global pandemic and its nonlinear and lingering effects on air pollution. But in other ways, this year was less ideal because TROPOMI pixel sizes changed in August 2019 from $7 \times 3.5 \text{ km}^2$ ($\sim 25 \text{ km}^2$) to $5.5 \times 3.5 \text{ km}^2$ ($\sim 19 \text{ km}^2$). The fraction of clear-sky pixels likely increased by 1 %–2 % after August 2019 as smaller pixel sizes can better “see around” clouds (Krijger et al., 2007). This probably did not meaningfully affect our analysis but is nonetheless a caveat of using 2019 data.

These results have repercussions for many applied studies that use satellite data to estimate surface NO₂ concentrations or NO_x emissions. First, for studies that estimate surface concentrations, it is important to ingest surface NO₂ measurements during cloudy (and nighttime) conditions in some capacity in order to appropriately estimate 24 h concentrations; most studies already do this. If one were to use the clear-sky satellite data coupled with only a chemical transport model as a transfer standard to convert the column measurement into a pseudo-surface “measurement”, this would underestimate annualized NO₂ concentration in most places. Unfortunately, there are many global regions with few or no surface measurements, so this is an important consideration when estimating surface NO₂ in these regions. But even if one were to ingest surface NO₂ during cloudy conditions, the spatial patterns of surface NO₂ during cloudy conditions may be slightly different than implied by the clear-sky satellite data. For example, we find that NO₂ surface concentrations under cloudy conditions are much larger in the Northeast US than the Southwest US, and a cloud-free satellite map does not capture this.

Second, for nitrogen oxide emissions estimates it is often assumed that anthropogenic emission rates are similar under cloudy and clear-sky conditions, but this is likely not the case in reality. Although we show that surface NO₂ concentrations are typically smaller under clear-skies, it is likely that anthropogenic NO_x emissions are actually larger under regionwide clear-skies during summer and winter due to the moderating impact of clouds on surface temperature and subsequent impacts on heating-ventilation-air conditioning (HVAC) usage/emissions (Abel et al., 2017). If we were able to better independently estimate tropospheric vertical column NO₂ during cloudy conditions, perhaps this could be investigated in the future.

Lastly, as satellite-derived NO₂ applications increase over the coming years, it is important to document its successes and shortcomings. We see this project as a first-step towards better accounting for the clear-sky bias of satellite NO₂ data. While future NO₂ applications may use geostationary data, such as TEMPO, which may suffer from a similar bias depending on the hour of the day, an advantage of geostationary satellite data is the ability to use multiple measurements per day before and just after the clouds. It might be possible to isolate a two-hour window (one with a cloud and one without) to get a better handle on the instantaneous versus long-term role of clouds affecting NO₂ concentrations.

This work also highlights the critical role that chemical transport models can play in satellite NO₂ applications. Errors in the model assumptions can hamstring many NO₂ applications. For example, using a model to infer NO₂ during cloudy conditions in the lack of clear-sky satellite data would yield significant errors. Therefore, future work should concurrently focus on acquiring and using sub-orbital measurements to diagnose errors related in simulating NO₂ in chemical transport models, so that they can be used as more robust transfer standards.

Code and data availability. TROPOMI NO₂ version 2.4 data (<https://doi.org/10.5270/S5P-9bnp8q8>, Copernicus Sentinel-5P, 2021) processed to 0.01° × 0.01° resolution (<https://doi.org/10.5067/MKJG22GUOD34>, Goldberg, 2024) and TEMPO NO₂ version 3 data (https://doi.org/10.5067/IS-40e/TEMPO/NO2_L3.003, NASA/LARC/SD/ASDC, 2024) can be freely downloaded from NASA Earthdata. EPA AQS surface NO₂ data can be downloaded from pre-generated files: https://aq5.epa.gov/aq5web/airdata/download_files.html (last access: 9 July 2025). ERA5 re-analysis hourly data on single levels (<https://doi.org/10.24381/cds.adbb2d47>, Hersbach et al., 2023) can be downloaded from Copernicus Climate Data Store (<https://cds.climate.copernicus.eu/#/!home>, last access: 28 December 2023). NOAA SURFAD data can be downloaded from: <https://gml.noaa.gov/grad/surfrad/sitepage.html> (last access: 20 February 2025). Output from the WRF-Chem simulation is available upon request. IDL code and processed data to create the figures can be accessed at: <https://doi.org/10.6084/m9.figshare.30376426>.

Supplement. The Supplement includes: (1) a spatial plot of the annual average of days with total cloud fraction less 0.5 as simulated by WRF-Chem during 08:00 a.m., 01:00 p.m., and 05:00 p.m. local time, (2) scatterplots of various 2019 annual averages of surface NO₂ and TROPOMI NO₂ measurements, (3) *j*NO₂ from WRF-Chem and an intercomparison with the NOAA SURFRAD network, (4) Tables of “cloudy vs. all days” and “all days vs. no clouds” analogous to Table 1 which shows “cloudy vs. no clouds”. The supplement related to this article is available online at <https://doi.org/10.5194/acp-25-16287-2025-supplement>.

Author contributions. DG, AC, SK, and SA developed the project design. JH and CL set-up and conducted the WRF-Chem simulations. DG downloaded and processed the TROPOMI NO₂, TEMPO NO₂, ERA5, and SURFRAD data and re-gridded all data to a standardized grid. MON helped to process the surface NO₂ data. DG developed all figures for the manuscript and wrote the paper. All authors edited the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. Preparation of this manuscript was funded by grants from the NOAA GeoXO program (1305M323PNRMA0668) and the NASA Health and Air Quality Applied Sciences Team (HAQAST) (80NSSC21K0511). NOAA Cooperative Agreement (NA17OAR4320101 and NA22OAR4320151) funded C. Lyu and J. He. The WRF-Chem simulation was supported by NOAA’s High Performance Computing Program. The authors would also like to thank Brian McDonald and Laura Judd for very helpful feedback during preparation of this manuscript.

Financial support. This research has been supported by the National Oceanic and Atmospheric Administration (grant nos. 1305M323PNRMA0668, NA17OAR432010, and NA22OAR4320151) and the National Aeronautics and Space Administration (grant no. 80NSSC21K0511 and 80NSSC23K1002).

Review statement. This paper was edited by Michel Van Roozendael and reviewed by two anonymous referees.

References

- Abel, D. W., Holloway, T., Kladar, R. M., Meier, P., Ahl, D., Harkey, M., and Patz, J.: Response of Power Plant Emissions to Ambient Temperature in the Eastern United States, *Environ. Sci. Technol.*, 51, 5838–5846, <https://doi.org/10.1021/acs.est.6b06201>, 2017.
- Acker, S., Holloway, T., and Harkey, M.: Satellite detection of NO₂ distributions using TROPOMI and TEMPO and comparison with ground-based concentration measurements, *Atmos. Chem. Phys.*, 25, 8271–8288, <https://doi.org/10.5194/acp-25-8271-2025>, 2025.
- Anenberg, S. C., Mohegh, A., Goldberg, D. L., Kerr, G. H., Brauer, M., Burkart, K., Hystad, P., Larkin, A., Wozniak, S., and Lamsal, L.: Long-term trends in urban NO₂ concentrations and associated paediatric asthma incidence: estimates from global datasets, *Lancet Planet Health*, 6, e49–e58, [https://doi.org/10.1016/S2542-5196\(21\)00255-2](https://doi.org/10.1016/S2542-5196(21)00255-2), 2022.
- Bechle, M. J., Millet, D. B., and Marshall, J. D.: National Spatiotemporal Exposure Surface for NO₂: Monthly Scaling of a Satellite-Derived Land-Use Regression, 2000–2010, *Environ. Sci. Technol.*, 49, 12297–12305, <https://doi.org/10.1021/acs.est.5b02882>, 2015.
- Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veefkind, J. P., Stammes, P., Huijnen, V., Kleipool, Q. L., Sneep, M., Claas, J., Leitão, J., Richter, A., Zhou, Y., and Brunner, D.: An improved tropospheric NO₂ column retrieval algorithm for the Ozone Monitoring Instrument, *Atmos. Meas. Tech.*, 4, 1905–1928, <https://doi.org/10.5194/amt-4-1905-2011>, 2011.
- Burnett, R. T., Stieb, D., Brook, J. R., Cakmak, S., Dales, R., Raizenne, M., Vincent, R., and Dann, T.: Associations between short-term changes in nitrogen dioxide and mortality in Canadian cities, *Arch. Environ. Health*, 59, 228–236, <https://doi.org/10.3200/AEOH.59.5.228-236>, 2004.
- Busca, G., Lietti, L., Ramis, G., and Berti, F.: Chemical and mechanistic aspects of the selective catalytic reduction of NO(x) by ammonia over oxide catalysts: A review, *Appl. Catal. B*, 18, 1–36, [https://doi.org/10.1016/S0926-3373\(98\)00040-X](https://doi.org/10.1016/S0926-3373(98)00040-X), 1998.
- Cao, E. L.: National ground-level NO₂ predictions via satellite imagery driven convolutional neural networks, *Front Environ. Sci.*, 11, <https://doi.org/10.3389/fenvs.2023.1285471>, 2023.
- Copernicus Sentinel-5P: TROPOMI Level 2 Nitrogen Dioxide total column products, Version 02, European Space Agency [data set], <https://doi.org/10.5270/S5P-9bnp8q8>, 2021.
- Crippa, M., Guizzardi, D., Pisoni, E., Solazzo, E., Guion, A., Muntean, M., Florczyk, A., Schiavina, M., Melchiorri, M., and Hutfilter, A. F.: Global anthropogenic emissions in urban areas: patterns, trends, and challenges, *Environmental Research Letters*, 16, 074033, <https://doi.org/10.1088/1748-9326/AC00E2>, 2021.
- de Foy, B., Krotkov, N. A., Bei, N., Herndon, S. C., Huey, L. G., Martínez, A.-P., Ruiz-Suárez, L. G., Wood, E. C., Zavala, M., and Molina, L. T.: Hit from both sides: tracking industrial and volcanic plumes in Mexico City with surface measurements and OMI SO₂ retrievals during the MILAGRO field campaign, *Atmos. Chem. Phys.*, 9, 9599–9617, <https://doi.org/10.5194/acp-9-9599-2009>, 2009.
- Demetillo, M. A. G., Navarro, A., Knowles, K. K., Fields, K. P., Geddes, J. A., Nowlan, C. R., Janz, S. J., Judd, L. M., Al-Saadi, J., Sun, K., McDonald, B. C., Diskin, G. S., and Pusede, S. E.: Observing Nitrogen Dioxide Air Pollution Inequality Using High-Spatial-Resolution Remote Sensing Measurements in Houston, Texas, *Environ. Sci. Technol.*, 54, 9882–9895, <https://doi.org/10.1021/acs.est.0c01864>, 2020.
- Dickerson, R. R., Anderson, D. C., and Ren, X.: On the use of data from commercial NO_x analyzers for air pollution studies, *Atmos. Environ.*, 214, 116873, <https://doi.org/10.1016/j.atmosenv.2019.116873>, 2019.
- Dressel, I. M., Demetillo, M. A. G., Judd, L. M., Janz, S. J., Fields, K. P., Sun, K., Fiore, A. M., McDonald, B. C., and Pusede, S. E.: Daily Satellite Observations of Nitrogen Dioxide Air Pollution Inequality in New York City, New York and Newark, New Jersey: Evaluation and Application, *Environ. Sci. Technol.*, 56, 15298–15311, <https://doi.org/10.1021/acs.est.2c02828>, 2022.
- Duncan, B. N., Anderson, D. C., Fiore, A. M., Joiner, J., Krotkov, N. A., Li, C., Millet, D. B., Nicely, J. M., Oman, L. D., St. Clair, J. M., Shutter, J. D., Souri, A. H., Strode, S. A., Weir, B., Wolfe, G. M., Worden, H. M., and Zhu, Q.: Opinion: Beyond global means – novel space-based approaches to indirectly constrain the concentrations of and trends and variations in the tropospheric hydroxyl radical (OH), *Atmos. Chem. Phys.*, 24, 13001–13023, <https://doi.org/10.5194/acp-24-13001-2024>, 2024.
- Eskes, H., van Geffen, J., Boersma, F., Eichman, K., Apituley, A., Pedernana, M., Sneep, M., Veefkind, J., and Loyola, D.: Sentinel-5 precursor/TROPOMI Level 2 Product User Manual Nitrogen dioxide, https://sentiwiki.copernicus.eu/_attachments/1673595/S5P-KNMI-L2-0021-MA-Sentinel-5P-TROPOMI-Level-2-Product-User-Manual-Nitrogen-Dioxide-2024-4.4.0.pdf (last access: 14 November 2025), 2022.
- Geddes, J. A., Murphy, J. G., O'Brien, J. M., and Celarier, E. A.: Biases in long-term NO₂ averages inferred from satellite observations due to cloud selection criteria, *Remote Sens. Environ.*, 124, 210–216, <https://doi.org/10.1016/j.rse.2012.05.008>, 2012.
- Ghahremanloo, M., Lops, Y., Choi, Y., and Yeganeh, B.: Deep Learning Estimation of Daily Ground-Level NO₂ Concentrations From Remote Sensing Data, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034925, <https://doi.org/10.1029/2021JD034925>, 2021.
- Ghahremanloo, M., Lops, Y., Choi, Y., Mousavinezhad, S., and Jung, J.: A Coupled Deep Learning Model for Estimating Surface NO₂ Levels from Remote Sensing Data: 15-Year Study Over the Contiguous United States, *Journal of Geophysical Research: Atmospheres*, e2022JD037010, <https://doi.org/10.1029/2022JD037010>, 2023.
- Glissenaar, I., Boersma, K. F., Angloul, I., Rijdsdijk, P., Verhoelst, T., Compernelle, S., Pinardi, G., Lambert, J.-C., Van Roozendael, M., and Eskes, H.: TROPOMI Level 3 tropospheric NO₂ dataset with advanced uncertainty analysis from the ESA CCI+ECV precursor project, *Earth Syst. Sci. Data*, 17, 4627–4650, <https://doi.org/10.5194/essd-17-4627-2025>, 2025.
- Goldberg, D.: HAQAST Sentinel-5P TROPOMI Nitrogen Dioxide (NO₂) CONUS Monthly Level 3 0.01 x 0.01 Degree Gridded Data V2.4, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC) [data set], <https://doi.org/10.5067/MKJG22GUOD34>, 2024.
- Goldberg, D. L., Anenberg, S. C., Kerr, G. H., Mohegh, A., Lu, Z., and Streets, D. G.: TROPOMI NO₂ in the United States: A Detailed Look at the Annual Averages, Weekly Cycles, Effects of Temperature, and Correlation With Sur-

- face NO₂ Concentrations, *Earths Future*, 9, e2020EF001665, <https://doi.org/10.1029/2020EF001665>, 2021.
- Goldberg, D. L., Harkey, M., de Foy, B., Judd, L., Johnson, J., Yarwood, G., and Holloway, T.: Evaluating NO_x emissions and their effect on O₃ production in Texas using TROPOMI NO₂ and HCHO, *Atmos. Chem. Phys.*, 22, 10875–10900, <https://doi.org/10.5194/acp-22-10875-2022>, 2022.
- Goldberg, D. L., Tao, M., Kerr, G. H., Ma, S., Tong, D. Q., Fiore, A. M., Dickens, A. F., Adelman, Z. E., and Anenberg, S. C.: Evaluating the spatial patterns of U.S. urban NO_x emissions using TROPOMI NO₂, *Remote Sens. Environ.*, 300, 113917, <https://doi.org/10.1016/j.rse.2023.113917>, 2024.
- Harkey, M. and Holloway, T.: Simulated Surface-Column NO₂ Connections for Satellite Applications, *Journal of Geophysical Research: Atmospheres*, 129, <https://doi.org/10.1029/2024JD041912>, 2024.
- He, J., Harkins, C., O'Dell, K., Li, M., Francoeur, C., Aikin, K. C., Anenberg, S., Baker, B., Brown, S. S., Coggon, M. M., Frost, G. J., Gilman, J. B., Kondragunta, S., Lamplugh, A., Lyu, C., Moon, Z., Pierce, B. R., Schwantes, R. H., Stockwell, C. E., Warneke, C., Yang, K., Nowlan, C. R., González Abad, G., and McDonald, B. C.: COVID-19 perturbation on US air quality and human health impact assessment, *PNAS Nexus*, 3, <https://doi.org/10.1093/pnasnexus/pgad483>, 2024.
- He, M. Z., Kinney, P. L., Li, T., Chen, C., Sun, Q., Ban, J., Wang, J., Liu, S., Goldsmith, J., and Kioumourtzoglou, M. A.: Short- and intermediate-term exposure to NO₂ and mortality: A multi-county analysis in China, *Environmental Pollution*, 261, 114165, <https://doi.org/10.1016/j.envpol.2020.114165>, 2020.
- Health Effects Institute: Systematic Review and Meta-analysis of Selected Health Effects of Long-Term Exposure to Traffic-Related Air Pollution, https://www.healtheffects.org/system/files/hej-special-report-23_6.pdf (last access: 14 November 2025), 2022.
- Hegarty, J. D., Lewis, J., McGrath-Spangler, E. L., Henderson, J., Scarino, A. J., DeCola, P., Ferrare, R., Hicks, M., Adams-Selin, R. D., and Welton, E. J.: Analysis of the Planetary Boundary Layer Height during DISCOVER-AQ Baltimore–Washington, D.C., with Lidar and High-Resolution WRF Modeling, *J. Appl. Meteorol. Climatol.*, 57, 2679–2696, <https://doi.org/10.1175/JAMC-D-18-0014.1>, 2018.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2023.
- Jin, X., Zhu, Q., and Cohen, R. C.: Direct estimates of biomass burning NO_x emissions and lifetimes using daily observations from TROPOMI, *Atmos. Chem. Phys.*, 21, 15569–15587, <https://doi.org/10.5194/acp-21-15569-2021>, 2021.
- Judd, L. M., Al-Saadi, J. A., Szykman, J. J., Valin, L. C., Janz, S. J., Kowalewski, M. G., Eskes, H. J., Veefkind, J. P., Cede, A., Mueller, M., Gebetsberger, M., Swap, R., Pierce, R. B., Nowlan, C. R., Abad, G. G., Nehrir, A., and Williams, D.: Evaluating Sentinel-5P TROPOMI tropospheric NO₂ column densities with airborne and Pandora spectrometers near New York City and Long Island Sound, *Atmos. Meas. Tech.*, 13, 6113–6140, <https://doi.org/10.5194/amt-13-6113-2020>, 2020.
- Kebabian, P. L., Wood, E. C., Herndon, S. C., and Freedman, A.: A practical alternative to chemiluminescence-based detection of nitrogen dioxide: Cavity attenuated phase shift spectroscopy, *Environ. Sci. Technol.*, 42, 6040–6045, <https://doi.org/10.1021/es703204j>, 2008.
- Kelly, K. E., Whitaker, J., Petty, A., Widmer, C., Dybwad, A., Sleeth, D., Martin, R. V., and Butterfield, A.: Ambient and laboratory evaluation of a low-cost particulate matter sensor, *Environmental Pollution*, 221, 491–500, <https://doi.org/10.1016/j.envpol.2016.12.039>, 2017.
- Kenagy, H. S., Sparks, T. L., Ebben, C. J., Wooldrige, P. J., Lopez-Hilfiker, F. D., Lee, B. H., Thornton, J. A., McDuffie, E. E., Fibiger, D. L., Brown, S. S., Montzka, D. D., Weinheimer, A. J., Schroder, J. C., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Dibb, J. E., Campos, T., Shah, V., Jaeglé, L., and Cohen, R. C.: NO_x Lifetime and NO_y Partitioning During WINTER, *Journal of Geophysical Research: Atmospheres*, 123, 9813–9827, <https://doi.org/10.1029/2018JD028736>, 2018.
- Kerr, G. H., Goldberg, D. L., Harris, M. H., Henderson, B. H., Hystad, P., Roy, A., and Anenberg, S. C.: Ethnoracial Disparities in Nitrogen Dioxide Pollution in the United States: Comparing Data Sets from Satellites, Models, and Monitors, *Environ. Sci. Technol.*, <https://doi.org/10.1021/acs.est.3c03999>, 2023.
- Khreis, H., Kelly, C., Tate, J., Parslow, R., Lucas, K., and Nieuwenhuijsen, M.: Exposure to traffic-related air pollution and risk of development of childhood asthma: A systematic review and meta-analysis, *Environ. Int.*, 100, 1–31, <https://doi.org/10.1016/j.envint.2016.11.012>, 2017.
- Kim, E. J., Holloway, T., Kokandakar, A., Harkey, M., Elkins, S., Goldberg, D. L., and Heck, C.: A Comparison of Regression Methods for Inferring Near-Surface NO₂ With Satellite Data, *Journal of Geophysical Research: Atmospheres*, 129, <https://doi.org/10.1029/2024JD040906>, 2024.
- Koltsakis, G. and Stamatelos, A.: Catalytic automotive exhaust aftertreatment, *Prog. Energy Combust. Sci.*, 23, 1–39, [https://doi.org/10.1016/s0360-1285\(97\)00003-8](https://doi.org/10.1016/s0360-1285(97)00003-8), 1997.
- Krijger, J. M., van Weele, M., Aben, I., and Frey, R.: Technical Note: The effect of sensor resolution on the number of cloud-free observations from space, *Atmos. Chem. Phys.*, 7, 2881–2891, <https://doi.org/10.5194/acp-7-2881-2007>, 2007.
- Kuhn, L., Beirle, S., Kumar, V., Osipov, S., Pozzer, A., Bösch, T., Kumar, R., and Wagner, T.: On the influence of vertical mixing, boundary layer schemes, and temporal emission profiles on tropospheric NO₂ in WRF-Chem – comparisons to in situ, satellite, and MAX-DOAS observations, *Atmos. Chem. Phys.*, 24, 185–217, <https://doi.org/10.5194/acp-24-185-2024>, 2024.

- Lambert, J.-C., Claas, J., Stein-Zweers, D., Ludewig, A., Loyola, D., Sneep, M., and Dehn, A.: Quarterly Validation Report of the Copernicus Sentinel-5 Precursor Operational Data Products #19, https://mpc-vdaf.tropomi.eu/index.php?option=com_vdaf&view=showReport&format=rawhtml&id=57 (last access: 14 November 2025), 2023.
- Lamsal, L. N., Martin, R. V., van Donkelaar, A., Steinbacher, M., Celarier, E. A., Bucsela, E. J., Dunlea, E. J., and Pinto, J. P.: Ground-level nitrogen dioxide concentrations inferred from the satellite-borne Ozone Monitoring Instrument, *Journal of Geophysical Research Atmospheres*, 113, 1–15, <https://doi.org/10.1029/2007JD009235>, 2008.
- Larkin, A., Anenberg, S., Goldberg, D. L., Moheg, A., Brauer, M., and Hystad, P.: A global spatial-temporal land use regression model for nitrogen dioxide air pollution, *Front. Environ. Sci.*, 11, 484, <https://doi.org/10.3389/FENVS.2023.1125979>, 2023.
- Laughner, J. L. and Cohen, R. C.: Direct observation of changing NO_x lifetime in North American cities, *Science*, 366, 723–727, <https://doi.org/10.1126/science.aax6832>, 2019.
- Levelt, P. F., Joiner, J., Tamminen, J., Veefkind, J. P., Bhartia, P. K., Stein Zweers, D. C., Duncan, B. N., Streets, D. G., Eskes, H., van der A, R., McLinden, C., Fioletov, V., Carn, S., de Laat, J., DeLand, M., Marchenko, S., McPeters, R., Ziemke, J., Fu, D., Liu, X., Pickering, K., Apituley, A., González Abad, G., Arola, A., Boersma, F., Chan Miller, C., Chance, K., de Graaf, M., Hakkarainen, J., Hassinen, S., Ialongo, I., Kleipool, Q., Krotkov, N., Li, C., Lamsal, L., Newman, P., Nowlan, C., Suleiman, R., Tilstra, L. G., Torres, O., Wang, H., and Wargan, K.: The Ozone Monitoring Instrument: overview of 14 years in space, *Atmos. Chem. Phys.*, 18, 5699–5745, <https://doi.org/10.5194/acp-18-5699-2018>, 2018.
- Lin, M., Horowitz, L. W., Hu, L., and Permar, W.: Reactive Nitrogen Partitioning Enhances the Contribution of Canadian Wild-fire Plumes to US Ozone Air Quality, *Geophys. Res. Lett.*, 51, <https://doi.org/10.1029/2024GL109369>, 2024.
- Liu, F., Beirle, S., Zhang, Q., Dörner, S., He, K., and Wagner, T.: NO_x lifetimes and emissions of cities and power plants in polluted background estimated by satellite observations, *Atmos. Chem. Phys.*, 16, 5283–5298, <https://doi.org/10.5194/acp-16-5283-2016>, 2016.
- Liu, S., Valks, P., Pinardi, G., Xu, J., Chan, K. L., Argyrouli, A., Lutz, R., Beirle, S., Khorsandi, E., Baier, F., Huijnen, V., Bais, A., Donner, S., Dörner, S., Gratsea, M., Hendrick, F., Karagkiozidis, D., Lange, K., PETERS, A. J. M., Remmers, J., Richter, A., Van Roozendaal, M., Wagner, T., Wenig, M., and Loyola, D. G.: An improved TROPOMI tropospheric NO₂ research product over Europe, *Atmos. Meas. Tech.*, 14, 7297–7327, <https://doi.org/10.5194/amt-14-7297-2021>, 2021.
- Liu, X., Wang, Y., Wasti, S., Li, W., Soleimanian, E., Flynn, J., Griggs, T., Alvarez, S., Sullivan, J. T., Roots, M., Twigg, L., Gronoff, G., Berkoff, T., Walter, P., Estes, M., Hair, J. W., Shingler, T., Scarino, A. J., Fenn, M., and Judd, L.: Evaluating WRF-GC v2.0 predictions of boundary layer height and vertical ozone profile during the 2021 TRACER-AQ campaign in Houston, Texas, *Geosci. Model Dev.*, 16, 5493–5514, <https://doi.org/10.5194/gmd-16-5493-2023>, 2023.
- Lorente, A., Folkert Boersma, K., Yu, H., Dörner, S., Hilboll, A., Richter, A., Liu, M., Lamsal, L. N., Barkley, M., De Smedt, I., Van Roozendaal, M., Wang, Y., Wagner, T., Beirle, S., Lin, J.-T., Krotkov, N., Stammes, P., Wang, P., Eskes, H. J., and Krol, M.: Structural uncertainty in air mass factor calculation for NO₂ and HCHO satellite retrievals, *Atmos. Meas. Tech.*, 10, 759–782, <https://doi.org/10.5194/amt-10-759-2017>, 2017.
- Martin, R. V., Brauer, M., van Donkelaar, A., Shaddick, G., Narain, U., and Dey, S.: No one knows which city has the highest concentration of fine particulate matter, *Atmos. Environ.*, 100040, <https://doi.org/10.1016/J.AEAOA.2019.100040>, 2019.
- Maruhashi, J., Mertens, M., Grewe, V., and Dedoussi, I. C.: A multi-method assessment of the regional sensitivities between flight altitude and short-term O₃ climate warming from aircraft NO_x emissions, *Environmental Research Letters*, <https://doi.org/10.1088/1748-9326/ad376a>, 2024.
- NASA/LARC/SD/ASDC: TEMPO gridded NO₂ tropospheric and stratospheric columns V03 (PROVISIONAL), NASA Langley Atmospheric Science Data Center DAAC [data set], https://doi.org/10.5067/IS-40e/TEMPO/NO2_L3.003, 2024.
- Nault, B. A., Laughner, J. L., Wooldridge, P. J., Crounse, J. D., Dibb, J., Diskin, G., Peischl, J., Podolske, J. R., Pollack, I. B., Ryerson, T. B., Scheuer, E., Wennberg, P. O., and Cohen, R. C.: Lightning NO_x Emissions: Reconciling Measured and Modeled Estimates With Updated NO_x Chemistry, *Geophys. Res. Lett.*, 44, 9479–9488, <https://doi.org/10.1002/2017GL074436>, 2017.
- Nawaz, M. O., Goldberg, D. L., Kerr, G. H., and Anenberg, S. C.: TROPOMI Satellite Data Reshape NO₂ Air Pollution Land-Use Regression Modeling Capabilities in the United States, *ACS ES&T Air*, <https://doi.org/10.1021/acsestair.4c00153>, 2025.
- Nawaz, M. O., Johnson, J., Yarwood, G., de Foy, B., Judd, L., and Goldberg, D. L.: An intercomparison of satellite, airborne, and ground-level observations with WRF–CAMx simulations of NO₂ columns over Houston, Texas, during the September 2021 TRACER-AQ campaign, *Atmos. Chem. Phys.*, 24, 6719–6741, <https://doi.org/10.5194/acp-24-6719-2024>, 2024.
- Nowlan, C. R., Abad, G. G., Liu, X., Wang, H., and Chance, K.: TEMPO Nitrogen Dioxide Retrieval Algorithm Theoretical Basis Document, <https://doi.org/10.5067/WX026254FI2U>, 2025.
- Palmer, P. I., Jacob, D. J., Chance, K. V., Martin, R. V., Spurr, R. J. D., Kurosu, T. P., Bey, I., Yantosca, R., Fiore, A. M., and Li, Q.: Air mass factor formulation for spectroscopic measurements from satellites: Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment, *Journal of Geophysical Research: Atmospheres*, 106, 14539–14550, <https://doi.org/10.1029/2000JD900772>, 2001.
- Platt, U.: Differential Optical Absorption Spectroscopy (DOAS), in: *Air monitoring by spectroscopic techniques*, Wiley-IEEE, 531, 27–84, 1994.
- Poraicu, C., Müller, J.-F., Stavrakou, T., Fonteyn, D., Tack, F., Deutsch, F., Laffineur, Q., Van Malderen, R., and Veldeman, N.: Cross-evaluating WRF-Chem v4.1.2, TROPOMI, APEX, and in situ NO₂ measurements over Antwerp, Belgium, *Geosci. Model Dev.*, 16, 479–508, <https://doi.org/10.5194/gmd-16-479-2023>, 2023.
- Rijsdijk, P., Eskes, H., Dingemans, A., Boersma, K. F., Sekiya, T., Miyazaki, K., and Houweling, S.: Quantifying uncertainties in satellite NO₂ superobservations for data assimilation and model evaluation, *Geosci. Model Dev.*, 18, 483–509, <https://doi.org/10.5194/gmd-18-483-2025>, 2025.
- Ryu, Y.-H., Hodzic, A., Barre, J., Descombes, G., and Minnis, P.: Quantifying errors in surface ozone predictions associated with

- clouds over the CONUS: a WRF-Chem modeling study using satellite cloud retrievals, *Atmos. Chem. Phys.*, 18, 7509–7525, <https://doi.org/10.5194/acp-18-7509-2018>, 2018.
- Sarwar, G., Hogrefe, C., Henderson, B. H., Mathur, R., Gilliam, R., Callaghan, A. B., Lee, J., and Carpenter, L. J.: Impact of particulate nitrate photolysis on air quality over the Northern Hemisphere, *Science of The Total Environment*, 917, 170406, <https://doi.org/10.1016/j.scitotenv.2024.170406>, 2024.
- Shah, V., Jacob, D. J., Li, K., Silvern, R. F., Zhai, S., Liu, M., Lin, J., and Zhang, Q.: Effect of changing NO_x lifetime on the seasonality and long-term trends of satellite-observed tropospheric NO₂ columns over China, *Atmos. Chem. Phys.*, 20, 1483–1495, <https://doi.org/10.5194/acp-20-1483-2020>, 2020.
- Shah, V., Keller, C. A., Knowland, K. E., Christiansen, A., Hu, L., Wang, H., Lu, X., Alexander, B., and Jacob, D. J.: Particulate Nitrate Photolysis as a Possible Driver of Rising Tropospheric Ozone, *Geophys. Res. Lett.*, 51, <https://doi.org/10.1029/2023GL107980>, 2024.
- Shetty, S., Schneider, P., Stebel, K., David Hamer, P., Kylling, A., and Koren Berntsen, T.: Estimating surface NO₂ concentrations over Europe using Sentinel-5P TROPOMI observations and Machine Learning, *Remote Sens Environ.*, 312, 114321, <https://doi.org/10.1016/j.rse.2024.114321>, 2024.
- Sullivan, D. M. and Krupnick, A.: Using Satellite Data to Fill the Gaps in the US Air Pollution Monitoring Network, NW, https://media.rff.org/documents/RFF20WP-18-21_0.pdf (last access: 14 November 2025), 2018.
- Sun, W., Tack, F., Clarisse, L., Schneider, R., Stavrou, T., and Van Roozendaal, M.: Inferring Surface NO₂ Over Western Europe: A Machine Learning Approach With Uncertainty Quantification, *Journal of Geophysical Research: Atmospheres*, 129, <https://doi.org/10.1029/2023JD040676>, 2024.
- Tack, F., Merlaud, A., Iordache, M.-D., Pinardi, G., Dimitropoulou, E., Eskes, H. J., Bomans, B., Veefkind, P., and Van Roozendaal, M.: Assessment of the TROPOMI tropospheric NO₂ product based on airborne APEX observations, *Atmos. Meas. Tech.*, 14, 615–646, <https://doi.org/10.5194/amt-14-615-2021>, 2021.
- Thornton, J. A., Wooldridge, P. J., and Cohen, R. C.: Atmospheric NO₂: In Situ Laser-Induced Fluorescence Detection at Parts per Trillion Mixing Ratios, *Anal. Chem.*, 72, 528–539, <https://doi.org/10.1021/ac9908905>, 2000.
- U.S. EPA: FY2023 and 2024 National Program Manager Guidance Monitoring Appendix, https://www.epa.gov/system/files/documents/2023-02/FY23andFY24NPMGuidance-MonitoringAppendix_020823.pdf (last access: 14 November 2025), 2023.
- Vandaele, A. C., Hermans, C., Simon, P. C., Carleer, M., Colin, R., Fally, S., Mérienne, M. F., Jenouvrier, A., and Coquart, B.: Measurements of the NO₂ absorption cross-section from 42 000 cm⁻¹ to 10 000 cm⁻¹ (238–1000 nm) at 220 K and 294 K, *J. Quant. Spectrosc. Ra.*, 59, 171–184, [https://doi.org/10.1016/S0022-4073\(97\)00168-4](https://doi.org/10.1016/S0022-4073(97)00168-4), 1998.
- van Geffen, J., Eskes, H. J., Boersma, K. F., Veefkind, J. P., Sneep, M., and ter Linden, M.: TROPOMI ATBD of the total and tropospheric NO₂ data products, https://sentiwiki.copernicus.eu/_attachments/1673595/S5P-KNMI-L2-0005-RP-Sentinel-5P-TROPOMI-ATBD-NO2-data-products-2024-2.8.0.pdf (last access: 14 November 2025), 2024.
- van Geffen, J., Boersma, K. F., Eskes, H., Sneep, M., ter Linden, M., Zara, M., and Veefkind, J. P.: S5P TROPOMI NO₂ slant column retrieval: method, stability, uncertainties and comparisons with OMI, *Atmos. Meas. Tech.*, 13, 1315–1335, <https://doi.org/10.5194/amt-13-1315-2020>, 2020.
- van Geffen, J., Eskes, H., Compernelle, S., Pinardi, G., Verhoelst, T., Lambert, J.-C., Sneep, M., ter Linden, M., Ludewig, A., Boersma, K. F., and Veefkind, J. P.: Sentinel-5P TROPOMI NO₂ retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-based data, *Atmos. Meas. Tech.*, 15, 2037–2060, <https://doi.org/10.5194/amt-15-2037-2022>, 2022.
- Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens. Environ.*, 120, 70–83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.
- Zare, A., Romer, P. S., Nguyen, T., Keutsch, F. N., Skog, K., and Cohen, R. C.: A comprehensive organic nitrate chemistry: insights into the lifetime of atmospheric organic nitrates, *Atmos. Chem. Phys.*, 18, 15419–15436, <https://doi.org/10.5194/acp-18-15419-2018>, 2018.