Atmos. Chem. Phys., 25, 16915–16943, 2025 https://doi.org/10.5194/acp-25-16915-2025 © Author(s) 2025. This work is distributed under the Creative Commons Attribution 4.0 License.





Parameterization adaption needed to unlock the benefits of increased resolution for the ITCZ in ICON

Clarissa A. Kroll¹, Robert C. Jnglin Wills¹, Luis Kornblueh², Ulrike Niemeier², and Andrea Schneidereit³

¹Institute for Atmospheric and Climate Science, ETH Zürich, Zurich, Switzerland
²Max Planck Institute for Meteorology, Hamburg, Germany
³Deutscher Wetterdienst, Offenbach am Main, Germany

Correspondence: Clarissa A. Kroll (clarissa.kroll@env.ethz.ch)

Received: 17 March 2025 – Discussion started: 25 March 2025 Revised: 17 October 2025 – Accepted: 20 October 2025 – Published: 26 November 2025

Abstract. The double Inter-Tropical Convergence Zone (double-ITCZ) bias has been a persistent tropical precipitation bias over many climate model generations. This motivates investigations of whether increasing resolution and discarding parameterizations improves the representation of the large-scale atmospheric circulation and ITCZ. In this work, we study the double-ITCZ bias in an ICON XPP resolution hierarchy spanning from parameterized to explicitly described deep convection within a consistent framework. We demonstrate that the double ITCZ persists across horizontal resolutions from 160 to 5 km in specified sea-surface temperature simulations, independent of deep-convective and non-orographic gravity wave parameterizations. Changes in the treatment of near-surface wind speed within the turbulence parameterization can reduce the bias. However, we highlight that a key driver of the double-ITCZ bias seems to lie in the insufficient moisture transport from the subtropics to the inner tropics. The resulting dry bias in tropical near-surface moisture reduces deep convection over the Warm Pool, leading to a weakened Walker circulation. These biases ultimately culminate in the double-ITCZ feature. Increasing the near-surface wind speed limiter improves tropical near-surface moisture but exacerbates the bias in the moisture source, increasing the inner-tropical contribution at the expense of the subtropics. This degrades the representation of the global circulation, energy balance, and teleconnections. Additionally, we show that parameter adjustments at low resolution are informative of the response to the same parameter adjustments at high resolution. Our findings showcase the benefits of models supporting a range of resolutions and underline the importance of continuing the development of non-discardable parameterizations.

1 Introduction

The hydrological cycle is an important manifestation of the global atmospheric circulation. Despite its importance, biases in the representation of precipitation within the Inter-Tropical Convergence Zone (ITCZ) have been a persistent challenge throughout many model generations of the Coupled Model Intercomparison Project (CMIP) (Tian and Dong, 2020). Among them, the double-ITCZ bias is the most prominent problem (Mechoso et al., 1995; Lin, 2007). It describes positive precipitation biases south of the equator in the eastern Pacific and Atlantic, as well as underestimated precipitation in the equatorial Pacific. While observations in-

dicate that a double-ITCZ feature can occur naturally in these regions at certain times of the year, preferentially in March (Magnusdottir and Wang, 2008; Gonzalez et al., 2025), this feature is much too strong and persistent in model simulations. In addition to its influence on regional precipitation biases, the double ITCZ can influence large-scale climate phenomena such as the El Niño-Southern Oscillation (Ham and Kug, 2014; Zhang et al., 2014). Therefore, identifying the causes of the double ITCZ is a key step towards improving climate models.

In order to understand the simulated double-ITCZ bias, it is helpful to consider the role of the ITCZ at the center of

the large-scale atmospheric meridional circulation that imports moisture into the tropics and exports energy to higher latitudes (Holton, 2004; Schneider et al., 2014). Subtropical moisture is transported within the trade winds into the inner tropics, where its convergence sustains the ITCZ (Peixoto and Oort, 1992). The moist static energy content is greater in the poleward flow in the upper troposphere, such that there is a net energy transport into the subtropics by the Hadley cells (Fig. 1a). Since the position and structure of the ITCZ and the distribution of the net energy input to the atmosphere are interrelated (Bischoff and Schneider, 2014; Ren and Zhou, 2024), changes in the distribution of precipitation will be reflected in the atmospheric energy budget and vice versa. For example, the ITCZ resides in the Northern Hemisphere in the annual-mean due to the difference in the net energy input between the Northern and Southern Hemispheres (Philander et al., 1996; Frierson et al., 2013; Marshall et al., 2014). This underlines that the double-ITCZ problem cannot be investigated as an isolated tropical phenomenon: Sub- and extratropical biases in the energy budget can also be sources of the problem (Kang et al., 2008; Hwang and Frierson, 2013; Adam et al., 2018; Kang et al., 2019), and tropical biases can likewise cause biases in the sub- and extratropics (Henderson et al., 2017; Dong et al., 2022; Feng et al., 2023).

It is critical to recognize the coupling between the double-ITCZ problem and biases in the energy budget also in the context of model tuning, because in many cases the first step of model evaluation is the energy balance of the model (Wild, 2020). Most model evaluation workflows focus first on global mean top-of-the-atmosphere (TOA) fluxes and potentially surface energy fluxes (Mauritsen et al., 2012), and then proceed to tune additional atmospheric fields and processes, for example, as outlined in Hourdin et al. (2017). This is done with the knowledge that there can be trade-offs between these different optimizations, making dedicated approaches to optimize across multiple processes necessary, for example focusing on the precipitation distribution and TOA fluxes within one framework (Zhao et al., 2018a, b). However, in general, most published model evaluation workflows provide little information on regional energy budgets that are mechanistically important for the large-scale circulation and precipitation distribution. This poses a problem: Sometimes improvements in global mean energy fluxes introduce compensating errors in regional energy fluxes and lead to deterioration of the large-scale circulation and precipitation distribution.

Next to model tuning choices, an additional possible source of tropical precipitation biases is the incorrect representation of convective processes in coarse-scale models. Discarding potentially error-prone convective parameterizations (Arakawa, 2004; Jones and Randall, 2011; Sherwood et al., 2014; Hardiman et al., 2015) has therefore been one of the avenues explored to address the double-ITCZ problem (Song and Zhang, 2018; Zhou et al., 2022; Ma et al., 2023). In this context, it is important to note that even at a hori-

zontal resolution of 5 km, the necessity of deep convective parameterizations is still disputed, with some studies showing improvements in atmospheric representation with the elimination of parameterizations (Vergara-Temprado et al., 2020) and others showing deterioration and insufficiently resolved processes (Clark et al., 2024). Additionally, although it is sometimes stated that high-resolution simulations are less sensitive to parameterization choices than lowresolution simulations, a number of parameterizations still remain – such as the cloud microphysics, non-orographic gravity wave, turbulence, and in some cases shallow convective parameterizations – all of which are important for the model's atmospheric circulation and energy budget (Hu et al., 2021; Hájková and Šácha, 2023). Nevertheless, high computational costs, in combination with the hope that the most relevant processes are now resolved, can lead to a shortening of the model tuning process in high-resolution simulations. A lack of systematic tuning could be one of the reasons why high-resolution convection-permitting simulations without parameterizations do not show improvements in skill for the tropical precipitation pattern compared to conventional CMIP models, while high-resolution simulations that still rely on parameterization can outperform both (Zhou et al., 2022; Schneider et al., 2024).

In this work, we investigate the double-ITCZ bias in a resolution hierarchy of the ICON model spanning from parameterization-supported 160 km to deep-convection-permitting 5 km horizontal resolution. We discuss the influence of modeling parameter choices over the resolution hierarchy using an otherwise identical setup to allow for a robust comparison. This endeavor has only been recently enabled by the newly developed XPP configuration of the ICON climate model (Früh et al., 2022; Niemeier et al., 2023; Müller et al., 2025b), which can be run efficiently employing GPUs at a wide range of horizontal resolutions (Giorgetta et al., 2022; Lapillonne et al., 2025; Dipankar et al., 2025) and, in contrast to ICON sapphire (Hohenegger et al., 2023), supports the usage of convective parameterizations. We focus on the following questions:

- 1. Resolution dependence of the double ITCZ and largescale climate biases. Can increased horizontal resolution improve the double ITCZ and large-scale climate biases? Are there common biases across resolutions? Where can resolution-dependent improvements be found? (addressed in Sect. 3.1)
- 2. Underlying mechanisms and targeted parameter adjustments. What are the underlying mechanisms leading to the double ITCZ and associated large-scale climate biases in ICON XPP and how do the chosen adjustments ameliorate it? (addressed in Sects. 3.2 and 4, summarized in Schematic Fig. 1)
- 3. Resolution and parameterization-(in)dependence of bias corrections. To the extent that there are common




Figure 1. Conceptual depiction of the main zonal and meridional circulations exchanging moisture and energy between the subtropics and the tropics as (a) observed, (b) in default ICON simulations, and (c) in U_{\min} adjusted ICON simulations. Observations show that the moisture within the innermost tropics originates from evaporation in tropical and subtropical areas. The moisture is imported into the innermost tropics by the trade winds, with small losses to the free troposphere. In the innermost tropics it is then transported to the main convective regions, i.e., the Warm Pool (WP) within the zonally anomalous Walker circulation. Energy is exported to the subtropics within the outward branches of the Hadley circulation. In conventional ICON simulations (b), a large portion of the subtropical moisture is lost due to too much vertical transport of moisture from the boundary layer to the free troposphere. This leads to a moisture deficit in the tropics, specifically over the WP, reducing deep convection, slowing down the Walker circulation and leading to spurious additional convective centers over the Eastern Pacific as too little moisture is transported to the WP within the boundary layer. In the U_{\min} adjusted ICON simulations (c) increased evaporation leads to an additional moisture source in the tropics, which resolves the dry bias over the WP. However, the adjustments increase the positive free-tropospheric moisture bias between 0 to 30°, reduce the trade wind strength and change the balance between innermost-tropical and subtropical sources of tropical moisture content. Green colors indicate high moisture content. Brown colors low moisture content.

(double-ITCZ and climate) biases, how can they be addressed and can the same adjustments be applied at various resolutions relying on all and a reduced set of parameterizations? (addressed in Sect. 3.3)

For parameter adjustments, we focus on the choice of a minimum surface wind speed threshold (U_{\min}) in the vertical turbulence scheme of ICON XPP (see Sect. 2.1 for details). The minimum wind speed limiter U_{\min} has been identified as a promising tuning candidate to address the double-ITCZ precipitation biases in low-wind regimes such as the Warm Pool (Segura et al., 2025). Additionally, it may also be resolution dependent, with generally increasing near-surface wind speeds at increased horizontal resolution (Jeevanjee, 2017; Iles et al., 2020; Paccini et al., 2021; Morris et al.,

2024), making it particularly interesting for a resolution sensitivity study.

In the following, we will show that precipitation biases are reduced with increases in resolution (Sect. 3.1) and with physically motivated parameter adjustments (Sect. 3.2.1), both in terms of the large-scale precipitation field and a simplified ITCZ metric. Then we discuss the underlying physical mechanisms of this improvement, identifying potential tradeoffs between the improved precipitation representation and other key quantities. We focus in particular on the impacts on the Hadley and Walker circulation (Sect. 3.2.2), global moisture sources (Sect. 3.2.3), moisture transport (Sect. 3.2.4) and the sensitivity of key global metrics to further tuning (Sect. 3.2.5). We outline why our experiments hint that the

actual origin of the precipitation biases does not reside in the tropics but in the subtropics (Sect. 3.2.6). For this purpose, we compare the reference observations, the ICON simulation with a double ITCZ, and the ICON simulation with tuning adjustments. The identified large-scale differences in the three scenarios will be visualized building up on Fig. 1. In a last step, we test the transferability of the gained knowledge with respect to tuning choices to higher resolution (Sect. 3.3).

2 Methods

2.1 Model simulations

2.1.1 Model setup

We perform simulations with the atmosphere model of the Icosahedral Nonhydrostatic Weather and Climate Model for Numerical Weather Prediction, ICON-NWP (Zängl et al., 2015; Prill et al., 2023), in the eXtended Prediction and Projection, XPP-configuration - previously: Seamless configuration (Früh et al., 2022; Niemeier et al., 2023; Müller et al., 2025b). The setup of ICON XPP allows users to conduct weather and climate studies within one consistent framework (Müller et al., 2025a). ICON XPP uses parameterizations for radiation (Hogan and Bozzo, 2018), cloud microphysics (Seifert, 2008), vertical diffusion (Mauritsen et al., 2007), convection (Tiedtke, 1989; Bechtold et al., 2008), subgrid-scale orographic drag (Lott and Miller, 1997) and non-orographic gravity wave drag (Orr et al., 2010). The atmosphere is coupled to the land model JSBACH (Reick et al., 2021).

The experiments cover a hierarchy of four horizontal resolutions: 160 km (R2B4), 80 km (R2B5), 40 km (R2B6), and 5 km (R2B9). This includes the horizontal resolution range of conventional CMIP-type models (160 km) up to the resolution of deep-convection-permitting models (5 km). For all resolutions, 150 vertical levels up to a model top of 75 km are employed with a constant layer thickness of 300 m from a height of 8.4 to 19 km. The prescribed constant layer thickness ensures that the altitude range of deep convection is fully captured, even in the inner tropics (Schmidt et al., 2024). When going from the 40 to the 5 km setups only the strongly resolution dependent non-orographic and orographic gravity wave parameters were adjusted. Two configurations are run at 5 km. The first configuration relies on the full suite of parameterizations. In the second configuration, the parameterization for subgrid non-orographic gravity waves, midlevel and deep convection are disabled. We will refer to this setup as the rParam simulation. In contrast to Hohenegger et al. (2023), the orographic gravity wave drag and shallowconvective parameterization are kept at all resolutions, as underlying processes are not sufficiently resolved.

In all experiments, we use prescribed 6-hourly climatological sea ice and sea-surface temperature (SST) fields interpolated from the monthly climatological values of the CMIP6

Forcing Datasets (input4MIPs, 1978–2020) as boundary conditions (Durack and Taylor, 2018). Prescribed climatological SSTs reduce the impact of interannual variability, such as the influence of ENSO events on precipitation. In addition, they isolate the effect of model biases in atmospheric processes from those in the SST representation. The simulations are initialized with data from the European Centre for Medium-Range Weather Forecasts ERA5 reanalysis (Hersbach et al., 2020) for 1 January 2004.

2.1.2 Experiments

Default and Umin adapted ICON

 U_{\min} , the near-surface wind speed limiter, has been suggested as a potential lever for precipitation biases over the Warm Pool region in the ICON sapphire model setup (Segura et al., 2025). In our study, we investigate potential consequences of using this limiter for the global circulation and climate in a resolution hierarchy of the ICON XPP setup. The near-surface wind speed limiter U_{\min} acts within the bulk flux formulation in the turbulence parameterization. In this framework, evaporation E is tied to wind speed U at the lowest atmospheric level via the bulk flux formula for evaporation

$$E = \rho_{\mathbf{a}} C_E U(q_{\mathbf{s}} - q_{\mathbf{a}}),\tag{1}$$

where ρ_a is the atmospheric density, q_s the surface specific humidity, q_a the near-surface atmospheric specific humidity, and C_E the bulk transfer coefficient for latent heat, which is inversely proportional to the Richardson number. The bulk flux formulation shows that the latent heat flux is directly proportional to the near-surface wind speed. The fix suggested by Segura et al. (2025) does not change Udirectly, but adapts a hard-coded lower limit for the nearsurface wind speeds. In ICON XPP's turbulence parameterization, the surface wind U used for the computation of the Richardson number has a lower prescribed threshold U_{\min} with $U = MAX(U_{min}, U)$. This lower limiter is used to account for the influence of subgrid-scale turbulence with the goal of increasing the turbulent fluxes in low-wind regimes. Increasing the default value of U_{\min} from 1 m s^{-1} decreases the Richardson number in low-wind regimes, e.g., in the Warm Pool, and leads to increased evaporation. The influence of changes in U_{\min} at various resolutions, including simulations with the entire set of parameterizations, has not been tested before. It is important to note that changes in U_{\min} can similarly impact turbulent fluxes of sensible heat and momentum in addition to the targeted latent heat flux. Potential consequences of the resulting changes in the momentum budget were not considered in Segura et al. (2025). Concretely, this means that the suggested increase in U_{\min} will increase the drag on near-surface winds. It is our aim to investigate whether there are associated negative influences on the circulation next to the positive effect for precipitation over the Warm Pool.

Description of experiment configurations

The simulations presented in this work are summarized in Table 1. We run five control simulations (CTL) in four different horizontal resolutions: 160, 80, 40 and 5 km. The simulations are run for six years for all coarser resolutions (160 to 40 km). Due to the very high computation demand, the two 5 km resolution setups - with full parameterization set and reduced parameterization set – are run for three years each instead of six years. Branching off at the fourth year of the 40 km simulations, two-year-long sensitivity (perturbation) studies are performed (hereafter labeled with PTB). In the PTB experiments, we vary the minimum surface wind speed limiter, U_{\min} , which is presented to the turbulence scheme (Table 1). The two U_{\min} settings that perform the best in representing the large-scale annual-mean precipitation, PTB-5 and PTB-6, with $U_{\min} = 5$ and $6 \,\mathrm{m \, s^{-1}}$, are chosen for further optimization. First, the wind speed is adapted for land and ocean separately. In PTB-5_1 and PTB-6_1, $U_{\text{min}} = 1 \text{ m s}^{-1}$ over land to account for the slower near-surface wind speeds. Second, in PTB-5_1t, PTB-6_1t, the model is retuned to reestablish a similar top-of-the-atmosphere (TOA) imbalance to CTL. Only the best performing U_{\min} setting, PTB-5_1 and the corresponding tuned settings PTB-5_1t, are tested in 3-month branch-off runs of the CTL-5 km and CTL-5 km-rParam simulation. For this purpose, MAM is chosen because it is the season with the largest double-ITCZ biases. To allow the model time to equilibrate with the changed settings, a onemonth spin-up is used. As the investigated horizontal resolutions differ by up to a factor of 32, all data sets are remapped for a fair comparison in the subsequent analysis. For this purpose, a 1° horizontal resolution is chosen. This resolution is close to the tropical 1.4° horizontal resolution of the 160 km simulation.

2.2 Reanalysis and observational data sets

The model output is evaluated against reanalysis and observational data sets. For the radiative fluxes, data from the "Clouds and the Earth's Radiant Energy System" (CERES; Doelling et al., 2013, 2016), an observational dataset for TOA energy fluxes, are used. The precipitation distribution is compared to the observational data sets "Integrated Multi-satellitE Retrievals for GPM" (IMERG; Huffman et al. (2010)) and "Global Precipitation Climatology Project" (GPCP; Huffman, 2021). In addition to these observational data sets, we show radiative fluxes from ERA5, (C3S, 2018; Simmons et al., 2020), noting that these values are not assimilated (Dee et al., 2011; Hersbach et al., 2017). This means that the reanalysis product might behave more like a conventional model for these variables and should only be used to complement the observational data. For latent heat flux, near-surface wind speeds, and near-surface specific humidity q_s , we refer to ERA5 as a primary reference, as it assimilates station-based humidity measurements (Simpson et al., 2024; Hersbach et al., 2020) and offers a homogeneous data set with values both over land and ocean. Whereas latent heat flux and near-surface wind speeds are direct outputs of ERA5, the values of q_s are computed (details in Appendix A). The ERA5 latent heat flux values are thought to have a positive bias (Martens et al., 2020; Song et al., 2021). Therefore, we also compare our data to the OAFlux dataset (Schneider et al., 2013; National Center for Atmospheric Research Staff, 2022), which integrates satellite retrievals and three atmospheric reanalysis. When comparing model data to observational data or ERA5 statistical significance is tested with a two-sided z-test at $\alpha = 0.1$. For the humidity and temperature fields, we correct for autocorrelation via adjustment of the net sample size based on lag-1 autocorrelation values of years 2004-2010 from ERA5 and the 40 km simulations (cf. Trenberth, 1984).

2.3 ITCZ evaluation metrics

To evaluate the large-scale ITCZ features, we use an energetically motivated framework, where precipitation biases are separated into hemispherically symmetric (Adam et al., 2016) and antisymmetric components (Hwang and Frierson, 2013). The symmetric component

$$E_p = \frac{\overline{P}_{2^{\circ} S - 2^{\circ} N}}{\overline{P}_{20^{\circ} S - 20^{\circ} N}} - 1 \tag{2}$$

is related to the net energy input near the equator. $E_p = 0$ would describe a state in which the precipitation average \overline{P} between 2° S and 2° N is equal to the precipitation average between 20° S and 20° N. The anti-symmetric component

$$A_{p} = \frac{\overline{P}_{0^{\circ} - 20^{\circ} N} - \overline{P}_{0^{\circ} - 20^{\circ} S}}{\overline{P}_{20^{\circ} S - 20^{\circ} N}},$$
(3)

is related to the cross-equatorial energy flux, which is influenced by asymmetries in the net energy input in the high latitudes of the two hemispheres. In both equations \overline{P} is the average precipitation in the region constrained by the two latitude boundaries referred to in the respective subscript.

2.4 Residence time in boundary layer based on moisture budget analysis

In order to assess biases in moisture transport, we evaluate the residence time of moisture within the boundary layer. Our analysis is based on a time-mean moisture budget (Peixóto and Oort, 1983; Peixoto and Oort, 1992) linking moisture flux convergence with precipitation and evaporation fluxes as

$$\overline{P} - \overline{E} = -\frac{1}{g} \int_{0}^{p_s} \nabla \cdot \overline{\beta_{\rho} u q} \, dp. \tag{4}$$

Table 1. Summary of Experiments: Experiment acronyms, employed resolutions, simulated timeframe, setting for the minimum horizontal wind speed over ocean/land and tuning. The control simulation (CTL) run for all resolutions and the best-performing perturbation experiment (PTB) for the large-scale precipitation representation are marked bold. The runs with the reduced set of parameterizations are labeled as rParam. Resolution is given in terms of the ICON grid used and the corresponding horizontal grid spacing.

| Experiment | Resolution | Simulated Timeframe | Surface Wind Speed Limiter $U_{\rm min}$ (Ocean/Land)/m s ⁻¹ | Tuned |
|-----------------|----------------------------|------------------------|---|--------------|
| PTB-0.5 | R2B6, 40 km | 2 years | 0.5/0.5 | |
| CTL | R2B{4,5,6}, 160, 80, 40 km | 6 years | 1.0/1.0 | \checkmark |
| | R2B9, 5 km | 3 years | 1.0/1.0 | \checkmark |
| CTL-rParam | R2B9, 5 km | 3 years | 1.0/1.0 | \checkmark |
| PTB-1.5 | R2B6, 40 km | 2 years | 1.5/1.5 | |
| PTB-2 | R2B6, 40 km | 2 years | 2.0/2.0 | |
| PTB-4 | R2B6, 40 km | 2 years | 4.0/4.0 | |
| PTB-5 | R2B6, 40 km | 2 years | 5.0/5.0 | |
| PTB-5_1 | R2B6, 40 km | 2 years | 5.0/1.0 | |
| PTB-5_1 | R2B9, 5 km | 2 MAM | 5.0/1.0 | |
| PTB-5_1-rParam | R2B9, 5 km | 2 MAM | 5.0/1.0 | |
| PTB-5_1t | R2B6, 40 km | 2 years | 5.0/1.0 | \checkmark |
| | R2B9, 5 km | 2 MAM | 5.0/1.0 | \checkmark |
| PTB-5_1t-rParam | R2B9, 5 km | 2 MAM | 5.0/1.0 | \checkmark |
| PTB-6 | R2B6, 40 km | 2 years | 6.0/6.0 | |
| PTB-6_1 | R2B6, 40 km | 2 years | 6.0/1.0 | |
| PTB-6_1t | R2B6, 40 km | 2 years | 6.0 / 1.0 | \checkmark |

In this equation $\overline{(\cdot)}$ denotes the annual-mean climatology, \boldsymbol{u} the horizontal wind vector, q the specific humidity, p is the pressure and g the gravitational constant. β_{ρ} is a Heaviside function indicating whether values are above or below the surface (cf. Boer, 1982). Based on Eq. (4), we then estimate the residence time $\tau_{R_{bl}}$ within the boundary layer by splitting the integral into a contribution from the boundary layer up to 850 hPa and the rest of atmosphere and rearranging:

$$\frac{1}{\tau_{R_{bl}}} = \left| \frac{-\frac{1}{g} \int_{850 \text{ hPa}}^{p_s} \nabla \cdot \overline{\beta_{\rho} u q} \, dp}{\frac{1}{g} \int_{850 \text{ hPa}}^{p_s} \overline{q} \, dp} \right|
= \left| \frac{\overline{P} - \overline{E} + \frac{1}{g} \int_{850 \text{ hPa}}^{850 \text{ hPa}} \nabla \cdot \overline{\beta_{\rho} u q} \, dp}{\frac{1}{g} \int_{850 \text{ hPa}}^{p_s} \overline{q} \, dp} \right|.$$
(5)

Dividing by the integrated moisture within the boundary layer helps control for the dependence of boundary-layer moisture flux on boundary-layer moisture.

2.5 Circulation analysis

We calculate the velocity potential χ at 200 hPa to assess the Walker and Hadley circulation strength following the approach by Tanaka et al. (2004) and Lu et al. (2007) (see Appendix B for derivation details). χ_{max} , the maximum of the zonal-mean velocity potential at 200 hPa within the 30° latitude band is used as a proxy for the Hadley circulation strength. As χ can only be determined with a constant offset due to the inverse Laplacian, we choose this constant so

that the global mean of χ is zero for consistency between the individual experiments. The Walker circulation is the zonally anomalous part of the circulation. Here, we employ χ^*_{max} , the maximum deviation of the annual mean χ in the 10° latitude band from its zonal mean, as a Walker circulation metric.

3 Results

3.1 Consistent large-scale precipitation biases across horizontal resolutions

To investigate the influence of horizontal resolution on the double-ITCZ bias and associated large-scale climate biases, we focus on CTL experiments with horizontal grid resolutions of 5 to 160 km employing all parameterizations. The two-year mean precipitation shows qualitatively similar biases compared to IMERG in all four resolutions of ICON (Fig. 2). Quantitatively, there is a consistent improvement with increased resolution in the global root mean square error (RMSE) with respect to IMERG from 1.55 mm d^{-1} (160 km) to 1.27 mm d⁻¹ (5 km). The large-scale patterns of precipitation bias persist with increasing resolution, but decrease in amplitude. Especially apparent in the tropics is the double-ITCZ feature over the Pacific and Atlantic, as well as the dry bias over the Maritime Continent. Areas of consistent improvement include coastal regions, for example, in the ITCZ on either side of South America or surrounding the Maritime Continent.




Figure 2. (a) Climatological IMERG precipitation distribution from 2004 to 2010 and (**b–e**) two-year mean precipitation bias with respect to IMERG for 160 km (**b**), 80 km (**c**), 40 km (**d**) and 5 km (**e**) setups. Statistically insignificant differences between IMERG and the experiments based on a two sided z-test at $\alpha = 0.1$ are grayed out in (**b**)–(**e**). The corresponding global RMSE in precipitation is stated beneath the panel for each resolution experiment. The global RMSE and mean error in near-surface specific humidity, calculated with respect to values derived from ERA5 reanalysis, is depicted in the inlays.

The tropical dry bias over the Warm Pool is tied to a lack of near-surface specific humidity throughout the entire tropics (average global values in inlays of Fig. 2; spatial distribution in Fig. C1). The lack of near-surface specific humidity affects the build-up of larger deep convective systems in the inner tropics and with it the vertical moisture distribution (Fig. 3). Consistent for all resolutions, the regions between 20° S and 20° N are too dry below 1 km in altitude, although a clear improvement with increasing horizontal resolution is apparent. The lack of near-surface moisture helps to explain

the lack of precipitation in convective areas close to the equator. In contrast to the inner tropics, the areas between 15 to 40° latitude exhibit a moist bias above 1 km in height, peaking at around 20° N and 20° S at a height of 2 km. The vertical structure of humidity biases indicates too strong vertical moisture flux out of the boundary layer. The drying of the boundary layer reduces the amount of moisture transported into the inner tropics by the trade winds (Fig. 1b). We will demonstrate this mechanism in more detail in Sect. 3.2.




Figure 3. (a–d) Two-year atmospheric specific humidity bias with respect to ERA5 (2004–2010) for 160 km (a), 80 km (b), 40 km (c) and 5 km setups (d). Statistically significant differences between ERA5 and the model data are shown, and insignificant regions are dashed out.

The consequences of the biases in the specific humidity distribution are also apparent in the zonal-mean temperature biases (Fig. 4, cf. Figs. 3 and C1). Across the resolutions, a near-surface warm bias and a cold bias in the tropical free troposphere persist. The near-surface warm bias could result from the dry bias in the near-surface layer. If the energy input into the lower atmosphere is similar in the CTL and PTB simulations, the same amount of energy would produce a warmer atmosphere under drier conditions. The colder free troposphere is consistent with Figs. 2 and 3: The lack of boundary layer moisture reduces deep convection over the Warm Pool. The missing latent heating in the free troposphere leads to a cooler atmosphere. As temperatures in the tropics are quite homogeneous, as explained by the Weak Temperature Gradient theory (Charney, 1963; Held and Hoskins, 1985; Bretherton and Smolarkiewicz, 1989), this bias propagates through the tropical region. The warm air band at 20–40° N coincides with the region influenced by the local topography of the Himalayas, where the too warm surface air is advected upwards (Pan et al., 2016).

Summarizing, we found that increased resolution alone does not resolve the double-ITCZ bias up to a resolution of 5 km. Although improvements can be found in the precipitation pattern, similar large-scale biases persist. From the biases, we gain a first understanding of the underlying problem: It appears that the negative bias in near-surface specific humidity is one of the causes of the precipitation bias. A negative bias in near-surface moisture inhibits deep convection in the Warm Pool, weakening the Walker circulation and allowing convection to develop the East Pacific ITCZ region. A




Figure 4. (a–d) Two-year atmospheric temperature bias with respect to ERA5 (2004 to 2010) for 160 km (a), 80 km (b), 40 km (c) and 5 km (d) setups. Statistically significant differences between ERA5 and the model data are shown, insignificant regions are dashed out.

too-weak Walker circulation reduces moisture transport from the East Pacific to the West Pacific and further weakens the Walker circulation. Furthermore, too-weak deep convection leads to a cold bias in the free troposphere, which reduces the temperature inversion also in distant regions, potentially leading to an additional positive feedback on the circulation by reducing the cloud cover in the subsidence regions. These resolution-independent ICON CTL simulation biases are conceptually summarized in Fig. 1b.

3.2 Sensitivity of large-scale precipitation biases to the surface wind threshold in the turbulence scheme

In this section, we investigate the efficacy of a parameter adjustment in addressing the double-ITCZ bias across resolutions in ICON. We test the U_{\min} parameter adjustment (Sect. 2.1, cf. Sect. 3.2.1) to enhance evaporation and resolve the near-surface dry bias over the Warm Pool region, with the goal of interrupting the circle of biases outlined in the previous paragraph (cf. Fig. 1). In this way, we use the impact of U_{\min} changes on the global climate system as a lens into the underlying mechanisms of the double-ITCZ bias. Specifically, we will analyze the impact on the global circulation in Sect. 3.2.2, explaining why the tropical circulation shows an strengthening despite increased surface drag coefficient. Section 3.2.3 explores how U_{\min} leads to an unintended shift in the origin of inner tropical moisture. In Sect. 3.2.4, we will demonstrate that ICON exhibits an exaggerated moisture transport out of the boundary layer, which can explain the deficit in near-surface moisture in the inner tropics. We discuss why this exaggerated vertical moisture transport necessitates high-biased evaporation (i.e. through increased $U_{\rm min}$) to maintain a realistic global circulation. In Sect. 3.2.5, we discuss the global impacts of the $U_{\rm min}$ changes and also how they change with retuning. Section 3.2.6 summarizes the impacts of $U_{\rm min}$ changes on the global dynamics and underlying mechanisms.

3.2.1 The choice of surface wind threshold

We use the 40 km resolution configuration to test the effects of varying the $U_{\rm min}$ threshold in the turbulence scheme. Starting from the default 1 m s⁻¹ setting in the CTL simulation, we change $U_{\rm min}$ to the values of {0.5, 1.5, 2, 4, 5, 6} m s⁻¹. The resulting two-year mean precipitation biases against IMERG are shown in the first two rows of Fig. 5. Indeed, RMSEs in both precipitation and near-surface specific humidity decrease with an increase in $U_{\rm min}$ up to 5 m s⁻¹ (PTB-5). Going beyond this value increases the RMSE in precipitation again and a positive precipitation bias develops over the Warm Pool in PTB-6.

Despite the decrease in the large-scale precipitation bias, precipitation biases over land remain or even worsen. Most apparent is the persistent dry bias over the islands in the Warm Pool region, but also an increase of the bias over South America and Africa, with increasing U_{\min} (Fig. 5a– d). Observational data show an average wind speed of below $2.5 \,\mathrm{m \, s^{-1}}$ at the surface over land, whereas the lower surface roughness over the ocean allows higher wind speeds with values of 6–7 m s⁻¹ (Fig. 6). High values of U_{\min} would therefore cut off more of the velocity distribution over land, leading to exaggerated evaporation over land and potentially drying of the soil. In the next experiments, we therefore distinguish between ocean and land when setting U_{\min} , going back to the CTL setting of 1 m s⁻¹ for land in PTB-5 1 and PTB-6 1. This change reverses the increasing dry bias over South America and improves the RMSE precipitation score even further (Fig. 5e, f).

3.2.2 The impact of the surface wind threshold: Hadley and Walker circulation

In a next step, we address concerns that increased U_{\min} might adversely affect the velocity distribution by increasing the surface drag coefficient on the near-surface velocities and correspondingly slowing down the circulation. A comparison of the probability density function of monthly mean surface wind speeds in CTL, PTB-5 and PTB-5_1 with ERA5 in Fig. 6 summarizes how the changes in U_{\min} feed back on the wind distribution in the tropics. Compared to ERA5, CTL exhibits a skew to smaller surface wind speeds, both over ocean and land. Against expectations based on increased surface drag coefficient, the skew to smaller near-surface wind speeds over ocean is reduced instead of exacerbated via adjustment of U_{\min} over the ocean in PTB-5 and PTB-5_1 (Fig. 6b). For the values over land (Fig. 6c), which are mainly

in the velocity bins from 0 to $2.5\,\mathrm{m\,s^{-1}}$, the main velocity peak remains essentially unchanged in location across all ICON simulations, although the probability of strong winds is reduced in PTB-5 relative to CTL.

An inspection of the spatially resolved near-surface wind fields and velocity potential of CTL, PTB-5 and PTB-5_1 compared to ERA5 in Fig. 7 gives more nuanced insights into the impact of U_{\min} changes on the atmospheric circulation. The choice of globally high U_{\min} in PTB-5 (Figs. 6, 7e) leads to the expected overall reduction in wind speeds on land. Over land, this reduction in overall wind speed compared to CTL is reverted when separate U_{\min} values are chosen for land and ocean in PTB-5_1 (Fig. 7g). Over ocean, a general slowdown of the Trades in the Southern Hemisphere, e.g., off the West Coast of South America, can be noted in all PTB experiments (Fig. 7e, g). This not only impacts moisture transport within the Trades ranging from 30° to the inner tropics, but may also interfere with the wind-evaporation-SST feedback and related teleconnections. The inner tropics are an exception to the general slow-down of the circulation, which would be expected considering the increase in surface drag coefficient.

Indeed, changes in the velocity potential at 200 hPa confirm changes in the Walker circulation (Fig. 7b, d, f, h and Table 2). In the ERA5 reanalysis (Fig. 7b), the velocity potential field shows a peak in divergence over the Warm Pool and convergence near the coast of South America. For the CTL configuration, the Walker circulation strength has only 80% of its strength in ERA5 (Fig. 7c, Table 2). This is in line with the reduced divergence both over the Warm Pool and South America, the locations of rising branches of the Walker circulation. As already suspected from the nearsurface wind plots, the U_{\min} changes improve the Walker circulation strength, such that it exhibits a similar strength as in ERA5. This is also apparent when comparing Fig. 7d and f, h, which show that the U_{\min} adjustment reduces the positive bias over the maritime continent. The reduced divergence over the Warm Pool area in CTL is almost entirely counterbalanced at the cost of an increased bias over South America, which shows a decreased divergence. In PTB5_1, the increased convergence over South America decreases again compared to PTB5, further reducing the velocity potential bias (Fig. 7f, h).

The Hadley circulation is generally too weak in all experiments (Table 2). In CTL, the Hadley circulation strength has 70 % of its strength in ERA5. In PTB5 and PTB5_1 it drops to 60 %. The differing response of the circulation in the innermost tropics and subtropical latitudes can be explained by the increased role of moisture in the inner tropics. More available near-surface moisture leads to increased deep convection, strengthening the Walker circulation and outweighing the increased drag on the near-surface winds. In the higher latitudes this effect is less important, leading to the slow-down of the Trades and the Hadley circulation.




Figure 5. Two-year mean large-scale precipitation bias with respect to IMERG averaged over 2004–2010 for different settings of the surface wind at 40 km resolution: PTB-0.5 (a), PTB-4 (b), PTB-5 (c), PTB-5_1 (e), PTB-5_1t (g), PTB-6 (d), PTB-6_1 (f) and PTB-6_1t (h). The corresponding global precipitation RMSE is stated beneath the panel for each sensitivity experiment. Statistically insignificant differences between IMERG and the experiments based on a two-sided z-test at $\alpha = 0.1$ are grayed out. The global RMSE and mean error in near-surface specific humidity, calculated with respect to values derived from ERA5 reanalysis, is depicted in the inlays.




Figure 6. Normalized probability density functions of the monthly mean near-surface wind speed distributions between 10° S to 10° N in the experiments CTL, PTB-5 and PTB-5_1. A comparison with ERA5 data is given for (a) the entire region, (b) over ocean only, and (c) over land only.

Table 2. Walker and Hadley circulation strengths for ERA5 as well as CTL, PTB-5 and PTB-5_1. Indices are calculated following the approach by Tanaka et al. (2004) as described in Sect. 2.5.

| Experiment | Walker circulation Strength $\chi*_{max}/10^7$ m ² s ⁻¹ | Hadley circulation Strength $\chi_{max}/10^6 \text{ m}^2 \text{ s}^{-1}$ |
|------------|---|--|
| ERA5 | 1.16 | 3.23 |
| CTL | 0.94 | 2.17 |
| PTB-5 | 1.27 | 1.98 |
| PTB-5_1 | 1.22 | 1.98 |

3.2.3 The impact of the surface wind threshold: Moisture sources

The changes in the spatial wind speed patterns (Fig. 7) have demonstrated that U_{\min} has the potential to substantially impact the general atmospheric circulation. It increases the Walker circulation strength, but leads to a reduction of the Hadley circulation strength, especially noticeable in the region of the Southern Hemisphere trade winds. The trade winds transport a substantial amount of moisture into the inner tropics. Therefore, the change in U_{\min} comes at the risk of changing the weighting of the inner-tropical versus the subtropical moisture sources. The analysis of the latent heat flux biases of CTL and PTB-5_1 with respect to ERA5 and OAFlux can give some indication of potential changes in the moisture sources (Fig. 8). When interpreting the biases, it is important to keep in mind that the latent heat flux in ERA5 likely has a positive bias (i.e. too strong evaporation, cf. Martens et al., 2020; Song et al., 2021) and is larger than in OAFlux (Fig. 8a, b). The biases against ERA5 will therefore tend to have a bias towards more negative values, although OAFlux is subject to its own observational biases such that there is no true ground truth. For consistency with the other plots, we primarily use ERA5 as reference, but we also show OAFlux for comparison.

CTL shows too little latent heat flux in the region of the ITCZ in the Pacific as well as in the Indian Ocean with

respect to ERA5 (Fig. 8c), whereas values compared to OAFlux almost perfectly capture the zonal-mean climatology. With the increase in U_{\min} (Fig. 8e–f), the low bias in the Warm Pool region is reverted to a positive latent heat flux bias in the tropical region, even compared to the already high values simulated by ERA5. Although the spatial contrast in evaporation bias reduces, negative evaporation biases emerge in the trade wind regions at the coasts of Central- and South-America, key source regions for moisture transported into the tropical Pacific. These reductions are obscured in the zonal mean, because subtropical regions with climatologically lower wind speeds, such as the region South-East of Australia, develop positive evaporation biases. All in all, these findings indicate a reduction of the subtropical moisture contribution to moisture in the Pacific inner tropics.

3.2.4 The impact of the surface wind threshold: Moisture transport

Next we investigate the potential role of vertical moisture transport out of the boundary layer at latitudes between 20 and 40° in reducing moisture transport by the trade winds into the inner tropics. Exaggerated vertical moisture transport out of the boundary layer would reduce trade-wind moisture transport and could lead to the negative near-surface humidity bias in the inner tropics (cf. Fig. 3). If this is the case, then strong evaporation in the innermost tropics is necessary




Figure 7. (a) ERA5 multi-year average near surface wind speeds for the years 2004–2010. Arrows visualizing the near-surface wind speed for ERA5 are superimposed to visualize the general circulation. (c, e, g) Two-year mean large-scale near-surface wind speed bias with respect to ERA5 for CTL and different settings of the near-surface wind speed limiter, U_{\min} , in 40 km resolution: PTB-5, PTB-5_1. (b) ERA5 multi-year average velocity potential χ at 200 hPa for the years 2004-2010. (d, f, h) Two-year mean bias of the CTL, PTB-5 and PTB-5_1 experiment with respect to the ERA5 velocity potential. Negative values of the velocity potential are found in regions of ascent and divergent motion; positive values in region of subsidence and convergence.




Figure 8. Latent heat flux in reanalysis and 40 km simulations. (a) ERA5 multi-year average latent heat flux for the years 2004–2010. (b) OAFlux multi-year average latent heat flux for the years 2004–2010. Two-year mean large-scale latent heat flux change with respect to ERA for CTL and PTB-5_1 (c-f). The left column shows global biases, the right column the zonal mean bias with respect to ERA5 (land and ocean) and OAFlux (ocean only).

to compensate for the lack of moisture transported into the tropics. However, this would also mean that the wind speed limiter fix addresses a downstream consequence and not the primary error in moisture transport.

Figure 9 quantifies this effect with the inverse residence time $\tau_{R_{bl}}$ of specific humidity within the boundary layer up to 850 hPa (Sect. 2.4). On a large scale, the $\tau_{R_{bl}}$ values simulated in CTL and PTB-5_1 are shorter compared to ERA5 (Fig. 9d–e), demonstrating a too fast loss of boundary layer moisture to the free troposphere. This transport bias is well aligned with positive specific humidity biases in the free troposphere in the mid latitudes (cf. Fig. 3). The fast loss of boundary layer moisture is most pronounced in the higher

latitudes around the poles, but tropical and midlatitude land as well as the trade wind region are also affected. In CTL, the weaker circulation manifests itself in a pronounced bias to longer residence times in the Northern ITCZ band over the Eastern Pacific, which is indicative of too weak moisture-flux divergence. With the U_{\min} adaptation, this bias is corrected at the expense of an overall shortening of residence times in the trade wind region. A quantitative analysis of $\tau_{R_{bl}}$ over the ocean only demonstrates that the probability density function of $\tau_{R_{bl}}$ is biased to shorter residence times without apparent improvements with the U_{\min} patch (Fig. 9c). The corresponding bias in the zonal distribution of $\tau_{R_{bl}}$ over the ocean compared to ERA5 confirms that shorter residence times dom-

inate over all latitudes in the ICON experiments (Fig. 9b). The U_{\min} adjustment does not correct the short $\tau_{R_{bl}}$ values in the tropics, where values were already too short. In the region at 10° N, where CTL exhibited long residence times, the adjustment overcompensates for the bias. In the subtropics the bias is slightly improved, but residence times remain too short. Together, these results illustrate that moisture is exported too quickly out of the boundary layer to the free troposphere in the ICON simulations, which leaves less moisture to be transported into the inner tropics. The change in U_{\min} does not directly correct for this problem, but combats the consequences by artificially increasing the moisture supply to the boundary layer, specifically in low-wind regimes in the tropics. A more physically motivated correction would need to increase the transport of subtropical moisture into the tropics rather than the subtropical free troposphere. The consequent increase in inner-tropical moisture source would likewise lead to a increase in the Walker circulation. This would naturally increase the near-surface wind speeds and compensate for the missing evaporation in the tropics.

3.2.5 The impact of the surface wind threshold: Global variables

As we have seen far-reaching impacts on the general atmospheric circulation when U_{\min} is increased, we investigate the impact of the altered U_{\min} on key global parameters in Fig. 10. The first parameter pair is the asymmetry index A_p and symmetry index E_p of the precipitation distribution. All sensitivity experiments show a positive A_p , consistent with the observational references IMERG and GPCP, indicating that the inter-hemispheric energy imbalance and precipitation contrast is captured correctly. E_p shows negative values, starting to increase only at a U_{\min} of $4 \,\mathrm{m \, s^{-1}}$. The negative E_p for U_{min} limiters smaller than 4 m s⁻¹ reflect the large dry bias over the Warm Pool. They indicates that the net energy input (NEI) to the atmosphere at the equator is most likely low biased. Since U_{\min} preferentially effects evaporation in warm regions where evaporation is climatologically high and in regions with low wind speeds, an increase in near-surface wind speed will have a more substantial effect in the innertropics, which would positively reflect in the E_p index, as is evident in the simulation scores.

The simulation scores confirm our earlier findings that the reduced dry bias in near-surface humidity with increasing U_{\min} comes at the cost of an increase in magnitude of the global latent heat flux bias compared to ERA5 (cf. Figs. 8 and C2 for spatial maps). Globally, there is already enough moisture supplied to the atmosphere in CTL, but there are biases in the moisture distribution associated with the global circulation.

The improvements in the ITCZ bias come at the cost of strong biases in the global annual-mean net top-of-the-atmosphere (TOA) imbalance which drops by $5 \,\mathrm{W} \,\mathrm{m}^{-2}$ for a $U_{\rm min}$ of $6 \,\mathrm{m} \,\mathrm{s}^{-1}$ (Fig. 10). The main contribution to the

changes in TOA imbalance arise from upward shortwave radiation. Due to the specified-SST setup, the increased bias in TOA imbalance can only have small effects on the surface temperature $T_{\rm s}$. Despite the negative TOA, $T_{\rm s}$ increases by around 1 K. This can be explained by the increase in free tropospheric moist static energy, as both specific humidity values and atmospheric temperature increase while the lapse rate over land does not change substantially (Zhang and Boos, 2023). The increase in $T_{\rm s}$ is reflected in the increase in outgoing long wave radiation. The tuning simulations 5-1t and 6-1t demonstrate that it is possible to readjust the radiative fluxes to expected values without strong deterioration of the improved precipitation distribution.

3.2.6 The impact of the surface wind threshold: Consequences on global dynamics

Within this section, we have answered the second research question group: The precipitation biases in coarse-scale 40 km simulations can be addressed by increasing evaporation via an increase in U_{\min} in the turbulence scheme. Mechanistically, the increase in U_{\min} should lead to an increase in the drag on the surface winds. However, in the inner tropics, the increase in U_{\min} also counterbalances the dry bias in the Warm Pool region through increased evaporation. This fuels deep convection and leads to an acceleration of the Walker circulation transporting moisture over the Eastern and Central Pacific into the main convective region. The increase in the Walker circulation strength counterbalances the increases in drag on the near-surface wind speeds in the inner tropics, leading to an increase in near-surface wind speed over ocean. In the subtropics, which do not benefit from the increase in Walker circulation strength, the increased drag on the surface winds leads to a reduction in the trade wind strength. U_{\min} compensates for three problems in the moisture transport:

- 1. for too strong vertical transport out of the boundary layer (cf. Fig. 9).
- 2. for the consequently too weak meridional equatorward near-surface specific humidity transport within the Trades from the subtropics, the main source region of inner tropical moisture (cf. Fig. 7, Table 2).
- 3. for the resulting dry bias in the tropics and too weak moisture transport by tropical circulations (cf. Fig. 3, Table 2).

However, by drawing the moisture primarily from the inner tropics without fixing the problem in the vertical transport in the higher latitudes, it acts more like a "patch" than a real fix. The improved representation of the ITCZ comes at the cost of a replacement of higher latitude (sub)tropical moisture sources by an increase in the inner-tropical moisture source, a reduction in the trade wind strength, and an increased global-mean latent heat flux bias (Fig. 1c).




Figure 9. Inverse residence time $\tau_{R_{bl}}$ within the boundary layer up to 850 hPa: (a) ERA5 for the reference years 2004–2010, and differences of two-year average of CTL (d) and PTB-5_1 (e) compared to ERA5. The gray lines mark the 40 and 20° latitudes. (b) Difference of zonal distribution of $\tau_{R_{bl}}$ of CTL and PTB-5_1 to ERA5 within the latitude band between 40° S and 40° N, restricted to ocean only. (c) Probability density function of $\tau_{R_{bl}}$ of ERA5, CTL and PTB-5_1 within the latitude band between 40° S and 40° N, restricted to ocean only.

3.3 Transferability to higher resolutions and reduced parameterization configurations

After understanding the sensitivity to U_{\min} changes in the coarser 40 km setup, we proceed to investigate whether the corresponding changes with U_{\min} also translate to higher 5 km resolution and additionally whether discarding parameterizations improves the representation of the mean atmospheric state. Due to computational constraints, we perform seasonal rather than year-long sensitivity tests. We focus on two MAM seasons, as this is the key season for the double-ITCZ bias (e.g., Si et al., 2021; Adam et al., 2018). A comparison of the precipitation bias for the CTL and PTB5-1 simulations in 40, 5 and 5 km-rParam is shown in Fig. 11. The 5 km setup with the full suite of parameterizations exhibits a less pronounced double-ITCZ feature in the East Pacific than rParam, its counterpart without gravity wave and deep convective parameterization (Fig. 11b, e; cf. Fig. 11c, f). Additionally, while the 5 km-Param setup reduces the globalmean near-surface specific humidity bias compared to the 40 km setup, the 5 km-rParam setup increases this bias. However, especially in the MAM average, it becomes apparent that the precipitation dry bias over the Warm Pool is stronger in the parameterized simulations (Fig. 11e; cf. Fig. 11f), with an RMSE comparable to the 40 km setup. This is also true for the dry bias over the islands in the Maritime Continent, which disappears in rParam but is still apparent in the 5 km setup with the full set of parameterizations.

In both 5 km configurations, the seasonal-average double-ITCZ bias is reduced by $U_{\rm min}$ changes, and their RMSE in seasonal-average precipitation is less than in the comparable 40 km simulations. The positive precipitation bias appearing in the Indian Ocean with the $U_{\rm min}$ adjustments is not as prominent at higher resolution (Fig. 11h, i; cf. Fig. 11g). In contrast to the 40 km setup, both 5 km setups also show strong improvements in the precipitation biases over the tropical Atlantic.

Across all simulations, the 40 and 5 km-Param setups show similar biases in the moisture distribution, but with reduced amplitude at increased resolution (Fig. 12). In the CTL simulations, the near-surface dry bias is much more pronounced in 5 km-rParam than in the 40 and 5 km-Param setups. In all setups, the increase in U_{\min} reduces the dry bias and changes the sign of the inner-tropical dry bias up to 3 km. In the setups with parameterizations, the U_{\min} fix counterbalances the near-surface specific humidity dry bias, while leading to a slight deterioration of the free tropospheric moisture bias at higher latitudes. In the 5 km-rParam setup, the nearsurface humidity dry bias can not be entirely counterbalanced by the U_{\min} fix, however, the free tropospheric moist bias does not increase as much. The remaining near-surface bias in PTB-5_1-rParam at 5 km resolution amounts to a percentage bias of less than 3 %, 15 % of the background values in 40 and 5 km. Overall, despite the differences described between the two resolutions, both 40 and 5 km exhibit the same symptoms of a too-strong vertical moisture transport from




Figure 10. Simulation scores for all 40 km sensitivity experiments as a function of prescribed minimum surface wind speed. The values for a U_{\min} of 1 m s⁻¹ were taken from the respective two years of the CTL simulation. Values for the asymmetric index A_p , symmetry index E_P as well as latent heat flux, near-surface specific humidity q_s , net top-of-the-atmosphere (TOA) imbalance, upwards radiative shortwave flux, outgoing long wave radiation (OLR) and global mean surface temperature T_S are shown. Each point depicts a one-year global average. For reference the corresponding values from IMERG, GPCP, ERA5 and CERES are depicted.

the boundary layer into the free troposphere between 20 to 40° latitude, which comes at the cost of reduced near-surface moisture transport into the tropics, and this remains the case in the setup without deep convective and non-orographic gravity wave parameterizations (rParam).

The increase in atmospheric specific humidity and deep convection in the inner tropics with modified U_{\min} is also visible in the zonal-mean temperature biases in Fig. 13. The tropical free troposphere warms in both resolutions as the convection ties the temperature profile to a different base-line moist adiabat with increased moisture supply. In the parameterized setups, the cold bias is overcompensated, resulting in

a warm bias. The 5 km-rParam exhibited a stronger cold bias than the parameterized setups. Here, the cold bias is reduced. A possible reason for the differences in the zonal-mean temperatures between the parameterized and non-parameterized simulations could be differences in the entrainment rates into the convective cores. The more pronounced cold bias in the 5 km-rParam setups suggests that more entrainment takes place in the simulations relying on explicitly described deep convection, resulting in an overall colder tropical free troposphere.

We also test the transferability of the tuning to higher resolution, using the same parameter adaptations for the 5 km




Figure 11. Large-scale precipitation bias with respect to IMERG in: 2-year averages from CTL simulations (**a–c**), two-MAM averages from CTL simulations (**d–f**) and two-MAM averages from PTB-5_1 simulations (**g–i**) for 40, 5 and 5 km-rParam. The corresponding global precipitation RMSE is indicated below the panel for each experiment. Statistically insignificant differences between IMERG and the experiments are shown based on a two-sided *z*-test at $\alpha = 0.1$, and insignificant regions are grayed out. The global RMSE and mean error in the near-surface specific humidity, calculated with respect to the values derived from the ERA5 reanalysis, are shown in the inlays.

setups as for 40 km. The simulation scores from this exercise are shown in Fig. 14. In many variables, 40 and 5 km behave similarly, although it is also evident that the 5 km simulations exhibit a greater spread (i.e., variability) than the coarser 40 km counterpart for some variables. In A_s all simulations perform well compared to IMERG, although the 5 km-rParam simulations exhibit a small positive deviation. The E_p score is negatively biased in the untuned version and improves for both resolutions with the U_{\min} adjustment. The improvement in near-surface specific humidity comes at the cost of an increased bias in latent heat flux. The TOA radiative imbalance shows increased variability in the 5 km simulations despite no clear drift in the three year control simulation from which these sensitivity experiments are branched off. This suggests that newly resolved processes enhance variability, both in SW up (suggesting a role of clouds) and OLR (suggesting a role of high clouds). The global-mean

near-surface temperature $T_{\rm s}$ is the only parameter in which different setups show opposing trends with adjustment of the $U_{\rm min}$ and tuning: warming for parameterized setups and cooling for 5 km-rParam. For the 40 and 5 km-rParam setups, the tuned versions are coming closer to the observational values. The 5 km simulations with parameterizations shows a deterioration in $T_{\rm s}$ bias.

Summarizing the main findings in this section, we demonstrated

- 1. that the U_{\min} adjustment to counteract the development of the double ITCZ is similarly effective in 5 km simulations as in 40 km simulations and
- 2. that the 40 km tuning is transferable to the 5 km-Param simulations.




Figure 12. Atmospheric specific humidity bias with respect to ERA5: all CTL for a 2 year average (**a–c**), all CTL for an average over two MAM seasons (**d–f**) and all PTB-5_1 for an average over two MAM seasons (**g–i**) for 40, 5 and 5 km-rParam. Statistically significant differences between ERA5 and the model data are shown, insignificant regions are hatched.

We also showed that the humidity and temperature fields change in a similar fashion to the adjustments of U_{\min} at different resolutions.

4 Discussion and Conclusions

In our study, we compared the representation of the precipitation as simulated by ICON over a resolution hierarchy spanning from CMIP-resolution up to km-scale. Although some small-scale improvements, such as a better representation of precipitation over land with more pronounced orography were found, increased resolution and the discard of deep convective and non-orographic gravity wave parameterization could not resolve persistent large-scale precipitation biases. The precipitation biases in our ICON resolution hierarchy arise from both atmospheric contributions and evaporation biases linked to a mismatch between prescribed SSTs and atmospheric conditions. Although our model setup is AMIPtype, it also expresses a double-ITCZ bias as seen across several CMIP generations (Tian et al., 2024). Additionally, we find a too-cold tropical free troposphere and a too-moist free troposphere in the subtropics. These biases are symptoms of a general misrepresentation of the exchange of energy and moisture between the inner- and outer tropics (Fig. 1a, cf. Fig. 1b). Again, similar temperature and humidity biases can be found in the CMIP models (Tian et al., 2013, 2024), suggesting that the described bias in moisture export from the subtropical boundary layer is not unique to the ICON model. To address the biases, we concentrated on the parameterizations that remain active at the km-scale, focusing on U_{\min} in the bulk flux formulation of the turbulence scheme. It is known that the bulk flux formulation can lead to biased results compared to observations in both very low and very high wind regimes, producing a high bias in latent heat release (Hsu et al., 2022). Additionally, adjusting U_{\min} yielded promising results in the Sapphire configuration of the ICON model (Segura et al., 2025). The usage of increased U_{\min} in the bulk flux formula resolved some of the precipitation biases in our ICON XPP simulations. This improvement came at the cost of a global overestimation of evaporative fluxes. Furthermore, a deterioration of precipitation over the land made a separate adjustment necessary. This was not seen in Segura et al. (2025) due to their focus on the double-ITCZ feature in the Warm Pool region. Furthermore, we showed that the U_{\min} adjustment led to similar changes at the kmscale as in our lower-resolution simulations. Our analysis also gives new insights into the mechanisms by which U_{\min} acts: By increasing evaporation, it counterbalances the low biases in the near-surface specific humidity in the tropics, enhancing deep convective activity, which increases free tro-




Figure 13. Zonal mean temperature bias with respect to ERA5: all CTL for a 2 year average (**a–c**), all CTL for an average over two MAM seasons (**d–f**) and all PTB-5_1 for an average over two MAM seasons (**g–i**) for 40, 5 and 5 km-rParam. Statistically significant differences between ERA5 and the model data are shown, insignificant regions are hatched.

pospheric temperatures and the temperature inversion in subsidence regions.

The fact that the model can not replicate the observed nearsurface wind distribution hints at more overarching problems in the global near-surface moisture advection. The increase in U_{\min} speed acts to address the too weak Walker circulation but does not address the lack of moisture transport from the higher (sub)tropical latitudes starting at 30° latitude, which is likely the origin of the tropical dry bias (Fig. 12c). This also means that the additional supply of near-surface moisture is drawn from the wrong area, i.e. the inner tropics instead of the subtropics. Our moisture budget analysis shows an exaggerated vertical moisture transport out of the boundary layer (cf. Fig. 9), which explains the deficit in moisture available for transport into the inner tropics. The study by Lang et al. (2023) gives further evidence that exaggerated vertical moisture transport out of the boundary layer is related to turbulent transport. The vertical moisture transport out of the boundary layer may be intensified if parameterizations such as the deep convective parameterization are discarded prematurely. Another source of the bias might also be the shallow convection; however, as similar biases were also encountered in the ICON sapphire configuration without the shallow convection scheme (Segura et al., 2025), a bias in the used turbulence schemes or in the dynamical core is more likely.

In summary, the main findings of this work are:

- Similar large-scale precipitation biases persist throughout the ICON resolution hierarchy up to km-scale resolution, although improvements over tropical islands in the Warm Pool region appear with increasing resolution and discard of parameterizations.
- 2. Precipitation biases can be reduced by adapting undiscardable parameterizations but only yield an overall improvement of the precipitation fields if ocean and land are treated separately. The direction of regional changes hold over different resolutions. The main driving mechanism is a strengthening of the Walker circulation by increasing the supply of available moisture for the deep convective regions. However, the Umin fix proposed by Segura et al. (2025) leads to additional drag on near-surface winds in higher latitudes. If the Umin limiter has to be used as a patch, it may be advisable to restrict it to the bulk flux formula for evaporation and not those for momentum exchange and sensible heat.
- 3. There are strong indications that the U_{\min} fix is not addressing the underlying core problem. A uniform wind threshold does not account for different regional circulation regimes and will lead to too high evaporation values in regions with low wind speed variations. Especially




Figure 14. Simulation scores for two MAM seasons of CTL, PTB-5_1 and PTB-5_1t sensitivity experiments in the 40, 5 and 5 km-rParam setup. The 40 km scores are shown in different shades of red, the 5 km setups in shades of green and the 5 km-rParam in different shades of blue. Values for the asymmetric index A_p , symmetry index E_p as well as latent heat flux, near-surface specific humidity q_s , net top-of-the-atmosphere (TOA) imbalance, upwards radiative shortwave flux, outgoing long wave radiation (OLR) and global mean near-surface temperature T_s are shown. Each point depicts a one season average. For reference the corresponding values from IMERG, GPCP, ERA5 and CERES are listed.

noticeable is also that the global latent heat flux and the 20 to 40° latitude free-tropospheric moisture bias are increased in the adjusted simulations. This demonstrates that there are problems with the source regions and transport of atmospheric moisture. The fix leads to a replacement of subtropical moisture by moisture originating from the inner tropics. Additionally, the increase in drag on the near-surface wind speeds leads to a slow down of the trade winds, such that the adjustment might compromise the representation of extra-tropical to tropical teleconnections. In a coupled climate simulation,

changes in U_{\min} could also critically influence ocean dynamics and vertical structure (e.g., lead to substantial cooling), potentially introducing severe large-scale biases.

- 4. Results indicate a two-fold faulty representation of the moisture transport in ICON, and potentially in the many other models with a double-ITCZ bias:
 - In the subtropics, the vertical moisture transport out of the boundary layer is too strong, leaving too little

moisture to be transported horizontally within the Trades into the tropics.

 In the tropics, the lack of moisture reduces deep convective activity, which in turn reduces temperature inversions and slows down the Walker circulation. The reduced zonally anomalous circulations favors the development of a second ITCZ branch as moisture is not transported into the core regions for deep convection.

The results of this study suggest that a physically motivated correction of the double-ITCZ bias in ICON (and potentailly other AMIP-style CMIP simulations) should focus on reducing exaggerated vertical moisture fluxes in the subtropics. In general, a two-fold climate modeling strategy would be greatly beneficial for the community: On the one side the advancement of resources for high-resolution simulations, on the other side, further development of parameterizations that are needed in both high- and coarse-resolution models. The persistence of large-scale biases across our ICON resolution hierarchy shows the potential of learning from both high- and low-resolution simulations. In the long term, scaleaware parameterizations or parameterizations with a seamless transition between convective and subgrid scale turbulent processes within one unified scheme (e.g., Tan et al., 2018) would help unify high- and low-resolution modeling strategies.

Appendix A: Computation of ERA5 near-surface specific humidity

The near-surface specific humidity q_s , is not included as direct output of ERA5. We derive it via

$$q_{\rm s} = \frac{0.622 \, p_{\rm v}}{p_{\rm s} - 0.378 \, p_{\rm v}}.\tag{A1}$$

using the vapour pressure p_v in hPa, which is calculated using ERA5 output of 2 m dew point temperature T_D in Celsius and surface pressure p_s in hPa according to the Magnus formula

$$p_{\rm v} = 6.112 \times 5 \exp\left(\frac{17.67 T_{\rm D}}{T_{\rm D} + 243.5 \,\rm K}\right).$$
 (A2)

Appendix B: Computation of the velocity potential

The velocity potential χ is obtained by computing the inverse Laplacian of the wind field divergence. Specifically, the wind field in the Helmholtz decomposition can be written as

$$\boldsymbol{u} = \nabla \times \boldsymbol{\psi} - \nabla \chi, \tag{B1}$$

where ψ is the streamfunction. The divergence of the wind field is directly related to χ via

$$\nabla \cdot \boldsymbol{u} = -\nabla^2 \chi, \tag{B2}$$

and $\chi_{l,m}$ in the spectral space can be obtained by calculating the inverse Laplacian in spectral space according to

$$\chi_{l,m} = -\frac{r^2 D_{l,m}}{l(l+1)},\tag{B3}$$

where $D_{l,m}$ are the spectral coefficients of the wind divergence $\nabla \cdot \boldsymbol{u}$, r is the earth's radius, l is the zonal wavenumber and m is the longitudinal wavenumber. We analyze χ after transforming back to grid space.

Appendix C: Near-surface humidity biases





Figure C1. Two-year mean large-scale surface specific humidity bias with respect to ERA5 (2004–2010) over the resolution hierarchy. Global mean root mean square errors are shown below the plots.






Figure C2. Two-year mean large-scale surface humdity bias with respect to ERA5 (2004–2010) for different settings of the surface wind in 40 km: PTB-0.5, PTB-4, PTB-5_1, PTB-5_1t, PTB-6, PTB-6_1 and PTB-6_1t. Contours show the ERA5 climatology. The corresponding global RMSE is stated beneath the panel for each sensitivity experiment.

Code and data availability. The ICON code was released in January 2024 under https://www.icon-model.org/news/news_open_ source_release (last access: 14 March 2025). The ICON XPP code release and source scripts are publicly available and published by Müller et al. (2025b). The model data and postprocessing scripts needed to replicate the work presented in this paper are available from a Zenodo archive: https://doi.org/10.5281/zenodo.17410280. The ERA5 reanalysis data set is available from the climate data store (https://cds.climate.copernicus.eu/, last access: 14 March 2025). IMERG is provided by NASA under https://gpm.nasa. gov/data/directory (last access: 14 March 2025). GPCP data sets are maintained by NOAA under https://psl.noaa.gov/data/gridded/ data.gpcp.html (last access: 14 March 2025). CERES data can be obtained from NASA under https://ceres.larc.nasa.gov/data/ (last access: 14 March 2025). The OAFlux data are maintained by NCAR under https://climatedataguide.ucar.edu/climate-data/ oaflux-objectively-analyzed-air-sea-fluxes-global-oceans (last access: 14 March 2025). For postprocessing and data analysis python and CDO were used; the latter can be downloaded from https: //code.mpimet.mpg.de/projects/cdo (last access: 14 March 2025).

Author contributions. CAK had the idea for the study, conceptualized it, acquired computational resources, performed the analysis and wrote the paper manuscript. Model simulations and tuning were done by UN in 160 km, by AS in 80 km and CAK in 40 and 5 km resolution. RJW provided input on many details of the analysis. LK advised on technical aspects on the model setup and provided the necessary input data. All authors, CAK, RJW, LK, UN and AS, contributed to the interpretation of the results and revised the paper manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. The authors specifically thank the ICON Seamless/ICON XPP Friends-circle and the XPP Precipitation task force for invaluable collaboration and support throughout the project year. This work greatly benefited from inspiring discussions with our colleagues Kristina Fröhlich, Karel Castro-Morales, Maike Ahlgrimm and Trang van Pham. It would not have been possible to run the ICON XPP code on GPU without Roland Wirth, who fixed some remaining GPU porting bugs. Annika Lauber, Jonas Jucker and Michael Jähn from the C2SM team at ETH have been a great discussion contacts for technical issues encountered around ICON. Computing and data storage resources for the 40 and 5 km simulations, were provided by the Swiss National Supercomputing Center (CSCS) in Lugano via the project: s1283 and lp67 as well as

the early-testers access to the vCluster Todi on the new Alps infrastructure. William Sawyer, Christoph Müller, and the technical staff at CSCS have been very supportive of the ICON community when transitioning to the new Alps computing infrastructure. For the R2B5 simulations, we thank the German Weather Services for providing computation resources on their system rts, for the R2B4 simulations, we thank the Deutsches Klima Rechenzentrum (DKRZ) for providing resources granted by its Scientific Steering Committee (WLA) under project ID bm0550. The authors thank the three anonymous reviewers for constructive and detailed comments on this work.

Financial support. This research has been supported by the ETH Postdoctoral Fellowship Program (CAK), the Swiss National Science Foundation Award PCEFP2 203376 (RJW), the research unit FOR 2820 VolImpact (grant no. 398006378) funded by the German Research Foundation (DFG) within the project VOLARC (UN) and DWD's "Innovation Programme for Applied Researches and Developments" IAFE ICON-Seamless VH 4.7 (AS).

Review statement. This paper was edited by Michael Byrne and reviewed by three anonymous referees.

References

Adam, O., Schneider, T., Brient, F., and Bischoff, T.: Relation of the double-ITCZ bias to the atmospheric energy budget in climate models, Geophysical Research Letters, 43, 7670–7677, https://doi.org/10.1002/2016GL069465, 2016.

Adam, O., Schneider, T., and Brient, F.: Regional and seasonal variations of the double-ITCZ bias in CMIP5 models, Climate Dynamics, 51, 101–117, https://doi.org/10.1007/s00382-017-3909-1, 2018.

Arakawa, A.: The Cumulus Parameterization Problem: Past, Present, and Future, Journal of Climate, 17, 2493–2525, 2004.

Bechtold, P., Köhler, M., Jung, T., Doblas-Reyes, F., Leutbecher, M., Rodwell, M. J., Vitart, F., and Balsamo, G.: Advances in simulating atmospheric variability with the ECMWF model: From synoptic to decadal time-scales, Quarterly Journal of the Royal Meteorological Society, 134, 1337–1351, https://doi.org/10.1002/qj.289, 2008.

Bischoff, T. and Schneider, T.: Energetic Constraints on the Position of the Intertropical Convergence Zone, Journal of Climate, 27, 4937–4951, https://doi.org/10.1175/JCLI-D-13-00650.1, 2014.

Boer, G. J.: Diagnostic Equations in Isobaric Coordinates, Monthly Weather Review, 110, 1801–1820, https://doi.org/10.1175/1520-0493(1982)110<1801:DEIIC>2.0.CO;2, 1982.

Bretherton, C. S. and Smolarkiewicz, P. K.: Gravity Waves, Compensating Subsidence and Detrainment around Cumulus Clouds, Journal of Atmospheric Sciences, 46, 740–759, https://doi.org/10.1175/1520-0469(1989)046<0740:GWCSAD>2.0.CO;2,1989.

C3S: ERA5 hourly data on single levels from 1940 to present, Climate Data Store [data set], https://doi.org/10.24381/CDS.ADBB2D47, 2018.

- Charney, J. G.: A Note on Large-Scale Motions in the Tropics, Journal of the Atmospheric Sciences, 20, 607–609, https://doi.org/10.1175/1520-0469(1963)020<0607:ANOLSM>2.0.CO;2, 1963.
- Clark, J. P., Lin, P., and Hill, S. A.: ITCZ Response to Disabling Parameterized Convection in Global Fixed-SST GFDL-AM4 Aquaplanet Simulations at 50 and 6 km Resolutions, Journal of Advances in Modeling Earth Systems, 16, e2023MS003968, https://doi.org/10.1029/2023MS003968, 2024.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, Quarterly Journal of the Royal Meteorological Society, 137, 553–597, https://doi.org/10.1002/qj.828, 2011.
- Dipankar, A., Bianco, M., Bukenberger, M., Ehrengruber, T., Farabullini, N., Gopal, A., Hupp, D., Jocksch, A., Kellerhals, S., Kroll, C. A., Lapillonne, X., Leclair, M., Luz, M., Müller, C., Ong, C. R., Osuna, C., Pothapakula, P., Röthlin, M., Sawyer, W., Serafini, G., Vogt, H., Weber, B., and Schulthess, T.: Toward Exascale Climate Modelling: A Python DSL Approach to ICON's (Icosahedral Non-hydrostatic) Dynamical Core (icon-exclaim v0.2.0), EGUsphere [preprint], https://doi.org/10.5194/egusphere-2025-4808, 2025.
- Doelling, D. R., Loeb, N. G., Keyes, D. F., Nordeen, M. L., Morstad, D., Nguyen, C., Wielicki, B. A., Young, D. F., and Sun, M.: Geostationary Enhanced Temporal Interpolation for CERES Flux Products, Journal of Atmospheric and Oceanic Technology, 30, 1072–1090, https://doi.org/10.1175/JTECH-D-12-00136.1, 2013.
- Doelling, D. R., Sun, M., Nguyen, L. T., Nordeen, M. L., Haney, C. O., Keyes, D. F., and Mlynczak, P. E.: Advances in Geostationary-Derived Longwave Fluxes for the CERES Synoptic (SYN1deg) Product, Journal of Atmospheric and Oceanic Technology, 33, 503–521, https://doi.org/10.1175/JTECH-D-15-0147.1, 2016.
- Dong, Y., Armour, K. C., Battisti, D. S., and Blanchard-Wrigglesworth, E.: Two-Way Teleconnections between the Southern Ocean and the Tropical Pacific via a Dynamic Feedback, Journal of Climate, 35, 6267–6282, https://doi.org/10.1175/JCLI-D-22-0080.1, 2022.
- Durack, P. J. and Taylor, K. E.: PCMDI AMIP SST and sea-ice boundary conditions version 1.1.4, Version 2018, Earth System Grid Federation [data set], https://doi.org/10.22033/ESGF/INPUT4MIPS.2204, 2018.
- Feng, X., Ding, Q., Wu, L., Jones, C., Wang, H., Bushuk, M., and Topál, D.: Comprehensive Representation of Tropical – Extratropical Teleconnections Obstructed by Tropical Pacific Convection Biases in CMIP6, Journal of Climate, 36, 7041–7059, https://doi.org/10.1175/JCLI-D-22-0523.1, 2023.
- Frierson, D. M. W., Hwang, Y.-T., Fučkar, N. S., Seager, R., Kang, S. M., Donohoe, A., Maroon, E. A., Liu, X., and Battisti, D. S.: Contribution of ocean overturning circulation to tropical rainfall

- peak in the Northern Hemisphere, Nature Geoscience, 6, 940–944, https://doi.org/10.1038/ngeo1987, 2013.
- Früh, B., Potthast, R., Müller, W., Korn, P., Brienen, S., Fröhlich, K., Helmert, J., Köhler, M., Lorenz, S., Pham, T. V., Pohlmann, H., Schlemmer, L., Schnur, R., Schulz, J.-P., Sgoff, C., Vogel, B., Wirth, R., and Zängl, G.: ICON-Seamless, the development of a novel Earth System Model based on ICON for time scales from weather to climate, EMS Annual Meeting 2022, Bonn, Germany, 5–9 September 2022, EMS2022-292, https://doi.org/10.5194/ems2022-292, 2022.
- Giorgetta, M. A., Sawyer, W., Lapillonne, X., Adamidis, P., Alexev, D., Clément, V., Dietlicher, R., Engels, J. F., Esch, M., Franke, H., Frauen, C., Hannah, W. M., Hillman, B. R., Kornblueh, L., Marti, P., Norman, M. R., Pincus, R., Rast, S., Reinert, D., Schnur, R., Schulzweida, U., and Stevens, B.: The ICON-A model for direct QBO simulations on GPUs (version icon-cscs:baf28a514), Geosci. Model Dev., 15, 6985–7016, https://doi.org/10.5194/gmd-15-6985-2022, 2022.
- Gonzalez, A. O., Fahrin, F., Magnusdottir, G., Kinsella, A., and Ganguly, I.: We need to simulate more double ITCZs and less southern ITCZs in reanalyses and coupled climate models, ESS Open Archive, https://doi.org/10.22541/essoar.174017095.57302520/v1, 2025.
- Hájková, D. and Šácha, P.: Parameterized orographic gravity wave drag and dynamical effects in CMIP6 models, Climate Dynamics, https://doi.org/10.1007/s00382-023-07021-0, 2023.
- Ham, Y.-G. and Kug, J.-S.: Effects of Pacific Intertropical Convergence Zone precipitation bias on ENSO phase transition, Environmental Research Letters, 9, 064008, https://doi.org/10.1088/1748-9326/9/6/064008, 2014.
- Hardiman, S. C., Boutle, I. A., Bushell, A. C., Butchart, N., Cullen, M. J. P., Field, P. R., Furtado, K., Manners, J. C., Milton, S. F., Morcrette, C., O'Connor, F. M., Shipway, B. J., Smith, C., Walters, D. N., Willett, M. R., Williams, K. D., Wood, N., Abraham, N. L., Keeble, J., Maycock, A. C., Thuburn, J., and Woodhouse, M. T.: Processes Controlling Tropical Tropopause Temperature and Stratospheric Water Vapor in Climate Models, Journal of Climate, 28, 6516–6535, https://doi.org/10.1175/JCLI-D-15-0075.1, 2015.
- Held, I. M. and Hoskins, B. J.: Large-Scale Eddies and the General Circulation of the Troposphere, in: Advances in Geophysics, vol. 28, 3–31, Elsevier, ISBN 978-0-12-018828-4, https://doi.org/10.1016/S0065-2687(08)60218-6, 1985.
- Henderson, S. A., Maloney, E. D., and Son, S.-W.: Madden–Julian Oscillation Pacific Teleconnections: The Impact of the Basic State and MJO Representation in General Circulation Models, Journal of Climate, 30, 4567–4587, https://doi.org/10.1175/JCLI-D-16-0789.1, 2017.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A.,
 Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M.,
 De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S. B., Hogan, R., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: Fifth generation of ECMWF atmospheric reanalyses of the

- global climate. Complete ERA5 from 1940: Fifth generation of ECMWF atmospheric reanalyses of the global climate, Copernicus Climate Change Service (C3S) Data Store (CDS) [data set], https://doi.org/10.24381/cds.143582cf, 2017.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, Quarterly Journal of the Royal Meteorological Society, 146, 1999–2049, https://doi.org/10.1002/qj.3803, 2020.
- Hogan, R. J. and Bozzo, A.: A Flexible and Efficient Radiation Scheme for the ECMWF Model, Journal of Advances in Modeling Earth Systems, 10, 1990–2008, https://doi.org/10.1029/2018MS001364, 2018.
- Hohenegger, C., Korn, P., Linardakis, L., Redler, R., Schnur, R., Adamidis, P., Bao, J., Bastin, S., Behravesh, M., Bergemann, M., Biercamp, J., Bockelmann, H., Brokopf, R., Brüggemann, N., Casaroli, L., Chegini, F., Datseris, G., Esch, M., George, G., Giorgetta, M., Gutjahr, O., Haak, H., Hanke, M., Ilyina, T., Jahns, T., Jungclaus, J., Kern, M., Klocke, D., Kluft, L., Kölling, T., Kornblueh, L., Kosukhin, S., Kroll, C., Lee, J., Mauritsen, T., Mehlmann, C., Mieslinger, T., Naumann, A. K., Paccini, L., Peinado, A., Praturi, D. S., Putrasahan, D., Rast, S., Riddick, T., Roeber, N., Schmidt, H., Schulzweida, U., Schütte, F., Segura, H., Shevchenko, R., Singh, V., Specht, M., Stephan, C. C., von Storch, J.-S., Vogel, R., Wengel, C., Winkler, M., Ziemen, F., Marotzke, J., and Stevens, B.: ICON-Sapphire: simulating the components of the Earth system and their interactions at kilometer and subkilometer scales, Geosci. Model Dev., 16, 779-811, https://doi.org/10.5194/gmd-16-779-2023, 2023.
- Holton, J. R.: An Introduction to Dynamic Meteorology, Academic Press, Amsterdam, Boston, 4th edn., ISBN 9780123848666, 2004
- Hourdin, F., Mauritsen, T., Gettelman, A., Golaz, J.-C., Balaji, V., Duan, Q., Folini, D., Ji, D., Klocke, D., Qian, Y., Rauser, F., Rio, C., Tomassini, L., Watanabe, M., and Williamson, D.: The Art and Science of Climate Model Tuning, Bulletin of the American Meteorological Society, 98, 589–602, https://doi.org/10.1175/BAMS-D-15-00135.1, 2017.
- Hsu, C., DeMott, C. A., Branson, M. D., Reeves Eyre, J., and Zeng, X.: Ocean Surface Flux Algorithm Effects on Tropical Indo-Pacific Intraseasonal Precipitation, Geophysical Research Letters, 49, e2021GL096968, https://doi.org/10.1029/2021GL096968, 2022.
- Hu, Z., Lamraoui, F., and Kuang, Z.: Influence of Upper-Troposphere Stratification and Cloud–Radiation Interaction on Convective Overshoots in the Tropical Tropopause Layer, Journal of the Atmospheric Sciences, 78, 2493–2509, https://doi.org/10.1175/JAS-D-20-0241.1, 2021.
- Huffman, G. J.: GPCP Precipitation Level 3 Monthly 0.5-Degree V3.2, GES DISC [data set], https://doi.org/10.5067/MEASURES/GPCP/DATA304, 2021.
- Huffman, G. J., Bolvin, D. T., Nelkin, E. J., and Stocker, J. T.: V06 IMERG Release Notes, NASA,

- https://gpm.nasa.gov/sites/default/files/2020-10/IMERG_ V06_release_notes_201006_0.pdf (last access: 13 March 2025), 2010.
- Hwang, Y.-T. and Frierson, D. M. W.: Link between the double-Intertropical Convergence Zone problem and cloud biases over the Southern Ocean, Proceedings of the National Academy of Sciences, 110, 4935–4940, https://doi.org/10.1073/pnas.1213302110, 2013.
- Iles, C. E., Vautard, R., Strachan, J., Joussaume, S., Eggen, B. R., and Hewitt, C. D.: The benefits of increasing resolution in global and regional climate simulations for European climate extremes, Geosci. Model Dev., 13, 5583–5607, https://doi.org/10.5194/gmd-13-5583-2020, 2020.
- Jeevanjee, N.: Vertical Velocity in the Gray Zone, Journal of Advances in Modeling Earth Systems, 9, 2304–2316, https://doi.org/10.1002/2017MS001059, 2017.
- Jones, T. R. and Randall, D. A.: Quantifying the limits of convective parameterizations, Journal of Geophysical Research: Atmospheres, 116, https://doi.org/10.1029/2010JD014913, 2011.
- Kang, S. M., Held, I. M., Frierson, D. M. W., and Zhao, M.: The Response of the ITCZ to Extratropical Thermal Forcing: Idealized Slab-Ocean Experiments with a GCM, Journal of Climate, 21, 3521–3532, https://doi.org/10.1175/2007JCLI2146.1, 2008.
- Kang, S. M., Hawcroft, M., Xiang, B., Hwang, Y.-T., Cazes, G., Codron, F., Crueger, T., Deser, C., Hodnebrog, ., Kim, H., Kim, J., Kosaka, Y., Losada, T., Mechoso, C. R., Myhre, G., Seland, ., Stevens, B., Watanabe, M., and Yu, S.: Extratropical–Tropical Interaction Model Intercomparison Project (Etin-Mip): Protocol and Initial Results, Bulletin of the American Meteorological Society, 100, 2589–2606, https://doi.org/10.1175/BAMS-D-18-0301.1, 2019.
- Lang, T., Naumann, A. K., Buehler, S. A., Stevens, B., Schmidt, H., and Aemisegger, F.: Sources of Uncertainty in Mid-Tropospheric Tropical Humidity in Global Storm-Resolving Simulations, Journal of Advances in Modeling Earth Systems, 15, e2022MS003443, https://doi.org/10.1029/2022MS003443, 2023.
- Lapillonne, X., Hupp, D., Gessler, F., Walser, A., Pauling, A.,
 Lauber, A., Cumming, B., Osuna, C., Müller, C., Merker, C.,
 Leuenberger, D., Leutwyler, D., Alexeev, D., Vollenweider, G.,
 Van Parys, G., Jucker, J., Jansing, L., Arpagaus, M., Induni,
 M., Jacob, M., Kraushaar, M., Jähn, M., Stellio, M., Fuhrer, O.,
 Baumann, P., Steiner, P., Kaufmann, P., Dietlicher, R., Müller,
 R., Kosukhin, S., Schulthess, T. C., Schättler, U., Cherkas,
 V., and Sawyer, W.: Operational numerical weather prediction
 with ICON on GPUs (version 2024.10), EGUsphere [preprint],
 https://doi.org/10.5194/egusphere-2025-3585, 2025.
- Lin, J.-L.: The Double-ITCZ Problem in IPCC AR4 Coupled GCMs: Ocean–Atmosphere Feedback Analysis, Journal of Climate, 20, 4497–4525, https://doi.org/10.1175/JCLI4272.1, 2007.
- Lott, F. and Miller, M. J.: A new subgrid-scale orographic drag parametrization: Its formulation and testing, Quarterly Journal of the Royal Meteorological Society, 123, 101–127, https://doi.org/10.1002/qj.49712353704, 1997.
- Lu, J., Vecchi, G. A., and Reichler, T.: Expansion of the Hadley cell under global warming, Geophysical Research Letters, 34, 2006GL028443, https://doi.org/10.1029/2006GL028443, 2007.
- Ma, X., Zhao, S., Zhang, H., and Wang, W.: The double-ITCZ problem in CMIP6 and the influences of deep convection and model

- resolution, International Journal of Climatology, 43, 2369–2390, https://doi.org/10.1002/joc.7980, 2023.
- Magnusdottir, G. and Wang, C.-C.: Intertropical Convergence Zones during the Active Season in Daily Data, Journal of the Atmospheric Sciences, 65, 2425–2436, https://doi.org/10.1175/2007JAS2518.1, 2008.
- Marshall, J., Donohoe, A., Ferreira, D., and McGee, D.: The ocean's role in setting the mean position of the Inter-Tropical Convergence Zone, Climate Dynamics, 42, 1967–1979, https://doi.org/10.1007/s00382-013-1767-z, 2014.
- Martens, B., Schumacher, D. L., Wouters, H., Muñoz-Sabater, J., Verhoest, N. E. C., and Miralles, D. G.: Evaluating the landsurface energy partitioning in ERA5, Geosci. Model Dev., 13, 4159–4181, https://doi.org/10.5194/gmd-13-4159-2020, 2020.
- Mauritsen, T., Svensson, G., Zilitinkevich, S. S., Esau, I., Enger, L., and Grisogono, B.: A Total Turbulent Energy Closure Model for Neutrally and Stably Stratified Atmospheric Boundary Layers, Journal of the Atmospheric Sciences, 64, 4113–4126, https://doi.org/10.1175/2007JAS2294.1, 2007.
- Mauritsen, T., Stevens, B., Roeckner, E., Crueger, T., Esch, M., Giorgetta, M., Haak, H., Jungclaus, J., Klocke, D., Matei, D., Mikolajewicz, U., Notz, D., Pincus, R., Schmidt, H., and Tomassini, L.: Tuning the climate of a global model, Journal of Advances in Modeling Earth Systems, 4, 2012MS000154, https://doi.org/10.1029/2012MS000154, 2012.
- Mechoso, C., Robertson, A., Barth, N., Davey, M., Delecluse, P., Gent, P., Ineson, S., Kirtman, B., Latif, M., Treut, H. L., Nagai, T., Neelin, J., Philander, S., Polcher, J., Schopf, P., Stockdale, T., Suarez, M., Terray, L., Thual, O., and Tribbia, J.: The Seasonal Cycle over the Tropical Pacific in Coupled Ocean–Atmosphere General Circulation Models, Monthly Weather Review, 123, 2825–2838, https://doi.org/10.1175/1520-0493(1995)123<2825:TSCOTT>2.0.CO;2, 1995.
- Morris, M., Kushner, P. J., Moore, G. W. K., and Mercan, O.: Resolution Dependence of Extreme Wind Speed Projections in the Great Lakes Region, Journal of Climate, 37, 3153–3171, https://doi.org/10.1175/JCLI-D-23-0547.1, 2024.
- Müller, W. A., Früh, B., Korn, P., Potthast, R., Baehr, J., Bettems, J.-M., Bölöni, G., Brienen, S., Fröhlich, K., Helmert, J., Jungclaus, J., Köhler, M., Lorenz, S., Schneidereit, A., Schnur, R., Schulz, J.-P., Schlemmer, L., Sgoff, C., Pham, T. V., Pohlmann, H., Vogel, B., Vogel, H., Wirth, R., Zaehle, S., Zängl, G., Stevens, B., and Marotzke, J.: ICON: Towards vertically integrated model configurations for numerical weather prediction, climate predictions and projections, Bulletin of the American Meteorological Society, https://doi.org/10.1175/BAMS-D-24-0042.1, 2025a.
- Müller, W. A., Lorenz, S., Pham, T. V., Schneidereit, A., Brokopf, R., Brovkin, V., Brüggemann, N., Chegini, F., Dommenget, D., Fröhlich, K., Früh, B., Gayler, V., Haak, H., Hagemann, S., Hanke, M., Ilyina, T., Jungclaus, J., Köhler, M., Korn, P., Kornblüh, L., Kroll, C., Krüger, J., Castro-Morales, K., Niemeier, U., Pohlmann, H., Polkova, I., Potthast, R., Riddick, T., Schlund, M., Stacke, T., Wirth, R., Yu, D., and Marotzke, J.: The ICON-based Earth System Model for Climate Predictions and Projections (ICON XPP v1.0), EGUsphere [preprint], https://doi.org/10.5194/egusphere-2025-2473, 2025b.
- National Center for Atmospheric Research Staff, E.: The Climate Data Guide: OAFlux: Objectively Analyzed air-sea Fluxes for the global oceans, National Center for Atmospheric

- Research, https://climatedataguide.ucar.edu/climate-data/oaflux-objectively-analyzed-air-sea-fluxes-global-oceans (last access: 13 March 2025), 2022.
- Niemeier, U., Wallis, S., Timmreck, C., Van Pham, T., and Von Savigny, C.: How the Hunga Tonga Hunga Ha'apai Water Vapor Cloud Impacts Its Transport Through the Stratosphere: Dynamical and Radiative Effects, Geophysical Research Letters, 50, e2023GL106482, https://doi.org/10.1029/2023GL106482, 2023.
- Orr, A., Bechtold, P., Scinocca, J., Ern, M., and Janiskova, M.: Improved Middle Atmosphere Climate and Forecasts in the ECMWF Model through a Nonorographic Gravity Wave Drag Parameterization, Journal of Climate, 23, 5905–5926, https://doi.org/10.1175/2010JCLI3490.1, 2010.
- Paccini, L., Hohenegger, C., and Stevens, B.: Explicit versus Parameterized Convection in Response to the Atlantic Meridional Mode, Journal of Climate, 34, 3343–3354, https://doi.org/10.1175/JCLI-D-20-0224.1, 2021.
- Pan, L. L., Honomichl, S. B., Kinnison, D. E., Abalos, M., Randel, W. J., Bergman, J. W., and Bian, J.: Transport of chemical tracers from the boundary layer to stratosphere associated with the dynamics of the Asian summer monsoon, Journal of Geophysical Research: Atmospheres, 121, https://doi.org/10.1002/2016JD025616, 2016.
- Peixóto, J. P. and Oort, A. H.: The Atmospheric Branch Of The Hydrological Cycle And Climate, in: Variations in the Global Water Budget, edited by: Street-Perrott, A., Beran, M., and Ratcliffe, R., Springer Netherlands, Dordrecht, 5–65, ISBN 978-94-009-6956-8 978-94-009-6954-4, https://doi.org/10.1007/978-94-009-6954-4
 2, 1983.
- Peixoto, J. P. and Oort, A. H.: Physics of Climate, American Institute of Physics, Melville, NY, USA, 520 pp., ISBN 978-0-88318-712-8, 1992.
- Philander, S. G. H., Gu, D., Lambert, G., Li, T., Halpern, D., Lau, N.-C., and Pacanowski, R. C.: Why the ITCZ Is Mostly North of the Equator, Journal of Climate, 9, 2958–2972, https://doi.org/10.1175/1520-0442(1996)009<2958:WTIIMN>2.0.CO;2, 1996.
- Prill, F., Reinert, D., and Zängl, G.: ICON Tutorial Working with the ICON Model, Deutscher Wetterdienst, Business Area "Research and Development", https://doi.org/10.5676/DWD_pub/nwv/icon_tutorial2023, 2023.
- Reick, C. H., Gayler, V., Goll, D., Hagemann, S., Heidkamp, M., Nabel, J. E. M. S., Raddatz, T., Roeckner, E., Schnur, R., and Wilkenskjeld, S.: JSBACH 3 The land component of the MPI Earth System Model: documentation of version 3.2, Max Planck Institute for Meteorology, Hamburg, 4990986, https://doi.org/10.17617/2.3279802, 2021.
- Ren, Z. and Zhou, T.: Understanding the alleviation of "Double-ITCZ" bias in CMIP6 models from the perspective of atmospheric energy balance, Climate Dynamics, https://doi.org/10.1007/s00382-024-07238-7, 2024.
- Schmidt, H., Rast, S., Bao, J., Cassim, A., Fang, S.-W., Jimenez-de la Cuesta, D., Keil, P., Kluft, L., Kroll, C., Lang, T., Niemeier, U., Schneidereit, A., Williams, A. I. L., and Stevens, B.: Effects of vertical grid spacing on the climate simulated in the ICON-Sapphire global storm-resolving model, Geosci. Model Dev., 17, 1563–1584, https://doi.org/10.5194/gmd-17-1563-2024, 2024.

- Schneider, D. P., Deser, C., Fasullo, J., and Trenberth, K. E.: Climate Data Guide Spurs Discovery and Understanding, Eos, Transactions American Geophysical Union, 94, 121–122, https://doi.org/10.1002/2013EO130001, 2013.
- Schneider, T., Bischoff, T., and Haug, G. H.: Migrations and dynamics of the intertropical convergence zone, Nature, 513, 45–53, https://doi.org/10.1038/nature13636, 2014.
- Schneider, T., Leung, L. R., and Wills, R. C. J.: Opinion: Optimizing climate models with process knowledge, resolution, and artificial intelligence, Atmos. Chem. Phys., 24, 7041–7062, https://doi.org/10.5194/acp-24-7041-2024, 2024.
- Segura, H., Bayley, C., Fievét, R., Glöckner, H., Günther, M., Kluft, L., Naumann, A. K., Ortega, S., Praturi, D. S., Rixen, M., Schmidt, H., Winkler, M., Hohenegger, C., and Stevens, B.: A Single Tropical Rainbelt in Global Storm-Resolving Models: The Role of Surface Heat Fluxes Over the Warm Pool, Journal of Advances in Modeling Earth Systems, 17, e2024MS004897, https://doi.org/10.1029/2024MS004897, 2025.
- Seifert, A.: A revised cloud microphysical parameterization for COSMO-LME, http://www.cosmo-model.org (last access: 19 March 2025), 2008.
- Sherwood, S. C., Bony, S., and Dufresne, J.-L.: Spread in model climate sensitivity traced to atmospheric convective mixing, Nature, 505, 37–42, https://doi.org/10.1038/nature12829, 2014.
- Si, W., Liu, H., Zhang, X., and Zhang, M.: Double Intertropical Convergence Zones in Coupled Ocean-Atmosphere Models: Progress in CMIP6, Geophysical Research Letters, 48, e2021GL094779, https://doi.org/10.1029/2021GL094779, 2021.
- Simmons, A., Soci, C., Nicolas, J., Bell, B., Berrisford, P., Dragani, R., Flemming, J., Haimberger, L., Healy, S., Hersbach, H., Horányi, A., Inness, A., Munoz-Sabater, J., Radu, R., and Schepers, D.: ERA5.1: Rerun of the Fifth generation of ECMWF atmospheric reanalyses of the global climate (2000–2006 only), Climate Data Store [data set], https://doi.org/10.24381/cds.143582cf, 2020.
- Simpson, I. R., McKinnon, K. A., Kennedy, D., Lawrence, D. M., Lehner, F., and Seager, R.: Observed humidity trends in dry regions contradict climate models, Proceedings of the National Academy of Sciences, 121, e2302480120, https://doi.org/10.1073/pnas.2302480120, 2024.
- Song, X. and Zhang, G. J.: The Roles of Convection Parameterization in the Formation of Double ITCZ Syndrome in the NCAR CESM: I. Atmospheric Processes, Journal of Advances in Modeling Earth Systems, 10, 842–866, https://doi.org/10.1002/2017MS001191, 2018.
- Song, X., Ning, C., Duan, Y., Wang, H., Li, C., Yang, Y., Liu, J., and Yu, W.: Observed Extreme Air–Sea Heat Flux Variations during Three Tropical Cyclones in the Tropical Southeastern Indian Ocean, Journal of Climate, 34, 3683–3705, https://doi.org/10.1175/JCLI-D-20-0170.1, 2021.
- Tan, Z., Kaul, C. M., Pressel, K. G., Cohen, Y., Schneider, T., and Teixeira, J.: An Extended Eddy-Diffusivity Mass-Flux Scheme for Unified Representation of Subgrid-Scale Turbulence and Convection, Journal of Advances in Modeling Earth Systems, 10, 770–800, https://doi.org/10.1002/2017MS001162, 2018.
- Tanaka, H. L., Ishizaki, N., and Kitoh, A.: Trend and interannual variability of Walker, monsoon and Hadley circulations defined by velocity potential in the upper troposphere,

- Tellus A: Dynamic Meteorology and Oceanography, 56, 250, https://doi.org/10.3402/tellusa.v56i3.14410, 2004.
- Tian, B. and Dong, X.: The Double-ITCZ Bias in CMIP3, CMIP5, and CMIP6 Models Based on Annual Mean Precipitation, Geophysical Research Letters, 47, e2020GL087232, https://doi.org/10.1029/2020GL087232, 2020.
- Tian, B., Fetzer, E. J., Kahn, B. H., Teixeira, J., Manning, E., and Hearty, T.: Evaluating CMIP5 models using AIRS tropospheric air temperature and specific humidity climatology, Journal of Geophysical Research: Atmospheres, 118, 114–134, https://doi.org/10.1029/2012JD018607, 2013.
- Tian, B., Fetzer, E. J., and Teixeira, J.: Assessing the Tropospheric Temperature and Humidity Simulations in CMIP3/5/6 Models Using the AIRS Obs4MIPs V2.1 Data, Journal of Geophysical Research: Atmospheres, 129, e2023JD040536, https://doi.org/10.1029/2023JD040536, 2024.
- Tiedtke, M.: A Comprehensive Mass Flux Scheme for Cumulus Parameterization in Large-Scale Models, Monthly Weather Review, 117, 1779–1800, https://doi.org/10.1175/1520-0493(1989)117<1779:ACMFSF>2.0.CO;2, 1989.
- Trenberth, K. E.: Some Effects of Finite Sample Size and Persistence Meteorological Statison Part I: Autocorrelations, Monthly Weather 2359-2368, https://doi.org/10.1175/1520-0493(1984)112<2359:SEOFSS>2.0.CO;2, 1984.
- Vergara-Temprado, J., Ban, N., Panosetti, D., Schlemmer, L., and Schär, C.: Climate Models Permit Convection at Much Coarser Resolutions Than Previously Considered, Journal of Climate, 33, 1915–1933, https://doi.org/10.1175/JCLI-D-19-0286.1, 2020.
- Wild, M.: The global energy balance as represented in CMIP6 climate models, Climate Dynamics, 55, 553–577, https://doi.org/10.1007/s00382-020-05282-7, 2020.
- Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOsahedral Non-hydrostatic) modelling framework of DWD and MPI-M: Description of the non-hydrostatic dynamical core, Quarterly Journal of the Royal Meteorological Society, 141, 563–579, https://doi.org/10.1002/qj.2378, 2015.
- Zhang, H., Deser, C., Clement, A., and Tomas, R.: Equatorial signatures of the Pacific Meridional Modes: Dependence on mean climate state, Geophysical Research Letters, 41, 568–574, https://doi.org/10.1002/2013GL058842, 2014.
- Zhang, Y. and Boos, W. R.: An upper bound for extreme temperatures over midlatitude land, Proceedings of the National Academy of Sciences, 120, e2215278120, https://doi.org/10.1073/pnas.2215278120, 2023.
- Zhao, M., Golaz, J., Held, I. M., Guo, H., Balaji, V., Benson, R., Chen, J., Chen, X., Donner, L. J., Dunne, J. P., Dunne, K., Durachta, J., Fan, S., Freidenreich, S. M., Garner, S. T., Ginoux, P., Harris, L. M., Horowitz, L. W., Krasting, J. P., Langenhorst, A. R., Liang, Z., Lin, P., Lin, S., Malyshev, S. L., Mason, E., Milly, P. C. D., Ming, Y., Naik, V., Paulot, F., Paynter, D., Phillipps, P., Radhakrishnan, A., Ramaswamy, V., Robinson, T., Schwarzkopf, D., Seman, C. J., Shevliakova, E., Shen, Z., Shin, H., Silvers, L. G., Wilson, J. R., Winton, M., Wittenberg, A. T., Wyman, B., and Xiang, B.: The GFDL Global Atmosphere and Land Model AM4.0/LM4.0: 2. Model Description, Sensitivity Studies, and Tuning Strategies, Journal of Advances in Modeling Earth Systems, 10, 735–769, https://doi.org/10.1002/2017MS001209, 2018a.

Zhao, M., Golaz, J., Held, I. M., Guo, H., Balaji, V., Benson, R., Chen, J., Chen, X., Donner, L. J., Dunne, J. P., Dunne, K., Durachta, J., Fan, S., Freidenreich, S. M., Garner, S. T., Ginoux, P., Harris, L. M., Horowitz, L. W., Krasting, J. P., Langenhorst, A. R., Liang, Z., Lin, P., Lin, S., Malyshev, S. L., Mason, E., Milly, P. C. D., Ming, Y., Naik, V., Paulot, F., Paynter, D., Phillipps, P., Radhakrishnan, A., Ramaswamy, V., Robinson, T., Schwarzkopf, D., Seman, C. J., Shevliakova, E., Shen, Z., Shin, H., Silvers, L. G., Wilson, J. R., Winton, M., Wittenberg, A. T., Wyman, B., and Xiang, B.: The GFDL Global Atmosphere and Land Model AM4.0/LM4.0: 1. Simulation Characteristics With Prescribed SSTs, Journal of Advances in Modeling Earth Systems, 10, 691–734, https://doi.org/10.1002/2017MS001208, 2018b.

Zhou, W., Leung, L. R., and Lu, J.: Linking Large-Scale Double-ITCZ Bias to Local-Scale Drizzling Bias in Climate Models, Journal of Climate, 35, 7965–7979, https://doi.org/10.1175/JCLI-D-22-0336.1, 2022.