



PISTON and CAMP²Ex observations of the fundamental modes of aerosol vertical variability in the Northwest Tropical Pacific and Maritime Continent's Monsoon

Jeffrey S. Reid¹, Robert E. Holz², Chris A. Hostetler³, Richard A. Ferrare³, Juli I. Rubin⁴, Elizabeth J. Thompson⁵, Susan C. van den Heever⁶, Corey G. Amiot⁷, Sharon P. Burton³, Joshua P. DiGangi³, Glenn S. Diskin³, Joshua H. Cossuth⁸, Daniel P. Eleuterio⁸, Edwin W. Eloranta², Ralph Kuehn², Willem J. Marais², Hal B. Maring⁹, Armin Sorooshian¹⁰, Kenneth L. Thornhill¹¹, Charles R. Trepte³, Jian Wang¹², Peng Xian¹, and Luke D. Ziemba³

¹Marine Meteorology Division, US Naval Research Laboratory, Monterey, CA 93943, USA

²Space Science and Engineering Center University of Wisconsin-Madison, Madison, WI 53715, USA

³NASA Langley Research Center, Hampton, VA 23681, USA

⁴Remote Sensing Division, US Naval Research Laboratory, Washington, DC 20375, USA

⁵NOAA Physical Sciences Lab, Boulder, CO 80305, USA

⁶Department of Atmospheric Science, Colorado State University, Fort Collins, CO 80523, USA

⁷NASA Postdoctoral Program, NASA Marshall Space Flight Center, Huntsville, AL 35805, USA

⁸Office of Naval Research, Code 322, Arlington, VA 22203-1995, USA

⁹NASA Headquarters (Retired), Washington, DC 20546-0001, USA

¹⁰Department of Chemical and Environmental Engineering, University of Arizona, Tucson, AZ 85721, USA

¹¹Science Systems and Applications, Inc. at NASA Langley Research Center, Hampton, Virginia, 23681, USA

¹²Center for Aerosol Science and Engineering, Department of Energy, Environmental and Chemical Engineering, Washington University in St. Louis, St. Louis, MO 63130, USA

Correspondence: Jeffrey S. Reid (jeffrey.s.reid20.civ@us.navy.mil)

Received: 3 June 2025 – Discussion started: 12 August 2025

Revised: 30 October 2025 – Accepted: 12 November 2025 – Published: 19 December 2025

Abstract. While most large-scale smoke advection occurs within the free troposphere, Maritime Continent smoke is suspected to be unique in its long-range, near-surface transport. Such a pathway likely creates strong gradients and uncertainties in interpreting satellite and model data on light extinction, air pollution, and cloud condensation nuclei. This paper documents High Spectral Resolution Lidar (HSRL) data from the 2019 ONR PISTON cruise and NASA CAMP2Ex flights that revealed Maritime Continent smoke and pollution transport pathways and heterogeneity around the Marine Atmospheric Boundary Layer (MABL) over thousands of kilometers. Observations showed that 95 % of integrated aerosol backscatter occurred below 2500 m altitude. The R/V *Sally Ride* observed 50th and 84th percentile aerosol backscatter altitudes at ~ 600 and ~ 1500 m respectively, regardless of aerosol loading. Peak backscatter values occurred within or near the MABL top, diminishing as we approached 2–3 km altitude, but with occasional plumes reaching the melting level at 4800 m. At monsoonal scales, aerosol models largely account for the observed directional wind shear that causes altitude-dependent particle transport: near-surface particles remain in the core monsoon flow around the MABL, while at lower latitudes, aerosol layers aloft advect more eastwardly. Around the MABL, however, significant cloud-scale variability exists due to fine-scale flow, halo-entrainment-detrainment, and cold pool phenomena. Backscatter enhancements beneath individual clouds, extending to the ocean surface, likely relate to MABL-free troposphere exchange and air-sea interaction. So while aerosol transport occurs near the surface, particle extinction heterogeneity must still be considered for in situ observations and satellite retrievals.

1 Introduction

Every continent hosts biomass burning receptor regimes from an ever-increasing number of large scale smoke events. Long-range transport of smoke is often facilitated by free tropospheric advection from either direct injection (e.g. super fires in Australia, North America, and boreal Asia; Kahn et al., 2008; Fromm et al., 2010; Sofiev et al., 2012; Paugam et al., 2016; Xian et al., 2022; Eck et al., 2023) or from “land plume” formation when the upper portions of a deep and polluted Planetary Boundary Layer (PBL)/lower free troposphere are acted upon by large scale vertical wind shear, often times along shorelines or mountain ranges (e.g. Sinha et al., 2004; Reid et al., 2013; Wang et al., 2013). Once in the free troposphere, or stratosphere for volcanic or pyro convection events, aerosol plumes are impacted by synoptic meteorological features, such as frontal activity, large-scale subsidence/lifting, tropospheric folds, etc. and transported tens of thousands of kilometers before being scavenged. Plumes even frequently reach the Arctic and Antarctic regions (e.g. Hu et al., 2013; Xian et al., 2022). An excellent example of this is the Southern Africa Aerosol system, with well documented elevated plumes that can be transported across the Atlantic (e.g. Transport and Atmospheric Chemistry Near the Equator-Atlantic- TRACE-A, Anderson et al., 1996; Observations of Aerosols above Clouds and their Interactions-ORACLES, Mallet et al., 2019; Southern African Regional Science Initiative-SAFARI-2000, Swap et al., 2002; The Cloud–Aerosol–Radiation Interaction and Forcing: Year 2017-CLARIFY, Haywood et al., 2021; Layered Atlantic Smoke Interactions with Clouds-LASIC, Dobracki et al., 2025) all the way to the Antarctic (e.g. Khan et al., 2019). Although these events are easily tracked by satellite sensors and models through Aerosol Optical Thickness (AOT), there is difficult-to-observe variability in particle vertical distribution. In particular, lower-level transport and free troposphere-PBL exchange result in the presence of anthropogenic aerosol particles in the remote Marine Atmospheric Boundary Layers (MABL). For Africa, such work ranges from Andreae (1983) to more recently Dobracki et al. (2025). Indeed, it has long been recognized that terrestrially generated cloud condensation nuclei (CCN) can dominate maritime nuclei and light scattering over much of the globe (e.g. Clarke et al. 1996, 2013; Quinn et al., 2017; Ross et al., 2018; Reid et al. 2022).

The Maritime Continent (MC), made up of the islands of Indonesia, Malaysia, East Timor, and the Philippines, has a distinct biomass burning and pollution transport phenomenology from other source regions, such as aforementioned Africa, as well as Boreal or South America. Smoke from small agricultural waste burns and smoldering peat fires have lower injection heights than fires in other regions (Labonne et al., 2007; Wang et al., 2013) and long-range

transport events appear to predominately be in the MABL and lower free troposphere (Campbell et al., 2013; Chew et al., 2013; Reid et al. 2013, 2023). While other major plumes in the world are ventilated into the free troposphere for long range transport with MABL transport as a small relative fraction, the MC’s “long-range-near-surface-maritime” plumes are in high enough concentrations to be easily tracked by satellite for over 7000 km (Reid et al., 2013, 2022; Ross et al., 2018). This transport scenario allows for the observation of particle evolution and lifetime through their interaction with an entirely different set of meteorological phenomena than what is more commonly studied in association with long range free troposphere or stratosphere aerosol transport. In particular, MC aerosol transport is often embedded within a convectively active monsoonal system with numerous cloud regimes (e.g. Reid et al., 2013, 2023; Ge et al., 2017; Hilario et al., 2021).

Copious anthropogenic emissions and numerous lower intensity fires are associated with shallow plume injection heights. Orographic effects, off-island smoke transport by the land breeze, and vertical wind shear within the boreal summer’s Southwest Monsoon (SWM) system have also been hypothesized to allow plumes to reach thousands of kilometers in length with the bulk of AOT being contributed by particles below 3 km in altitude (Reid et al., 2013, 2015, 2016; Wang et al., 2013; Ge et al., 2017; Hilario et al. 2021). While major burning events of the MC can occur at any time of year with a sufficient dry spell (Reid et al., 2013), boreal summertime drought conditions such as those related to El Niño can lead to the draining and significant burning of peatlands well past the monsoon transition (e.g. Nichol 1998; Field and Shen, 2008; Reid et al., 2012, 2023; Goldstein et al. 2020). Such events result in the highest monthly average AOT values recorded by the Aerosol Robotic Network (AERONET; Eck et al., 2019; Reid et al., 2023). Similar to biomass burning plumes, MC megacity pollution can also be transported 1000 km or more (Cruz et al., 2019; Braun et al., 2020; Hilario et al., 2020; Lorenzo et al., 2023).

The particle transport from the MC, peninsular SE Asia, and southern China through the South China, Sulu, and Philippine Seas, to their eventual annihilation in the West Pacific monsoonal trough lends itself to being a natural laboratory of aerosol lifecycle and cloud-related process studies, including entrainment, detrainment, microphysics/chemistry, and scavenging. At the same time, the nature of regional particle diversity and evolution makes the MC’s monsoon system a venue to explore new remote sensing and modeling technologies. In 2019, the Office of Naval Research (ONR) Propagation of IntraSeasonal Oscillations (PISTON) Intensive Observation Period (IOP) on the R/V *Sally Ride* was conducted in partnership with the NASA Cloud, Aerosol and Monsoon Processes Philippines Experiment (CAMP²Ex) with its P-3 and Stratton Park Engineer-

ing Company (SPEC) Lear 35 aircraft in the vicinity of the Philippines during the Southwest Monsoon (SWM) to North-east Monsoon (NEM) transition (Reid et al., 2023). The combined resources of these two missions allowed for the first time detailed observations of MC, East Asia, and maritime aerosol transport characteristics, particle evolution, and cloud process outcomes. Included in the missions were three High Spectral Resolution Lidars (HSRL): (1) a CAMP²Ex-sponsored Langley Research Center (LaRC) HSRL-2 instrument on the NASA P3 aircraft; (2) a PISTON sponsored University of Wisconsin-Madison Space Sciences and Engineering Center HSRL (UW-HSRL) deployed to the *Sally Ride* in the Philippine Sea 600 km east-northeast of Manila; and (3) A second NASA CALIPSO sponsored UW-HSRL (henceforth MO-HSRL) deployed to the Manila Observatory, Metro Manila, Philippines. Combined, CAMP²Ex and PISTON allowed for the observation of volumetric variability of lower troposphere aerosol transport across the region (Hilario et al., 2021; Edwards et al., 2022; Reid et al., 2023).

It was clear during the CAMP²Ex and PISTON deployment that moisture, aerosol, and hence light extinction can exhibit strong variability on relatively short spatial and temporal scales. As noted in Su et al. (2008), Trickett and Di Girolamo (2009); Eck et al. (2014), Reid et al. (2019) and the review by Marshak et al. (2021), aerosol properties can have sharply different properties around clouds in association with so-called “Halos” (Radke and Hobbs, 1991). However, given the locality and speed of key evolving cloud properties coupled with sampling limitations from aircraft and ship, it is challenging to analyze specific physical processes. An aircraft can measure properties at a single point and time to high precision, but only a few hundred meters or seconds away the environment can be quite different. The ship can provide a very detailed bottom up-view, but waits for the atmosphere to pass over it. As the start of further investigation to understand how MABL and cloud heterogeneity are related to coupled aerosol-cloud, meteorological, and remote sensing impacts, this paper documents the nature of SWM and NEM aerosol particle layering and heterogeneity phenomena during 2019 PISTON/CAMP²Ex campaigns. This review of the different “Animals in the coupled aerosol-cloud-ocean zoo” is necessary before we can efficiently tease out the intertwined roles of individual processes that work together to create intricate patterns in light extinction in and around the MABL. This first analysis largely relies on NASA P-3 and PISTON R/V *Sally Ride* HSRL assets to identify types and ranges of scales of the maritime environment’s aerosol structures in preparation for more detailed follow-on studies.

In this survey, we make use of the strengths of different instrument datasets to paint a holistic picture of aerosol variability in the Northwest Tropical Pacific’s (NWTP) monsoon environment. We begin with an overview of available assets (Sect. 2), followed by a brief large-scale survey of aerosol vertical profiles from models and in situ measurements from the P-3 (Sect. 3). After this overview, we provide exam-

ple environments measured by the airborne LaRC HSRL-2 (Sect. 4), limited to the analysis of three core environments (NWTP marine, MC biomass burning, and Asian pollution). The lidar observations show significant complexity in aerosol features. We use these examples to lay the groundwork for a time-series analysis from the R/V *Sally Ride* (Sect. 5) with further examples highlighting specific lidar-diagnosed aerosol features. Finally, we summarize findings and discuss implications in our conclusions (Sect. 6).

2 Mission overview, observations, and this investigation design

A full description of the CAMP²Ex field mission and its relationship to PISTON is provided by Reid et al. (2023). A brief summary on relevant topics drawn from this paper is provided here. The NASA CAMP²Ex experiment was conducted out of Clark International Airport, Philippines with (a) 19 flights between 25 August–5 October 2019 of the NASA P-3 as the primary platform for remote sensing, state, and composition hosting instruments including the HSRL-2 (Hair et al., 2008, Burton et al., 2018); and (b) 11 SPEC Lear 35 flights from 7–29 September 2019 for sampling deep convection (not used here). P-3 research flight days as well airmass sources were published by Hilario et al. (2021) via the application of HYSPLIT systematically along flight trajectories. They identified a series of subcontinental sources, including used here the remote Western Pacific (WP); the Maritime Continent of Sumatra, the Malay Peninsulas, and Borneo (MC); East Asian Pollution (EA), largely originating from the Pearl River Delta; and indeterminate mixtures (MX). A synopsis of the MABL and Lower Free Troposphere (LFT) airmass origins are listed in Table 1. Within the CAMP²Ex study period, PISTON was conducted with the R/V *Sally Ride* on station ~ 600 km northeast of Manila 6–25 September 2019, transiting to and from Taiwan. Of interest to this study regarding *Sally Ride* data, was the zenith-pointing and scanning University of Wisconsin Space Science and Engineering Center (SSEC) HSRL (Eloranta 2005; Reid et al., 2017), SEAPOL C-band dual-polarization Doppler radar (Rutledge et al., 2019), and the NOAA PSL air-sea flux and near-surface meteorological and ocean time series (Fairall et al., 1996, 1997, 2003, Edson et al., 2013, Sobel et al., 2021). Finally, the NASA Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) mission funded the long-term deployment of another, zenith pointing, SSEC HSRL at the Manila Observatory between June 2019–September 2020. Discussion of these urban observations is forthcoming in works.

2.1 Model and satellite products

To provide context to observations we utilize satellite products generated by SSEC (outlined in Reid et al., 2023) and the Navy Aerosol Analysis and Prediction System Reanal-

Table 1. P3 flight summary including the local day for the flight (GMT is +8 h). Properties are taken from surface to > 5 km profiles over ocean. Included is (a) Marine Atmospheric Boundary Layer (MABL; 100–400 m) and Lower free troposphere (LFT 1200–1600 m). Approximate source regions as extracted from Hilario et al., (2021): BB-Biomass burning from Borneo and Sumatra; EA-East Asia. MC-Maritime Continent; WP-Western Pacific. MX-Mixture/Other. (b) Carbon monoxide values for the MABL and 1.4 km representing the lower free troposphere measured from Picarro Cavity Ring Down Spectrometer (c) Volume concentration of fine and coarse mode particles measured from the Laser Aerosol Spectrometer (LAS; $0.1 < d_p < 1 \mu\text{m}$ and $1 < d_p < 3 \mu\text{m}$). (d) Estimated sub-micron dry light extinction as the sum of TSI 3 λ nephelometer and Radiance Research Particle Soot/Absorption Photometer (PSAP).

RF	Local Day	MABL Source	Lower Free Trop Source	MABL:LFT CO (ppb)	MABL:LFT Fine LAS Vol $0.1 < d_p < 1 \mu\text{m}$ ($\mu\text{m}^3 \text{cm}^{-3}$)	MABL:LFT Coarse LAS Vol $1 < d_p < 3 \mu\text{m}$ ($\mu\text{m}^3 \text{cm}^{-3}$)	MABL:LFT Dry Sub micron Light Extinction (532 nm Mm $^{-1}$)
1	8/25/2019*	MC	WP	108 : 86	2.5 : 0.9	2.3 : 1.0	31 : 9
2	8/27/2019	MX	MX	91 : 85	2.0 : 0.7	2.1 : 0.6	22 : 9
3	8/30/2019*	MC	MC	110 : 96	3.0 : 1.1	1.7 : 0.6	31 : 31
4	8/31/2019*	MC	MC	102 : 88	2.5 : 1.7	1.7 : 1.7	28 : 21
5	9/4/2019	MC	MC	121 : 104	5.5 : 3.4	2.2 : 2.2	54 : 31
6	9/7/2019*	BB	BB	242 : 133	13.0 : 4.6	4.8 : 2.1	166 : 44
7 [#]	9/9/2019*	BB	WP	175 : 112	7.5 : 1.4	2.9 : 0.8	83 : 13
8	9/14/2019*	WP	MX	74 : 79	3.5 : 3.2	1.0 : 0.8	39 : 32
9	9/16/2019*	BB	BB	678 : 374	44.0 : 23	15.3 : 8.0	698 : 351
10	9/17/2019*	BB	WP	312 : 194	17.0 : 6.7	6.8 : 2.7	224 : 74
11	9/20/2019*	EA	EA	138 : 139	4.5 : 2.6	3.9 : 1.8	54 : 20
12	9/22/2019*	EA	EA	181 : 153	7.0 : 3.9	3.9 : 2.2	82 : 38
13 [#]	9/24/2019	EA	EA	166 : 154	5.0 : 3.1	2.9 : 2.1	54 : 32
14 [#]	9/25/2019*	EA	EA	149 : 143	4.0 : 2.2	2.5 : 1.2	36 : 21
15	9/28/2019	WP	WP	77 : 76	0.4 : 0.1	0.7 : 0.1	5 : 3
16	9/29/2019	EA	WP	80 : 73	1.5 : 0.3	1.5 : 0.4	18 : 3
17	10/2/2019*	EA	EA	122 : 177	5.0 : 8.4	2.0 : 2.1	56 : 101
18	10/4/2019*	MX	MX	161 : 100	9.0 : 2.6	1.6 : 0.7	91 : 24
19	10/5/2019	WP	WP	75 : 74	0.5 : 0.1	0.6 : 0.2	4 : 1

* Takeoff before 00:00 UTC on this day and hence the data filename is for the day before the local day. [#] Flight with R/V *Sally Ride* overpass.

ysis (NAAPS-RA; Lynch et al., 2016 and Edwards et al., 2022). P-3 and Lear 35 aircraft tracks as well as *Sally Ride* positions and SEAPOL C-band weather radar (Rutledge et al., 2019) plots overlaid on imagery and a variety of cloud and aerosol products used here were generated on the field mission instance of NASA Worldview hosted at the UW SSEC. Six-hourly NAAPS data of mass ratios, extinctions, and AOTs of smoke, sea salt, dust and combined anthropogenic/biogenic fine species (ABF), as well as associated Navy Global Environmental Model (NAVGEM; Hogan et al., 2014) meteorology (see Sect. 7).

2.2 P-3 payload

A review of the full complement of instruments on the NASA P-3 aircraft is provided in the Supplement of Reid et al. (2023). Data used here include atmospheric state (e.g. temperature, water vapor, wind, etc.) from the P-3's Vaisala dropsondes, as well as P-3 forward camera data. Mission-wide integrated plots of particle vertical profiles from the Laser Aerosol Spectrometer (LAS; Froyd et al., 2019, Moore et al., 2021) and High-Resolution Time-of-Flight Aerosol Mass

Spectrometer (HR-ToF-AMS; DeCarlo et al., 2008, Crosbie et al., 2022) in Sect. 3 to provide an overview of the vertical profiles of key constituents for marine, biomass burning, and Asian pollution flights.

The focus of airborne observations in this overview paper is on the HSRL-2 (Hair et al., 2008, Burton et al., 2012, 2015, 2018; Ferrare et al., 2023) to demonstrate specific vertically-resolved aerosol phenomena. A limitation of backscatter lidars is that the measurement is fundamentally ill constrained, requiring assumptions regarding the aerosol/cloud scattering characteristics to retrieve the particulate backscatter and extinction. The HSRL technique takes advantage of the molecular Doppler broadening due to molecular Brownian motion to separate the molecular return from particulate (aerosol/cloud) backscatter (Eloranta, 2005). The ratio of total to molecular backscatter leads to a calibrated particulate backscatter signal as a function of range, and the slope of molecular signal relates directly to the particulate extinction. Compared to other lidar types, the HSRL provides excellent daytime sensitivity due to its improved solar background rejection resulting in more tractable range dependent error propagation than more commonly used elastic

backscatter or Raman systems. As noted in Reid et al. (2017) and clearly demonstrated in this current work, while the vertical profile of extinction derived from the HSRL is a valuable product, the calibrated aerosol backscatter product from HSRL systems has greater vertical and temporal resolution, precision, and accuracy. Thus, for the topics examined here, the focus is more towards the application of aerosol backscatter products.

The LaRC HSRL is what is termed a $3\beta + 2\alpha + 3\delta$ system, providing calibrated aerosol backscatter and extinction (α) at 355 and 532 nm, and attenuated backscatter (β) and depolarization (δ) at 355, 532, and 1064 nm (i.e. including both aerosol and molecular signal). The HSRL-2 acquires profiles at a rate of 200 profiles per second. Profiles are accumulated to 100-profile averages in hardware, making the finest available temporal profile spacing of 0.5 s. To provide adequate signal-to-noise ratios, level 2 aerosol backscatter and depolarization products are further averaged over 10 s, and the native 1.25 m vertical bins are averaged to 30 m. At the P-3's flight speed of $\sim 150 \text{ m s}^{-1}$, the effective horizontal resolution is 1500 m. Extinction and lidar ratio estimates are also reported at 10 s/30 m but require a longer 60 s integration (7500 m on center-at flight speed) and with a 300 m vertical window. Other derived products include estimates of planetary boundary layer (PBL) heights, which use the vertically resolved profile measurements of aerosol backscatter, and then secondarily derived aerosol mixed layer heights during the daytime (Scarino et al., 2014).

2.3 R/V *Sally Ride* payload

The ONR PISTON campaign sent the R/V *Sally Ride* in cooperation with CAMP²Ex to study air-sea interaction and convection and their relationship to Madden Julian Oscillation's summertime manifestation, the Boreal Summer Intraseasonal Oscillation (BSISO; e.g. Jiang et al., 2004). The R/V *Sally Ride* was thus equipped with a complement of turbulent flux, ocean sensing, and radar systems (overviewed by Sobel et al., 2021 for the 2018 cruise; all but the underway CTD data were repeated in the 2019 cruise). Data were continuously collected during the cruise while it was in international waters. Of particular interest here is the deployment of the UW-HSRL configured for both zenith pointing and occasional scans from the surface (0° horizontal) up to 17° in the MBL. This HSRL is very similar in nature to the one used in the NASA SEAC⁴RS project described in Reid et al. (2017) with the addition of the horizontal scanning capability, and the analysis here will follow a similar methodology. This HSRL provides aerosol backscatter at 532 nm and elastic backscatter at 1064 nm. Because the MABL, its entrainment zone, and lower free troposphere are within the region where the UW-HSRL suffers from incomplete overlap of the transmitted beam and receiver field of view, reliable lidar ratios in the area of interest are laborious to obtain. The focus here is on aerosol backscatter, and when needed, extinction

estimated with the lidar ratios estimated from the HSRL-2. An ongoing study is underway on the HSRL's side scanning capabilities to develop more reliable lidar ratios in the MABL to mitigate the uncertainties of the near field (0–5 km) where the receiver is out of focus.

Mission-wide figures and statistics at times utilize a 300 s cloud screened product at 30 m vertical resolution. Considering advection for a typical 8 m s^{-1} wind speed, 30 and 300 s averaging is equivalent to 240 and 2400 m horizontal resolutions, respectively. This system was determined to have a precision in aerosol backscatter of better than 2×10^{-7} and $2 \times 10^{-8} (\text{m sr})^{-1}$ for these temporal averages respectively, more than an order of magnitude lower than Rayleigh backscatter (1.16×10^{-5} , 7.7×10^{-6} , and $4.4 \times 10^{-6} (\text{m sr})^{-1}$, at 1, 5, and 10 km). For specific cases, resolution of up to 5 s and 15 m are shown. In addition to the UW-HSRL, in this paper we make use of Vaisala radiosondes released every 3 h and mean meteorology measured by the NOAA PSL bow mast sensors.

2.4 Cross comparison between the P-3 and *Sally Ride* HSRLs

Because of how the UW-HSRL scan pattern was set and to maintain safety for the P-3 and *Sally Ride*, there were no direct overflights with contiguous data on the two platforms. The closest the P-3 and *Sally Ride* came to joint lidar observations for these cases was on the order of 3–5 km in the horizontal. A full discussion of the combined dataset is part of an ongoing analysis. However, here in the methods section it is worth briefly comparing nearby data when the P-3 flew near the *Sally Ride* to demonstrate consistency in aerosol backscatter values. The flight on 24 September observed a combination of maritime and Asian polluted air masses. Nevertheless, the comparison of HSRL-2 data from the P-3's "top down" and UW-HSRL *Sally Ride*'s "bottom up" points of view is quite enlightening regarding the nature of lower free troposphere aerosol variability with overall good agreement of the calibrated backscatter between the instruments.

On 24 September 2019 [Research Flight (RF) 13] the P-3 passed 4 km just to the south of the *Sally Ride*, in an area devoid of clouds (Fig. 1a). At the surface, winds were NE at 6 m s^{-1} , with a relative humidity (RH) of 74 %, and the air-sea temperature difference was slightly unstable at 0.5°C . These conditions are such that we do not expect MABL roll structures to form, and the near absence of clouds inhibits MABL free troposphere exchange (Lemone, 1973; Salesky et al., 2017). In every sense, this should be the most uniform of environments, and that is what is observed. In Fig. 1b, we compare P-3 HSRL-2 in the standard 10 s increments in small markers centered on the closest position to the *Sally Ride*. For the *Sally Ride*'s UW-HSRL, we chose 5 s samples around a 2 min window when the lidar was pointed vertically. Data for the two platforms match exceptionally well in the MABL ML at $\sim 2.3 (\text{M m sr})^{-1}$ at the surface and with




Figure 1. Overview of when the NASA P3 flew in proximity to the *Sally Ride* on 24 September at $\sim 01:15$ UTC. Included is (a) the nearest RGB image from Himiwari 8, with the *Sally Ride* location as a yellow circle and the P3 track in blue; (b) aerosol backscatter associated with the nearest approach of the P3 including within a 2 min window for selected samples when the SSEC HSRL is pointed vertically and within 3 nearest cloud free samples from the P3's HSRL-2.

the SSEC-HSRL particulate backscatter within the range of HSRL-2 values. Both HSRLs also show a typical increase in backscatter in the MABL presumed due to humidity, as well as good agreement in the height of the inversion at ~ 550 – 600 m. The strength of the backscatter at the top of the inversion varies more significantly due to strong non-linearities in hygroscopicity at high RH values (as discussed later in this paper), but the variability in both the P-3 and *Sally Ride* platforms overlap. Above the inversion is a lesser elevated layer between 800 – 1400 m. Here too there is some variability, but again the individual P-3 profiles and *Sally Ride* profiles overlap.

3 Results 1: regional aerosol properties and transport

Here we provide brief context to the regional environment explored by HSRL data in the following sections. An overview of the CAMP²Ex/PISTON mission meteorology and aerosol fields is provided in Reid et al. (2023) and its associated Supplement. An overview of transport patterns for individual CAMP²Ex flights is provided in Hilario et al. (2021), and more regional overviews of the vertical variability in aerosol, thermodynamic and wind parameters in the region can be found in Xian et al. (2013), Bukowski et al. (2017), and Reid et al. (2012, 2023). In short, the region is dominated in the boreal summer months by the southwesterly flow of the SWM, followed by a monsoon transition to the NEM in late September to early October. For the CAMP²Ex/PISTON study season, the monsoon transition happened abruptly between 20–22 September 2019, about two weeks earlier than the climatological median. The large-scale aerosol transport pattern is elucidated in Fig. 2 where




Figure 2. Overview of the CAMP²Ex/PISTON study area for the (a) Southwest and (b) Northeast Monsoon periods of study. Included is the base average Aerosol Optical Thickness (AOT) from the NAAPS reanalysis, with (black arrows) NAVGEM surface winds; and (magenta) 3000 m AGL winds. Overlaid are CAMP²Ex flight numbers. Included are the NAAPS reanalysis smoke wet deposition fluxes over the study region for these two periods. Included in (a) in dotted orange are the cross-section meridians shown in Fig. 3. Included on (c) are geographic locations referenced in this paper.

the average NAAPS-RA AOT is plotted with surface (black) and 700 hPa/ ~ 3 km (magenta) winds for SWM and NEM CAMP²Ex mission days, in (a) and (b) respectively. Also included are the P-3 flight numbers located in the primary areas of study. To map receptor locations for biomass burning emissions, corresponding wet deposition fields of the smoke species in NAAPS-RA are provided for the two monsoon periods in Fig. 2c and d along with geographic notations referenced throughout this paper.

Biomass burning and pollution emissions of the MC are dominated by Indonesia, Malaysia and Singapore. During the SWM, these emissions are carried north through the South

China and Sulu Seas. Smoke wet deposition is at a maximum along the western coast of the Philippine archipelago islands due to upstream convection induced by the islands. The Philippines then adds its own emissions to the airmass, as they are further transported and scavenged in the NWTP trough farther east. Smoke was observed from southern Japan to Guam after 4000+ km of transport. Notable in Fig. 2a is directional vertical wind shear around the Philippines, indicated by the difference in the surface and 700 hPa winds. Smoke near the surface is typically advected from Borneo and Sumatra towards Luzon, whereas air in the LFT has a stronger westerly component. As discussed in Reid et al. (2013) and Bukowski et al. (2017), air above the middle troposphere overall reverses direction to northeasterly. Because of directional wind shear, back trajectories for air above the MABL are more likely to originate from Peninsular Southeast Asia than Borneo and Sumatra.

As part of the transition to the NEM, the monsoonal trough orientates from southeast to northwest in the NWTP to a more zonal orientation through Indonesia into the Indian Ocean. An opposite flow pattern to the SWM is then established (Fig. 2b). Notable is that emissions from East Asia are transported along the coast to the southwest as well as some residual smoke in and around Mindanao. Biomass burning emissions from Indonesia, Malaysia and Singapore are advected into the Indian Ocean where they are scavenged in the wintertime monsoonal trough. In both monsoon periods, vertical directional wind shear influences the vertical distribution of aerosol particles in the region. As discussed in Reid et al. (2013) and shown specifically for CAMP²Ex in the trajectory analysis of Hilario et al. (2021), during the SWM the MABL related air masses more typically originate from Peninsular Southeast Asia such as from Malaysia, Thailand, Cambodia and southern Vietnam.

The impact of vertical wind shear on fine mode aerosol vertical distribution is provided in Fig. 3, that shows the 119.5 and 127.5° meridional cross section of NAAPS-RA simulated fine mode mass concentration and horizontal winds for periods examined in this paper. Rows (a) and (b) represent 7 and 17 September 2019, the two largest sampled biomass burning events, respectively; row (c) is a mixed marine case where the P-3 and *Sally Ride* operated together; row (d) is a clean marine case north of Luzon sampled by the P-3; and row (e) is the most significant Asian pollution case sampled. These events match the general flow and AOT pattern as shown in Fig. 2a and b with the meridional transects indicated as dashed orange lines, differing only in more westward or eastward axes of the plumes. Examining the cross-sections, we see what one would expect for a sheared environment. For the two biomass burning cases (a and b), smoke has a latitudinal dependence in altitude, with elevated LFT smoke around 5–12° N, followed by the MABL dominated transport north of 15°. These plume areas are clearly aligned with areas of wind speed and directional shear in the vertical. Comparing the western to the eastern side of the Philippines,

the move towards higher aerosol altitudes for lower latitudes is also visible. For the mixed (c), clean marine (d), and most polluted Asian (e) cases of 24, 27 September, and 2 October 2019, respectively, we also see consistently higher altitudes in the aerosol plumes to the south and east.

While Fig. 2 provides an overview of the mean pattern, daily flow pattern can be quite complicated (e.g. Fig. 3), modulated by monsoonal surges and breaks driven by such factors as convection patterns and tropical cyclone activity. Consequently, many aerosol environments are advected into the study area from around the region, with more variability around Luzon. As outlined in Hilario et al. (2021) and Reid et al. (2023), CAMP²Ex/PISTON was able to monitor MC burning and pollution, Asian pollution, and cleaner marine conditions. For example, off Borneo air was advected well within the southwesterly monsoonal flow, whereas near Luzon, Philippines, air masses originated from all around the region depending on the state/strength of the monsoonal flows. Based on the back-trajectory analysis of Hilario et al. (2021), we segregated P-3 profiles based on their MABL and LFT origin. The dominant aerosol sources for the MABL (~ 100 to 400 m) and the free troposphere just above the boundary layer clouds (between 1200–1600 m) are included in Table 1. The average P-3 profiles, excluding cloud sampling, are categorized as: (1) cleaner Western Pacific air advected from the east (WP) on RF8, 15, and 19; (2) two significant biomass burning outbreaks (BB) that were sampled on consecutive days on either side of the Philippines for RF6–7, 9–10; (3) other Maritime Continent and Southeast Asia outflow (MC) on RF1, 3–5; (4) East Asian sources, predominantly from China (EA) on RF11–14, 16, 17; and (5) mixed and/or indeterminate origins around the Philippines, including the Manila superplume and ship emissions – (Mixed and/or other, MX) on RF2, 18. Key parameters provided at the two levels for each flight are CO, submicron aerosol volume from the LAS, coarse mode geometric aerosol volume derived from the LAS, and dry light extinction at 532 nm derived from the 3λ nephelometer and PSAP. Average profiles of these and other key constituents are shown in Fig. 4, calculated as the mean of all offshore datapoints by flight by the indicated RF numbers, as the mean of flight means. Included are (a) fine and coarse mode volume concentrations as calculated by the LAS and APS; (b) non-volatile and volatile condensation nuclei concentration; (c) AMS organics and sulfate mass concentrations; and (d) dry and liquid water light extinction at 532 nm calculated from the nephelometers, PSAP, and hygroscopicity measurements. Also included is “excess” CO, where the lowest value of CO measured during the mission (70 ppb) is subtracted from the mean profiles. Mission-wide, flight averaged peak particle indicators are always below 1200 m and often in the lowest 400 m (i.e. in the MABL’s mixed layer). Although local exceptions exist as shown below in lidar data, nucleation processes were observed associated with free troposphere cloud detrainment, and associated primary particle scavenging and nucleation precursor




Figure 3. Height cross-sections of NAAPS-RA fine mode aerosol mass concentrations along the 120.5 and 127.5° meridians for cases explored in this paper. Overlaid is the NAVGEM wind vectors pointing towards the direction towards (i.e. wind up is towards N/southerly wind, towards right is East/westerly wind). Indicator lines for these meridians are shown in Fig. 1a. Values below $0.1 \mu\text{g m}^{-3}$ are white.

detrainment (e.g. Xiao et al., 2023 and similar to Reid et al., 2019). Specific regimes are discussed in the subsections below with the regions of flight concentration listed by flight numbers on Fig. 2.

3.1 Western Pacific

Western Pacific air was well sampled once in the SWM (RF8) and twice in the NEM (RF15, 19). Back trajectories from Hilario et al. (2021) suggested that for these cases air masses sampled by the P-3 spent more than 7 d over the ocean originating east of the Philippines. If we consider the Western Pacific as the region’s “background environment”, it is unsurprising particle volumes/masses/extinctions are overwhelmingly below 2 km in altitude, with more significant coarse mode contributions from sea spray as one approaches the MABL (mixed layer inversion was typically $\sim 450\text{--}640\text{ m}$; see Sects. 4 and 5). CO was quite low, ranging from 74–77 ppb for all three flights. Given the aircraft

inlet’s aerodynamic cut point of $5.0 \mu\text{m}$ aerodynamic diameter, these coarse particle volumes are certainly underestimated. A slight fine mode aerosol particle enhancement is found just above the typical MABL inversion height of 450–550 m in the fine mode LAS and AMS sulfate and some organics (sea salt species are not measured). This peak coincides with a modest CO enhancement. At the same time, CN concentrations do not peak as fine particles, with particles with nonvolatile cores also making an increasing fraction of counts towards the surface, suggesting secondary production processes dominating particle sources in the middle free troposphere as discussed in detail by Xiao et al. (2023). All of these findings are suggestive of extraordinarily long-lived aerosol species or secondary production originating from distant anthropogenic sources being embedded in the “background environment” (e.g. see discussion below and Clarke et al., 2013). Fine mode light extinction values are also low, and are 40 % contributed by hygroscopic water uptake—again expected for high relative humidity found in




Figure 4. Mission average profiles of key aerosol parameters based on their primary marine boundary layer source. Included are columns (left to right) more pristine marine sources from the Western Pacific; two major biomass burning outbreaks; Sources from the Maritime Continent and Southeast Asia; East Asia; and complex mixtures of local and transported pollution. Parameters include (top to bottom), row (a) estimated dry geometric volumes for sub and super microns sampled through the P3s inlet; (b) Nonvolatile core and completely volatile CN concentration for particles > 10 nm; (c) AMS inferred submicron organics and sulfate; (d) Fine mode light extinction at 532 nm for dry particles and inferred water derived from the PSAP and dry and ambient nephelometer; (e) Excess CO above 70 ppb.

the MABL ($\sim 70\%$ – 80% RH at the surface, saturation at the MABL Mixed Layer top). Some indications of occasionally observed and highly scavenged detrainment layers are also visible in the free troposphere (e.g. Xiao et al., 2023; Hilario et al., 2025 Fig. S10).

3.2 Biomass burning

In contrast to the *near* pristine Western Pacific air masses, CAMP²Ex sampled two exceptional Borneo smoke outflow events, each on consecutive days on either side of the Philippines (RF6–7; 9–10). The 2019 burning season was one of the largest on record, with AOTs seasonally nearing and at time exceeding the mammoth 1998 and 2015 El Niño induced events (Eck et al. 2019; Reid et al., 2023 in the Supplement). Such AOTs have not been observed since 2019 at the writing of this paper. Yet, the 2019 season was marked by neutral but warming El Niño/Southern Oscillation (ESNO) conditions followed by a modern record Indian Ocean Dipole (IOP). The season nevertheless followed the emissions and transport that are typical of previous major events, including strong September through October emissions. Examining Fig. 4, we see as previously noted all major indicators of plume transport (Particle size, AMS, CN, light extinction, CO) placing smoke near the surface with indications of lower free troposphere plumes. Unsurprisingly, smoke was dominated by organic species in the AMS with mass and particle volume overwhelmingly below 1600 m for all cases, from off of Borneo to the east of the Philippines. In comparison, MABL mixed layer heights were on the order of 500 m. On one flight, RF9 (16 September 2019 local day), an isolated entrainment layer was visible at 1800–2200 m, a case shown in more detail in Sect. 4.

Also notable is that while these flights monitored exceptional levels of fine mode smoke particles, the super-micron particle volumes were non-negligible. We hypothesize these coarse mode particles come from two sources. First, some coarse mode particles are a natural part of the burning emissions, including dust and ash (Reid et al., 2005). Second, major biomass burning outbreaks are typically part of a monsoon enhancement that exhibit stronger near surface wind speeds and may generate sea salt particles through white-capping.

One key area of continued investigation by the science team is smoke particle hygroscopicity. In cases of heavy smoke, fine mode particles were measured to have light scattering hygroscopic growth factors less than one (i.e. $f(80) < 1$). That is, hydrated particles scatter less than their dry counterparts. This is not an uncommon observation, thought to be a result of possibly chain aggregates collapsing, volatilization effects, and refractive index modification while hydrating (Shingler et al., 2016; Lorenza et al., 2025). However, as shown in the lidar profiles, overall particles are clearly hygroscopic. There has been much internal discussion as to what such hygroscopicity measurements mean and how they com-

pare to bulk lidar observations (which include total fine and coarse, including any monsoon generated sea salt) as well as the potential for measurement artifacts. While reconciliation of these observations is outside the scope of this paper, this topic is discussed further in Sects. 4 and 5.

3.3 Other maritime continent observations

While major biomass burning events were observed toward the end of the SWM, the P-3 flew multiple cases off Borneo when fire prevalence was low (RF1, 3, 4, 5). These cases are likely a mixture of some burning, and also regional anthropogenic emissions (domestic fuel, industry, petrochemical, ship emissions, etc.; e.g. Hilario et al., 2020). For these cases from the MC, we see a similar pattern of low-level transport as with biomass burning with much lower and more distributed particle concentrations to ~ 3000 m. The sulfate fraction is significantly enhanced over biomass burning with slight enhancement in hygroscopicity, demonstrating a mixture of biomass burning and industrial contributions. Further, the coarse mode volume is a larger fraction of particles overall, although the absolute magnitude of coarse mode for the MC cases is half of what was present in the biomass burning cases. Also notable is that for the MC and biomass burning overall, there are elevated CO values for altitudes above where particle concentrations fall off. This can be reasonably explained as a result of long-range transport from the rest of SE and South Asia (e.g. Reid et al. 2015) and detrainment of aerosol scavenged air.

3.4 East Asia

In comparison to biomass burning and other Maritime Continent emissions, the East Asian sourced flights showed the first indications of a significant land plume transport phenomenology. East Asian sources were sampled on RF11, 12, 13, 14, 16, 17 around Luzon and they showed higher particle concentrations above the mixed layer indicative of land plume formation. These flights are the only ones from land sources where sulfate dominates fine mode aerosol particle masses over organics. As one would then expect, higher hygroscopicity was exhibited from other land sources as well. Also notable are CO and volatile CN enhancements in the lower free troposphere.

3.5 Mixed environments

Finally, two flights can be categorized as mixed environments (RF2 and 18), that were an amalgam of local Philippine sources, including the Manila Super Plume and well-aged Asian and Maritime Continent sources including ship emissions. These mixed cases were just that, variable in chemistry and vertical profile, although nevertheless dominated by particles < 3 km in altitude but with high CN volatility fraction and an enhancement in CN concentrations. Notable for both

cases are the enhancements in CN concentrations in or just above the MBL as well as higher CN concentrations aloft. For RF2, this is in part due to observed ship emissions in the area. For RF18, urban emissions in the presence of enhanced radiation led to explosive nucleation (e.g. Reid et al., 2016).

4 Results 2: example regional aerosol environment as monitored from the P-3 HSRL-2

Of the 19 CAMP²Ex research flights, several stand out as having easily describable maritime environments that demonstrate commonly observed aerosol features. Here we examine (a) the background marine cases of the Western Pacific (RF15 and 19); (b) the biomass burning smoke outbreaks (RF6, 9, and 10); and (c) East Asian pollution (RF17). Excluded in these cases is an evaluation of the many multi-spectral lidar products, as well as more complex environments, mixed maritime continent outflow, and local pollution of Philippines; these are all topics that are quite nuanced and require separate, more in-depth investigations. For this section, we extracted cases with good P-3 HSRL-2 viewing conditions to the surface from altitude and successful 532 nm light extinction in the vicinity of dropsondes and in situ profile data that typified the nature of aerosol layering phenomenon (Table 2). Even so, because of the high frequency variability shown here and in Sect. 5, it is quite difficult to perform true closure between the lidar profiles, sondes and in situ profiles. Nevertheless, these cases represent more idealized conditions of different classes of aerosol environments. Provided in Fig. 5 are 30 s average aerosol backscatter clean marine and polluted cases, in (a) and (b) respectively (the average of three consecutive 10 s profiles from the standard LaRC HSRL-2 product). Corresponding light extinction and lidar ratio products for the same 30 s product window are shown in (c, d) and (e, f), respectively. It should be remembered that lidar extinction and lidar ratio in the HSRL-2 archived data products are computed on a 60 s running average reported on a 10 s grid, making a 30 s product average shown here to have an effective smoothing of 90 s (~ 114 km horizontal center-weighted average assuming the P-3 flying at 300 kts). As the clean marine conditions are near the noise threshold for extinction and lidar ratio, lidar products in Fig. 5 have been further boxcar smoothed to 90 m in the vertical, making a 360 m center-weighted product. Finally, the relative humidity fields from nearby dropsondes are provided in (g) and (h).

4.1 NWTP background marine

We begin the analysis of P-3 observations of “clean background marine” environments that often come to mind as being representative of the Western Pacific. As this is the first example shown in Sect. 4, we go into more detail here than subsequent environments. As noted in Clarke et al. (1996, 2003, 2013), Reid et al. (2022), and Sect. 3.1 above, the




Figure 5. Example LaRC HSRL-2 profiles at 532 nm of common phenomena for clean marine (a, c, e) and polluted (b, d, f) marine conditions. Included are 30 s (~ 4.5 km) aerosol backscatter, and extracted 1.5 min (~ 14 km) aerosol extinction and lidar ratio with additional 90 m boxcar smoothing. Also included are nearby dropsonde releases (g, h).

idea of “background” or “pristine” marine can be problematic as there is often some terrestrial influence on aerosol populations in the remotest of oceans. Indeed, enhanced CO₂ as an indicator of anthropogenic influence, was found just above the MABL (Fig. 4). Nevertheless, after reviewing flight data, two flights immediately stand out as being closest to nominal “background marine” conditions observed during CAMP²Ex: RF15 (28 September 2019 local date) and 19 (5 October 2019 local date). RF8 is excluded as it has some orographic influence of small islands along the aircraft track. RF15 and 19 were conducted post monsoon transition and these flights experienced excellent fair-weather subtrop-

Table 2. Summary properties of representative marine environments studied in detail here using the NASA P3, including focus area latitude and longitude, winds at key levels, Marine Boundary Mixed Layer Inversion Height (MBLH) from aircraft soundings and the HSRL-2 product, 532 nm Aerosol Optical Thickness from the HSRL, and the fraction of that AOT within the Marine Boundary Layer's Mixed Layer. Sounding Mixed layer heights provide a maximum/minimum range in meters based on local variability in relative humidity and potential temperature during the airborne profile. NA stands for not available.

	Lat/Long (° N/E)	200 m Wind (ms ⁻¹ @°)	1500 m Wind (ms ⁻¹ @°)	3000 m Wind (ms ⁻¹ @°)	Sounding MBLH (m)	HSRL-2 MBLH (m)	HSRL-2 532 AOT	Fraction AOT < MLH
Clean Marine								
RF15 #1 9/27 22:06 UTC	20.85/122.75	7.5@004	7.6@025	1.5@061	460/400	450	0.09	0.38
RF15 #2 9/27 22:22 UTC	20.55/123.75	4.6@042	5.3@057	2.5@046	675/400	450	0.06	0.44
RF15 #3 9/28 00:48 UTC	20.60/125.50	6.0@099	NA	NA	675/340	475	0.03*	0.35
Biomass Burning								
RF06 #1 9/7 03:34 UTC	16.75/119.75	12.5@231	12.1@240	11.9@216	560/350	700	0.30	0.40
RF06 #2 9/7 06:22 UTC	12.50/120.5	10.7@208	11.4@244	9.1@237	550/550	350	0.31	0.28
RF09 #1 9/16 02:59 UTC	8.65/118.70	8.7@208	12.5@245	13.0@251	625/625	630	1.00	0.37
RF10 #1 9/17 00:05 UTC	13.80/125.45	11.12@217	17.5@253	13.8@256	575/575	645	0.59	0.55
Asia Pollution								
RF17 #1 10/2 02:26 UTC	19.5/120.50	0.2@256	0.8@132	3.2@229	340/340	450	0.37	0.24

ical conditions while capturing key MABL and cloud detraining phenomena. Visible images with flight tracks and 925 hPa NAVGEM winds are provided in Fig. 6 (a; Terra MODIS and (b; SNPP VIIRS) for RF15 and RF19 cases, respectively. Subsequent panels provide P-3 forward camera images and HSRL-2 aerosol backscatter curtains. It is clear from the Fig. 5 images that for “fair weather marine” conditions there are numerous macroscopic MABL cloud organization or “textures” for similarly low to moderate wind conditions. Back trajectories from Hilario et al. (2021) for these flights reveal airmasses originating from the Western Pacific. Table 2 has estimated AOT at 532 nm (t_{532}) from the HSRL-2 processing algorithm ranging from ~ 0.03 to 0.09. These are within one standard deviation of what sun photometers observe in typical remote ocean (e.g. Reid et al., 2022). MABL inversion height estimates varied in the 450–675 m range.

RF15 (28 September, 2019 local day) represented a period of easterly MABL winds ahead of Typhoon Mitag, which was propagating into the northeast quadrant of our operations area (Fig. 6a). The P-3 sampled an array of macroscopic cumulus cloud organizational structures of cloud properties before sunrise to mid-morning. The flight started with a pre-dawn Aeolus satellite wind lidar underpass after which the P-3 sampled two distinct regimes: (1) Between points 1 and 2, across an area of some crosswind oriented weak cloud development; and (2) A similar length line west of Point 3 in an area of scattered MABL clouds, largely subpixel in the MODIS imagery. A third area of investigation by the P-3 to the north was similar to the second and thus not dealt with here. In the two regimes shown in Fig. 6a, MABL winds were largely northeasterly to easterly, with cloud fea-

tures and macroscopic textures aligned north-south/cross-wind. NAVGEM and dropsondes suggest convergence across the convective line, with winds $\sim 7.5 \text{ ms}^{-1}$ @25° on the eastern side below 850 hPa, 5 ms^{-1} @50° on the western side below 850 hPa, and diminishing winds at and beyond 700 hPa. Examining geostationary data, this cloud band was shed from TC Mitag 24 h earlier while it was in a tropical depression stage. Moisture profiles (Fig. 5g) show the typical increase in RH in the ML from the surface to the inversion base at ~ 400 m, followed by a moist layer to 1700 m, and then some drying aloft (as is often observed in tropical maritime regions). For the segment between points 1 and 2, the P-3 cut across a feature of *cumulus mediocris* (Cu med) with cloud top heights to 2–3 km and thin detraining *altocumulus* (Ac) clouds, associated aerosol layers, and outflow remnants on the western side between 500 and 1500–1800 m (Fig. 6 a marker 1; Fig. 6c and f). As noted in Table 2, the western “downwind” side of this convective area had HSRL-2 derived 532 nm AOTs (t_{532}) of 0.09 vs. 0.06 on the eastern side. The western side AOT and backscatter enhancement is consistent with detraining, although long range continental transport from Asia cannot be completely ruled out. Nevertheless, small areas of enhanced particulate backscatter (i.e. so called “halos”; Radke and Hobbs, 1991) are occasionally visible around clouds on the eastern side of the convective line.

The second RF15 domain around Point 3 had appreciably less convective development than RF15a. The region hosted small fields of *Cumulus Humilis* (Cu hum) with limited development, largely 100–200 m deep, with one cloud top height observed reaching 1 km (Fig. 6d and g). And yet,




Figure 6. HSRL-2 aerosol backscatter profiles for two research flights focusing on background marine conditions (RF15, takeoff 27 September 2019; RF19 takeoff 5 October 2019). (a) and (b), Terra MODIS RGB with P3 flight tracks and curtain segments labeled; (c–e) selected forward camera at the start of lidar segments used here; (f–i) HSRL-2 calibrated aerosol backscatter with cloud screening for flight segments marked on (a) and (b) – all with colorbar noted in (g) and (j) 2 Hz zoom of the box in (i) with its own linear color bar. Blacked out areas of lidar profiles are due to the presence of opaque clouds.

dropsonde RH fields show only a slightly drier environment just above the MABL than in the convergence zone between points 1 and 2. This air mass originated from Mitag just behind the convective line sampled between Points 1 and 2. Consequently, the derived τ_{532} product generated as part of the standard LaRC processing was low and matched Point 2 with a value of 0.06. Even though convective development was less than the other domain, about 1/3 of light extinction and aerosol backscatter was below cloud base at approximately 425 m, with limited detrainment to 1 km. What is highly notable, however, is that one plume was observed between 800–1400 m at the ~ 100 km marker. This is likely a cloud halo/residual from an evaporating cloud.

The second example day of a Western Pacific marine environment, RF19, differentiates itself from RF15 in the sampled region's cloud organization and development associated with flow around a mesoscale convective system (MCS; Fig. 6b and e). Like RF15, near surface winds were roughly in the $6\text{--}8\text{ ms}^{-1}$ range, with areas of *Cu Hu* and *Cu Med* to 3 km in height with prevalent Ac (Fig. 6e, h, and i). However, for these cases the macroscopic cloud pattern is along wind instead of crosswind, an indicator of MABL roll structures (e.g. Moeng and Sullivan 1994; Salesky et al., 2017; Park et al., 2022). Moisture profiles were slightly drier in the ML than the RF15 case, especially for the second profile. RF19 sampled in both the cross and along wind directions (versus only cross wind in for RF15). t_{532} was also similar to RF15 ($\sim 0.06\text{--}0.09$; Table 2) with lidar ratios at ~ 30 sr in the mixed layer, and increasing with height (Fig. 5).

These cases provide us with some spatial perspectives of clean marine conditions. Both flights sampled areas low in AOT. While the HSRL-2 measurements of AOT have compared very well with Sun photometer AOT measurements, the instrument is near the signal-to-noise extinction derivation capabilities for the airborne HSRL-2 (Sawamura et al., 2017). Indeed, examining 30 s average aerosol backscatter profiles (Fig. 5a), we easily observe well-known characteristics of a boundary layer. Backscatter increases in the mixed layer due to adiabatic cooling and likewise increases in RH, as demonstrated in the associated dropsondes. When backscatter reaches a maximum coincident with potential temperature at the top of the mixed layer, roughly at cloud base, backscatter then also asymptotically falls with altitude with punctuations of thin aerosol layers and Ac clouds. However, light extinction and lidar ratio (Fig. 5c and e) are difficult to interpret, even with the additional averaging imposed.

Despite the noise in the derived extinction, we can still estimate the nature of ambient light extinction through the aerosol backscatter product. Within the MABL, backscatter from these background days can vary by a factor of two. With the exception of RF15 Point 1 (e.g. ahead of the weak convective line), backscatter values are all fairly similar above the MABL. While aerosol backscatter is dominated by the MABL, as is aerosol mass as shown Fig. 4, the fainter aerosol backscatter in the free troposphere integrates spatially to appreciable values. Indeed, overall, 25 %–45 % of the estimated extinction based on an assumed constant lidar ratio for these cases is within the MABL Mixed Layer (Table 2). If we assume the increasing lidar ratio with altitude is also due to the increase in RH near the top of the MBL (e.g. Fig. 5e), then this fraction diminishes further—although establishing the veracity of the derived lidar ratio under these clean conditions requires a study in and of its own. Nevertheless, the role of spatially extensive but low extinction aerosol environments has been problematic to the lidar community for some time. Noteworthy is the exclusion of such environments in CALIOP retrievals with a subsequent retrieval bias (e.g. Toth et al., 2018).

Figure 6 also shows that overall the height of pronounced aerosol backscatter is often, but not always, related to the height of local cloud tops. This is clearly visible in both RF15 cases, as well as RF19 area 2. Indeed, even with cloud tops to 3 km in RF15b, a clear delineation in aerosol backscatter is visible. And yet, for RF19 area 1 the opposite is true, as there is a minimum in aerosol backscatter between 2–3 km. Such collocated enhancements or dearth of aerosol particles may well be related to changing natures of cloud to detrainment for precipitating vs. non-precipitating cells—either through scavenging or through differences in the detrainment processes themselves.

Fine detrainment layer structure and cloud halos are also clearly visible in the individual profiles (Fig. 5) and curtain plots (Fig. 6f–i). Thin aerosol layers may be produced by nearby cumulus cloud development, or generated by deep convection and differential advection further afield. The nature of detrained aerosol layers in and around cumulus clouds is quite apparent with the HSRL-2 curtains in Fig. 6f–i. The deeper Cu fields for RF15 Area 1 have more visible detrainment layers aloft, while cloud fields with limited vertical development (e.g. RF15 Area 3) do not. Nevertheless one isolated cloud residual is visible at 1 km in altitude. However, this is not a universal observation, as deeper convection was present in RF19a to the same heights as RF15 Area 1, but did not show as strong detrainment features. One hypothesis explaining this difference is related to the forcing of the cloud fields. The cross-flow cloud feature in RF15a may be forced by slight near-surface convergence, leading to net detrainment aloft. Whereas for the RF19a case, cloud streets are a result of shear and differences in buoyancy (Lemone, 1973; Moeng and Sullivan, 1994). And yet, for RF19 Area 2, Ac were prevalent in that RF19 area, as were more visible detrainment layers. One notable difference between RF19a and b is sampling. The RF19 Area 1 sampled more cross wind, whereas RF19 area 2 was more along wind. As this has implications for MABL sampling (e.g. Park et al., 2022), it is probably fair to hypothesize along wind/cross wind sampling bias plays a role in MABL cloud detrainment studies as well. Other factors potentially impacting detrainment, and hence the formation of Ac and associated aerosol layers, include environmental static stability and relative humidity, the vertical structure of cloud buoyancy, in-cloud vertical pressure gradients, and evaporation along cloud edges (e.g. Moser and Lasher-Trapp 2017; Savre 2022; Morrison 2025), and should be further explored.

Finally, in addition to MABL cloud detrainment variability, there is also significant vertical structure in the mixing layer itself, shown in figures such as Fig. 5. As noted above, we expect an increase in aerosol backscatter with height in the mixed layer due to simple adiabatic cooling, leading to increases in RH, and hence hygroscopic growth. However, updrafts under clouds with subsequent in-between drier downdrafts (e.g. Reid et al., 2017) and associated MABL roll structures (Park et al., 2022) also result in alternating higher

and lower scattering columns. Examining the 10 s data in Fig. 6g and i in particular, we can see horizontal oscillation in backscatter in the lowest 500 m. Given the P-3 was flying at 170 m s^{-1} , a 10 s sample is 1.7 km, not too far off the common 2.5–3 km distance between the roll structures comprising cloud streets. If we plot backscatter at the instrument native resolution of 0.5 s (noting that 10 s is still considered the full independent data rate of the instrument, Fig. 6j), the enhanced backscatter column under clouds becomes much clearer. Oscillating areas of enhanced backscatter of about a factor of 2 extend around the cloud to the surface. Although the independent signal sampling period is 10 s, the symmetry of these features around the clouds suggests they are physical and not an electronic signal problem. We hypothesize that these features are related to updrafts and the nonlinear nature of a hygroscopic growth curve and vertical exchange of air under and between clouds (e.g. as in the terrestrial PBL described in Kunkel et al., 1977 and Reid et al., 2017). Such variability is examined in much more detail in Sect. 5.

4.2 Biomass burning outbreaks

In this subsection, we move from the very cleanest environments sampled by the P-3 during CAMP²Ex to the most polluted. The P-3 sampled the 2019 season's two major SWM smoke outbreaks, both related to monsoonal enhancements associated with tropical cyclone development in the NWTP—a common source of monsoonal perturbation (Reid et al., 2012). The first enhancement was driven by the passage of Typhoon Lingling from the east of the Philippines up through north of Taiwan and into North Korea. Smoke from Central Kalimantan, Borneo was sampled just to the west of Luzon on RF6 after $\sim 1600 \text{ km}$ of transport (Fig. 7a local day 7 September 2023) within a well-defined wind enhancement of 12 m s^{-1} . t_{532} from the HSRL-2 was on the order of ~ 0.3 (Table 2). This plume was transported in active warm convection with cloud tops at 3–5 km transitioning to deep convection due to orographic enhancement along the Luzon coast. A second smoke outbreak occurred 9 d later by the influence of multiple tropical disturbances that eventually formed into Typhoon Tapah. This outbreak was sampled on subsequent days: RF9 sampling 150 km north of Borneo where t_{532} were measured > 1 after $\sim 1000 \text{ km}$ of transport from Central Kalimantan (Fig. 7b local day 16 September 2023); and RF10 (Fig. 7c local day 17 September 2023) sampling just east of Luzon with t_{532} measured at 0.6 after an additional 1000 km of transport. While the RF9 and 10 were roughly one day apart in transport given the MBL winds at 10 m s^{-1} , we are reluctant to call these two flights a true Lagrangian pair. Nevertheless, some consistencies and differences are indicative of evolution processes.

In contrast to the background marine conditions, biomass burning outbreaks sampled during CAMP²Ex were well above signal to noise limitations for the HSRL-2 for extinction measurements. Figure 5b and d show extinction profiles




Figure 7. Same as Fig. 4 but for HSRL-2 aerosol backscatter profiles for three research flights focusing on major biomass burning smoke outbreaks (RF6, takeoff 6 September 2019; RF9 takeoff 15 September 2019, 2019; and RF10 takeoff 16 September 2019). (a–c) Terra MODIS RGB with P3 flight tracks and curtain segments labelled; (d–f) selected forward camera at the start of lidar segments used here. (g–j) HSRL-2 calibrated aerosol backscatter with cloud screening for flight segments marked in (a–c). Marked in red arrows on (g) and (h) is a contrast in notable differences in the aerosol vertical distribution inside and outside the convective region. Likewise arrows in (i) and (j) notable differences between boundary layer profiles sampled just off of Borneo and after 1000 km of transport to the north east demonstrating how shear results in separating the plume into a MABL and lower free troposphere plume. Finally, in (k) is a zoomed 0.5 s sample of the purple box in (j) showing individual role feature induced backscatter enhancements slanted due to the presence of wind shear.

that largely match aerosol backscatter structure quite well, although smoothed due to its 300 m native sampling window. Lidar ratios are also more stable than marine counterparts (Fig. 4f) and vertical variability can be resolved—albeit nevertheless with some potential noise. In all but one case (RF6 #2, 06:35 UTC; see below), we find the typical aerosol backscatter profile peaking at the top of the 450–650 m high mixed layer. Also embedded in the mean profiles are many geomet-

rically thin, but high backscatter aerosol layers that can persist for 10–100 s of kilometers as well as what appears to be other detrainment events and residual cloud halos. Frequent oscillations at the top of the smoke layers observed in RF9 are indicative of gravity waves (Fig. 7i).

Examining these three flights together, there are notable similarities and a few key differences between lidar backscatter profiles shown for the RF6 and RF9–10 cases. In all cases, over 98 % of measured aerosol backscatter was at altitudes below 3 km (Fig. 5) and subjected to low-level transport. These cases also demonstrate the nature of cloud processing and detrainment of aerosol particles in the presence of vertical wind shear. The RF6 profiles have the same t_{532} at ~ 0.3 , but exhibit vastly different profile distributions. RF6 #1 being MABL dominated, and RF6 #2 showing one of the few cases having a very low fraction of extinction in the mixed layer (28 %). These profiles for RF6 and 7 were taken within and just outside the area of convection associated with the monsoon enhancement (Fig. 7a). Track RF6 #1 with its MABL dominated smoke transport (note red arrow) was collected in a cloud free slot in between convective lines with prevalent *Ac* and noticeable detrainment to 3 km (Fig. 7d and g). Even near the surface, there is considerable variability in profiles and layer strengths. In contrast, track RF6 b was on the south side of the monsoon enhancement cutting northwards into convection. In Fig. 7h, we can clearly see a shift from an aerosol environment dominated by smoke above the MABL on the southeast side of the main convective area (note red arrow, distance marker 0–10 km) to profiles within the convection (distance of 65 km) that are more indicative of MABL transport as was seen in RF6 a. We initially hypothesized that the smoke layer at 1200 m may be a result of detrainment and shear from the eastern side of monsoonal enhancement transporting the smoke. However, this shear hypothesis is not supported by the available dropsonde data. Alternatively, this enhancement at RF6 #2 may not be an enhancement at all, rather, smoke may be efficiently scavenged in the convective area, and thus the lower free troposphere is left with many optically thin detrainment layers. Or, as discussed in Sect. 4.4 may be a result of large scale cold pool lifting.

The pair of flights on RF9 and 10 are notable in that unlike RF6, they were not embedded in a region of highly active convection, but rather in more scattered congestus and deep convection. Long transects at altitude on RF9 allowed for good HSRL-2 viewing conditions to fully sample the environment. Significant variability in aerosol profiles is visible in the lowest kilometer, with multiple aerosol layers visible that are 30–100 km in length but only a few hundred meters in depth. While the smoke layer is largely capped at 3 km, very thin aerosol layers were visible up to 4.5 km, the approximate 0 °C level. After an additional 1000 km of transport, the center of the plume was observed again the next day during RF10. While some detrainment is visible to 3 km, the majority of extinction is within the mixed layer. This is perhaps the mission's best example of how a change in vertical

profile can be a result of slight changes in wind shear. Winds from the surface were 13 ms^{-1} from the southwest, while winds aloft increased to 15 ms^{-1} west-southeast with smoke being preferentially advected more eastward.

Like the marine environment, biomass burning smoke cases all exhibited systematically increasing lidar ratios with altitude (Fig. 5f). This results in noticeable shifts in light extinction weighting upwards relative to aerosol backscatter. From a lidar data interpretation point of view, these changes can have significant implications for estimating aerosol budget and transport. In the case of RF9 #1 00:13 UTC, this increase in lidar ratio shifts the maximum extinction from the top of the mixed layer to just above. Thin aerosol layer enhancements above MBL are also enhanced by the growth of lidar ratio with height. A second area of interest is notable differences in the lidar ratios for the two monsoon enhancement events. Typically, smoke has higher lidar ratios $\sim 60 \text{ sr}$ or above (Burton et al., 2012), as was seen for the RF9 and 10 flights of local day 16–17 September 2019. However, for the first monsoon enhancement of 7 September (RF6) case, lidar ratios are a third to one-half of that. Surface winds for RF6 were $\sim 12 \text{ ms}^{-1}$ and over a long maritime fetch; perhaps lower lidar ratio of 30–40 sr are a result of mixing with coarse mode sea salt. Yet this explanation is not entirely satisfactory, as the RF9 and 10 pair exhibited high winds too and the ratio of fine to coarse mode volume in the MABL from the LAS is also the same at 0.36 (Table 1). A second possibility is the lower lidar ratios for RF6 are simply a result of the lower overall AOTs for that event, and hence a greater coarse/giant mode fraction that is not sampled in the P-3's inlets. Lastly, we pose that perhaps these lower lidar ratios in part have to do with differences in the nature of cloud processing of smoke for the air mass sampled in RF6. Indeed, this region was quite active with warm multi-thermal convection (see analysis in Reid et al., 2023). Nevertheless, this difference is highlighted here to demonstrate that even for what one would classify as a “major biomass burning event” interpretation of the lidar data is not straightforward. Higher order analysis of this comparison is underway.

Increasing values in aerosol backscatter and extinction profiles indicate increases in extinction with height in the MABL suggesting increasing water-uptake due to enhanced particle hygroscopicity. This is contrary to onboard in situ hygroscopicity measurements in the LARGE package visible in Fig. 4 and noted in Reid et al. (2023) as needing further investigation. Possible reasons for this difference may be physical, such as the role of sea salt (i.e. hygroscopicity measurements of only those penetrating particles through the P-3 inlet vs. the clearly hygroscopic nature of particles within the ambient lidar beam) or some measurement artifact related to particle drying and dehydration process in the instrumentation. This is discussed to some extent by Lorenzo et al. (2025) and commented on further in Sect. 5

Lastly, notable are individual updrafts and cloud halos as in the clean marine case. With surface winds in monsoon en-

hancements at 12 ms^{-1} , convective roll structures are also visible in 2 Hz (90 m) aerosol backscatter data (Fig. 7k). As with background marine, oscillating pillars of higher and lower aerosol backscatter are observed $\sim 2\text{--}3 \text{ km}$ apart. However, these oscillations are lower in magnitude than the maritime counterparts-perhaps due to less active convection or lower hygroscopicity of the smoke particles. Given the higher winds associated with monsoon enhancements, there is notable evidence of wind shear-aerosol backscatter in tilted convective lines rather than vertical pillars.

4.3 Asian air pollution

In comparison to heavy biomass burning smoke, RF17 sampled northwest of Luzon the most significant Asian pollution case advected into the region on 2 October 2019 (Fig. 8a). Post-monsoon, winds over the northern South China and Philippine Sea were quite light and variable (Table 2). Leading up to this flight, pollution from China, including from the Pearl River Delta region was transported to the southeast towards Luzon. RF17 occurred right between a shift in the winds to more northerly conditions, with speeds being variable and $< 1 \text{ ms}^{-1}$ winds from the surface to 2 km. North of Luzon, there were fields of very shallow Cu Hum with thicknesses of 50–200 m (Fig. 8b), although slightly deeper convection was in the vicinity to the south. RF17 fits a land plume model of stratified aerosol features (Fig. 8c) with high sulfate to organic ratios (Fig. 4). Above the typical mixed layer peak in aerosol backscatter at 450 m, a broader land plume is visible from 800–2000 m. Embedded are aerosol backscatter enhancements of just a few hundred meters deep, but up to 50 km long. Lidar ratios for this pollution case were even higher than for biomass burning, being over 60 sr at the surface and increasing to over 80 sr by mid plume (e.g. Fig. 5f). Examining the 2 Hz data (Fig. 8d), we do not find strong updrafts as in the earlier cases of biomass burning or marine conditions-understandable given the very low wind speeds and most limited cloud development. Nevertheless, individual cloud halos are still visible with less scattering downdrafts between clouds.

4.4 Cold pool

The final airborne case presented here is the sampling of an evolving cold pool on RF7 around the 9 September 2019 01:00–04:00 UTC timeframe. Unlike previous cases where the P-3 provided large scale surveys of regional environments, this cold pool case demonstrates perhaps CAMP²Ex's best example of P-3 sampling an evolving environment. Describing the complexity of this case with the P-3's remote sensors and dropsondes is far outside the scope of this paper and is covered in much more detail in a future work. However, in the context of this CAMP²Ex's aerosol “animals in the zoo” paper it is worth giving a brief overview here.




Figure 8. Same as Fig. 4 but for HSRL-2 aerosol backscatter profiles for the RF17 research flight focusing on air pollution from East Asia. (Take-off 1 October 2019). (a) Terra MODIS RGB with P3 flight tracks and curtain segments labelled; (b) forward camera at the start of lidar segments used here. (c) HSRL-2 calibrated aerosol backscatter with cloud screening for the flight segments marked on (a).

After RF7 takeoff, the P-3 was directed to the northeast of Luzon where an isolated thunderstorm began forming at 00:00 UTC. After arriving at 01:00 UTC, the P-3 surveyed cloud features around thunderstorm outflow, followed by a 50 km east-west cross section along a 10 km wide clear air slot above the cold pool that began to develop between the storm core and cold pool's leading edge due to the outward propagation of the cold pool itself (Fig. 9a and c). Forty minutes later the cold pool propagated another 10 km and the P-3 performed a northwest to southeast cross section. Given the complexity of the scene, here we use the 0.5 s HSRL-2 aerosol backscatter for the two transects (Fig. 9e and f) as well as estimated surface wind retrievals from the Advanced Microwave Precipitation Radiometer (AMPR; Amiot et al., 2021; Lang et al., 2021) microwave data for context (Fig. 9g and h). Dropsondes were scattered over the region.

Overall, the RF7 observations conceptually match an almost idealized cold pool case (Tompkins 2001; Drager and van den Heever 2017). Geostationary imagery estimated propagation speed of the arc cloud away from the storm center was $\sim 5\text{--}6\text{ m s}^{-1}$. For the P-3's first crossing at 1:49Z, the arc cloud tops reached $\sim 2\text{ km}$ (distance 8 km on transect), with a congestus to 3.5 km in the middle of the cold pool (distance 35 km) that from satellite imagery may have been from a secondary pulse from the storm. Retrieved surface winds from AMPR qualitatively show significant increases




Figure 9. Similar as Fig. 4 but for the cold pool case on RF7 around 9 September 2019 for two flight segments for 01:39–01:24 and 02:19–02:24 UTC. (a, b) Satellite image with path corresponding to 0–50 km; (c, d) forward camera; (e, f) 2 Hz HSRL-2 aerosol backscatter; (g, h) 20 Hz microwave derived wind speed. Marked under (g) and are where the P3 is outside and inside the cold pool, as well as the location of a potential secondary cold pool forming.

and heterogeneity under the early stages of the arc cloud, rapidly diminishing after it weakened significantly (Fig. 9g and h, respectively). Optically thin altocumulus clouds are also visible in the HSRL-2 as well as the forward and downward camera images. Forty minutes later for the second transect, the cold pool arc cloud had visibly started to dissipate, with cloud tops at only 1 km. Scattered alto cumulus remain visible in the HSRL-2 and video footage.

What makes this case so useful in the context of this paper is that these observations provide more of a mechanistic view of the elevated aerosol layers observed leaving the convective outflow in the RF6 b case provided in Fig. 7. Nominally, our working hypothesis is that smoke is transported long distances in and around the MABL. With the help of vertical wind shear, convectively pumped smoke can be transported in different horizontal directions near the surface. Here, we can observe not just cumulus cloud convective pumping, but the injection of cleaner free troposphere air into the MABL by thunderstorm downdrafts. Indeed, outside the cold pools we find 532 nm AOTs at 0.14–0.18 reducing to 0.05 within (Table 2). We see high aerosol backscatter outside the cold pools (distance $< 5\text{ km}$ for 01:39 case, and $> 30\text{ km}$ for 02:19 case) with AOTs on the order of 0.14–0.18 (Table 2). Behind the cold pool, however, cool density current flow pushed the

aerosol layer upwards creating an elevated layer and overall diminished AOTs to ~ 0.05 . Additional scavenging within the arc cloud may also further reduce aerosol backscatter. Noteworthy is how detraining altocumulus clouds are associated with the tops of aerosol layers, as described in terrestrial counterparts by Reid et al. (2019). This lifting of aerosol particles may also explain the enhanced smoke layers aloft for the RF6 biomass burning case. Also notable is enhanced near surface aerosol backscatter immediately behind the arc cloud in Fig. 9e, we hypothesize due to wind generated sea spray associated with the arc cloud. This is discussed further for a *Sally Ride* case in Sect. 5.2.4. Finally, complex interactions between aerosol loading, precipitation amounts and size distributions, environmental conditions, cold pool intensity, and ultimately cold pool lofting of particles, have been noted in a number of idealized modeling studies (e.g. Grant and van den Heever, 2015). These co-located CAMP²Ex observations of aerosols and cold pools will assist in further unravelling these interactions.

5 Results 3: PBL and vertical variability from the *Sally Ride* HSRL

The R/V *Sally Ride* with its upward pointing UW-HSRL gives a different perspective from the P-3's downward pointing HSRL2. While the NASA P-3 provides an excellent platform for surveying a region at any given time and is configured to provide estimates of the lidar ratio in the MABL, the near-continuous *Sally Ride* measurements observe aerosol vertical distribution within the MABL and MABL variability. The HSRL provides calibrated backscatter and depolarization to within 150 m of the instrument, however retrievals of extinction and lidar ratio are problematic for the up-looking observations due to the receiver being out of focus close to the instrument (< 5 km). Further, the *Sally Ride*'s strategic position has the advantage of long-term monitoring just upwind of maximum convection around the NWTP monsoonal trough, with the disadvantage of westerly shear advecting upstream smoke above the MABL to the east (e.g. Sect. 3). This said, smoke aloft is then much more likely to have been placed there by more local convection. In this section, we examine the time series data and cloud related perturbations of aerosol backscatter.

5.1 Period overview

Sally Ride started collecting lidar data on 4 September 2019 on its way from Taiwan and was on station 7 September in the NWTP just at the start of the first of two major Indonesian smoke outbreaks. Figure 10 provides an overview of cloud and precipitation screened aerosol backscatter taken over the course of the *Sally Ride* deployment along with bow mast meteorology and radiosonde winds. Figure 10a provides a time series of aerosol backscatter to 5 km, with a log scale for aerosol backscatter. Focusing on the MABL in

Fig. 10b, we replot Fig. 10a on a log vertical scale but a linear aerosol backscatter scale. Overlaid are estimated hourly cloud bases for the lowest 10 % of detected clouds from the onboard ceilometer as a standard. Given the high sensitivity of the HSRL, defining the backscatter threshold that differentiates between aerosol and cloud is needed given the system resolves the increase in backscatter through this transition. Nevertheless, the ceilometer cloud base is a fair first approximation to MABL mixed layer depth. To help with interpretation, a “Pseudo Extinction” scale is included by assuming a lidar ratio of 50 sr, the middle of the range in Fig. 4e and f. In clean marine conditions where a lidar ratio of ~ 20 – 25 sr (e.g. Fig. 4e) may be more appropriate, resulting in an overestimate of extinction by a factor of 2 by using this scale. Conversely, for pollution cases where a lidar ratio may reach 60 sr, the use of 50 would underestimate extinction by 20 %. Thus, this depiction should be considered only for qualitative scaling purposes for biomass burning and pollution and more details will be provided in a subsequent paper on MABL lidar ratios. Using integrated aerosol backscatter, we likewise provide pseudo AOT for cloud free profiles to 7 km are provided in Fig. 10c. Included is Pseudo extinction for 200 m altitude as a mid-mixed layer benchmark. Meteorological data from the *Sally Ride*'s bow mast is provided in Fig. 10d – Precipitation, e – 10 m temperature, sea surface skin temperature and relative humidity and f 10 m winds and directions. Also included are extracted g wind speed and h direction at 250, 1500, and 3000 m from the 3 hourly radiosonde releases.

Significant aerosol events were observed throughout the cruise. The largest aerosol backscatter values are overwhelmingly within MABL's mixed layer, and peaking at and around cloud base. The line plot of integrated and 250 m backscatter/pseudo extinction (Fig. 10c), estimated 532 nm maximum extinction coefficients were associated with major smoke events on the order of 0.3 – 0.5 km^{-1} with RH < 85 %, nominally a visibility of only 6–4 km, respectively, associated with the Koschmieder coefficient of 1.9 (Griffing, 1980; Husar et al., 2000). These are similar to the P-3-HSRL2 for smoke flights. Corresponding 532 nm Pseudo AOT for cloud free paths to 7 km for events range from 0.3–0.6. The 1 Mm^{-1} sr^{-1} line, indicated by the blue to green transition in the colorbars, is a good qualitative indicator of polluted conditions. This level is at approximately 1–2 km, with isolated events at 2–3 km and the highest notable plumes detected on 17 September to the 0°C level at ~ 5 km. This was confirmed with personal observations on the ship. Again, given a choice of 50 sr as a lidar ratio, this estimated extinction and AOT is probably underestimated by ~ 0.05 – 0.1 km^{-1} and 0.06–0.12, respectively. Conversely, pseudo extinctions and AOTs were as low as 0.1 km^{-1} and 0.10, but again the true values were likely half these values. Thus, overall, we expect estimated AOTs and extinction coefficients to be under-dispersive.

Estimated MABL cloud base heights throughout the cruise ranged from a low of 300 m to a high of 800 m, with a typ-




Figure 10. September 2019 timeseries of UW-HSRL aerosol backscatter and surface meteorology from the R/V *Sally Ride* bow mast. Meteorology data began when the *Sally Ride* arrived on station on 6 September 2019, (a) Cloud screened aerosol backscatter at 532 nm on a log scale colorbar. Included is a “pseudo extinction” for scaling based on an assumed lidar ratio of 50 sr. This will likely underestimate clean marine conditions by a factor of two, and underestimate extinction for fine dominated environments by $\sim 20\%$. (b) same as (a) but on a log vertical scale and linear aerosol backscatter scale. Overlaid in red is estimated cloud base; (c) For reference, pseudo Aerosol Optical Depth (AOT) and extinction at 200 m using an assumed lidar ratio of 50 sr, (d) Surface rain rate; (e) 2 m temperature and relative humidity and sea surface skin temperature; (f) 10 m windspeed and direction; (g) Radiosonde derived wind speed at 250, 1500, and 3000 m; (h) Radiosonde derived wind direction at 250, 1500, and 3000 m.

ical value of ~ 450 m during the SWM, rising to ~ 650 m during the NWM. Examination of Fig. 10b and as discussed further on cloud edges are frequently observed to be associated with an overall enhanced aerosol backscatter to 1500–3000 m, with occasional altocumulus layers on top.

Examining the different panels of Fig. 10 as a whole, one can see significant aerosol and meteorology variability at both low and high frequencies. Aerosol backscatter in Fig. 10b has a distinct striping nature well beyond what can be attributed to precipitation or very high relative humidity, with events arriving and departing quite suddenly. Even within individual aerosol events aerosol backscatter/pseudo extinction often has a $\sim 20\%$ – 30% scatter in variability. While one would think that being in the Southwest Monsoon there would be continuous rainfall, in reality precipitation is in scattered cells, leading to short deluges and occasional light stratiform precipitation (Fig. 10d). For indicators of monsoonal state (Fig. 10e–h) – southwest monsoon also showed variations in strength, building though the cruise with an abrupt transition to a winter northeast monsoon mode on 22 September. A flow reversal to easterly winds on 10–13 September was due to a tropical depression to the east bringing additional variability in aerosol backscatter. Temperature and relative humidity also tracked monsoonal strength, with slight diurnal cycles punctuated by occasional cold pools and convective development (Fig. 10f). The air sea temperature differences were consistently unstable; typically skin SST minus 10 m air temperature was 5°C , punctuated by cold pools reducing the still unstable value in magnitude to 3.5°C . There were occasional very slightly stable periods characterized by skin SST minus air temperature values of -0.5°C , likely due to advection from Asia (e.g. just 18–19, and 21 September out of the total 22 d cruise time series). Comparing monsoonal state back to the lidar profiles, aerosol events track monsoon enhancements in wind. Directional vertical wind shear is also at times noticeable. This leads to the aforementioned shearing phenomena that reinforces a low-level transport pattern.

We summarized the observed variability in aerosol backscatter by performing a series of statistics for 3 h increments in Fig. 11. Over the course of the deployment, cloud and precipitation free aerosol backscatter spanned almost a factor of 5 (Fig. 11a), with Western Pacific air sampled at $1.8 (\text{Mm sr})^{-1}$ during a monsoon break on 10 September, and during the NEM on 24 September. Mean values reached $9 (\text{Mm sr})^{-1}$ for aforementioned monsoonal enhancements, or an extinction of roughly 0.45 km^{-1} if one assumes a 50 sr^{-1} lidar ratio. It is notable that there is also significant variability in backscatter deviations, being relatively low in the NWM monsoon breaks and NEM, and increasing in the monsoon enhancements. As with the P-3 observations, the highest backscatter values are in the MABL, with the 50th% in integrated aerosol backscatter being at 400–700 m (Fig. 11b), and the 84th % a kilometer above that-reaching a high of 2 km mid mission. If we take the MABLs mixed layer

to be cloud base taken as the lowest 10 % of clouds from the ceilometer plus 200 m for cloud entrainment/detrainment zone, 25 %–60 % of integrated aerosol backscatter is with the MABL's Mixed layer, averaging 50 % (Fig. 11c). Also as noted in the P-3 analyses using the HSRL-2's 0.5 s data and in Kunkel et al. (1977) and Reid et al. (2017), aerosol backscatter enhancements exist under clouds owing to up-drafts. We can in part quantify this variability by taking the ratio of aerosol backscatter at an altitude of 200 m for those UW-HSRL samples with observations attenuated in the lowest 800 m representing cloudy conditions, to those with adequate sample to 5 km representing clear (Fig. 11d). Almost all samples show ratios above 1, with 10 %–20 % enhancements being typical.

5.2 Example periods of aerosol variability

It is helpful to examine short periods of the *Sally Ride* data to better interpret the long time series presented in Sect. 5.1. Indeed, like performed for the P-3, here we examine the nature of MABL variability for different aerosol regimes, but with a bottom-up instead of top-down perspective. The SSEC HSRL typically had a scan pattern between vertical stare and horizontal scan, repeating every 5.5 min. This made isolating individual cloud features difficult. However, there were several periods of vertical stare of several hours that allow for more detailed analysis. Whereas Fig. 10 images were made with 30 s/30 m data where available, for these long periods of vertical stare, data were reprocessed for continuous 5 s, 7.5 m vertical resolution. At nominal 10 m s^{-1} wind speeds, data are functionally ~ 50 m in horizontal resolution, as opposed to resolution ≥ 1 km for the P-3 and *Sally Ride* baseline observations.

5.2.1 Fair weather MBL

Based on Fig. 11a, one of the lowest aerosol backscatter periods in magnitude and variability observed on *Sally Ride* was the post monsoon transition to the NWM at the end of the cruise, 24–25 September 2019. Nominally we consider this mostly Western Pacific “fair weather marine” conditions sampled by *Sally Ride*, given its low backscatter conditions and moderate regional northeasterly winds. No precipitation was detected in and around the vessel at this time. Nevertheless, given the analysis of Hilario et al. (2021) and Sect. 3, some East Asian influence cannot be categorically excluded. Indeed, earlier that day, the P-3 was in the vicinity of the *Sally Ride* and a lidar ratio of ~ 40 sr was found in the region, indicative of a mixed environment of fine and coarse particles. Retrievals of AOT at 550 nm for Terra/Aqua MODIS and at 532 nm for the LaRC HSRL for this period were also on the order of 0.15–0.2-above what is more typical for clean marine conditions (e.g. Reid et al., 2023). Because this is our most ideal “fair weather marine” baseline case, we go into more detail on the nature of the shipboard observa-




Figure 11. Marine Boundary and lower free troposphere statistics of UW-HSRL aerosol backscatter from the R/V *Sally Ride* based on 3 h increments. **(a)** Average and standard deviation of 532 nm aerosol backscatter taken at 200 m (mid mixed layer) with a second scale of “pseudo extinction” assuming a lidar ratio of 50 sr^{-1} ; **(b)** 50th and 84th percentile of integrated aerosol backscatter taken from cloud free samples to 5 km; **(c)** Percentage of integrated aerosol backscatter within the Marine Atmospheric Boundary Layer’s (MABL) mixed layer taken as MABL cloud base; **(d)** The ratio of aerosol backscatter taken at 200 m for samples under MABL clouds and those that are cloud free.

tions (Fig. 12). Subsequent Sects. 5.2.2 and 5.2.3 cover more briefly highly wind sheared and heavy smoke environments.

Figure 12 provides a synopsis of a two-hour period from 24 September 2019 22:00 UTC to 25 September 2019 00:00 UTC. Aerosol backscatter time series data (Fig. 12a) were collected in an area of scattered cumulus clouds adjacent to northeasterly MCS outflow (Fig. 12b). Notable is the frequent striping nature in aerosol backscatter below the fair Cu (red colors with backscatter $> 10 \text{ Mm}^{-1} \text{ sr}^{-1}$). Enhanced backscatter features under clouds are consistent with cloud updrafts, atmospheric roll phenomena, and associated particle transport and hygroscopicity (Kunkel et al., 1977). Particle hygroscopic growth is further indicated in the linear depolarization (Fig. 12c) as more water content leads to more spherical particles relative to a refractory core with shrinking volume fraction; areas of enhanced backscatter have lower depolarization in the mixed layer. Overall, while these depolarization values in the mixed layer are quite low, it is nevertheless supportive of this being a “mixed” environment of both sea salt and a combination of Asian pollution and perhaps residual smoke from the Maritime Continent. The lowest depolarizations are below cloud base. However, once a cloud forms, particles grow to sufficient size as cloud droplets that they depolarize again. Above cloud level, even in clear air, particles have dried somewhat and higher depolarization appears.

The striping patterns in aerosol backscatter and linear depolarization are quite vivid and can be more readily quantified in the generation of time series (d) including 150 m, the altitude at which we are most confident in near surface returns, and 500, 50–100 m below cloud base. As expected, higher altitudes in the ML exhibit higher backscatter. At 150 m, the amplitude in aerosol backscatter varies between $2\text{--}3 \text{ Mm}^{-1} \text{ sr}^{-1}$, 20 % standard deviation around the mean and at 500 m 1.6 to $> 8 \text{ (Mm sr}^{-1}\text{)}$, or almost a factor of two standard deviations around the mean. Notably, at 500 m values in aerosol backscatter frequently drop below the 150 m values. These oscillations are consistent with Fig. 12d in that they are most often associated with clouds/between clouds and extend from cloud base to the surface.

Clearly, these areas of aerosol backscatter enhancement extend to near the surface, modulated by the presence or absence of clouds. Individual events are also much greater than what would be expected from the bulk analysis presented in Fig. 12d. On this particular day, the frequency of cloud observation was 5 min or roughly $\sim 2.7 \text{ km}$ assuming a mean wind speed of $\sim 9 \text{ m s}^{-1}$. At the 5 s sample level, correlations of aerosol backscatter between 150–500 m are only moderate ($r = 0.42$; $r^2 = 0.21$), in part because of the significant non-linear behavior below cloud base. Performing 30 s averages, correlations improve to $r = 0.60$; or $r^2 = 0.36$.




Figure 12. A selected contiguous timeseries of lidar data on 24 September 2019 during the more maritime Northwest monsoon. While winds are from the North Tropical Western Pacific, there is still likely some influence from Asia. Included are (a) Aerosol backscatter up to 1000 m for 22:00–00:00 UTC; (b) AHI imager RGB of the region, with the *Sally Ride* location and surface wind barb; (c) Linear depolarization ratio timeseries; (d) Aerosol backscatter timeseries from 150 and 500 m; (e) Time series of bow mast/surface relative humidity, temperature and wind speed on the *Sally Ride*; (f) At the time of the 23:27 UTC radiosonde release aerosol backscatter, and sonde relative humidity (RH), potential temperature (θ), and equivalent potential temperature (θ_e); (g) and (h) Aerosol backscatter and linear depolarization (same color bar as a and c) for 90 s just after the radiosonde release. Marked is the radiosonde release (magenta line), and notable areas of positive correlations of aerosol backscatter and temperature/anticorrelation with wind speed (red arrows).

While the lowest altitude we are very confident in lidar observations is 150 m, there are indications of these enhancements reaching ship level. The time series of *Sally Ride* 1 min relative humidity, temperature and surface wind speed from the bow mast show the telltale signs of sweep and ejection events associated with variations in aerosol backscatter (Fig. 12e). Marked as red arrows between Fig. 12d and e are notable isolated peaks in backscatter, and these correspond with increases in temperature and decreases in horizontal wind—consistent with a surface ejection event (positive sensible heat flux, negative moment flux). Corresponding adjacent drops in temperatures and increases in wind speed likewise are a result of large-scale downdrafts (so-called sweeps). Humidity for this cruise was measured with an aspirated system with a response time of nearly 30 s, and values were subject to long period variability. Thus RH does not correlate as well as temperature and wind, but nevertheless, small amplitude

peaks in RH likewise correlated with temperature and anticorrelate with wind. While a full analysis of the relationship between surface air sea fluxes and their relation to individual clouds is outside the scope of this survey paper, the preliminary evidence strongly suggests such connections between surface fluxes and individual clouds. Indeed, the strong amplitude in 500 m aerosol backscatter in Fig. 12d shows how structured variation exists around cloud base. This leads to questions about sampling and the distinct “modal” nature of aerosol backscatter below and between clouds.

Examining above the ML in the MABL’s entrainment zone, we also see signs of inhomogeneity. There are sharp delineations in aerosol backscatter and depolarization around cloud level. Areas of particle aerosol enhancement are also decorrelated with aerosol depolarization above 600 m, a sign of particle detrainment. This said, some of the increased depolarization may be due to the extremely clean environment

of the free troposphere. We can examine this further with radiosonde data. The challenge with applying radiosonde data to high-resolution lidar analysis is observations can be aliased; we cannot exactly match up the sonde with the lidar-wind shear and the evolving MABL. However, the sondes are quite useful in describing the overall environment. Marked under Fig. 12a is the radiosonde release time of 23:26 UTC. In Fig. 12f we provide the aerosol backscatter at that time, +5, -15 min, and the 2 h average of contiguous data, along with the radiosondes wind speed, relative humidity, potential temperature, and equivalent potential temperature profiles. While the mixed layer top is often taken as cloud base and derived as the first peak in backscatter or at some threshold value, presumably at a peak in RH, we can see in Fig. 12a that such definitions can be problematic. The inversion “entrainment zone” of mixed layer free troposphere exchange is somewhat ambiguous, extending well into and above the mixed layer; again noting the detraining aerosol plumes are quite visible in Fig. 12a and b. Profiles in backscatter are quite variable depending on if it is taken under or between a cloud, with a 2 h average smoothing these states together. It is also noteworthy that the lower values of depolarization extend well beyond the peaks in aerosol backscatter. Comparing these profiles to radiosonde profiles is likewise problematic, because it can take up to 90–120 s in a sheared environment to reach 500 m. Nevertheless, we do observe some “better behaved” profiles. RH systematically increases to 98.8 % at 580 m while potential temperature and equivalent potential temperature stay roughly constant, except for some near-surface instability below 40 m. At the top of the mixed layer, RH and equivalent potential temperature rapidly drop, and potential temperature rapidly increases. This is consistent with conditions supporting a moist MABL in fair weather conditions. While wind shear is visible in the lowest few hundred meters, the winds aloft are fairly uniform at 9 m s^{-1} with wind directions ranging from 40–60°.

Finally, we can examine all of the data above in the context of a single cloud. Figure 12g and h show aerosol backscatter and linear depolarization around a faint cloud observed a few minutes after the radiosonde release. Clearly visible are an inner updraft core of $\sim 250 \text{ m}$ in diameter, of high backscatter and lower depolarization surrounded by a 100 m thick structured “halo” of related cloud influence. Notable are some areas of enhanced backscatter protruding beyond and below the cloud.

5.2.2 Convectively active marine environment

We can compare the “fair weather marine conditions” with a more convectively active regime. Around 12 September, the *Sally Ride* was in relatively clear air surrounded by tropical depression Marilyn and two other MCSs. In effect, the *Sally Ride* was within the middle of the monsoon trough. Figure 13 provides similar information as Fig. 12 for 04:00–06:00 UTC on 12 September 2019. During this period while being sur-

rounded by convective systems and precipitating fair weather clouds, the *Sally Ride* sampled a series of shallow convective lines advecting from depression Marilyn to the northeast. Unlike more typical roll structures that are aligned parallel to the wind shear vector (longitudinal roll clouds), these cloud features were oriented transverse to the prevailing MABL wind shear vector (transverse roll clouds). Despite the convective nature of the region, no precipitation was detected at the ship for this short investigation period. Comparing to our fair-weather case, we see significant structural differences in aerosol backscatter (Fig. 13a) and depolarization (Fig. 13b), areas of aerosol backscatter enhancement are not evenly spaced, but in distinct areas associated with cloud line passage. Under clouds, high frequency variability still exists. As air is advecting from tropical depression Marilyn with greater wind speeds, enhanced sea salt related backscatter is likely well above the 24 September case. With the absence of any continental influence, depolarization is likewise greatly diminished (Fig. 13b), almost undetectable. Areas of aerosol detraining between 700–1200 m are more strongly evident. Variability in aerosol backscatter at 405 m (50 m below nominal cloud base) relative to 150 m is far greater than the 24 September case, but backscatter at levels throughout the ML are nevertheless correlated (Fig. 13d). Likewise, peaks in the anticorrelation in surface wind and temperature appear to be related to backscatter enhancements (Fig. 13e) although at longer temporal periods than the 24 September case as well. In this environment, correlation to RH are less dramatic.

One of the interesting aspects of the 12 September case is its greater wind speeds and vertical wind shear. The 05:29 UTC radiosonde release occurred during the passing of a line of clouds. Included in Fig. 13f are 5 coincident 5 s data collections as the cloud passed over the site. The first profile at 05:29:15 UTC was just as a cloud exited, with a peak backscatter of $200 \text{ Mm}^{-1} \text{ sr}^{-1}$ at 600 m. Five seconds later or $\sim 75 \text{ m}$, backscatter was down to $20 (\text{Mm sr})^{-1}$ reaching a minimum of $8 (\text{Mm sr})^{-1}$. Progressively, 15 more seconds after that with a peak in backscatter decreased in altitude and magnitude to 400 m, resolving an exchange event. Notable in the associated soundings are not only higher winds than the 24 September case, but more vertical wind shear – increasing from 8 m s^{-1} at the surface to 16 m s^{-1} at cloud level. Nonlinearities in the relative humidity, potential temperature, and equivalent potential temperature plots show evidence of highly localized mixing. Indeed, if we examine a short five-minute period of lidar data around the radiosonde release (Fig. 13g aerosol backscatter and (h) linear depolarization) we can clearly see the relationship between wind shear and the slanted aerosol backscatter returns as well as the presence of a detraining layer at 700 m just above the cloud level enhancement.

Because of wind shear, the correlations between 150 m and cloud base are more difficult to quantify, although we can qualitatively see them at larger scales. Taken at 5 s resolution,




Figure 13. Similar to Fig. 12 but for a selected contiguous timeseries of lidar data on 12 September 2019 examining outflow from a tropical system. Included are (a) Aerosol Backscatter up to 1000 m for 22:00–00:00 UTC; (b) AHI imager RGB of the region, with the *Sally Ride* location and surface wind barb; (c) Linear depolarization ratio timeseries; (d) Aerosol backscatter timeseries for 150 and 500 m; (e) Time series of bow mast/surface relative humidity, temperature and winds speed on the *Sally Ride*; (f) At the time of the 23:27 UTC radiosonde release aerosol backscatter, and sonde relative humidity (RH), potential temperature (θ), and equivalent potential temperature (θ_e); (g) and (h) Aerosol backscatter and linear depolarization (same color bar as a and c) for 5 min just after the radiosonde release. Marked is the radiosonde release (magenta line), and notable areas of positive correlations of aerosol backscatter and temperature/anticorrelation with wind speed (red arrows).

for the 2 h period 150 and 400 m aerosol backscatter correlations are difficult to compute as cloud edge effects generate significant variability (Fig. 13d). But if we exclude all cases where 405 m backscatter is higher than, say, $20 \text{ Mm}^{-1} \text{ sr}^{-1}$, correlations are still reasonable—even with shear (slope of 1.75, $r = 0.54$; $r^2 = 0.3$). Performing a lag correlation results in a slight increase to an r^2 of 0.32 after 25 s. Averaging to 30 s sees significant improvement, with a slope now of 3.5 ($r = 0.8$; $r^2 = 0.66$).

5.2.3 Heavily polluted marine

Finally, we can quickly contrast the cleaner cases of 12 and 24 September and to the most polluted period sampled by the *Sally Ride* during PISTON, ~ 16 –18 September 2019. The P-3 flew 16 and 17 September, and the *Sally Ride* was forced to disengage to the north to avoid severe weather.

The early morning of 18 September the *Sally Ride* started to transit back to its station when it intercepted the thick polluted air mass as it entered a tropical system (Fig. 14, similar format to Fig. 12 but note significant scale changes). Like the more-pristine days, while we see similar features of enhanced backscatter and suppressed linear depolarization under clouds, the situation overall is much more complex. The monsoon enhancement comes with even stronger winds and wind shear than the previous cases reviewed. Notable detrainment layers with altocumulus formation were observed for the last 30 min of the time series. Backscatter also correlates vertical within the ML but with a lesser increase with altitude (~ 1.3 from 150 to 500 m or lowest observation level to 50 below cloud base)—consistent with less hygroscopic smoke vs. more sea salt and sulfate for the cleaner marine case in Sects. 5.2.1 and 5.5.2 – but with stronger outliers. Positive correlations between backscatter and temperatures,




Figure 14. Similar to Fig. 12 but for selected contiguous timeseries of lidar data on 18 September 2019 during the most polluted period samples during PISTON. Note significant scale changes from the previous figures. Included are (a) Aerosol backscatter up to 1000 m for 22:00–00:00 UTC; (b) AHIMAGER RGB of the region, with the *Sally Ride* location and surface wind barb; (c) Linear depolarization ratio timeseries; (d) Aerosol backscatter timeseries for 150 and 500 m; (e) Time series of bow mast/surface relative humidity, temperature and winds speed on the *Sally Ride*; (f) At the time of the 23:27 UTC radiosonde release aerosol backscatter, and sonde relative humidity (RH), potential temperature (θ), and equivalent potential temperature (θ_e); (g) and (h) Aerosol backscatter and linear depolarization (same color bar as a and c) for 5 min just after the radiosonde release.

and anticorrelation with wind speed exist, but also on a larger scale, in association with what may be more organized cloud-related circulations. Perturbations to RH, if they exist, are below detectable limits. The sounding is not as well behaved as the previous, with multiple detrainment levels visible; even variables like RH that are typically monotonic in the mixed layer show significant variation. Mixed layer height is at 600 m, and cloud tops in the lidar data correspond well to a second inversion at 1000 m. Indeed, for the one clearly “cloud free” case near radiosonde release aerosol backscatter was higher aloft along with many fine detrainment layers. In nearly every sense the polluted marine environment is much more complex; it is difficult to find an individual cloud (e.g. like Fig. 13g and h). Spatial inhomogeneity and the effect of wind shear is significant in both backscatter and linear depolarization, and even within the 2 h period different regimes are visible.

5.2.4 Cold pools

As noted in Sect. 4.4 transient cold pool events may be an important driver of aerosol lifting out of the MABL and into the free troposphere. Based on the *Sally Ride*’s near-surface air temperature time series in Fig. 10, six rapid temperature drops of greater than 1°C collocated with sharp near-surface wind speed increases were observed during the cruise. Of these six, two occurred while the SSEC HSRL was viewing vertically, between 16:00–23:00 UTC on 20 September 2019. The first of these two, between 16:00 and 08:00 UTC, had superior viewing conditions and is used as an example in Fig. 15 following a similar figure format as Figs. 12–14. However, unlike previous figures given the dynamic range of the observed lidar signals, log scales are used.

The aerosol backscatter profiles for a ninety-minute period are provided in Fig. 14a. At the beginning of the time




Figure 15. Similar to Fig. 8 but for two transects across an evolving cold pool for RF7 around 20 September 2019 between 16:00 and 17:30 UTC. Included are (a) Aerosol backscatter up to 1000 m; (b) AH1 thermal infrared image of the region, with the *Sally Ride* location and surface wind barb; (c) Linear depolarization ratio timeseries; (d) Aerosol backscatter timeseries for 150 and 450 m; (e) Time series of bow mast/surface relative humidity, temperature and winds speed on the *Sally Ride*; (f) At the time of the 17:29 UTC radiosonde release aerosol backscatter, and sonde relative humidity (RH), potential temperature (θ), and equivalent potential temperature (θ_e); (g) aerosol backscatter (same color bar as a) for 10 min after cold pool passage starting at 16:30 UTC.

series, we see similar aerosol backscatter features to the convective environment in Figs. 12a and 13a. However, notable is a heavy rain shaft at 16:20 UTC, followed by that storm's cold pool with very clean air above 200 m at 16:30 UTC. Another minor event was observed just 15 min later. This cold pool occurred at night within one of Typhoon Tapa's convective arms, and only IR imagery is available (Fig. 15b). Like the previous active convection and clean (presumably sea salt dominated) conditions, there is virtually no linear depolarization signal associated with aerosol particles, although the rain shafts are clearly delineated around and above 0.2 resulting from multiple scattering within the precipitation. While overall, comparisons of nearer surface (e.g. 150 m) vs. mixed layer top are similar to other cases, near the cold pool boundaries vertical and horizontal gradients in aerosol backscatter between the near surface and upper mixed layer environments can be quite dramatic (Fig. 15d).

The influence of the cold pool is well described by *Sally Ride*'s observation of a drop in air temperature and sharp

increase in wind speed and RH (Fig. 13e). Given the presence of rain shafts next to clear air, aerosol backscatter signals span two orders of magnitude in this data segment. Ultimately, the near cold pool environment results in more varied aerosol and thermodynamic profiles compared to more benign conditions. Included in Fig. 15f are example aerosol backscatter profiles within and outside of the cold pool near radiosonde release. Within the cold pool (16:34 and 16:35Z) aerosol backscatter profiles are essentially static in the MABL, except for a near surface aerosol backscatter enhancement. After the atmosphere had recovered, the MABL sounding and aerosol profiles were much more typical.

Magnifying on the cold pool area (Fig. 15g and h), strong enhancements of aerosol backscatter very near the surface are associated with the enhanced winds along the cold pool boundary. As noted in the P-3 case in Sect. 4.4 and in similar observations of aerosol particles in similar convective regimes in the South China Sea (Reid et al., 2015), these near

surface backscatter enhancements may be due to wind driven sea spray-essentially sea salt haboobs.

6 Conclusions and implications

The purpose of this paper is to outline the modes of variability in aerosol vertical profiles over the North Western Tropical Pacific (NWTP) using the combined assets of the ONR Propagation of Intraseasonal Oscillations (PISTON) R/V *Sally Ride* stationed East of Luzon and the NASA Cloud, Aerosol and Monsoon Processes Philippines Experiment (CAMP²Ex) P-3 flying over the Sulu, South China and Philippine Seas. These cooperative experiments were deployed in the late boreal summer of 2019, and fortuitously captured the monsoonal transition from the large-scale Southwest to Northeast flow. Consequently, the platforms observed a wide variety of aerosol sources, including significant biomass burning activity and pollution emissions from the Maritime Continent (MC), Asian pollution, and pristine air from the NWTP and convective outflow. The focus in the paper is on both the P-3's and *Sally Ride*'s High Spectral Resolution Lidars (HSRLs) to outline the “animals in the zoo” of aerosol vertical phenomenology as an introduction to the rationale for needed efforts in understanding how aerosol inhomogeneity relates to the heterogeneous nature of aerosol transport/exchange/transformation, radiation, and flux/exchange fields. Throughout, we note areas of ongoing work that reference this overview paper.

As has been noted in aforementioned work (e.g. Hilario et al., 2021; Edwards et al., 2022; Reid et al., 2023; Xiao et al., 2023) and presented in more detail here, P-3 in situ and *Sally Ride* HSRL observations highlight the MC's distinct monsoon driven lower free troposphere and long-range transport phenomenology. As a consequence, the larger NWTP's Marine Atmospheric Boundary Layer (MABL) and lower free troposphere are impacted by a very wide range of aerosol properties. Unlike most subcontinental to continental scale aerosol events where long-range transport is facilitated by the formation of land plumes and advection by free tropospheric winds, the 2019 PISTON and CAMP²Ex missions verified the dominance of MABL and lower free troposphere aerosol transport pathways of thousands of kilometers with 532 nm Aerosol Optical Thickness (AOT) at times above 1. Background marine, Indonesian biomass burning outbreaks, MC emissions and Asian pollution cases were observed to have 95 % of integrated aerosol backscatter below 3 km in altitude in both P-3 and *Sally Ride* observations. Twenty to fifty percent of total aerosol backscatter and extinction are contributed by the MABL's 400–650 m high Mixed Layer (ML), with the remainder mostly falling off with altitude in the free troposphere, with intermittent cloud detrainment layers. Conversely, aerosol backscatter is quite low in the middle and upper free troposphere. All three cases of P-3 observations near the *Sally Ride* point to good comparisons of aerosol

backscatter profiles between the instruments in the MABL overall, but demonstrate significant variability in spatial scale of aerosol layering in the lower free troposphere. These different “top down” vs. “bottom up” views agree in the nature of lower-level transport, but there are noticeable differences in the benchmark calculation of fraction of aerosol backscatter that is in the boundary layer. Closure studies are underway.

There are several key physics findings in the analysis that are related to scale, from the large-scale monsoon to the role of convection and the importance of individual updrafts.

1. *Monsoonal Maintenance*: In this paper, we expanded on Edwards et al. (2022) analysis of the Navy Aerosol Analysis and Prediction System Reanalysis (NAAPS-RA) simulations. At the largest monsoonal scale, what maintains the primary low-level aerosol transport process is a combination of low injection heights of smoke and pollution off of the MC islands (as noted in Wang et al., 2013) coupled with veering directional wind shear with height. Thus, particle transport naturally bifurcates in altitude, with the near surface particles remaining in and around the MABL during long range transport in the SWM flow. Any aerosol layers aloft from either enhanced injection or convective pumping would overall be preferentially advected more eastward than those in the MABL. As demonstrated in Bukowski et al. (2017), above the melting level (~ 4.8 km) monsoonal flow reverses direction completely to northeasterly. The P-3 observations clearly observed a reduction in aerosol layering aloft between just offshore of Borneo and 1000 km to the northeast. The absence of strong aerosol layering above 3 km suggests a combination of low level free tropospheric inversions capping sometimes active warm convection and efficient wet scavenging as contributors to keeping aerosol layering heights low. Further, even though the region is convectively active, it is only through major tropical features or significant coastally forced convection that major scavenging takes place. Consequently, fine mode particle lifetimes in or just above MABL can be extraordinarily long lived in this SWM system lasting over 4000 km. This work supports the long-standing suggestions by Clarke et al. (1996, 2013) and Quinn et al. (2017) that note that even in the most remote marine areas, cloud condensation nuclei populations can be heavily influenced by terrestrial sources. However, while Clark et al.'s observations suggest mixing from the free troposphere into the MABL, this present work shows that particle can be exceptionally long lived in the MABL only to be scavenged in significant storm features. Because these primary maintenance mechanisms are at the large scale, the global models replicate this long-range transport quite well as shown in the NAAPS-RA. The extent that smoke as

CCN may inhibit precipitation and thus *significantly reduce* scavenging is still an open question for CAMP²Ex,

- Lower free troposphere layering*: While the large-scale flow helps sustain long-range low-level aerosol transport, in our discussions with other regional scientists investigators are still at times surprised by the dominance of the MABL and its mixed layer over the lower free troposphere for the highest particle concentrations and aerosol backscatter. With so much convection, it is argued that more detrainment above the MABL and scavenging within the MABL should be expected. Indeed, only once in this analysis (the P-3 RF6 case on 6 September, Fig. 5h) did we find a situation of maximum particle concentrations above the MABL. But even here it might be explainable again as a combination of vertical wind shear to the east of an area of active warm convection capped by a strong mid-level inversion and potential cold pool lofting. However, it is noteworthy that while the highest concentrations and maximum aerosol backscatter are in the shallow MABL, the integral of particle mass species and backscatter is still often on par with the lower free troposphere, punctuated by frequently observed and remarkably thin (< 100 m) aerosol layers, demonstrating significant convective detrainment. In cases adjacent to detraining clouds, detrainment layers are also associated with altocumulus. Further, both the P-3 and *Sally Ride* HSRL observations show how intricate the detrainment process is in and around cumulus clouds. This likely explains some deficiencies in sub-grid parameterizations in describing lower free troposphere aerosol loading as identified in Edwards et al. (2022). The absence of strong aerosol layers in the middle to upper free troposphere coupled with enhanced CO is suggestive of efficient wet scavenging for those cells that puncture lower-level inversions (e.g. Hilario et al., 2025). Although as Xiao et al. (2023) discusses, there is significant nucleation potential of detrained precursors aloft.
- MABL Exchange*: One of the most striking findings of this survey is how variable aerosol backscatter, depolarization, and presumably light extinction are in the MABL and its associated mixed layer. Local enhancements in aerosol backscatter of 30 %–100 % were found underneath clouds. Just as observed by the ground observations by Kunkel et al. (1977) and the SSEC-HSRL deployment at Huntsville AL for the terrestrial NASA Study of Emissions and Atmospheric Composition, Clouds and Climate by Regional Surveys campaign (SEAC⁴RS; Reid et al., 2017), there are persistent enhancements in aerosol backscatter and decreases in linear depolarization due to moist updrafts under clouds. Cloud halos relative to large detrainment layers are also clear. The difference for the marine environment measured in PISTON/CAMP²Ex vs. terrestrial observations

is shallowness of the boundary layer inversion, ~ 450 – 650 m over water vs. 1.5 – 2 km over land. Consequently, these under-cloud backscatter enhancements reach the ocean surface, even visibly correlated to surface winds and temperature, and presumably fluxes. High temporal resolution observations from the *Sally Ride* clearly demonstrate the formation of cloud halos and air mass exchange around clouds. Such features are also observed in P-3 HSRL-2 0.5 s data at the edges of clouds, although with the aircraft flying at 150 m s⁻¹ fine features are washed out. Mixed layer backscatter enhancements are more prevalent for higher large-scale MABL wind speeds (> 7 m s⁻¹) where MABL roll features and associated cloud streets form. Such features are diminished for lower wind speeds or stable conditions-which unfortunately existed during all three periods of P-3-*Sally Ride* joint observations. As expected for periods of higher MABL wind shear, aerosol backscatter enhancement features are likewise noticeably sheared (e.g. Fig. 13g). Nevertheless, the lidar observations clearly link cloud base to the air-sea interface. Increased aerosol backscatter is locally related to reduced wind speed and increased temperature associated with an ejection event, or ejection; decreased backscatter is associated with decreased temperature and increased wind (i.e. sweep event). Further studies are needed to establish the extent these backscatter enhancements are associated with surface fluxes, MABL/free troposphere exchange, and moisture convergence.

- Observational signal to noise in the HSRL*: The great advantage of the HSRL systems over other lidars is its ability to very accurately derive calibrated aerosol backscatter both day and night from the ratio of total to molecular backscatter. Consequently, even in the most pristine conditions, both HSRLs had excellent signal to noise performance yielding profiles for aerosol backscatter signals with precisions that are a small fraction of the Rayleigh signal. Error propagation is also much more localized than more common elastic backscatter lidar systems. It is for these reasons that there are active proponents of HSRL development for satellites, as evidenced with the recent EarthCARE launch. However, extinction and lidar ratio are calculated from the gradient in the molecular signal and are thus much more sensitive to noise. This is quite apparent in Fig. 4 when comparing backscatter vs. extinction products for typically marine to polluted conditions. The 10 s reported extinction products are over-sampled in a boxcar 60 s window leading to functionally a 10–12 km product (versus 1.75 km 10 s backscatter product), and in the vertical the 15 m reported product is itself computed over a 300 m vertical window, that we in addition performed additional averaging. In the context of large scale polluted conditions, the HSRL

provides a valuable degree of freedom in the derivation of lidar ratio. However, even with longer integration times and lengths, extinction profiles for typical conditions are nevertheless quite noisy as are associated lidar ratios for typical marine conditions and key exchange features are washed out. For the highest 532 nm AOTs measured for the clean marine conditions, RF15a on the western side of convection, lidar ratios in the 20–25 sr⁻¹ range are consistent with expected coarse mode, spherical sea-salt droplet dominated conditions (Burton et al., 2012), lending confidence to these measurements. Nevertheless, the lidar extinction profile, while still showing a mixed layer dominance, does not show the clean transitions as observed in the backscatter due to the coarser resolution and higher noise profile for the more tenuous aerosol. For other cases where aerosol backscatter in the MABL is a half (e.g. RF15 2) to a tenth (RF19 a) of the polluted cases, extinction profiles and lidar ratios are much more noisy and difficult to interpret—although the lidar ratio is still overall close to marine environments at 30 sr. Although aloft lidar ratios steadily rise, perhaps due to a lower prevalence of sea salt and higher prevalence of a fine mode (e.g. sea salt may have been preferentially scavenged/longer range transport of a fine mode/particle nucleation) or as a low extinction related artifact. Applying even a mean lidar ratio of 30 sr to a cleaner aerosol backscatter profile, 532 nm AOTs are much lower than what the extinction retrieval would suggest, down as low as 0.03 for RF19-1. What this suggests is that for cleaner or even more typical conditions and fine features, algorithm development should be more heavily weighted on projection of lidar ratios onto aerosol backscatter than specifically the extinction products more directly.

5. *3D variability and radiative effects*: The combined PISTON/CAMP²Ex observations are spawning studies focused on the features and physical processes related to the scales outlined above. Remote sensing will no doubt play a prominent role in quantifying such processes. Historically, cloud and aerosol populations are measured in a 2-dimensional product (e.g. cloud top heights, AOT, etc). With the recent advancement of higher-level technologies (e.g. polarimetry; hyperspectral; HSRL in space, better geostationary, etc), additional complexity will no doubt be able to be monitored. However, PISTON/CAMP²Ex demonstrated that just because aerosol transport is near the surface it is not necessarily simple. In fact, the opposite is true—the fine scales associated with near surface aerosol and cloud features have challenges beyond large mid and upper-level clouds and plumes. Despite the prevalence of attempts to correlate AOT with boundary layer aerosol mass or CCN populations, skill in such a regression or machine learning is highly localized. We find that

as expected, use of AOT as an indicator of particle concentrations in the MABL or cloud detrainment is problematic. Despite the highest concentrations of particles being in the MABL, just as in Reid et al. (2017) over land, the variability in AOT contributions above the MABL is what often dominates AOT variability (e.g. Fig. 10b). Low AOTs can still have extraordinarily high aerosol backscatter in the MABL and vice versa. Further, backscatter enhancements under clouds add a potential 3-dimensional effect for non-nadir optical paths, becoming even more complex for high wind shear in and around the MABL. All of this points to potential challenges of extracting aerosol information in and around clouds. Indeed, as pointed out in Marshak et al. (2021) there are significant radiance enhancements due to added radiative effects—notably cloud reflections. Spencer et al. (2019) showed AOT enhancements immediately adjacent to clouds, but Cloud Physics Lidar data they used was unable to resolve or penetrate under or immediately adjacent to clouds. It requires further study to determine under what circumstance uncertainties may be larger for any slant path application such as polarimetry or if the complexity of the system simply cancels out. Similar morphological considerations also need to be considered for clouds. A review of downward video with examples in Figs. 5 and 6 suggest aerosol detrainment is often associated with altocumulus clouds. These vertically thin but horizontally expansive clouds may be an appreciable fraction of what is perceived by satellites as cumulus clouds and are also difficult to replicate in all but the highest resolution LES models.

6. *Higher order inversions and hygroscopicity*: Questions regarding 3D radiative transfer immediately lead to questions of higher order retrievals, such as size, index of refraction and hygroscopicity. Particle hygroscopicity is a particularly important parameter to understand as water uptake in high relative humidity environments like the MABL is what largely modulates size, refractive index, and ultimately scattering. Notable from the onset of analysis of P-3 data was the finding of negative hygroscopicity within the in-situ instrumentation. The second question is to what extent we can estimate aerosol hygroscopicity from lidar profiles. Indeed, it has been previously hypothesized by Feingold and Morley (2003) that by assuming saturation at cloud base and an adiabatic lapse rate that then aerosol hygroscopicity can be inverted. More formal theoretical investigations on extraction of particle microphysics from lidar and polarimetry are also optimistic (Kahnert and Kanngießer, 2024). However, the authors here have long experimented with the SEAC⁴RS ground SSEC HSRL and CAMP²Ex P-3 HSRL-2 data for such an extraction and papers are forthcoming. To be sure, some-

thing can be said about particle hygroscopicity and other particle properties. Overall, RH increases systematically as expected in the mixed layer. Particle backscatter increases and linear depolarization decreases under clouds. Clearly, the particles observed in Figs. 6 and 13 are hygroscopic. However, there are several considerations to make an inference to quantitative data. First, the lidars and other remote sensors observe the ensemble of aerosol particles regardless of size. In a marine environment, there is always a mixture of particles with hygroscopic sea salt that must be controlled. Second, dry hygroscopic particles grow rapidly with RH. However, once hydrated, particles retain their moisture as they dry, known as hysteresis (e.g. Carrico et al., 2003). This “upper” hygroscopic curve is more uncertain, and shows less sensitivity to RH. In and around the MABL, it is often assumed that particles represent the upper part of the hygroscopic hysteresis curve. However, this is not always a good assumption; Ferrare et al. (2023) has noted that depolarizing sea salt is sometimes found around MABL in areas of dry air aloft. As particles never fully dry out (RH even above the MABL is 75 % as opposed to ~ 35 % often required for efflorescence) we are not currently in a position to compare quantitatively with the airborne measurements. Further, as particles grow and shrink, their lidar ratio should also change. Finally, we know from our findings that there is heterogeneity in the MABL itself and assuming a simple adiabatic lapse rate with altitude and uniform moisture might also be problematic. While as a whole the balloon sounding is “well-behaved” and depicts the MABL as we expect, it is nevertheless a single profile in an inhomogeneous and sheared environment. We do not know specifically if the sounding is predominately in an up-draft or downdraft region, or even the extent humidity changes both vertically and horizontally especially between clouds when airborne or spaceborne lidar observations are made. But, we do know below cloud base for even sea salt dominated environments that aerosol backscatter between clouds can be larger at 200 m than just below cloud base (e.g. Figs. 11–13). In comparisons, “bottom up” observations from ship samples in an Eulerian fashion suggest that small changes in distance or direction can result in differences in environments (Park et al., 2022). What is perhaps most uncertain, however, is the nature of hygroscopic growth, the hysteresis for these particles, and associated changes in lidar ratio that must be parsed within the ensemble of particles in the beam.

Data availability. The PISTON *Sally Ride* dataset can be found at the NASA Langley Atmospheric Data Center at <https://doi.org/10.5067/SUBORBITAL/PISTON2019-ONR-NOAA/RVSALLYRIDE/DATA001> (Moum, 2020) or at the NASA

Langley Airborne Science Data Repository (<https://www-air.larc.nasa.gov/cgi-bin/ArcView/camp2ex?RV-SALLY-RIDE=1>, last access: 2 December 2025). PISTON context can be gleaned from the mission site: <https://onrpiston.colostate.edu/> (last access: 2 December 2025). *Sally Ride* HSRL data can be visualized at: <https://hsrl.ssec.wisc.edu/> (last access: 2 December 2025). The CAMP²Ex data doi is <https://doi.org/10.5067/SUBORBITAL/CAMP2EX2018/DATA001> (Reid and Czech, 2019; Chen et al., 2020) with access available at <https://www-air.larc.nasa.gov/missions/camp2ex/index.html> (last access: 2 December 2025). Additional descriptive and satellite data can be found on the University of Wisconsin Geoworldview site (<https://geoworldview.ssec.wisc.edu/>, last access: 2 December 2025). Model data can be found on the NRL Global Oceans Data Assimilation Experiment site <https://usgodae.org> (last access: 2 December 2025).

Author contributions. Author JSR conceived the study, performed the bulk analysis, and was lead writer of this paper. Authors JIR and PX conducted NAAPS analysis, data processing and aided in the shear analysis. Authors CAH, RAF, and SPB participated in the field experiment (HSRL-2 team) and contributed to reviewing and editing of this paper. Authors REH, EWE, WJM, and RK developed the HSRL technology, calibrated/deployed/processed the SSECHSRL measurements for the deployment. Author EJT contributed to reviewing and editing of this paper, led data archiving of all PISTON datasets, and led data collection and processing of the shipboard near-surface oceanic and meteorological, air-sea flux, and ceilometer data. Authors HBM, DPE, and JHC provided financial and science support. Author SCvdH participated in the field experiment (PI for dropsondes; P-3 flight scientist), was a CAMP²Ex white paper co-author, and contributed to reviewing and editing of this paper. Other coauthors that provided data and editing include CGA (AMPR), GSD and JPD (Trace Gas) AS (AMS), KLT (State), JW (FIMS), LDZ (LARGE).

Competing interests. At least one of the (co-)authors is a member of the editorial board of *Atmospheric Chemistry and Physics*. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Special issue statement. This article is part of the special issue “Cloud, Aerosol and Monsoon Processes Philippines Experiment (CAMP²Ex) (ACP/AMT inter-journal SI)”. It is not associated with a conference.

Acknowledgements. CAMP²Ex could only be performed with the persistent effort and the cooperation of many Philippine entities including the Manila Observatory, PAGASA, the Department of Foreign Affairs (DFA), Philippine Department of Science and Technology (DOST), Civil Aviation Authorities of the Philippines (CAAP), and Armed Forces Philippines (AFP). We are also grateful to the US Embassy in Manila Economics and Science Department for shepherding the CAMP²Ex mission over many years of preparation. For PISTON, we thank collaborators in Taiwan, particularly Prof. Sen Jan and colleagues at National Taiwan University, and in Palau. The PISTON and CAMP²Ex deployments were an enormous logistical undertaking. Team members are most appreciative to the University of California San Diego Crew of *Sally Ride*, the NASA Wallops flight facility, and the NASA Earth Science Project Office.

Financial support. This research has been supported by the Office of Naval Research (grant no. N0001424WX02457), the NASA Headquarters (grant no. 80HQTR21T0089), and the US Naval Research Laboratory Base Program.

Review statement. This paper was edited by Geraint Vaughan and reviewed by two anonymous referees.

References

- Amiot, C. G., Biswas, S. K., Lang, T. J., and Duncan, D. I.: Dual-polarization deconvolution and geophysical retrievals from the Advanced Microwave Precipitation Radiometer during OLYMP-EX/RADEX, *J. Atmos. Oceanic Technol.*, 38, 607–628, <https://doi.org/10.1175/JTECH-D-19-0218.1>, 2021.
- Anderson, B. E., Grant, W. B., Gregory, G. L., Browell, E. V., Collins Jr., J. E., Sachse, G. W., Bagwell, D. R., Hudgins, C. H., Blake, D. R., and Blake, N. J.: Aerosols from biomass burning over the tropical South Atlantic region: distributions and impacts, *J. Geophys. Res.*, 101, 24117–24137, <https://doi.org/10.1029/96JD00717>, 1996.
- Andreae, M. O.: Soot carbon and excess fine potassium: long-range transport of combustion-derived aerosols, *Science*, 220, 1148–1151, <https://doi.org/10.1126/science.220.4602.1148>, 1983.
- Braun, R. A., Aghdam, M. A., Bañaga, P. A., Betito, G., Cambaliza, M. O., Cruz, M. T., Lorenzo, G. R., MacDonald, A. B., Simpas, J. B., Stahl, C., and Sorooshian, A.: Long-range aerosol transport and impacts on size-resolved aerosol composition in Metro Manila, Philippines, *Atmos. Chem. Phys.*, 20, 2387–2405, <https://doi.org/10.5194/acp-20-2387-2020>, 2020.
- Bukowski, J., Posselt, D. J., Reid, J. S., and Atwood, S. A.: Modes of vertical thermodynamic and wind variability over the Maritime Continent, *Atmos. Chem. Phys.*, 17, 4611–4626, <https://doi.org/10.5194/acp-17-4611-2017>, 2017.
- Burton, S. P., Ferrare, R. A., Hostetler, C. A., Hair, J. W., Rogers, R. R., Obland, M. D., Butler, C. F., Cook, A. L., Harper, D. B., and Froyd, K. D.: Aerosol classification using airborne High Spectral Resolution Lidar measurements – methodology and examples, *Atmos. Meas. Tech.*, 5, 73–98, <https://doi.org/10.5194/amt-5-73-2012>, 2012.
- Burton, S. P., Hair, J. W., Kahnert, M., Ferrare, R. A., Hostetler, C. A., Cook, A. L., Harper, D. B., Berkoff, T. A., Seaman, S. T., Collins, J. E., Fenn, M. A., and Rogers, R. R.: Observations of the spectral dependence of linear particle depolarization ratio of aerosols using NASA Langley airborne High Spectral Resolution Lidar, *Atmos. Chem. Phys.*, 15, 13453–13473, <https://doi.org/10.5194/acp-15-13453-2015>, 2015.
- Burton, S. P., Hostetler, C. A., Cook, A. L., Hair, J. W., Seaman, T., Scola, S., Harper, D. B., Smith, J. A., Fenn, M. A., Ferrare, R. A., Saide, P. E., Chemyakin, E. V., and Müller, D.: “Calibration of a high spectral resolution lidar using a Michelson interferometer, with data examples from ORACLE S.” *Appl. Optics*, 57, 6061–6075, 2018.
- Campbell, J. R. Reid, J. S. Westphal, D. L. Zhang, J. Tackett, J. L. Chew, B. N. Welton, E. J. Shimizu A., and Sugimoto, N.: Characterizing aerosol particle composition and the vertical profile of extinction and linear depolarization over Southeast Asia and the Maritime Continent: the 2007–2009 view from CALIOP, *Atmos. Res.*, 122, 520–543, <https://doi.org/10.1016/j.atmosres.2012.05.007>, 2013.
- Carrico, C. M., Kus, P., Rood, M. J., Quinn, P. K., and Bates T. S.: Mixtures of pollution, dust, sea salt, and volcanic aerosol during ACE-Asia: radiative properties as a function of relative humidity, *J. Geophys. Res.*, 108, 8650, <https://doi.org/10.1029/2003JD003405>, 2003.
- Chew, B. N. Campbell, J. R. V., Salinas, S. V. Chang, C. W. Reid, J. S. Welton, E. J. Holben, B. N., and Liew, S. C.: Aerosol particle vertical distributions and optical properties over Singapore, *Atmos. Environ.*, 79, 599–613, <https://doi.org/10.1016/j.atmosenv.2013.06.026>, 2013.
- Clarke, A. D., Ki, Z., and Litchy, M.: Aerosol dynamics in the equatorial Pacific marine boundary layer: microphysics, diurnal cycles and entrainment, *Geophys. Res. Lett.*, 23, 733–736, <https://doi.org/10.1029/96GL00778>, 1996.
- Clarke, A. D., Freitag, S., Simpson, R. M. C., Hudson, J. G., Howell, S. G., Brekhovskikh, V. L., Campos, T., Kapustin, V. N., and Zhou, J.: Free troposphere as a major source of CCN for the equatorial Pacific boundary layer: long-range transport and teleconnections, *Atmos. Chem. Phys.*, 13, 7511–7529, <https://doi.org/10.5194/acp-13-7511-2013>, 2013.
- Crosbie, E., Ziemba, L. D., Shook, M. A., Robinson, C. E., Winstead, E. L., Thornhill, K. L., Braun, R. A., MacDonald, A. B., Stahl, C., Sorooshian, A., van den Heever, S. C., DiGangi, J. P., Diskin, G. S., Woods, S., Bañaga, P., Brown, M. D., Gallo, F., Hilario, M. R. A., Jordan, C. E., Leung, G. R., Moore, R. H., Sanchez, K. J., Shingler, T. J., and Wiggins, E. B.: Measurement report: Closure analysis of aerosol–cloud composition in tropical maritime warm convection, *Atmos. Chem. Phys.*, 22, 13269–13302, <https://doi.org/10.5194/acp-22-13269-2022>, 2022.
- Cruz, M. T., Bañaga, P. A., Betito, G., Braun, R. A., Stahl, C., Aghdam, M. A., Cambaliza, M. O., Dadashazar, H., Hilario, M. R., Lorenzo, G. R., Ma, L., MacDonald, A. B., Pabroa, P. C., Yee, J. R., Simpas, J. B., and Sorooshian, A.: Size-resolved composition and morphology of particulate matter during the southwest monsoon in Metro Manila, Philippines, *Atmos. Chem. Phys.*, 19, 10675–10696, <https://doi.org/10.5194/acp-19-10675-2019>, 2019.
- DeCarlo, P. F., Dunlea, E. J., Kimmel, J. R., Aiken, A. C., Sueper, D., Crouse, J., Wennberg, P. O., Emmons, L., Shi-

- nozuka, Y., Clarke, A., Zhou, J., Tomlinson, J., Collins, D. R., Knapp, D., Weinheimer, A. J., Montzka, D. D., Campos, T., and Jimenez, J. L.: Fast airborne aerosol size and chemistry measurements above Mexico City and Central Mexico during the MILAGRO campaign, *Atmos. Chem. Phys.*, 8, 4027–4048, <https://doi.org/10.5194/acp-8-4027-2008>, 2008.
- Dobracki, A., Lewis, E. R., Sedlacek III, A. J., Tatro, T., Zawad-owicz, M. A., and Zuidema, P.: Burning conditions and trans- portation pathways determine biomass-burning aerosol prop- erties in the Ascension Island marine boundary layer, *Atmos. Chem. Phys.*, 25, 2333–2363, <https://doi.org/10.5194/acp-25-2333-2025>, 2025.
- Drager, A. J. and van den Heever, S. C.: Characterizing convective cold pools, *J. Adv. Model. Earth Sy.*, 9, 1091–1115, 2017.
- Eck, T. F., Holben, B. N., Reid, J. S., Arola, A., Ferrare, R. A., Hostetler, C. A., Crumeyrolle, S. N., Berkoff, T. A., Welton, E. J., Lolli, S., Lyapustin, A., Wang, Y., Schafer, J. S., Giles, D. M., Anderson, B. E., Thornhill, K. L., Minnis, P., Pickering, K. E., Loughner, C. P., Smirnov, A., and Sinyuk, A.: Observations of rapid aerosol optical depth enhancements in the vicinity of pol- luted cumulus clouds, *Atmos. Chem. Phys.*, 14, 11633–11656, <https://doi.org/10.5194/acp-14-11633-2014>, 2014.
- Eck, T. F., Holben, B. N., Giles, D. M., Slutsker, I., Sinyuk, A., Schafer, J. S., Smirnov, A., Sorokin, M., Reid, J. S., Sayer, A. M., Hsu, N. C., Shi, Y. R., Levy, R. C., Lyapustin, A., Rah- man, M. A., Liew, S. C., Salinas Cortijo, S. V., Li, T., Kalber- matter, D., Keong, K. L., Yuggotomo, M. E., Aditya, F., Mo- hamad, M., Mahmud, M., Chong, T. K., Lim, H.-S., Choon, Y. E., Deranadyan, G., Kusumaningtyas, S. D. A., and Aldrian, E.: AERONET remotely sensed measurements and retrievals of biomass burning aerosol optical properties during the 2015 In- donesian burning season, *J. Geophys. Res.-Atmos.*, 124, 4722–4740. <https://doi.org/10.1029/2018JD030182>, 2019.
- Eck, T. F., Holben, B. N., Reid, J. S., Sinyuk, A., Giles, D. M., Arola, A., Slutsker, I., Schafer, J. S., Sorokin, M. G., Smirnov, A., LaRosa, A. D., Kraft, J., Reid, E. A., O'Neill, N. T., Welton, E. J. Menendez, A. R. (2023) The extreme forest fires in California/Oregon in 2020: Aerosol optical and physical properties and comparisons of aged versus fresh smoke, *Atmos. Environ.*, 305, 119798, <https://doi.org/10.1016/j.atmosenv.2023.119798>, 2023.
- Edson, J. B., Raju, V. S., Weller, R. A., Bigorre, S., Pluedde- mann, A., Fairall, C. W., Miller, S., Mahrt, L., Vickers, D., and Hersbach, H.: On the exchange of momentum over the open ocean, *J. Phys. Oceanogr.*, 43, 1589–1610, 2013.
- Edwards, E.-L., Reid, J. S., Xian, P., Burton, S. P., Cook, A. L., Crosbie, E. C., Fenn, M. A., Ferrare, R. A., Freeman, S. W., Hair, J. W., Harper, D. B., Hostetler, C. A., Robinson, C. E., Scarino, A. J., Shook, M. A., Sokolowsky, G. A., van den Heever, S. C., Winstead, E. L., Woods, S., Ziemba, L. D., and Sorooshian, A.: Assessment of NAAPS-RA performance in Maritime Southeast Asia during CAMP²Ex, *Atmos. Chem. Phys.*, 22, 12961–12983, <https://doi.org/10.5194/acp-22-12961-2022>, 2022.
- Eloranta, E. W.: High spectral resolution lidar, in: Lidar: Range Resolved Optical Remote Sensing of the Atmosphere, edited by: Weitkamp, K., Springer, New York, 143–163, https://doi.org/10.1007/0-387-25101-4_5, 2005.
- Fairall, C. W., Bradley, E. F., Godfrey, J. S., Wick, G. A. Ed- son, J. B., and Young, G. S.: Cool-skin and warm-layer effects on sea surface temperature, *J. Geophys. Res.*, 101, 1295–1308, <https://doi.org/10.1029/95JC03190>, 1996.
- Fairall, C. W., White, A. B., Edson, J. B., and Hare, J. E.: Integrated shipboard measurements of the marine boundary layer, *J. Atmos. Ocean. Tech.*, 14, 338–359, 1997.
- Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Ed- son, J. B.: Bulk parameterization of air-sea fluxes: updates and verification for the COARE algorithm, *J. Climate*, 16, 571–591, 2003.
- Feingold, G. and Morley, B.: Aerosol hygroscopic prop- erties as measured by lidar and comparison with in situ measurements, *J. Geophys. Res.*, 108, 4327, <https://doi.org/10.1029/2002JD002842>, 2003.
- Ferrare, R., Hair, J., Hostetler, C., Shingler, T., Burton, S. P., Fenn, M., Clayton, M., Scarino, A. J., Harper, D., Seaman, S., Cook, A., Crosbie, E., Winstead, E., Ziemba, L., Thorn- hill, L., Robinson, C., Moore, R., Vaughan, M., Sorooshian, A., Schlosser, J. S., Liu, H., Zhang, B., Diskin, G., DiGangi, J., Nowak, J., Choi, Y., Zuidema, P., and Chellappan, S.: Airborne HSRL-2 measurements of elevated aerosol depolarization asso- ciated with non-spherical sea salt, *Frontiers in Remote Sensing*, 4, <https://doi.org/10.3389/frsen.2023.1143944>, 2023.
- Field, R. D., and Shen, S. S. P.: Predictability of carbon emis- sions from biomass burning in Indonesia, *J. Geophys. Res.*, 113, G04024, <https://doi.org/10.1029/2008JG000694>, 2008.
- Fromm, M., Lindsey, D. T., Servranckx, R. Yue, G. E. Trickl, T. Sica, R. Doucet, P. and Godin-Beekmann, S.: The untold story of pyrocumulonimbus, *B. Am. Meteorol. Soc.*, 91, 1193–1210 <https://doi.org/10.1175/2010BAMS3004.1>, 2010.
- Froyd, K. D., Murphy, D. M., Brock, C. A., Campuzano-Jost, P., Dibb, J. E., Jimenez, J.-L., Kupc, A., Middlebrook, A. M., Schill, G. P., Thornhill, K. L., Williamson, C. J., Wilson, J. C., and Ziemba, L. D.: A new method to quantify mineral dust and other aerosol species from aircraft platforms using single- particle mass spectrometry, *Atmos. Meas. Tech.*, 12, 6209–6239, <https://doi.org/10.5194/amt-12-6209-2019>, 2019.
- Ge, C. J., Wang, J. S., Reid, D. J., Posselt, P., Xian, and Hyer, E.: Mesoscale modeling of smoke transport from equa- torial Southeast Asian Maritime Continent to the Philip- pines: first comparison of ensemble analysis with in situ observations, *J. Geophys. Res.-Atmos.*, 122, 5380–5398, <https://doi.org/10.1002/2016JD026241>, 2017.
- Goldstein, J. E., Graham, L., Ansori, S., Vetruta, Y., Thomas, A., Ap- plegate, G., Vayda, A. P., Saharjo, B. H., and Cochrane, M. A.: Beyond slash-and-burn: the roles of human activities, altered hydrology and fuels in peat fires in Central Kalimantan, In- donesia, *Singapore Journal of Tropical Geography*, 41, 190–208, <https://doi.org/10.1111/sjtg.12319>, 2020.
- Grant, L. D., and van den Heever, S. C.: Cold pool and precipitation responses to aerosol loading: modulation by dry layers, *J. Atmos. Sci.*, 72, 1398–1408, 2015.
- Griffing, G. W.: Relations between the prevailing visibility, neph- elometer scattering coefficient and sunphotometer turbidity co- efficient, *Atmos. Environ.*, 14, 577–584, 1980.
- Hair, J. W., Hostetler, C. A., Cook, A. L., Harper, D. B., Fer- rare, R. A., Mack, T. L., Welch, W., Izquierdo, L. R., and Hovis F. E.: Airborne high spectral resolution lidar for pro- filing aerosol optical properties, *Appl. Optics*, 47, 6734–6752, <https://doi.org/10.1364/AO.47.006734>, 2008.

- Haywood, J. M., Abel, S. J., Barrett, P. A., Bellouin, N., Blyth, A., Bower, K. N., Brooks, M., Carslaw, K., Che, H., Coe, H., Cotterell, M. I., Crawford, I., Cui, Z., Davies, N., Dingley, B., Field, P., Formenti, P., Gordon, H., de Graaf, M., Herbert, R., Johnson, B., Jones, A. C., Langridge, J. M., Malavelle, F., Partridge, D. G., Peers, F., Redemann, J., Stier, P., Szpek, K., Taylor, J. W., Watson-Parris, D., Wood, R., Wu, H., and Zuidema, P.: The CLoud–Aerosol–Radiation Interaction and Forcing: Year 2017 (CLARIFY-2017) measurement campaign, *Atmos. Chem. Phys.*, 21, 1049–1084, <https://doi.org/10.5194/acp-21-1049-2021>, 2021.
- Hilario, M. R. A., Cruz, M. T., Cambaliza, M. O. L., Reid, J. S., Xian, P., Simpas, J. B., Lagrosas, N. D., Uy, S. N. Y., Cliff, S., and Zhao, Y.: Investigating size-segregated sources of elemental composition of particulate matter in the South China Sea during the 2011 *Vasco* cruise, *Atmos. Chem. Phys.*, 20, 1255–1276, <https://doi.org/10.5194/acp-20-1255-2020>, 2020.
- Hilario, M. R. A., Crosbie, E., Shook, M., Reid, J. S., Cambaliza, M. O. L., Simpas, J. B. B., Ziemba, L., DiGangi, J. P., Diskin, G. S., Nguyen, P., Turk, F. J., Winstead, E., Robinson, C. E., Wang, J., Zhang, J., Wang, Y., Yoon, S., Flynn, J., Alvarez, S. L., Behrangi, A., and Sorooshian, A.: Measurement report: Long-range transport patterns into the tropical northwest Pacific during the CAMP²Ex aircraft campaign: chemical composition, size distributions, and the impact of convection, *Atmos. Chem. Phys.*, 21, 3777–3802, <https://doi.org/10.5194/acp-21-3777-2021>, 2021.
- Hilario, M. R. A., Barth, M., Bennett, R., Crosbie, E., DiGangi, J. P., Diskin, G. S., Lorenzo, G. R., Rutledge, S., Martin, M. Y., and Ziemba, L.: Quantifying scavenging efficiencies of different aerosol species and size-resolved volume concentrations in tropical convective clouds over the West Pacific, *J. Atmos. Sci.*, 82, 267–282, <https://doi.org/10.1175/JAS-D-24-0064.1>, 2025.
- Hogan, T. F., Liu, M., Ridout, J. A., Peng, M. S., Whitcomb, T. R., Ruston, B. C., Reynolds, C. A., Eckermann, S. D., Moskaitis, J. R., Baker, N. L., McCormack, J. P., Viner, K. C., McLay, J. G., Flatau, M. K., Xu, L., Chen, C., and Chang, S. W.: The Navy Global Environmental Model, *Oceanography* 27, 116–125, <https://doi.org/10.5670/oceanog.2014.73>, 2014.
- Hu, Q. H., Xie, Z. Q., Wang, X. M., Kang, H., and Zhang, P.: Levoglucosan indicates high levels of biomass burning aerosols over oceans from the Arctic to Antarctic, *Sci. Rep.*, 3, 3119, <https://doi.org/10.1038/srep03119>, 2013.
- Husar, R. B., Husar, J. D., Martin L.: Distribution of continental surface aerosol extinction based on visual range data *Atmos. Environ.*, 34, 5067–5078, 2000.
- Jiang, X., Li, T., and Wang B.: Structures and mechanisms of the northward propagating boreal summer intraseasonal oscillation, *J. Climate*, 17, 1022–1039, [https://doi.org/10.1175/1520-0442\(2004\)017<1022:SAMOTN>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<1022:SAMOTN>2.0.CO;2), 2004.
- Kahn, R. A., Chen, Y., Nelson, D. L., Leung, F.-Y., Li, Q., Diner, D. J., and Logan, J. A.: Wildfire smoke injection heights: two perspectives from space, *Geophys. Res. Lett.*, 35, L04809, <https://doi.org/10.1029/2007GL032165>, 2008.
- Kahnert, M. and Kanngießer, F.: Optical characterization of marine aerosols using a morphologically realistic model with varying water content: implications for lidar applications and passive polarimetric remote sensing, *Geophys. Res. Lett.*, 51, e2023GL107541. <https://doi.org/10.1029/2023GL107541>, 2024.
- Khan, A. L., Klein, A. G., Katich, J. M., and Xian, P.: Local emissions and regional wildfires influence refractory black carbon observations near Palmer Station, Antarctica, *Front. Earth Sci.*, 7, 1–8, <https://doi.org/10.3389/feart.2019.00049>, 2019.
- Kunkel, K. E., Eloranta, E. W., and Shipley, S. T.: Lidar observations of the convective boundary layer, *J. Appl. Meteorol.*, 16, 1306–1311, [https://doi.org/10.1175/1520-0450\(1977\)016<1306:LOOTCB>2.0.CO;2](https://doi.org/10.1175/1520-0450(1977)016<1306:LOOTCB>2.0.CO;2), 1977.
- Labonne, M., Bréon, F.-M., and Chevallier, F.: Injection height of biomass burning aerosols as seen from a spaceborne lidar, *Geophys. Res. Lett.*, 34, L11806, <https://doi.org/10.1029/2007GL029311>, 2007.
- Lang, T., Amiot, C., and Biswas, S.: AMPR CAMP2Ex, calibrated and quality-controlled flight dataset, level 2B, revision 1, NASA Langley Research Center [data set], <https://www-air.larc.nasa.gov/cgi-bin/ArcView/camp2ex> (last access: 2 December 2025), 2021.
- Lemone, M. A.: Structure and dynamics of horizontal roll vortices in planetary boundary-layer, *J. Atmos. Sci.*, 30, 1077–1091, [https://doi.org/10.1175/1520-0469\(1973\)030<1077:Tsadoh>2.0.Co;2](https://doi.org/10.1175/1520-0469(1973)030<1077:Tsadoh>2.0.Co;2), 1973.
- Lorenzo, G. R., Arellano, A. F., Cambaliza, M. O., Castro, C., Cruz, M. T., DiGangi, J. P., Gacal, G. F., Hilario, M. R. A., Lagrosas, N., Ong, H. J., Simpas, J. B., Uy, S. N., and Sorooshian, A.: An emerging aerosol climatology via remote sensing over Metro Manila, the Philippines, *Atmos. Chem. Phys.*, 23, 10579–10608, <https://doi.org/10.5194/acp-23-10579-2023>, 2023.
- Lorenzo, G. R., Ziemba, L. D., Arellano, A. F., Barth, M. C., Crosbie, E. C., DiGangi, J. P., Diskin, G. S., Ferrare, R., Hilario, M. R. A., Shook, M. A., Tilmes, S., Wang, J., Xiao, Q., Zhang, J., and Sorooshian, A.: Measurement report: Characterization of aerosol hygroscopicity over Southeast Asia during the NASA CAMP2Ex campaign, *Atmos. Chem. Phys.*, 25, 5469–5495, <https://doi.org/10.5194/acp-25-5469-2025>, 2025.
- Lynch, P., Reid, J. S., Westphal, D. L., Zhang, J., Hogan, T. F., Hyer, E. J., Curtis, C. A., Hegg, D. A., Shi, Y., Campbell, J. R., Rubin, J. I., Sessions, W. R., Turk, F. J., and Walker, A. L.: An 11-year global gridded aerosol optical thickness reanalysis (v1.0) for atmospheric and climate sciences, *Geosci. Model Dev.*, 9, 1489–1522, <https://doi.org/10.5194/gmd-9-1489-2016>, 2016.
- Mallet, M., Nabat, P., Zuidema, P., Redemann, J., Sayer, A. M., Stengel, M., Schmidt, S., Cochrane, S., Burton, S., Ferrare, R., Meyer, K., Saide, P., Jethva, H., Torres, O., Wood, R., Saint Martin, D., Roehrig, R., Hsu, C., and Formenti, P.: Simulation of the transport, vertical distribution, optical properties and radiative impact of smoke aerosols with the ALADIN regional climate model during the ORACLES-2016 and LASIC experiments, *Atmos. Chem. Phys.*, 19, 4963–4990, <https://doi.org/10.5194/acp-19-4963-2019>, 2019.
- Marshak, A., Ackerman, A., da Silva, A. M., Eck, T., Holben, B., Kahn, R., Kleidman, R., Knobelspie, K., Levy, R., Lyapustin, A., Oreopoulos, L., Remer, L., Torres, O., Várnai, T., Wen, G., and Yorks, J.: Aerosol Properties in Cloudy Environments from Remote Sensing Observations: A Review of the Current State of Knowledge, *B. Am. Meteorol. Soc.*, 102, E2177–E2197, <https://doi.org/10.1175/BAMS-D-20-0225.1>, 2021.
- Moeng, C. H. and Sullivan, P. P.: A comparison of shear-driven and buoyancy-driven planetary boundary-layer Flows,

- J. Atmos. Sci., 51, 999–1022, [https://doi.org/10.1175/1520-0469\(1994\)051<0999:Acosab>2.0.Co;2](https://doi.org/10.1175/1520-0469(1994)051<0999:Acosab>2.0.Co;2), 1994.
- Moore, R. H., Wiggins, E. B., Ahern, A. T., Zimmerman, S., Montgomery, L., Campuzano Jost, P., Robinson, C. E., Ziemba, L. D., Winstead, E. L., Anderson, B. E., Brock, C. A., Brown, M. D., Chen, G., Crosbie, E. C., Guo, H., Jimenez, J. L., Jordan, C. E., Lyu, M., Nault, B. A., Rothfuss, N. E., Sanchez, K. J., Schueneman, M., Shingler, T. J., Shook, M. A., Thornhill, K. L., Wagner, N. L., and Wang, J.: Sizing response of the Ultra-High Sensitivity Aerosol Spectrometer (UHSAS) and Laser Aerosol Spectrometer (LAS) to changes in submicron aerosol composition and refractive index, *Atmos. Meas. Tech.*, 14, 4517–4542, <https://doi.org/10.5194/amt-14-4517-2021>, 2021.
- Morrison, H.: Entrainment, detrainment, and dilution of dry and moist atmospheric thermals, *J. Atmos. Sci.*, 82, 361–389, 2025.
- Moser, D. H. and Lasher-Trapp, S.: The influence of successive thermals on entrainment and dilution in a simulated cumulus congestus, *J. Atmos. Sci.*, 74, 375–392, 2017.
- Moum, J.: PISTON 2019 Research Vessel (RV) Sally Ride Data, NASA Langley Atmospheric Science Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/SUBORBITAL/PISTON2019-ONR-NOAA/RVSALLYRIDE/DATA001>, 2020.
- Nichol, J.: Smoke haze in Southeast Asia: a predictable recurrence, *Atmos. Environ.*, 32, 2715–2716, 1998.
- Park, H. J., Reid, J. S., Freire, L. S., Jackson, C., and Richter, D. H.: In situ particle sampling relationships to surface and turbulent fluxes using large eddy simulations with Lagrangian particles, *Atmos. Meas. Tech.*, 15, 7171–7194, <https://doi.org/10.5194/amt-15-7171-2022>, 2022.
- Paugam, R., Wooster, M., Freitas, S., and Val Martin, M.: A review of approaches to estimate wildfire plume injection height within large-scale atmospheric chemical transport models, *Atmos. Chem. Phys.*, 16, 907–925, <https://doi.org/10.5194/acp-16-907-2016>, 2016.
- Quinn, P., Coffman, D., Johnson, J., Upchurch, L. M., Bates, T. S.: Small fraction of marine cloud condensation nuclei made up of sea spray aerosol, *Nature Geosci.*, 10, 674–679, <https://doi.org/10.1038/ngeo3003>, 2017.
- Radke, L. F. and Hobbs, P. V.: Humidity and particle fields around some small cumulus clouds, *J. Atmos. Sci.*, 48, 1190–1193, 1991.
- Reid, J. and Czech, E.: Clouds, Aerosol and Monsoon Processes-Philippines Experiment Data, NASA Langley Atmospheric Science Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/SUBORBITAL/CAMP2EX2018/DATA001>, 2019.
- Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmos. Chem. Phys.*, 5, 799–825, <https://doi.org/10.5194/acp-5-799-2005>, 2005.
- Reid, J. S., Hyer, E. J., Johnson, R. S., Holben, B. N., Yokelson, R. J., Zhang, J., Campbell, J. R., Christopher, S. A., Di Girolamo, L., Giglio, L., Holz, R. E., Kearney, C., Miettinen, J., Reid, E. A., Turk, F. J., Wang, J., Xian, P., Zhao, G., Balasubramanian, R., Chew, B. N., Janjai, S., Lagrosas, N., Lestari, P., Lin, N. H., Mahmud, M., Nguyen, A. X., Norris, B., Oanh, N. T. K., Oo, M., Salinas, S. V., Welton, E. J., and Liew, S. C.: Observing and understanding the Southeast Asian aerosol system by remote sensing: an initial review and analysis for the Seven Southeast Asian Studies (7SEAS) program, *Atmos. Res.*, 122, 403–468, <https://doi.org/10.1016/j.atmosres.2012.06.005>, 2013.
- Reid, J. S., Lagrosas, N. D., Jonsson, H. H., Reid, E. A., Sessions, W. R., Simpas, J. B., Uy, S. N., Boyd, T. J., Atwood, S. A., Blake, D. R., Campbell, J. R., Cliff, S. S., Holben, B. N., Holz, R. E., Hyer, E. J., Lynch, P., Meinardi, S., Posselt, D. J., Richardson, K. A., Salinas, S. V., Smirnov, A., Wang, Q., Yu, L., and Zhang, J.: Observations of the temporal variability in aerosol properties and their relationships to meteorology in the summer monsoonal South China Sea/East Sea: the scale-dependent role of monsoonal flows, the Madden–Julian Oscillation, tropical cyclones, squall lines and cold pools, *Atmos. Chem. Phys.*, 15, 1745–1768, <https://doi.org/10.5194/acp-15-1745-2015>, 2015.
- Reid, J. S., Lagrosas, N. D., Jonsson, H. H., Reid, E. A., Atwood, S. A., Boyd, T. J., Ghate, V. P., Xian, P., Posselt, D. J., Simpas, J. B., Uy, S. N., Zaiger, K., Blake, D. R., Bucholtz, A., Campbell, J. R., Chew, B. N., Cliff, S. S., Holben, B. N., Holz, R. E., Hyer, E. J., Kreidenweis, S. M., Kuciauskas, A. P., Lolli, S., Oo, M., Perry, K. D., Salinas, S. V., Sessions, W. R., Smirnov, A., Walker, A. L., Wang, Q., Yu, L., Zhang, J., and Zhao, Y.: Aerosol meteorology of Maritime Continent for the 2012 7SEAS southwest monsoon intensive study – Part 2: Philippine receptor observations of fine-scale aerosol behavior, *Atmos. Chem. Phys.*, 16, 14057–14078, <https://doi.org/10.5194/acp-16-14057-2016>, 2016.
- Reid, J. S., Kuehn, R. E., Holz, R. E., Elooranta, E. W., Kaku, K. C., Kuang, S., Newchurch, M. J., Thompson, A. M., Trepte, C. R., Zhang, J., Atwood, S. A., Hand, J. L., Holben, B. N., Minnis, P., and Posselt, D. J.: ground-based high spectral resolution lidar observation of aerosol vertical distribution in the summertime Southeast United States, *J. Geophys. Res.-Atmos.*, 122, 2970–3004, <https://doi.org/10.1002/2016JD025798>, 2017.
- Reid, J. S., Posselt, D. J., Kaku, K., Holz, R. A., Chen, G., Elooranta, E. W., Kuehn, R. E., Woods, S., Zhang, J., Anderson, B., Bui, T. P., Diskin, G. S., Minnis, P., Newchurch, M. J., Tanelli, S., Trepte, C. R., Thornhill, K. L., and Ziemba, L. D.: Observations and hypotheses related to low to middle free tropospheric aerosol, water vapor and altocumulus cloud layers within convective weather regimes: a SEAC⁴RS case study, *Atmos. Chem. Phys.*, 19, 11413–11442, <https://doi.org/10.5194/acp-19-11413-2019>, 2019.
- Reid, J. S., Gumber, A., Zhang, J., Holz, R. E., Rubin, J. I., Xian, P., Smirnov, A., Eck, T. F., O’Neill, N. T., Levy, R. C., Reid, E. A., Colarco, P. R., Benedetti, A., and Tanaka, T.: A coupled evaluation of operational MODIS and model aerosol products for maritime environments using sun photometry: evaluation of the fine and coarse mode, *Remote Sens.-Basel*, 14, 2978, <https://doi.org/10.3390/rs14132978>, 2022.
- Reid, J. S., Maring, H. B., Narisma, G. T., van den Heever, S., Girolamo, L., Ferrare, R., Lawson, P., Mace, G. G., Simpas, J. B., Tanelli, S., Ziemba, L., van Diedenhoven, B., Bruintjes, R., Bucholtz, A., Cairns, B., Cambaliza, M. O., Chen, G., Diskin, G. S., Flynn, J. H., Hostetler, C. A., Holz, R. E., Lang, T. J., Schmidt, K. S., Smith, G., Sorooshian, A., Thompson, E. J., Thornhill, K. L., Trepte, C., Wang, J., Woods, S., Yoon, S., Alexandrov, M., Alvarez, S., Amiot, C. G., Bennett, J. R., Brooks, M., Burton, S. P., Cayanan, E., Chen, H., Collow, A., Crosbie, E., DaSilva, A., DiGangi, J. P., Flagg, D. D., Freeman, S. W., Fu, D., Fukada, E., Hilario, M. R. A., Hong, Y., Hristova-Veleva, S. M., Kuehn, R.,

- Kowch, R. S., Leung, G. R., Loveridge, J., Meyer, K., Miller, R. M., Montes, M. J., Moum, J. N., Nenes, A., Nesbitt, S. W., Norgren, M., Nowottnick, E. P., Rauber, R. M., Reid, E. A., Rutledge, S., Schlosser, J. S., Sekiyama, T. T., Shook, M. A., Sokolowsky, G. A., Starnes, S. A., Tanaka, T. Y., Wasilewski, A., Xian, P., Xiao, Q., Xu, Z., and Zavaleta, J.: The Coupling Between Tropical Meteorology, Aerosol Lifecycle, Convection, and Radiation during the Cloud, Aerosol and Monsoon Processes Philippines Experiment (CAMP2Ex), *B. Am. Meteorol. Soc.*, 104, E1179–E1205, <https://doi.org/10.1175/BAMS-D-21-0285.1>, 2023.
- Ross, A. D., Holz, R. E., Quinn, G., Reid, J. S., Xian, P., Turk, F. J., and Posselt, D. J.: Exploring the first aerosol indirect effect over Southeast Asia using a 10-year collocated MODIS, CALIOP, and model dataset, *Atmos. Chem. Phys.*, 18, 12747–12764, <https://doi.org/10.5194/acp-18-12747-2018>, 2018.
- Rutledge, S. A., Chandrasekar, V., Fuchs, B., George, J., Junyent, F., Dolan, B., Kennedy, P. C., and Drushka, K.: SEA-POL Goes to Sea, *B. Am. Meteorol. Soc.*, 100, 2285–2301, <https://doi.org/10.1175/BAMS-D-18-0233.1>, 2019.
- Salesky, S. T., Chamecki, M., and Bou-Zeid, E.: On the nature of the transition between roll and cellular organization in the convective boundary layer, *Bound.-Lay. Meteorol.*, 163, 41–68, <https://doi.org/10.1007/s10546-016-0220-3>, 2017.
- Savre, J.: What controls local entrainment and detrainment rates in simulated shallow convection?, *J. Atmos. Sci.*, 79, 3065–3082, 2022.
- Sawamura, P., Moore, R. H., Burton, S. P., Chemyakin, E., Müller, D., Kolgotin, A., Ferrare, R. A., Hostetler, C. A., Ziemba, L. D., Beyersdorf, A. J., and Anderson, B. E.: HSRL-2 aerosol optical measurements and microphysical retrievals vs. airborne in situ measurements during DISCOVER-AQ 2013: an intercomparison study, *Atmos. Chem. Phys.*, 17, 7229–7243, <https://doi.org/10.5194/acp-17-7229-2017>, 2017.
- Scarino, A. J., Obland, M. D., Fast, J. D., Burton, S. P., Ferrare, R. A., Hostetler, C. A., Berg, L. K., Lefer, B., Haman, C., Hair, J. W., Rogers, R. R., Butler, C., Cook, A. L., and Harper, D. B.: Comparison of mixed layer heights from airborne high spectral resolution lidar, ground-based measurements, and the WRF-Chem model during CalNex and CARES, *Atmos. Chem. Phys.*, 14, 5547–5560, <https://doi.org/10.5194/acp-14-5547-2014>, 2014.
- Shingler, T., Sorooshian, A., Ortega, A., Crosbie, E., Wonaschütz, A., Perring, A. E., Beyersdorf, A., Ziemba, L., Jimenez, J. L., Campuzano-Jost, P., Mikoviny, T., Wisthaler, A., Russell, L. M.: Ambient observations of hygroscopic growth factor and $f(\text{RH})$ below 1: case studies from surface and airborne measurements, *J. Geophys. Res.-Atmos.*, 121, 13661–13677, <https://doi.org/10.1002/2016JD025471>, 2016.
- Sinha, P., Jaeglé, L., Hobbs, P. V., and Liang, Q.: Transport of biomass burning emissions from southern Africa, *J. Geophys. Res.*, 109, D20204, <https://doi.org/10.1029/2004JD005044>, 2004.
- Sobel, A. H., Sprintall, J., Maloney, E. D., Martin, Z. K., Wang, S., de Szoeko, S. P., Trabling, B. C., and Rutledge, S. A.: Large-scale state and evolution of the atmosphere and ocean during PISTON 2018, *J. Climate*, 34, 5017–5035, <https://doi.org/10.1175/JCLI-D-20-0517.1>, 2021.
- Sofiev, M., Ermakova, T., and Vankevich, R.: Evaluation of the smoke-injection height from wild-land fires using remote-sensing data, *Atmos. Chem. Phys.*, 12, 1995–2006, <https://doi.org/10.5194/acp-12-1995-2012>, 2012.
- Spencer, R. S., Levy, R. C., Remer, L. A., Mattoo, S., Arnold, G. T., Hlavka, D. L., Meyer, K. G., Marshak, A., Wilcox, E. M., and Platnick, S. E.: Exploring aerosols near clouds with high-spatial-resolution aircraft remote sensing during SEAC⁴RS, *J. Geophys. Res.-Atmos.*, 124, 2148–2173, <https://doi.org/10.1029/2018JD028989>, 2019.
- Su, W., Schuster, G. L., Loeb, N. G., Rogers, R. R., Ferrare, R. A., Hostetler, C. A., Hair, J. W., and Obland, M. D.: Aerosol and cloud interaction observed from high spectral resolution lidar data, *J. Geophys. Res.*, 113, D24202, <https://doi.org/10.1029/2008JD010588>, 2008.
- Swap, R. J., Annegarn, H. J., Suttles, J. T., Haywood, J., Helmlinger, M. C., Hely, C., Hobbs, P. V., Holben, B. N., Ji, J., King, M. D., Landmann, T., Maenhaut, W., Otter, L., Pak, B., Piketh, S. J., Platnick, S., Privette, J., Roy, D., Thompson, A. M., Ward, D., and Yokelson, R.: The Southern African Regional Science Initiative (SAFARI 2000): overview of the dry season field campaign, *South African J. Sci.*, 98, 125–130, 2002.
- Trickett, J. L. and Di Girolamo, L.: Enhanced aerosol backscatter adjacent to tropical trade wind clouds revealed by satellite-based lidar, *Geophys. Res. Lett.*, 36, L14804, <https://doi.org/10.1029/2009GL039264>, 2009.
- Tompkins, A. M.: Organization of tropical convection in low vertical wind shears: the role of cold pools, *J. Atmos. Sci.*, 58, 16501672, [https://doi.org/10.1175/1520-0469\(2001\)058<1650:OOTCIL>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<1650:OOTCIL>2.0.CO;2), 2001.
- Toth, T. D., Campbell, J. R., Reid, J. S., Tackett, J. L., Vaughan, M. A., Zhang, J., and Marquis, J. W.: Minimum aerosol layer detection sensitivities and their subsequent impacts on aerosol optical thickness retrievals in CALIPSO level 2 data products, *Atmos. Meas. Tech.*, 11, 499–514, <https://doi.org/10.5194/amt-11-499-2018>, 2018.
- Xian, P., Zhang, J., O'Neill, N. T., Reid, J. S., Toth, T. D., Sorenson, B., Hyer, E. J., Campbell, J. R., and Ranjbar, K.: Arctic spring and summertime aerosol optical depth baseline from long-term observations and model reanalyses – Part 2: Statistics of extreme AOD events, and implications for the impact of regional biomass burning processes, *Atmos. Chem. Phys.*, 22, 9949–9967, <https://doi.org/10.5194/acp-22-9949-2022>, 2022.
- Xiao, Q., Zhang, J., Wang, Y., Ziemba, L. D., Crosbie, E., Winstead, E. L., Robinson, C. E., DiGangi, J. P., Diskin, G. S., Reid, J. S., Schmidt, K. S., Sorooshian, A., Hilario, M. R. A., Woods, S., Lawson, P., Starnes, S. A., and Wang, J.: New particle formation in the tropical free troposphere during CAMP²Ex: statistics and impact of emission sources, convective activity, and synoptic conditions, *Atmos. Chem. Phys.*, 23, 9853–9871, <https://doi.org/10.5194/acp-23-9853-2023>, 2023.
- Wang, J., Gei, C., Yang, Z., Hyer, E., Reid, J. S., Chew, B. N., and Mahmud, M.: Mesoscale modeling of smoke transport over the South Asian maritime continent: vertical distributions and topographic effect, *Atmos. Res.*, 122, 486–503, 2013.