



## A WRF-Chem study of the greenhouse gas column and in situ surface mole fractions observed at Xianghe, China – Part 2: Sensitivity of carbon dioxide (CO<sub>2</sub>) simulations to critical model parameters

Sieglinde Callewaert<sup>1,4</sup>, Minqiang Zhou<sup>2,1</sup>, Bavo Langerock<sup>1</sup>, Pucai Wang<sup>3</sup>, Ting Wang<sup>3</sup>, Emmanuel Mahieu<sup>4</sup>, and Martine De Mazière<sup>1</sup>

<sup>1</sup>Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

<sup>2</sup>State Key Laboratory of Atmospheric Environment and Extreme Meteorology, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

<sup>3</sup>CNRC & LAGEO, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

<sup>4</sup>UR SPHERES, Department of Astrophysics, Geophysics and Oceanography, University of Liège, Liège, Belgium

**Correspondence:** Sieglinde Callewaert (sieglinde.callewaert@aeronomie.be) and Minqiang Zhou (minqiang.zhou@aeronomie.be)

Received: 13 August 2025 – Discussion started: 4 September 2025

Revised: 15 December 2025 – Accepted: 7 January 2026 – Published: 20 January 2026

**Abstract.** Understanding the variability and sources of atmospheric CO<sub>2</sub> is essential for improving greenhouse gas monitoring and model performance. This study investigates temporal CO<sub>2</sub> variability at the Xianghe site in China, which hosts both remote sensed (TCCON-affiliated) and in situ (PICARRO) observations. Using the Weather Research and Forecast model coupled with Chemistry, in its greenhouse gas option (WRF-GHG), we performed a one-year simulation of surface and column-averaged CO<sub>2</sub> mole fractions, evaluated model performance and conducted sensitivity experiments to assess the influence of key model configuration choices. The model captured the temporal variability of column-averaged mole fraction of CO<sub>2</sub> (XCO<sub>2</sub>) reasonably well ( $r = 0.7$ ), although a persistent bias in background values was found. A July 2019 heatwave case study further demonstrated the model's ability to reproduce a synoptically driven anomaly. Near the surface, performance was good during afternoon hours ( $r = 0.75$ , MBE = 2.44 ppm), nighttime mole fractions were overestimated (MBE = 7.86 ppm), resulting in an exaggerated diurnal amplitude. Sensitivity tests revealed that land cover data, vertical emission profiles, and adjusted VPRM-parameters (Vegetation Photosynthesis and Respiration Model) can significantly influence modeled mole fractions, particularly at night. Tracer analysis identified industry and energy as dominant sources, while biospheric fluxes introduced seasonal variability – acting as a moderate sink in summer for XCO<sub>2</sub> and a net source in most months near the surface. These findings demonstrate the utility of WRF-GHG for interpreting temporal patterns and sectoral contributions to CO<sub>2</sub> variability at Xianghe, while emphasizing the importance of careful model configuration to ensure reliable simulations.

## 1 Introduction

Climate change is one of the most pressing global challenges, and carbon dioxide (CO<sub>2</sub>) is its primary driver due to its long atmospheric lifetime and rising atmospheric abundance (Masson-Delmotte et al., 2021). Understanding how atmospheric CO<sub>2</sub> levels vary over time and space is essential for detecting long-term trends, distinguishing natural fluctuations from anthropogenic signals, and deepening our insight into the carbon cycle and its interactions with the atmosphere. Observational records are key to unraveling local and regional carbon budgets and assessing the effectiveness of mitigation strategies. To fully interpret such observations, especially in complex environments, we rely on atmospheric transport models, which provide spatial and temporal context and help disentangle the observed CO<sub>2</sub> signal into contributions from different sources and processes. As the world's largest fossil CO<sub>2</sub> emitter (Friedlingstein et al., 2025), our study focuses on China – a country whose vast and densely populated regions, strong industrial activity, and ecological diversity make it a complex but highly relevant area for atmospheric CO<sub>2</sub> research. In this context, a ground-based remote sensing instrument was installed in 2018 at the Xianghe site, a suburban location approximately 50 km southeast of Beijing. The Fourier Transform Infrared (FTIR) spectrometer provides high-precision column-averaged CO<sub>2</sub> mole fractions and is part of the global Total Carbon Column Observing Network (TCCON). Complementing this, a PICARRO Cavity Ring-Down Spectroscopy (CRDS) analyzer measures near-surface CO<sub>2</sub> mole fractions at 60 m above ground level. This unique combination of collocated column and in situ observations – to our knowledge currently the only such setup in China – offers a valuable opportunity to study both local and regional CO<sub>2</sub> signals and to evaluate model performance for different levels of the atmosphere.

Previous studies by Yang et al. (2020, 2021) provided initial insights into the seasonal and diurnal variability of both column-averaged and near-surface CO<sub>2</sub> mole fractions at Xianghe. Their work highlighted the strong influence of local and regional emissions, as well as planetary boundary layer dynamics, on observed CO<sub>2</sub> levels. However, these analyses were either purely observational or relied on coarser-resolution model products such as CarbonTracker, which are limited in their ability to resolve mesoscale variability. Furthermore, the two observation types were not jointly analyzed within a high-resolution modeling framework, leaving room for a more detailed and integrated approach. To gain a deeper understanding of the processes shaping the observed CO<sub>2</sub> mole fractions at Xianghe, we apply the high-resolution WRF-GHG model in this work, a specific configuration of the widely used WRF-Chem model tailored for greenhouse gas simulations (Beck et al., 2011). The current study is part of a broader research effort investigating multiple greenhouse gases at the site. While a companion paper

has already presented the results for CH<sub>4</sub> (Callewaert et al., 2025), the present work focuses exclusively on CO<sub>2</sub>.

The WRF-GHG model was originally developed to address the limitations of coarse-resolution global models by providing a more detailed representation of CO<sub>2</sub> transport, surface flux exchanges, and meteorological processes at the mesoscale. Thanks to its coupling with the Vegetation Photosynthesis and Respiration Model (VPRM), WRF-GHG has demonstrated strong capabilities in simulating biogenic CO<sub>2</sub> fluxes (NEE, net ecosystem exchange) and atmospheric dynamics. It has been successfully applied across a range of environments, from rural areas influenced by sea-breeze circulations (Ahmadov et al., 2007, 2009, here referred to as WRF-VPRM) to urban regions with complex emission patterns and boundary layer processes (Feng et al., 2016; Park et al., 2018; Zhao et al., 2019). Further, the model has been evaluated against in situ, tower, aircraft and satellite data during large-scale campaigns such as ACT-America in the US (Hu et al., 2020) and KORUS-AQ in South-Korea (Park et al., 2020), showing its ability to reasonably capture spatiotemporal variability of CO<sub>2</sub>. In China, WRF-GHG has been used to study CO<sub>2</sub> fluxes and atmospheric mole fractions on a national scale and to explore the role of biospheric and anthropogenic sources (Dong et al., 2021; Ballav et al., 2020). Li et al. (2020) evaluated WRF-GHG against tower observations in northeast China, showing the model could capture seasonal trends and episodic enhancements, despite underestimating diurnal variability and respiration fluxes.

Our study uses WRF-GHG to investigate the main drivers of observed temporal variations at Xianghe and to evaluate the model's ability to reproduce these patterns, identifying key sources of error where relevant. The model's tracer framework further allows us to disentangle the contributions of anthropogenic, biogenic, and meteorological processes to simulated CO<sub>2</sub> levels. The structure of the paper is as follows: Sect. 2 describes the observations, model configuration, and the design of additional model sensitivity experiments. Section 3 presents the results, including model performance, tracer-based analyses and sensitivity experiments. Section 4 discusses some of the results in more detail, while Sect. 5 summarizes the conclusions and provides an outlook.

## 2 Methods

### 2.1 Observations at Xianghe site

We use observational data from the atmospheric monitoring station situated in Xianghe county (39.7536°N, 116.96155°E; 30 m a.s.l.). This site is located in a suburban part of the Beijing-Tianjin-Hebei (BTH) region in northern China. The town center of Xianghe lies approximately 2 km to the east, while the major metropolitan areas of Beijing and Tianjin are situated roughly 50 km to the northwest and 70 km to the south-southeast, respectively (Fig. 1b). The

dominant vegetation in the surrounding area consists of cropland.

Continuous atmospheric measurements have been conducted at the Xianghe observatory by the Institute of Atmospheric Physics (IAP), Chinese Academy of Sciences (CAS), since 1974. FTIR solar absorption measurements have been performed since June 2018, from the roof of the observatory by a Bruker IFS 125HR. This ground-based remote sensing instrument records spectra in the infrared range and is part of the TCCON network (Wunch et al., 2011; Zhou et al., 2022), providing data on total column-averaged dry air mole fractions for gases such as CO<sub>2</sub>, CH<sub>4</sub>, and CO (noted as XCO<sub>2</sub>, XCH<sub>4</sub> and XCO, respectively). The current study employs the GGG2020 data product (Laughner et al., 2024). Observations are typically taken every 5 to 20 min, depending on weather conditions and instrument status. TCCON measurements are exclusively performed under clear skies. The uncertainty associated with the XCO<sub>2</sub> measurements is approximately 0.5 ppm. Further details regarding the instrument and the retrieval methodology can be found in Yang et al. (2020).

In addition to the FTIR measurements, in situ measurements of CO<sub>2</sub> and CH<sub>4</sub> mole fractions have been conducted since June 2018 using a PICARRO cavity ring-down spectroscopy G2301 analyzer. This instrument draws air from an inlet situated on a 60 m tower. A more comprehensive description of this measurement setup is available in Yang et al. (2021). The measurement uncertainty for CO<sub>2</sub> with this instrument is about 0.06 ppm. The data used in this study were converted to align with the WMO CO<sub>2</sub> X2019 scale (Hall et al., 2021).

## 2.2 WRF-GHG model simulations

We make use of the WRF-GHG model simulations elaborated in Part 1 of this work (Callewaert et al., 2025), and provide a brief summary here for completeness. The simulations were performed using the Weather Research and Forecasting model with Chemistry (WRF-Chem v4.1.5; Grell et al., 2005; Skamarock et al., 2019; Fast et al., 2006) in its greenhouse gas configuration, called WRF-GHG (Beck et al., 2011). This Eulerian transport model simulates three-dimensional greenhouse gas mole fractions simultaneously with meteorological fields, without accounting for chemical reactions. The model setup includes three nested domains with horizontal resolutions of 27, 9, and 3 km (Fig. 1a), and 60 vertical levels extending from the surface up to 50 hPa. There are 11 layers in the lowest 2 km, with a layer thickness ranging from about 50 m near the surface, to 400 m above 2 km.

Anthropogenic CO<sub>2</sub> emissions were obtained from CAMS-GLOB-ANT v5.3 (Granier et al., 2019; Soulé et al., 2024) and temporally disaggregated using CAMS-TEMPO profiles (Guevara et al., 2021). The original 11 source sectors were aggregated into four broader categories and included in separate tracers: energy, industry, transportation, and residen-

tial & waste. Biomass burning emissions were taken from the FINN v2.5 inventory (Wiedinmyer et al., 2011), and ocean-atmosphere CO<sub>2</sub> fluxes were prescribed based on the climatology from Landschützer et al. (2017). Finally, net biogenic CO<sub>2</sub> fluxes were calculated online using VPRM (Mahadevan et al., 2008; Ahmadov et al., 2007), driven by WRF-GHG meteorology and MODIS surface reflectance data, with ecosystem-specific VPRM parameters from Li et al. (2020) and land cover information from SYNMAP (Jung et al., 2006). Meteorological fields (e.g., wind, temperature, humidity) were driven by hourly data from the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 hourly data (0.25° × 0.25°; Hersbach et al., 2023a, b). Daily restarts were performed at 00:00 UTC, with model initialization at 18:00 UTC the day before to allow for a 6-h spin-up, stabilizing the simulation. For tracer fields, mole fractions at 00:00 UTC were copied from the previous day's simulation to maintain consistency. The initial and lateral boundary conditions for CO<sub>2</sub> were prescribed using the 3-hourly Copernicus Atmosphere Monitoring Service (CAMS) global reanalysis (EGG4, Agustí-Panareda et al., 2023).


The final simulated CO<sub>2</sub> field is composed of the sum of several tracers that track contributions from individual sources. These include a background tracer (reflecting the evolution of initial and lateral boundary conditions from CAMS), as well as tracers for energy, industry, residential, transportation, ocean, biomass burning, and biogenic fluxes.

WRF-GHG was run from 15 August 2018 to 1 September 2019. However, the first two weeks were regarded as a spin-up phase, so the analysis in Sect. 3.1 is made on one full year of data: from 1 September 2018 until 1 September 2019. The complete data set can be accessed on <https://doi.org/10.18758/P34WJEW2> (Callewaert, 2023).

To enable comparison with the observations, model data from the grid cell containing the measurement site are extracted. For near-surface observations, the model profile is interpolated to the altitude of the instrument, while for column measurements the model output is smoothed with the FTIR retrieval's a priori profile and averaging kernel, after being extended above the model top with the FTIR a priori profile. The hourly model output represents instantaneous values, as do the observational measurements. To align the datasets temporally, the observations are averaged around each model output time step – using a ±15-min window. Further details are provided in Part 1 (Callewaert et al., 2025).

## 2.3 Sensitivity experiment design

A series of sensitivity experiments was conducted to assess the impact of key model assumptions on surface CO<sub>2</sub> fluxes, such as the treatment of emission heights, land cover classification, and biogenic flux parameterizations. Four two-week periods were selected for these sensitivity simulations, spanning from the 15–29 March, May, July, and December. These months were identified as being most critical for simulating



**Figure 1.** (a) Location of the WRF-GHG domains, with horizontal resolutions of 27 km (d01), 9 km (d02) and 3 km (d03). All domains have 60 (hybrid) vertical levels extending from the surface up to 50 hPa. (b) Terrain map including the largest cities in the region of Xianghe, roughly corresponding to d03. The location of the Xianghe site is indicated by the red triangle in both maps. Figure taken from Callewaert et al. (2025). ©OpenStreetMap.

the diurnal CO<sub>2</sub> cycle while representing different seasons. Although the specific dates within each month (15–29) were chosen somewhat arbitrarily, they were applied consistently across all four months to ensure comparability. Four different simulation experiments (BASE, PROF, LC, PARAM) were performed over these four periods to isolate the impact of three model assumptions (see Table 1):

– **Emission height.** To assess the impact on simulated in situ CO<sub>2</sub> mole fractions at Xianghe of the height at which anthropogenic emissions are released in the atmosphere (all at the lowest model level near the surface, or according to sector-specific vertical profiles), we applied the vertical profiles for point sources from Brunner et al. (2019) to the CAMS-GLOB-ANT sector-specific CO<sub>2</sub> emissions. For the fluxes in the 'industrial processes' sector, we used the average of the profiles of SNAP 3 (Combustion in manufacturing industry) and SNAP 4 (Production processes). Note that we do not make a distinction between area and point sources as in Brunner et al. (2019), as this information is not available for our study region. Profile emissions were included in all but the BASE experiment: PROF, LC and PARAM.

– **Land cover classification.** The net biogenic CO<sub>2</sub> fluxes are calculated online in WRF-GHG as the weighted average of the Net Ecosystem Exchange (NEE) for eight vegetation classes (evergreen trees, deciduous trees, mixed trees, shrubland, savanna, cropland, grassland and non-vegetated land) (Mahadevan et al., 2008). As a default, the SYNMAP land cover map is used to calculate the vegetation fraction for every model grid cell. To assess the impact of this classification, we prepare the VPRM model input files for the LC and PARAM experiments with the global 100-m Copernicus Dynamic Land Cover Collection 3 (epoch 2019) (Buchhorn et al., 2020), using the pyVPRM python package (Glauch et al., 2025), allowing for an updated and

**Table 1.** Overview of the model configuration for the sensitivity experiments. Note that the Glauch parameter table does not include values for the "Savanna" class, consistent with its absence in the Copernicus land cover map.

Experiment name	Emission height	Land cover map	VPRM parameters
BASE	SFC	SYNMAP	Li table
PROF	PROF	SYNMAP	Li table
LC	PROF	Copernicus LC	Li table
PARAM	PROF	Copernicus LC	Glauch table

higher-resolution representation of vegetation types in the domain.

– **VPRM parameterization.** The VPRM-calculated NEE can be tuned for different regions around the globe by specifying four empirical parameters ( $\alpha$ ,  $\beta$ ,  $\lambda$  and  $PAR_0$ ) per vegetation class. These parameter tables are a mandatory input to the WRF-GHG model and can be calibrated using a network of eddy flux tower sites, representing the different vegetation classes in the region, or taken from literature. Due to the lack of a dedicated calibration study in China, we applied the table from Li et al. (2020) in the one-year simulations, and the BASE and LC experiments. Seo et al. (2024) reported the lowest RMSE in East Asia using these parameter values, relative to the default US settings and those of Dayalu et al. (2018). To evaluate the impact of these parameters at Xianghe, we conducted an experiment (PARAM) with an alternative parameter table, optimized over Europe by Glauch et al. (2025). The exact parameter values used in each experiment are provided in Table A1.

By comparing the results from the four sensitivity experiments, the influence of individual model components can be isolated. The role of vertical emission distribution can be

assessed by comparing the BASE and PROF experiments. Similarly, the impact of land cover classification is assessed by comparing the PROF and LC simulations, which differ only in the land cover dataset used. Finally, the effect of VPRM parameterization can be evaluated by comparing LC and PARAM, which share the same land cover input but differ in the VPRM parameter table. This approach enables a systematic investigation of potential model deficiencies affecting the representation of CO<sub>2</sub> at Xianghe. Note that the BASE experiment corresponds exactly with the settings of the one-year simulation described above.

### 3 Results

#### 3.1 Evaluation of one-year simulation

##### 3.1.1 Timeseries and statistical comparison

Table 2 summarizes the comparison between simulated and observed CO<sub>2</sub> mole fractions at Xianghe. Overall, WRF-GHG demonstrates a reasonable accuracy in replicating these measurements: the XCO<sub>2</sub> observations are slightly underestimated, with a mean bias error of  $-1.43 (\pm 1.99)$  ppm and a Pearson correlation coefficient of 0.70. Note that the XCO<sub>2</sub> time series was de-seasonalized before calculating the correlation coefficient in order to remove the effect of the seasonal variation. After applying a bias correction to the modeled values, the XCO<sub>2</sub> MBE decreases to  $-0.86 (\pm 1.57)$  ppm (corrected values shown in parentheses in Table 2), and the RMSE improves from 2.45 to 1.80 ppm, while the correlation remains unchanged. Details of the bias correction are provided in Sect. 3.1.2, and the resulting time series are shown in Fig. 2 (and Fig. A1).

The data near the surface has been divided into afternoon (12:00–15:00 LT), nighttime (22:00–04:00 LT) and morning (08:00–12:00 LT) periods to assess model performance under different boundary layer conditions. Indeed, WRF-GHG shows a smaller bias ( $-2.44$  ppm) during the afternoon, when the lower atmosphere is well-mixed, compared to nighttime (7.86 ppm). Additionally, the MBE differs in sign between the two periods: near-surface CO<sub>2</sub> levels tend to be underestimated by the model in the afternoon but overestimated at night. Except for the moderate correlation observed for in situ CO<sub>2</sub> during nighttime (0.60), WRF-GHG achieves relatively high correlation coefficients ( $\geq 0.7$ ) for other CO<sub>2</sub> data, indicating satisfactory model performance. Overall, the bias correction has only a minor influence on the comparison with near-surface mole fractions, where the effect on RMSE and correlation coefficients are negligible ( $< 1.2\%$  and  $< 0.01$ , respectively).

Finally, the XCO<sub>2</sub> time series in Fig. 2a reveals a notable spike between 20–29 July (highlighted in gray), interrupting the general decline associated with northern hemispheric photosynthetic uptake during the growing season, from May onwards. A dedicated analysis of this July XCO<sub>2</sub> event is

provided in Sect. 4.2. Note that there is a gap in the in situ CO<sub>2</sub> time series during this period due to instrument malfunctions (Yang et al., 2021).

##### 3.1.2 Correction of background bias

Our WRF-GHG simulations underestimate XCO<sub>2</sub> by approximately 2 ppm until May 2019, after which the negative bias diminishes (see Fig. A1a, c). This bias likely originates from a similar error in the background data, inaccuracies in representing the actual sources and sinks in the region, or a combination of both.

The CAMS validation report (Ramonet et al., 2021) presents “a very good agreement for all (TCCON) sites”, suggesting that the CAMS reanalysis that is driving the WRF-GHG simulations is of good quality without known biases. However, their criteria for what constitutes “very good” appears to be relatively mild (within  $\pm 2$  ppm). Moreover, the Xianghe site was not included in this report and the accompanying figure does not provide very detailed information. Therefore, we reproduced their analysis for several TCCON sites at similar latitudes for the period of our interest (September 2018–September 2019): Karlsruhe (49.1° N), Orleans (48.0° N), Garmisch (47.5° N), Park Falls (45.9° N), Rikubetsu (43.5° N), Lamont (36.6° N), Tsukuba (36.0° N), Edwards (35.0° N), Pasadena (34.1° N), Saga (33.2° N), and Hefei (31.9° N). The results of this analysis are presented in Fig. 3.


We find an underestimation of the CAMS reanalysis XCO<sub>2</sub> at all TCCON sites between 30–50° N (except Pasadena) from October 2018 until May 2019. More specifically for Xianghe, monthly mean errors range from  $-2.20 (\pm 1.3)$  ppm in January 2019 to  $3.38 (\pm 1.28)$  ppm in July 2019, which is of a similar magnitude as the bias found with WRF-GHG (where the monthly mean differences with respect to the TCCON site of Xianghe range from  $-2.53 (\pm 1.7)$  ppm in December 2018 to  $1.28 (\pm 1.57)$  ppm in July 2019).

Therefore, we assume that the error pattern detected in the XCO<sub>2</sub> time series is primarily the result of the same pattern in the background information. Moreover, this bias pattern is not found in the in situ CO<sub>2</sub> time series, likely because the relative contribution from the background to the in situ mole fractions is smaller than it is to the column data.

To account for the systematic bias introduced by the background values, we applied a bias correction to the WRF-GHG simulations. Specifically, we subtract the monthly mean difference between CAMS and TCCON XCO<sub>2</sub>, averaged across all TCCON sites located between 30–50° N (excluding Pasadena due to outlier behavior), from the model’s background tracer. The resulting improvements in model performance are summarized in Table 2 between parentheses.

**Table 2.** Statistics of the model-data comparison of the ground-based CO<sub>2</sub> observations at the Xianghe site from 1 September 2018 until 1 September 2019. We present the mean bias error (MBE), root mean square error (RMSE) and Pearson correlation coefficient (CORR). The MBE and RMSE are given in ppm. For in situ observations, the data is split in afternoon (12:00–15:00 LT), night (22:00–04:00 LT) and morning transition (08:00–12:00 LT) hours. The bias corrected model values are given between brackets.

	insitu CO <sub>2</sub>			XCO <sub>2</sub>
	afternoon	night	morning transition	
MBE	−3.12 (−2.44)	7.18 (7.86)	−0.73 (−0.04)	−1.43 (−0.86)
RMSE	15.35 (15.27)	23.77 (24.04)	22.04 (22.15)	2.45 (1.80)
CORR	0.75 (0.76)	0.60 (0.60)	0.69 (0.69)	0.70 (0.70)



**Figure 2.** Time series of the observed (black) and simulated (red) (a) XCO<sub>2</sub> and (b) insitu CO<sub>2</sub> mole fractions at the Xianghe site. Panels (c) and (d) show the differences between (smoothed) WRF-GHG simulations and observations for XCO<sub>2</sub> and in situ CO<sub>2</sub>, respectively. Data points are hourly, if available. The red data points in (b) and (d) represent the monthly mean differences. A bias correction was applied to the WRF-GHG values.

### 3.2 Sector contributions to observed mole fractions

WRF-GHG tracks all fluxes in separate tracers, enabling the decomposition of the total simulated CO<sub>2</sub> mole fractions at Xianghe into contributions from different source sectors. Figure 4 shows the monthly mean values, while additionally the median and interquartile ranges are presented in Table 3. Note that all simulated hours were used for this analysis, not just the ones coinciding with observations.

The main sectors contributing to the modeled CO<sub>2</sub> variability at Xianghe are energy, industry, and the biosphere. For XCO<sub>2</sub>, we find median values of 0.85 and 0.63 ppm for the energy and industry sectors, respectively. Furthermore, the biosphere significantly influences the column-averaged CO<sub>2</sub> values, where it acts as a sink from April to September with a median value of −0.77 ppm during this period. During the rest of the year, the biogenic tracer acts as a small source (median value of 0.22 ppm).


Near the surface, median enhancements of in situ CO<sub>2</sub> mole fractions are 6.85 and 5.69 ppm for the energy and industry sectors, respectively. The biosphere generally acts as a source throughout the year, with a median contribution of

2.69 ppm, except in August, when it becomes a significant sink of −6.76 ppm.


Next to the three dominant sectors (biosphere, industry, and energy), transportation and also residential sources have a smaller but still relevant influence on the Xianghe data. During winter, the contribution of residential sources increases, where the highest values for the column simulations are found in February (median of 0.45 ppm) while near the surface this occurs in January (4.28 ppm). This peak aligns with heightened residential emissions in winter, driven by increased heating demands correlated with air temperature (Guevara et al., 2021). Finally, no relevant impact was found from biomass burning and the ocean. Overall, the total tracer enhancement for the in situ mole fractions is about ten times greater than that of the column-averaged values.

### 3.3 Diurnal cycle analysis of in situ data

The planetary boundary layer (PBL) plays a crucial role in regulating near-surface CO<sub>2</sub> mole fractions. Figure 6 displays the diurnal variation of the PBL height as simulated by WRF-GHG, along with the corresponding CO<sub>2</sub> mole fractions near the surface (both simulated and observed).



**Figure 3.** Monthly mean difference (in ppm) between CAMS reanalysis model and TCCON XCO<sub>2</sub> between 30–50° N over the simulation period of this study.



**Figure 4.** Monthly mean tracer contributions above the background for (a) XCO<sub>2</sub> and (b) in situ CO<sub>2</sub> simulated mole fractions at Xianghe.

During the day, solar radiation promotes turbulent mixing, leading to a deepening of the PBL and the dilution of near-surface CO<sub>2</sub>. The PBL reaches its maximum height around 15:00 local time (LT), coinciding with the lowest surface CO<sub>2</sub> mole fractions. Conversely, during the night, radiative cooling leads to the formation of a stable, shallow PBL, trapping CO<sub>2</sub> near the surface and causing mole fractions to rise. As the sun rises and the PBL height begins to increase again, the CO<sub>2</sub> mole fractions drop, giving rise to a characteristic diurnal cycle.

Indeed, the lowest values are observed between 14:00 and 16:00 LT, with a minimum of 421.32 (hourly median value, with an interquartile range of 415.80–431.76) ppm at 16:00 LT. In the early morning, the observed CO<sub>2</sub> mole fractions show a distinct peak at 07:00 LT, reaching 443.42 (428.00–459.32) ppm. WRF-GHG successfully captures the general shape of this diurnal cycle, but discrepancies remain in the amplitude and timing. The model slightly underestimates daytime mole fractions, with a minimum value that is 1.22 ppm lower and occurs one hour earlier than the ob-


servations (at 15:00 LT). The peak CO<sub>2</sub> mole fractions in WRF-GHG are also reached at 07:00 LT, but overestimated by 3.36 ppm. Furthermore, this peak is less distinct as in the observations, where the model remains relatively stable between 03:00 and 08:00 LT. This results in an overestimation of the diurnal amplitude by approximately 4.58 ppm. Such a nighttime overestimation was not observed for CH<sub>4</sub> at the same site (Callewaert et al., 2025), suggesting that the bias is more likely related to the surface fluxes of CO<sub>2</sub> than to PBL dynamics.

### 3.4 Sensitivity experiments

Several sources of uncertainty may affect the accuracy of the simulated anthropogenic CO<sub>2</sub> fluxes and NEE in WRF-GHG. First of all, the parameters used in VPRM in this study are based on Li et al. (2020), who optimized them for ecosystems in the United States. Applying these values to China likely introduces regional mismatches, as differences in dominant species, climate conditions, and land use history can significantly alter ecosystem carbon dynamics even

**Table 3.** Statistics of the total simulated CO<sub>2</sub> mole fractions and the different tracer contributions over the complete simulation period. Q1 and Q3 represent the first and third quartile, respectively, between which 50 % of the data fall.

	XCO <sub>2</sub> (ppm)				in situ CO <sub>2</sub> (ppm)			
	Q1	median	mean	Q3	Q1	median	mean	Q3
Total	408.32	412.11	411.37	414.11	419.44	430.93	437.86	450.14
Background	407.2	410.21	409.54	412.12	397.63	412.42	411.08	414.69
Biomass burning	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Energy	0.36	0.85	1.07	1.53	2.74	6.85	10.51	14.06
Residential	0.03	0.06	0.17	0.17	0.30	0.65	1.88	1.95
Industry	0.24	0.63	0.79	1.14	2.70	5.69	8.53	10.51
Transportation	0.08	0.16	0.18	0.25	0.81	1.73	2.43	3.19
Biosphere	-0.77	0.04	-0.38	0.31	-0.01	2.36	3.44	7.39
Ocean	-0.00	-0.00	-0.00	-0.00	-0.01	-0.00	-0.01	-0.00
Total tracers	0.33	1.23	1.82	2.80	7.36	18.99	26.78	38.30




**Figure 5.** Map of the mean CO<sub>2</sub> flux (mol km<sup>-2</sup> h<sup>-1</sup>) in WRF-GHG domain d03 during the entire simulation period from September 2018 until September 2019, for the most important sectors. Remark that the panels have different color scales. The location of the Xianghe site is indicated by a blue cross.

within the same vegetation class (Mahadevan et al., 2008; Seo et al., 2024). Moreover, the linear formulation of the respiration term in VPRM has been identified as a source of potential bias (Dong et al., 2021; Hu et al., 2021). A third concern is the land cover classification. VPRM uses the SYN-MAP product (Jung et al., 2006), which is a 1-km global land cover map that classifies the area around Xianghe as 100 % cropland. While broadly consistent with the regional land use, this dataset does not account for increasing urbanization during the last decades. In WRF-GHG, built-up areas are assigned zero NEE, so their omission could contribute to the observed nighttime overestimation of respiration and daytime photosynthetic uptake.

Further, the representation of anthropogenic emission heights may also affect the modeled surface mole fractions. In this study, all anthropogenic CO<sub>2</sub> emissions are released in the lowest model layer, which simplifies reality. Especially for sectors such as energy and industry, this is a crude approximation, since facilities like power plants typically emit at elevated stacks. Previous work by Brunner et al. (2019) has shown that ignoring the vertical distribution of emissions can lead to overestimation of near-surface mole fractions.

Finally, while it is well known that uncertainties in simulating planetary boundary layer (PBL) dynamics can substantially affect near-surface CO<sub>2</sub> mole fractions, the influence of different PBL parameterization schemes is not explored in the current sensitivity experiments. WRF-GHG offers sev-



**Figure 6.** Hourly median and interquartile range of the (a) simulated planetary boundary layer height, and observed and simulated surface (b) CO<sub>2</sub> mole fraction at Xianghe.

eral PBL schemes, some of which were tested and discussed in Part 1 of this work (Callewaert et al., 2025). Here, the focus is instead on evaluating how model configuration choices related to CO<sub>2</sub> fluxes impact the simulated mole fractions.

### 3.4.1 Emission height sensitivity

To evaluate the impact of emission injection height on simulated CO<sub>2</sub> mole fractions at Xianghe, we compare the BASE and PROF sensitivity experiments (see Sect. 2.3). Figure 7 presents the median diurnal cycle of the anthropogenic CO<sub>2</sub> tracer across the four 14-d simulation periods (top), together with the total simulated and observed CO<sub>2</sub> mole fractions (bottom). Similarly, Table 4 provides a summary of the impact on the anthropogenic tracer.

Simulations using elevated emission profiles (PROF) consistently yield lower near-surface CO<sub>2</sub> mole fractions than those with surface-only emissions (BASE), with the most pronounced differences occurring during nighttime and the morning transition. The largest reduction is observed in December, where mean nighttime mole fractions in the PROF simulation are 16.92 ppm lower than in BASE.


An overview of the key statistical performance metrics with respect to the observations at Xianghe is given in Table 5. In December and March, BASE shows large positive mean biases (MBE), primarily driven by nighttime overestimation; this bias is substantially reduced in PROF. By contrast, in May and July the absolute MBE increases and changes sign from positive to negative. The use of elevated emissions leads to a reduction in the RMSE compared to the surface-only configuration in all periods, while the correlation coefficient remains largely unaffected. The use of elevated emission profiles also has a pronounced effect on the simulated diurnal amplitude of near-surface CO<sub>2</sub>. Compared to the surface-only configuration, which strongly overestimates the amplitude, the more realistic vertical distribution

results in a better agreement with observations – particularly in March and July. In March, for example, the amplitude overestimation is reduced from 22.73 ppm to just 1.74 ppm, and in July from 14.16 to –5.96 ppm. The implementation of elevated anthropogenic emissions has a minimal effect on the column-averaged XCO<sub>2</sub> mole fractions at Xianghe, see statistical metrics in Table A2.

### 3.4.2 Biogenic flux and land cover sensitivity

To evaluate the impact of land cover representation on the VPRM-calculated CO<sub>2</sub> fluxes, we compare the PROF and LC sensitivity experiments. In the PROF simulation, the WRF-GHG grid cell containing the Xianghe site is classified as 100 % cropland using the SYNMAP dataset. In contrast, the Copernicus Dynamic Land Cover data, used in the LC experiment, classifies the same cell as 68.88 % cropland, 23.5 % no vegetation (representing urban surfaces, water, ice, rocks, etc.), 3.45 % mixed forest, 2.03 % wetland, 1.5 % shrubland, and 0.64 % grassland. A comparison of the two land cover datasets over the innermost WRF-GHG domain (d03) is shown in Fig. A4. Due to its higher spatial resolution and more up-to-date information, the Copernicus dataset introduces more heterogeneous vegetation fractions and especially reduces the cropland fraction while increasing the urban land category compared to SYNMAP.

The top panels of Fig. 8 present the median diurnal cycles of NEE, at Xianghe across the four 14-d simulation periods. Differences between the experiments are negligible in winter months (December and March), whereas during May and July, the LC simulation exhibits both reduced daytime CO<sub>2</sub> uptake and lower nighttime respiration compared to PROF. This change in NEE is reflected in the simulated biogenic CO<sub>2</sub> tracer at Xianghe (Fig. 8e–h, Table 6). However, we notice that the magnitude of the difference between LC and PROF is more pronounced at night. For instance, in July, the



**Figure 7.** Median diurnal cycle of in situ CO<sub>2</sub> mole fractions (ppm) at Xianghe. The solid red line presents the simulated values of the PROF sensitivity experiment (using vertical profiles for the anthropogenic emissions), while the dashed blue line represents the simulated CO<sub>2</sub> values using only surface emissions (SFC). Observations are plotted in black.

**Table 4.** Mean and standard deviation (in ppm) of the total anthropogenic CO<sub>2</sub> tracer contribution (sum of industry, energy, transportation and residential tracer) to near-surface mole fractions at Xianghe for the BASE and PROF sensitivity experiments, and simulation period. “All” indicates that all simulated hours (00:00–23:00 LT) were used to calculate the metrics, in contrast to “Afternoon” (12:00–15:00 LT), “Night” (22:00–04:00 LT) and “Morning” (08:00–12:00 LT).

		December 2018	March 2019	May 2019	July 2019
All	BASE	34.32 ± 33.64	18.94 ± 18.74	15.59 ± 14.35	24.25 ± 14.78
	PROF	20.37 ± 20.47	9.51 ± 7.76	8.5 ± 6.16	13.98 ± 5.32
Afternoon	BASE	17.13 ± 17.23	9.12 ± 9.07	8.94 ± 7.44	15.53 ± 9.99
	PROF	11.82 ± 11.10	6.54 ± 5.73	7.09 ± 5.24	12.06 ± 4.85
Night	BASE	41.43 ± 35.36	23.17 ± 16.14	18.41 ± 15.23	27.57 ± 11.99
	PROF	24.51 ± 22.15	10.80 ± 7.02	8.89 ± 5.60	15.07 ± 5.62
Morning	BASE	31.07 ± 31.45	22.75 ± 27.99	19.36 ± 14.53	27.43 ± 14.45
	PROF	14.30 ± 10.16	10.27 ± 10.25	10.83 ± 6.31	14.77 ± 4.19

mean ( $\pm$  standard deviation) difference in the biogenic tracer is  $-3.91$  ( $\pm 2.13$ ) ppm during nighttime, while the daytime difference is only  $-1.41$  ( $\pm 1.26$ ) ppm.

To assess the impact of the VPRM parameterization on CO<sub>2</sub> simulations at Xianghe, we compare the LC and PARAM experiments. Both simulations use the 100-m Copernicus Dynamic Land Cover dataset but differ in their VPRM parameter tables (see Table 1). The comparison reveals that PARAM systematically produces more positive NEE than LC, both during daytime and nighttime. The change in NEE is reflected in the biogenic CO<sub>2</sub> tracer mole fractions at Xianghe. In July, for example, tracer values in PARAM are on average  $10.61$  ( $\pm 5.03$ ) ppm higher than in LC. This difference is again more pro-


nounced at night ( $11.48 \pm 4.27$  ppm) than during the afternoon ( $8.01 \pm 3.09$  ppm).

A summary of the model performance of the different experiments is shown in Table 5. Generally, the PARAM experiment yields the best agreement with observations. Across all months, PARAM shows the highest correlation coefficients and the lowest MBE and RMSE, with the exception of March. In that month, the LC experiment outperforms PARAM, with a smaller MBE ( $-0.2$  ppm vs.  $4.61$  ppm) and RMSE ( $10.73$  ppm vs.  $13.09$  ppm).

The various VPRM inputs have only a minor influence on the column-averaged XCO<sub>2</sub> mole fractions at Xianghe, as indicated by the statistical metrics in Table A2.

**Table 5.** Overview of statistical metrics (MBE, RMSE and CORR) for the different sensitivity experiments BASE, PROF, LC and PARAM with respect to the observations, per simulation period. Values for MBE and RMSE are given in ppm.

		MBE				RMSE				CORR			
		Dec	Mar	May	Jul	Dec	Mar	May	Jul	Dec	Mar	May	Jul
All	BASE	10.04	10.80	0.69	1.99	31.90	22.41	17.22	19.22	0.55	0.47	0.61	0.64
	PROF	-3.91	1.38	-6.40	-8.27	23.51	11.47	16.08	16.82	0.53	0.49	0.62	0.67
	LC	-3.92	-0.20	-8.04	-9.72	23.52	10.73	17.20	17.30	0.53	0.48	0.58	0.69
	PARAM	-1.82	4.61	0.34	0.82	23.60	13.09	13.79	13.64	0.55	0.50	0.68	0.73
Afternoon	BASE	-3.96	-0.51	-2.67	-6.70	21.59	10.49	12.22	18.76	0.58	0.63	0.12	-0.11
	PROF	-9.28	-3.09	-4.53	-10.17	24.42	8.88	11.92	17.11	0.47	0.66	0.11	-0.03
	LC	-9.29	-3.80	-4.54	-8.77	24.43	8.85	11.90	15.91	0.47	0.66	0.13	0.03
	PARAM	-8.53	-0.23	2.32	-0.84	23.84	9.75	11.00	13.70	0.50	0.63	0.23	0.10
Night	BASE	17.00	17.75	6.38	7.64	32.42	23.81	19.21	20.71	0.69	0.55	0.54	0.35
	PROF	0.08	5.37	-3.15	-4.86	18.57	11.14	11.69	16.12	0.72	0.49	0.70	0.43
	LC	0.06	3.17	-6.56	-8.77	18.58	9.51	14.06	17.26	0.72	0.49	0.61	0.46
	PARAM	3.00	8.90	2.56	2.64	19.38	13.78	11.15	14.47	0.73	0.48	0.73	0.54
Morning	BASE	-0.79	11.49	-1.44	-3.51	32.79	31.13	19.83	19.34	0.46	0.39	0.39	0.52
	PROF	-17.57	-1.00	-9.98	-16.17	30.43	14.28	20.44	21.50	0.60	0.47	0.35	0.57
	LC	-17.58	-2.60	-10.59	-15.97	30.45	13.85	20.72	21.10	0.59	0.45	0.33	0.59
	PARAM	-15.66	2.21	-0.66	-3.87	29.27	15.18	17.09	14.44	0.58	0.50	0.46	0.62



**Figure 8.** Median diurnal cycle of NEE (a–d), biogenic CO<sub>2</sub> tracer contribution (e–h) to the near surface CO<sub>2</sub> mole fractions (i–l) at Xianghe for the different simulation periods (columns). Different curves (colors) represent different sensitivity experiments PROF, LC and PARAM. Observations are plotted in black in the bottom row.

**Table 6.** Mean and standard deviation of the biogenic CO<sub>2</sub> tracer at Xianghe for the different sensitivity experiments BASE, LC and PARAM, per simulation period. “All” indicates all simulated hours (00:00–23:00 LT) were used to calculate the metrics, in contrast to “Afternoon” (12:00–15:00 LT), “Night” (22:00–04:00 LT), and “Morning” (08:00–12:00 LT).

		December 2018	March 2019	May 2019	July 2019
All	PROF	2.29 ± 2.67	6.25 ± 5.60	3.79 ± 9.35	8.07 ± 12.89
	LC	2.28 ± 2.66	4.67 ± 4.19	2.15 ± 7.91	6.62 ± 10.04
	PARAM	4.38 ± 4.91	9.49 ± 7.54	10.53 ± 10.09	17.24 ± 13.85
Afternoon	PROF	1.24 ± 1.72	3.34 ± 4.64	−2.62 ± 3.02	−3.19 ± 4.44
	LC	1.23 ± 1.70	2.63 ± 3.60	−2.62 ± 4.08	−1.78 ± 3.51
	PARAM	1.99 ± 2.91	6.20 ± 7.13	4.23 ± 3.11	6.23 ± 5.32
Night	PROF	3.03 ± 3.30	8.25 ± 5.01	9.91 ± 9.26	16.68 ± 9.92
	LC	3.02 ± 3.28	6.05 ± 3.55	6.50 ± 8.38	12.77 ± 8.11
	PARAM	5.96 ± 5.84	11.78 ± 6.57	15.62 ± 10.11	24.25 ± 11.18
Morning	PROF	1.88 ± 2.51	6.49 ± 6.78	0.42 ± 6.35	4.60 ± 9.35
	LC	1.87 ± 2.50	4.89 ± 5.21	−0.19 ± 5.39	4.80 ± 7.92
	PARAM	3.78 ± 4.90	9.70 ± 9.12	9.75 ± 8.73	17.00 ± 12.16

## 4 Discussion

### 4.1 Sector contributions: differences between in situ CO<sub>2</sub> and XCO<sub>2</sub>

In Sect. 3.2 we compared relative tracer contributions in the WRF-GHG simulations for near-surface CO<sub>2</sub> and column-averaged XCO<sub>2</sub>, and found a notable difference in the biogenic contribution between the two. To analyze this difference, Fig. 9 presents mean vertical profiles of the simulated CO<sub>2</sub> tracers at Xianghe.

The profiles show that the dominant tracer signals in WRF-GHG are limited to the lowest 4 km of the column. Panel (a) reveals a large monthly variability in the vertical distribution of the biogenic tracer. From May through September the biogenic signal at Xianghe is generally negative through much of the column but positive in the lowest levels. Near-surface values (below 200 m) are, on average, positive in all months except August. This pattern is consistent with Fig. 4, which shows a negative biosphere contribution in August for in situ CO<sub>2</sub> while XCO<sub>2</sub> indicates a biospheric sink across May–September. These vertical profiles indicate that the difference can be linked to two factors: the different sensitivities of the measurement techniques and Xianghe’s location relative to strong land sinks. In situ observations are typically more sensitive to local fluxes (i.e. from urban areas and cropland), which are a net source for most months (except August) as calculated by VPRM (see Fig. 5). In contrast, column measurements (XCO<sub>2</sub>) integrate the entire atmosphere and are sensitive to fluxes on a larger scale: in this case the forested mountains roughly 50 km north and 90 km east of Xianghe (see Fig. 5), producing a sink over Summer.

Panel (b) of Fig. 9 shows mean profiles of all tracers averaged over the full simulation period. Unlike the biogenic tracer, the industry, residential, and energy tracers are posi-


tive at all heights and exhibit a strong near-surface maximum that decays exponentially with altitude. Because these anthropogenic tracers do not change sign with height, their relative contributions are similar for both in situ CO<sub>2</sub> and XCO<sub>2</sub>.

### 4.2 July XCO<sub>2</sub> anomaly case study

A notable spike in XCO<sub>2</sub> levels is observed between 20–29 July (see Fig. 2a), diverging from the typical decreasing trend of XCO<sub>2</sub> from May to September. We will focus on the model simulations between 7 July and 30 August 2019 to explain the causes of this XCO<sub>2</sub> summer spike, as WRF-GHG correlates well with the observations during this period (correlation coefficient of 0.84).

As shown in Fig. 10a, the total simulated XCO<sub>2</sub> increases from  $406.30 \pm 1.97$  ppm before the summer spike (7–19 July) to  $408.23 \pm 1.67$  ppm during the spike (20–29 July), then decreases to  $405.01 \pm 1.63$  ppm afterward (30 July–30 August). These values represent the mean and standard deviation of all hourly WRF-GHG simulated XCO<sub>2</sub> values during each period. Figure 10 shows the simulated background and tracer contributions during this period. Figure 10a shows that the background XCO<sub>2</sub> remains relatively constant in July ( $406.65 \pm 0.93$  ppm), and decreases to  $405.63 \pm 1.19$  ppm in August. It further clearly indicates a negative contribution of the tracers before and after the summer spike to a positive enhancement during the spike period. Looking at the different tracers in Fig. 10b, we see that it is mainly the biogenic tracer that has a different behavior in the spike period compared to the periods before and after. Thus, the increase in XCO<sub>2</sub> between 20–29 July is mainly linked to a weaker biogenic sink ( $−0.86 \pm 1.04$  ppm) compared to the periods before ( $−3.56 \pm 1.44$  ppm) and after ( $−2.19 \pm 1.39$  ppm).

Further analysis reveals that during the spike, a heatwave with surface temperatures up to 39 °C occurred, together with



**Figure 9.** Vertical distribution of simulated CO<sub>2</sub> tracers in WRF-GHG at Xianghe up to 12 km altitude (a) at monthly scale for the biogenic tracer and (b) annual scale for all tracers.

800 hPa winds predominantly from the west (see Fig. 10a and c). The biogenic tracer also shows increased values across a large vertical extent in the troposphere (Fig. 10c), indicating advection from other regions. Synoptic maps (Figs. A2 and A3) show that at the onset of the event, on 20 July 2019, a warm air mass arrived from the northwest. This air mass, originating from the Gobi Desert and grasslands in Inner Mongolia, both areas that are characterized by sparse vegetation and elevated temperatures, carried a lack of biogenic signal and coincides with the jump in the biogenic XCO<sub>2</sub> tracer.

Additionally, the mean NEE around Xianghe, as calculated by VPRM, is slightly higher between 20 and 29 July (average of  $-5941 \text{ mol km}^{-2} \text{ h}^{-1}$  over domain d03) compared to the periods before and after (respectively  $-9153$  and  $-12\,785 \text{ mol km}^{-2} \text{ h}^{-1}$ ). In VPRM, the respiration component is linearly dependent on surface temperature, and the gross ecosystem exchange also has a temperature dependency representing the temperature sensitivity of photosynthesis, with CO<sub>2</sub> uptake decreasing at temperatures higher than optimal (Mahadevan et al., 2008). Indeed, it has been shown that extreme temperatures impact CO<sub>2</sub> fluxes (Xu et al., 2020; Ramonet et al., 2020; Gupta et al., 2021).


Therefore, we conclude that the spike was caused by an atmospheric circulation anomaly resulting in the advection of a warm air mass with high biogenic CO<sub>2</sub> levels, followed by locally reduced photosynthesis and increased respiration due to the resulting hot temperatures.

### 4.3 Sensitivity of near-surface CO<sub>2</sub> simulations to model configuration choices

#### 4.3.1 Emission height

In Sect. 3.4.1, we showed that applying vertical profiles to anthropogenic emissions improved the near-surface CO<sub>2</sub> simulations at Xianghe, substantially reducing the observed nighttime overestimation in the BASE experiment. These findings are consistent with Brunner et al. (2019) and Peng et al. (2023), who highlighted the critical role of emission height in determining near-surface CO<sub>2</sub> mole fractions, particularly under weak mixing conditions. The strong impact on nighttime simulations at Xianghe is likely driven by the proximity of strong point sources (Fig. 5a, b), where the effective release height determines whether emissions remain trapped below or mix above the shallow nocturnal boundary layer.

Nevertheless, some discrepancies with observations remain. In May, for instance, BASE agrees better with observations than PROF, despite May being selected due to poor agreement in the one-year BASE simulation. This apparent contradiction reflects both the short (two-week) duration of the sensitivity runs and the temporal variability of model performance: the main mismatch in BASE occurred in early May, which was not included in the experiments. A full-year sensitivity study would better capture meteorological variability and provide a more robust evaluation, but was not feasible here. The poorer performance of PROF in the second half of May, and the larger MBE in July, remain unexplained, and suggest that further assessment is needed to



**Figure 10.** Simulated time series of XCO<sub>2</sub> at Xianghe from 7 July to 30 August 2019, with the spike period highlighted in all panels. Daily mean (a) background tracer (cyan triangles) and total tracers (red diamonds) from WRF-GHG at Xianghe, and TCCON values (black dots). Error bars represent the standard deviation of the daily mean. Daily mean 800 hPa wind direction is indicated by wind barbs at the bottom. (b) Time series of different tracer contributions at Xianghe, with hourly values shown as thin lines and points for TCCON observation times. (c) Color coded vertical profiles of the biogenic CO<sub>2</sub> contributions (left y-axis) shown in red and blue, and surface temperature (right y-axis) in black.

determine whether the vertical profiles from Brunner et al. (2019) are appropriate for China and consistent with the sectoral and spatial patterns of the emission inventory used in this study.

#### 4.3.2 Land cover representation

Replacing the land-cover dataset with the Copernicus product systematically reduced simulated NEE and nighttime CO<sub>2</sub> mole fractions at Xianghe. This weakening of the biospheric signal is consistent with the larger fraction of non-vegetated land in the Copernicus compared to SYNMAP,

leading to smaller VPRM-driven fluxes. The differences are negligible in December and March, when biospheric activity is minimal. While implementing the Copernicus map does not uniformly improve agreement with observations and slightly worsens performance in some periods, it provides a more realistic land-cover representation and is therefore recommended for future regional experiments.

#### 4.3.3 VPRM parameterization

Applying the VPRM parameters from Glauch et al. (2025) produces higher NEE, primarily due to larger  $\alpha$  and  $\beta$  values

that enhance respiration rates (Table A1). Gross ecosystem exchange (GEE) remains similar between the two configurations, indicating that the net effect is primarily driven by increased respiration rather than changes in photosynthetic uptake. Across all experiments and periods, the PARAM configuration shows the closest agreement with observations, though residual discrepancies suggest that additional model errors remain.

A key limitation is the absence of a standardized VPRM parameter set optimized for China. The only known regional calibration, by Dayalu et al. (2018), introduces seasonal crop subtypes but requires detailed, time-varying land-cover input that is not readily available. Moreover, Seo et al. (2024) reported that the parameter values adopted here (Li et al., 2020) outperform those of Dayalu et al. (2018) over East Asia, supporting our choice.

Recent studies also show that including soil moisture effects in VPRM can improve simulated NEE (Gourdji et al., 2022; Segura-Barrero et al., 2025). However, implementing such modifications would require additional region-specific parameters and could introduce further uncertainty. While the Glauch et al. (2025) parameter set reduces model–observation errors, future work should focus on dedicated VPRM calibration using multi-site eddy-covariance data to develop parameter sets representative of Chinese ecosystems.

#### 4.3.4 Remaining sources of uncertainty

Several factors that were not explicitly tested in this study may still contribute to the remaining biases in simulated near-surface CO<sub>2</sub>. One potential source of uncertainty in many regional modeling studies is the coupling between planetary boundary layer (PBL) dynamics and biogenic CO<sub>2</sub> fluxes, a relationship often referred to as the atmospheric CO<sub>2</sub> *rectifier effect* (Larson and Volkmer, 2008). Since both processes are driven by solar radiation, they interact nonlinearly throughout the diurnal cycle: daytime CO<sub>2</sub> minima result from enhanced turbulent mixing and photosynthetic uptake, whereas nighttime maxima arise from stable stratification and ecosystem respiration. Consequently, accurate simulations require both realistic net ecosystem exchange (NEE) and reliable PBL dynamics. However, in our case, the companion study on CH<sub>4</sub> did not reveal any systematic biases in the mean diurnal cycle, suggesting that PBL processes are reasonably well represented here and unlikely to be the dominant cause of the remaining CO<sub>2</sub> discrepancies. Nevertheless, it is well known that the choice of PBL schemes can have a substantial influence on simulated tracer concentrations (Yu et al., 2022; Kretschmer et al., 2014; Feng et al., 2019; Díaz-Isaac et al., 2018), and uncertainties are generally amplified under weak mixing conditions, such as during the night (Maier et al., 2022). Minor deviations in modeled turbulence or nocturnal stability could therefore still contribute to the observed nighttime biases.

Another contributing factor is the vertical representation of the atmosphere near the surface. Since the observations are collected at 60 m above ground, the simulated CO<sub>2</sub> fields were interpolated to this height. However, in our configuration, the difference in simulated nighttime CO<sub>2</sub> between the two lowest model layers (about 50 and 64 m thick) can reach 20 ppm. This implies that combining an interpolation with too coarse a resolution may introduce errors of several ppm. Increasing vertical resolution within the PBL would help to reduce these artifacts. Overall, the accuracy of near-surface CO<sub>2</sub> simulations depends on the interplay between several model components, including emission height profiles, land cover representation, biogenic flux parameterization, and PBL scheme choice. Improving each of these aspects not only reduces biases but most of all enhances the physical realism of the modeled processes, ultimately leading to more reliable simulations of surface–atmosphere CO<sub>2</sub> exchanges.

## 5 Conclusions

This study is the second part of a broader investigation into greenhouse gas variability and model performance at the Xianghe site, following earlier work focused on CH<sub>4</sub>. Here, we shift the focus to CO<sub>2</sub>, aiming to better understand the observed variability through source attribution and model simulations. Using the WRF-GHG model, we performed a one-year simulation of both surface and column-averaged CO<sub>2</sub>, evaluated model performance against FTIR and in situ observations, and carried out sensitivity experiments to assess the impact of key model settings.

Model evaluation against FTIR observations at Xianghe shows that WRF-GHG is capable of capturing the temporal variability in column-averaged CO<sub>2</sub> (XCO<sub>2</sub>), with a correlation coefficient of 0.7. However, a systematic bias was identified in the model's background CO<sub>2</sub> values from CAMS, with a negative offset exceeding 2 ppm between September and May. After applying a bias correction based on monthly mean CAMS–TCCON differences, the mean bias error was reduced to  $-0.86$  ppm. These findings underscore the importance of accurate boundary conditions when simulating XCO<sub>2</sub>, particularly due to the long atmospheric lifetime of CO<sub>2</sub> and the relatively small contribution of regional emissions to the total column. In our simulations, emissions within the model domain contributed only  $\sim 1.82$  ppm to XCO<sub>2</sub>, making the column signal highly sensitive to background mole fractions. Furthermore, the model successfully captured a strong positive anomaly observed in July 2019, attributed to the advection of a warm, CO<sub>2</sub>-rich air mass. This case study illustrates the value of combining transport and mole fraction diagnostics for interpreting episodic events in column data and highlights the dominant role of synoptic meteorology in driving short-term variability in XCO<sub>2</sub>.


For near-surface CO<sub>2</sub> mole fractions, WRF-GHG shows good agreement with afternoon observations at Xianghe, achieving a correlation coefficient of 0.75 and a mean bias error of  $-2.44$  ppm after bias correction. In contrast to XCO<sub>2</sub>, near-surface CO<sub>2</sub> was more strongly influenced by local sources, with a mean tracer enhancement of 26.78 ppm, resulting in a smaller relative importance of boundary condition errors. Nighttime mole fractions are consistently overestimated, with a mean bias of 7.86 ppm and a lower correlation of 0.60. These discrepancies are reflected in the diurnal cycle: while the model captures the overall structure driven by planetary boundary layer dynamics, it overestimates the daily amplitude of 22.1 ppm by 4.58 ppm. Likely causes include inaccuracies in NEE and the vertical distribution of anthropogenic emissions.

Additional sensitivity experiments show that applying vertical emission profiles and a more recent land cover map can reduce nighttime CO<sub>2</sub> mole fractions and improve agreement with observations. Adjustments to the VPRM vegetation parameters substantially affected near-surface mole fractions, with differences up to 10 ppm, underscoring the critical role of appropriate parameter selection—especially in the absence of a standardized VPRM configuration for China.

Tracer analysis confirms that the industry and energy sectors are the dominant contributors to CO<sub>2</sub> levels at Xianghe, while the biosphere plays a secondary, seasonal role. For XCO<sub>2</sub>, the biosphere acts as a sink from April to September ( $-0.77$  ppm on average) and a weak source in the remaining months ( $+0.22$  ppm). At the surface, biospheric uptake is only seen in August ( $-6.76$  ppm), while respiration dominates the rest of the year ( $+2.69$  ppm on average). These differences illustrate the greater local sensitivity of in situ measurements compared to column observations and the varying spatial influence of different source types.

Overall, this study demonstrates the value of using a modeling framework like WRF-GHG to interpret both temporal and sectoral variations in surface and column CO<sub>2</sub> observations. It also highlights that model accuracy is strongly dependent on appropriate configuration choices, including the representation of boundary conditions, vertical emission profiles, and biogenic flux parameterizations. Addressing these factors is essential for improving simulations and supporting more accurate source attribution of observed CO<sub>2</sub> variability.

## Appendix A: Additional tables and figures




**Figure A1.** Same as Fig. 2 but showing the original (not bias corrected) model values.

**Table A1.** VPRM parameter values for different vegetation classes.


		Evergreen forest	Deciduous forest	Mixed forest	Shrubland	Savanna	Cropland	Grassland	Wetland
Li table	PAR <sub>0</sub>	745.306	514.13	419.5	590.7	600	1074.9	717.1	392.666
	$\lambda$	0.13	0.1	0.1	0.18	0.18	0.085	0.115	0.1377
	$\alpha$	0.1247	0.092	0.2	0.0634	0.2	0.13	0.0515	0.0779
	$\beta$	0.2496	0.8430	0.27248	0.2684	0.3376	0.542	-0.0986	0.0902
Glauch table	PAR <sub>0</sub>	521.9	500.8	451.1	444.1		960.8	443.4	399.7
	$\lambda$	0.13	0.13	0.14	0.1		0.09	0.22	0.12
	$\alpha$	0.21	0.23	0.19	0.08		0.17	0.27	0.3
	$\beta$	1.15	1.26	0.93	0.56		1.14	1.63	-0.39

**Table A2.** Same as Table 5 but for XCO<sub>2</sub>.


		MBE				RMSE				CORR			
		Dec	Mar	May	Jul	Dec	Mar	May	Jul	Dec	Mar	May	Jul
All	BASE	-3.15	-2.15	-1.45	0.37	4.07	2.36	1.8	1.36	0.6	0.67	0.38	0.59
	PROF	-3.55	-2.22	-1.56	0.02	4.29	2.41	1.89	1.09	0.63	0.69	0.35	0.71
	LC	-3.56	-2.27	-1.44	0.14	4.29	2.45	1.8	1.07	0.63	0.67	0.33	0.73
	PARAM	-3.56	-1.8	-0.44	1.6	4.29	2.13	1.14	1.97	0.63	0.66	0.39	0.71



**Figure A2.** Maps of eastern China at 800 hPa for 19–23 July 2019, 07:00 UTC (15:00 LT). Panels in the first column show potential temperature (K, in color) with wind barbs and geopotential height contour lines at 800 hPa (contour interval every 20 m); panels in the second column show biogenic XCO<sub>2</sub> enhancements (ppm, in color) with the same wind vectors and contours; the third column shows the NEE. The Xianghe site is marked by a black star.



**Figure A3.** Same as Fig. A2 but over the period 24–28 July 2019.



**Figure A4.** Dominant VPRM vegetation category in WRF-GHG domain d03 (3 km) for (a) 1-km SYNMAP and (b) 100-m Copernicus Dynamic Land Cover Collection 3 (epoch 2019). The black cross indicates the location of the Xianghe site.

**Code and data availability.** The ERA5 and CAMS reanalysis data set <https://doi.org/10.24381/cds.bd0915c6> and <https://doi.org/10.24381/cds.adbb2d47> (Hersbach et al., 2023a, b), used as input for the WRF-GHG simulations, was downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store (2022). The CAMS-GLOB-ANT v5.3 emissions (<https://doi.org/10.24380/D0BN-KX16>, Granier et al., 2019; <https://doi.org/10.5194/essd-16-2261-2024>, Soulie et al., 2024) and temporal profiles CAMS-GLOB-TEMPO v3.1 (<https://doi.org/10.5194/essd-13-367-2021>, Guevara et al., 2021) are archived and distributed through the Emissions of atmospheric Compounds and Compilation of Ancillary Data (ECCAD) platform. The WRF-Chem model code is distributed by NCAR (<https://doi.org/10.5065/D6MK6B4K>, NCAR, 2020). The WRF-GHG simulation output created in the context of this study can be accessed on <https://doi.org/10.18758/P34WJEW2> (Callewaert, 2023). The TCCON data were obtained from the TCCON Data Archive hosted by CaltechDATA at <https://doi.org/10.14291/tccon.ggg2020.xianghe01.R0> (Zhou et al., 2022), while the surface observations at Xianghe were received through private communication with the co-authors.

**Author contributions.** SC made the model simulations and performed the formal analysis, investigation and visualization. The research was conceptualized by SC, MDM and EM and supervised by MDM and EM. MZ, TW and PW have provided the observational in situ data at Xianghe. BL supported with computing tools to correctly compare the model with TCCON data. SC prepared the initial draft of this manuscript while it was reviewed and edited by MZ, BL, TW, MDM, EM and PW.

**Competing interests.** The contact author has declared that none of the authors has any competing interests.

**Disclaimer.** The results contain modified Copernicus Climate Change Service information 2022. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains.

Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

**Special issue statement.** This article is part of the special issue “Greenhouse gas monitoring in the Asia-Pacific region (ACP/AMT/GMD inter-journal SI)”. It is not associated with a conference.

**Acknowledgements.** We would like to thank all staff at the Xianghe site for operating the FTIR and PICARRO measurements. Emmanuel Mahieu is a research director with the F.R.S.-FNRS. The authors acknowledge all providers of observational data and emission inventories. We thank the IT team at BIRA-IASB for their support on data storage and HPC maintenance. Christophe Gerbig, Roberto Kretschmer, and Thomas Koch (MPI BGC) are thanked for distributing the VPRM preprocessor code. Finally, we are grateful for fruitful discussions with Jean-François Müller (BIRA-IASB) and Bernard Heinesch (ULiège).

**Financial support.** This work is supported by the National Key Research and Development Program of China (grant nos. 2023YFB3907500, 2023YFB3907505). This research has been supported by the Belgian Federal Government and the Belgian

tax exemption law for promoting scientific research (Art. 275/3 of CIR92).

**Review statement.** This paper was edited by Chris Wilson and reviewed by Sha Feng and one anonymous referee.

## References

Agustí-Panareda, A., Barré, J., Massart, S., Inness, A., Aben, I., Ades, M., Baier, B. C., Balsamo, G., Borsdorff, T., Bousserez, N., Boussetta, S., Buchwitz, M., Cantarello, L., Crevoisier, C., Engelen, R., Eskes, H., Flemming, J., Garrigues, S., Hasekamp, O., Huijnen, V., Jones, L., Kipling, Z., Langerock, B., McNorton, J., Meilhac, N., Noël, S., Parrington, M., Peuch, V.-H., Ramonet, M., Razinger, M., Reuter, M., Ribas, R., Suttie, M., Sweeney, C., Tarniewicz, J., and Wu, L.: Technical note: The CAMS greenhouse gas reanalysis from 2003 to 2020, *Atmospheric Chemistry and Physics*, 23, 3829–3859, <https://doi.org/10.5194/acp-23-3829-2023>, 2023.

Ahmadov, R., Gerbig, C., Kretschmer, R., Koerner, S., Neininger, B., Dolman, A. J., and Sarrat, C.: Mesoscale Covariance of Transport and CO<sub>2</sub> Fluxes: Evidence from Observations and Simulations Using the WRF-VPRM Coupled Atmosphere-Biosphere Model, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2007JD008552>, 2007.

Ahmadov, R., Gerbig, C., Kretschmer, R., Körner, S., Rödenbeck, C., Bousquet, P., and Ramonet, M.: Comparing high resolution WRF-VPRM simulations and two global CO<sub>2</sub> transport models with coastal tower measurements of CO<sub>2</sub>, *Biogeosciences*, 6, 807–817, <https://doi.org/10.5194/bg-6-807-2009>, 2009.

Ballav, S., Naja, M., Patra, P. K., Machida, T., and Mukai, H.: Assessment of Spatio-Temporal Distribution of CO<sub>2</sub> over Greater Asia Using the WRF-CO<sub>2</sub> Model, *Journal of Earth System Science*, 129, 80, <https://doi.org/10.1007/s12040-020-1352-x>, 2020.

Beck, V., Koch, T., Kretschmer, R., Marshall, J., Ahmadov, R., Gerbig, C., Pillai, D., and Heimann, M.: The WRF Greenhouse Gas Model (WRF-GHG), Tech. rep., iSSN 1615-7400, 2011.

Brunner, D., Kuhlmann, G., Marshall, J., Clément, V., Fuhrer, O., Broquet, G., Löscher, A., and Meijer, Y.: Accounting for the vertical distribution of emissions in atmospheric CO<sub>2</sub> simulations, *Atmospheric Chemistry and Physics*, 19, 4541–4559, <https://doi.org/10.5194/acp-19-4541-2019>, 2019.

Buchhorn, M., Smets, B., Bertels, L., Roo, B. D., Lesiv, M., Tsendbazar, N.-E., Herold, M., and Fritz, S.: Copernicus Global Land Service: Land Cover 100 m: Collection 3: Epoch 2019: Globe, <https://doi.org/10.2909/c6377c6e-76cc-4d03-8330-628a03693042>, 2020.

Callewaert, S.: WRF-Chem Simulations of CO<sub>2</sub>, CH<sub>4</sub> and CO around Xianghe, China, BIRA-IASB [data set], <https://doi.org/10.18758/P34WJEW2>, 2023.

Callewaert, S., Zhou, M., Langerock, B., Wang, P., Wang, T., Mahieu, E., and De Mazière, M.: A WRF-Chem study of the greenhouse gas column and in situ surface concentrations observed in Xianghe, China – Part 1: Methane (CH<sub>4</sub>), *Atmospheric Chemistry and Physics*, 25, 9519–9544, <https://doi.org/10.5194/acp-25-9519-2025>, 2025.

Dayalu, A., Munger, J. W., Wofsy, S. C., Wang, Y., Nehrkorn, T., Zhao, Y., McElroy, M. B., Nielsen, C. P., and Luus, K.: Assessing biotic contributions to CO<sub>2</sub> fluxes in northern China using the Vegetation, Photosynthesis and Respiration Model (VPRM-CHINA) and observations from 2005 to 2009, *Biogeosciences*, 15, 6713–6729, <https://doi.org/10.5194/bg-15-6713-2018>, 2018.

Díaz-Isaac, L. I., Lauvaux, T., and Davis, K. J.: Impact of physical parameterizations and initial conditions on simulated atmospheric transport and CO<sub>2</sub> mole fractions in the US Midwest, *Atmospheric Chemistry and Physics*, 18, 14813–14835, <https://doi.org/10.5194/acp-18-14813-2018>, 2018.

Dong, X., Yue, M., Jiang, Y., Hu, X.-M., Ma, Q., Pu, J., and Zhou, G.: Analysis of CO<sub>2</sub> spatio-temporal variations in China using a weather–biosphere online coupled model, *Atmospheric Chemistry and Physics*, 21, 7217–7233, <https://doi.org/10.5194/acp-21-7217-2021>, 2021.

Fast, J. D., Gustafson Jr., W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of Ozone, Particulates, and Aerosol Direct Radiative Forcing in the Vicinity of Houston Using a Fully Coupled Meteorology-Chemistry-Aerosol Model, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006721>, 2006.

Feng, S., Lauvaux, T., Newman, S., Rao, P., Ahmadov, R., Deng, A., Díaz-Isaac, L. I., Duren, R. M., Fischer, M. L., Gerbig, C., Gurney, K. R., Huang, J., Jeong, S., Li, Z., Miller, C. E., O’Keffe, D., Patarasuk, R., Sander, S. P., Song, Y., Wong, K. W., and Yung, Y. L.: Los Angeles megacity: a high-resolution land–atmosphere modelling system for urban CO<sub>2</sub> emissions, *Atmospheric Chemistry and Physics*, 16, 9019–9045, <https://doi.org/10.5194/acp-16-9019-2016>, 2016.

Feng, S., Lauvaux, T., Davis, K. J., Keller, K., Zhou, Y., Williams, C., Schuh, A. E., Liu, J., and Baker, I.: Seasonal Characteristics of Model Uncertainties From Biogenic Fluxes, Transport, and Large-Scale Boundary Inflow in Atmospheric CO<sub>2</sub> Simulations Over North America, *Journal of Geophysical Research: Atmospheres*, 124, 14325–14346, <https://doi.org/10.1029/2019JD031165>, 2019.

Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Landschützer, P., Le Quéré, C., Li, H., Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshakel, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Arneth, A., Arora, V., Bates, N. R., Becker, M., Bellouin, N., Berghoff, C. F., Bittig, H. C., Bopp, L., Cadule, P., Campbell, K., Chamberlain, M. A., Chandra, N., Chevallier, F., Chini, L. P., Colligan, T., Decayeux, J., Djutchouang, L. M., Dou, X., Duran Rojas, C., Enyo, K., Evans, W., Fay, A. R., Feely, R. A., Ford, D. J., Foster, A., Gasser, T., Gehlen, M., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A. K., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Kato, E., Keeling, R. F., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Lan, X., Lauvset, S. K., Lefèvre, N., Liu, Z., Liu, J., Ma, L., Maksyutov, S., Marland, G., Mayot, N., McGuire, P. C., Metzl, N., Monacci, N. M., Morgan, E. J., Nakao, S.-I., Neill, C., Niwa, Y., Nützel, T., Olivier, L., Ono, T., Palmer, P. I., Pierrot, D., Qin, Z., Resplandy, L., Roobaert, A., Rosan, T. M., Rödenbeck, C., Schwinger, J., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Steinhoff, T., Sun, Q., Sutton, A. J., Séférian, R., Takao, S., Tatebe, H., Tian, H., Tilbrook, B., Torres, O., Tourigny, E., Tsujino, H.,

Tubiello, F., van der Werf, G., Wanninkhof, R., Wang, X., Yang, D., Yang, X., Yu, Z., Yuan, W., Yue, X., Zaehle, S., Zeng, N., and Zeng, J.: Global Carbon Budget 2024, *Earth System Science Data*, 17, 965–1039, <https://doi.org/10.5194/essd-17-965-2025>, 2025.

Glauch, T., Marshall, J., Gerbig, C., Botía, S., Gałkowski, M., Vardag, S. N., and Butz, A.: *pyVPRM*: A next-generation Vegetation Photosynthesis and Respiration Model for the post-MODIS era, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-3692>, 2025.

Gourdji, S. M., Karion, A., Lopez-Coto, I., Ghosh, S., Mueller, K. L., Zhou, Y., Williams, C. A., Baker, I. T., Haynes, K. D., and Whetstone, J. R.: A Modified Vegetation Photosynthesis and Respiration Model (VPRM) for the Eastern USA and Canada, Evaluated With Comparison to Atmospheric Observations and Other Biospheric Models, *Journal of Geophysical Research: Biogeosciences*, 127, <https://doi.org/10.1029/2021JG006290>, 2022.

Granier, C., Darras, S., Denier van der Gon, H., Doubalova, J., Elguindi, N., Galle, B., Gauss, M., Guevara, M., Jalkanen, J.-P., Kuenen, J., Liousse, C., Quack, B., Simpson, D., and Sindelarova, K.: The Copernicus Atmosphere Monitoring Service Global and Regional Emissions (April 2019 Version), Copernicus Atmospheric Monitoring Service, <https://doi.org/10.24380/D0BN-KX16>, 2019.

Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully Coupled “Online” Chemistry within the WRF Model, *Atmospheric Environment*, 39, 6957–6975, <https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.

Guevara, M., Jorba, O., Tena, C., Denier van der Gon, H., Kuenen, J., Elguindi, N., Darras, S., Granier, C., and Pérez García-Pando, C.: Copernicus Atmosphere Monitoring Service TEMPOral profiles (CAMS-TEMPO): global and European emission temporal profile maps for atmospheric chemistry modelling, *Earth System Science Data*, 13, 367–404, <https://doi.org/10.5194/essd-13-367-2021>, 2021.

Gupta, S., Tiwari, Y. K., Revadekar, J. V., Burman, P. K. D., Chakraborty, S., and Gnanamoorthy, P.: An Intensification of Atmospheric CO<sub>2</sub> Concentrations Due to the Surface Temperature Extremes in India, *Meteorology and Atmospheric Physics*, 133, 1647–1659, <https://doi.org/10.1007/s00703-021-00834-w>, 2021.

Hall, B. D., Crotwell, A. M., Kitzis, D. R., Mefford, T., Miller, B. R., Schibig, M. F., and Tans, P. P.: Revision of the World Meteorological Organization Global Atmosphere Watch (WMO/GAW) CO<sub>2</sub> calibration scale, *Atmospheric Measurement Techniques*, 14, 3015–3032, <https://doi.org/10.5194/amt-14-3015-2021>, 2021.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Munoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 Hourly Data on Pressure Levels from 1940 to Present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.bd0915c6>, 2023a.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Munoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 Hourly Data on Single Levels from 1940 to Present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2023b.

Hu, X.-M., Crowell, S., Wang, Q., Zhang, Y., Davis, K. J., Xue, M., Xiao, X., Moore, B., Wu, X., Choi, Y., and DiGangi, J. P.: Dynamical Downscaling of CO<sub>2</sub> in 2016 Over the Contiguous United States Using WRF-VPRM, a Weather-Biosphere-Online-Coupled Model, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001875, <https://doi.org/10.1029/2019MS001875>, 2020.

Hu, X.-M., Gourdji, S. M., Davis, K. J., Wang, Q., Zhang, Y., Xue, M., Feng, S., Moore, B., and Crowell, S. M. R.: Implementation of Improved Parameterization of Terrestrial Flux in WRF-VPRM Improves the Simulation of Nighttime CO<sub>2</sub> Peaks and a Daytime CO<sub>2</sub> Band Ahead of a Cold Front, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD034362, <https://doi.org/10.1029/2020JD034362>, 2021.

Jung, M., Henkel, K., Herold, M., and Churkina, G.: Exploiting Synergies of Global Land Cover Products for Carbon Cycle Modeling, *Remote Sensing of Environment*, 101, 534–553, <https://doi.org/10.1016/j.rse.2006.01.020>, 2006.

Kretschmer, R., Gerbig, C., Karstens, U., Biavati, G., Vermeulen, A., Vogel, F., Hammer, S., and Totsche, K. U.: Impact of optimized mixing heights on simulated regional atmospheric transport of CO<sub>2</sub>, *Atmospheric Chemistry and Physics*, 14, 7149–7172, <https://doi.org/10.5194/acp-14-7149-2014>, 2014.

Landschützer, P., Gruber, N., and Bakker, D. C. E.: An Observation-Based Global Monthly Gridded Sea Surface pCO<sub>2</sub> Product from 1982 Onward and Its Monthly Climatology (NCEI Accession 0160558), NOAA National Centers for Environmental Information [data set], <https://doi.org/10.7289/v5z899n6>, 2017.

Larson, V. E. and Volkmer, H.: An Idealized Model of the One-Dimensional Carbon Dioxide Rectifier Effect, *Tellus B: Chemical and Physical Meteorology*, 60, 525, <https://doi.org/10.1111/j.1600-0889.2008.00368.x>, 2008.

Laughner, J. L., Toon, G. C., Mendonca, J., Petri, C., Roche, S., Wunch, D., Blavier, J.-F., Griffith, D. W. T., Heikkinen, P., Keeling, R. F., Kiel, M., Kivi, R., Roehl, C. M., Stephens, B. B., Baier, B. C., Chen, H., Choi, Y., Deutscher, N. M., DiGangi, J. P., Gross, J., Herkommmer, B., Jeseck, P., Laemmel, T., Lan, X., McGee, E., McKain, K., Miller, J., Morino, I., Notholt, J., Ohyama, H., Pollard, D. F., Rettinger, M., Riris, H., Rousogenous, C., Sha, M. K., Shiomi, K., Strong, K., Sussmann, R., Té, Y., Velazco, V. A., Wofsy, S. C., Zhou, M., and Wennberg, P. O.: The Total Carbon Column Observing Network’s GGG2020 data version, *Earth System Science Data*, 16, 2197–2260, <https://doi.org/10.5194/essd-16-2197-2024>, 2024.

Li, X., Hu, X.-M., Cai, C., Jia, Q., Zhang, Y., Liu, J., Xue, M., Xu, J., Wen, R., and Crowell, S. M. R.: Terrestrial CO<sub>2</sub> Fluxes, Concentrations, Sources and Budget in Northeast China: Observational and Modeling Studies, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031686, <https://doi.org/10.1029/2019JD031686>, 2020.

Mahadevan, P., Wofsy, S. C., Matross, D. M., Xiao, X., Dunn, A. L., Lin, J. C., Gerbig, C., Munger, J. W., Chow, V. Y., and Gottlieb, E. W.: A Satellite-Based Biosphere Parameterization for Net Ecosystem CO<sub>2</sub> Exchange: Vegetation Photosynthesis and Respiration Model (VPRM), *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2006GB002735>, 2008.

Maier, F., Gerbig, C., Levin, I., Super, I., Marshall, J., and Hammer, S.: Effects of point source emission heights in WRF-STILT: a step towards exploiting nocturnal observations in

models, *Geoscientific Model Development*, 15, 5391–5406, <https://doi.org/10.5194/gmd-15-5391-2022>, 2022.

Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou (eds.), A. B.: IPCC, 2021: Climate Change 2021: The Physical Science Basis, Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Tech. rep., Cambridge University Press, <https://doi.org/10.1017/9781009157896>, 2021.

NCAR: The Weather Research and Forecasting Model, Github [code], <https://github.com/wrf-model/WRF> (last access: 21 April 2020), 2020.

Park, C., Gerbig, C., Newman, S., Ahmadov, R., Feng, S., Gurney, K. R., Carmichael, G. R., Park, S.-Y., Lee, H.-W., Goulden, M., Stutz, J., Peischl, J., and Ryerson, T.: CO<sub>2</sub> Transport, Variability, and Budget over the Southern California Air Basin Using the High-Resolution WRF-VPRM Model during the CalNex 2010 Campaign, *Journal of Applied Meteorology and Climatology*, 57, 1337–1352, <https://doi.org/10.1175/JAMC-D-17-0358.1>, 2018.

Park, C., Park, S.-Y., Gurney, K. R., Gerbig, C., DiGangi, J. P., Choi, Y., and Lee, H. W.: Numerical Simulation of Atmospheric CO<sub>2</sub> Concentration and Flux over the Korean Peninsula Using WRF-VPRM Model during Korus-AQ 2016 Campaign, *PLOS ONE*, 15, e0228106, <https://doi.org/10.1371/journal.pone.0228106>, 2020.

Peng, Y., Hu, C., Ai, X., Li, Y., Gao, L., Liu, H., Zhang, J., and Xiao, W.: Improvements of Simulating Urban Atmospheric CO<sub>2</sub> Concentration by Coupling with Emission Height and Dynamic Boundary Layer Variations in WRF-STILT Model, *Atmosphere*, 14, 223, <https://doi.org/10.3390/atmos14020223>, 2023.

Ramonet, M., Ciais, P., Apadula, F., Bartyzel, J., Bastos, A., Bergamaschi, P., Blanc, P. E., Brunner, D., Caracciolo di Torchiarolo, L., Calzolari, F., Chen, H., Chmura, L., Colomb, A., Conil, S., Cristofanelli, P., Cuevas, E., Curcoll, R., Delmotte, M., di Sarra, A., Emmenegger, L., Forster, G., Frumau, A., Gerbig, C., Gheusi, F., Hammer, S., Haszpra, L., Hatakka, J., Hazan, L., Heliasz, M., Henne, S., Hensen, A., Hermansen, O., Keronen, P., Kivi, R., Komíková, K., Kubistin, D., Laurent, O., Laurila, T., Lavric, J. V., Lehner, I., Lehtinen, K. E. J., Leskinen, A., Leuenberger, M., Levin, I., Lindauer, M., Lopez, M., Myhre, C. L., Mammarella, I., Manca, G., Manning, A., Marek, M. V., Marklund, P., Martin, D., Meinhardt, F., Mihalopoulos, N., Mölder, M., Morgui, J. A., Necki, J., O'Doherty, S., O'Dowd, C., Ottosson, M., Philippon, C., Piacentino, S., Pichon, J. M., Plass-Duelmer, C., Resovsky, A., Rivier, L., Rodó, X., Sha, M. K., Scheeren, H. A., Sferlazzo, D., Spain, T. G., Stanley, K. M., Steinbacher, M., Trisolino, P., Vermeulen, A., Vítková, G., Weyrauch, D., Xueref-Remy, I., Yala, K., and Yver Kwok, C.: The Fingerprint of the Summer 2018 Drought in Europe on Ground-Based Atmospheric CO<sub>2</sub> Measurements, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375, 20190513, <https://doi.org/10.1098/rstb.2019.0513>, 2020.

Ramonet, M., Langerock, B., Warneke, T., and Eskes, H.: Validation Report of the CAMS Greenhouse Gas Global Reanalysis, Years 2003–2020, KNMI, <https://doi.org/10.24380/438C-4597>, 2021.

Segura-Barrero, R., Lauvaux, T., Lian, J., Ciais, P., Badia, A., Ventura, S., Bazzi, H., Abbessi, E., Fu, Z., Xiao, J., Li, X., and Villalba, G.: Heat and Drought Events Alter Biogenic Capacity to Balance CO<sub>2</sub> Budget in South-Western Europe, *Global Biogeochemical Cycles*, 39, e2024GB008163, <https://doi.org/10.1029/2024GB008163>, 2025.

Seo, M.-G., Kim, H. M., and Kim, D.-H.: Effect of Atmospheric Conditions and VPRM Parameters on High-Resolution Regional CO<sub>2</sub> Simulations over East Asia, *Theoretical and Applied Climatology*, 155, 859–877, <https://doi.org/10.1007/s00704-023-04663-2>, 2024.

Skamarock, C., Klemp, B., Dudhia, J., Gill, O., Liu, Z., Berner, J., Wang, W., Powers, G., Duda, G., Barker, D., and Huang, X.-y.: A Description of the Advanced Research WRF Model Version 4.1, Technical report, National Center for Atmospheric Research, <https://doi.org/10.5065/1dfh-6p97>, 2019.

Soulie, A., Granier, C., Darras, S., Zilberman, N., Doumbia, T., Guevara, M., Jalkanen, J.-P., Keita, S., Liousse, C., Crippa, M., Guizzardi, D., Hoesly, R., and Smith, S. J.: Global anthropogenic emissions (CAMS-GLOB-ANT) for the Copernicus Atmosphere Monitoring Service simulations of air quality forecasts and reanalyses, *Earth System Science Data*, 16, 2261–2279, <https://doi.org/10.5194/essd-16-2261-2024>, 2024.

Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): A High Resolution Global Model to Estimate the Emissions from Open Burning, *Geoscientific Model Development*, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.

Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column Observing Network, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369, 2087–2112, <https://doi.org/10.1098/rsta.2010.0240>, 2011.

Xu, H., Xiao, J., and Zhang, Z.: Heatwave Effects on Gross Primary Production of Northern Mid-Latitude Ecosystems, *Environmental Research Letters*, 15, 074027, <https://doi.org/10.1088/1748-9326/ab8760>, 2020.

Yang, Y., Zhou, M., Langerock, B., Sha, M. K., Hermans, C., Wang, T., Ji, D., Vigouroux, C., Kumps, N., Wang, G., De Mazière, M., and Wang, P.: New ground-based Fourier-transform near-infrared solar absorption measurements of XCO<sub>2</sub>, XCH<sub>4</sub> and XCO at Xianghe, China, *Earth System Science Data*, 12, 1679–1696, <https://doi.org/10.5194/essd-12-1679-2020>, 2020.

Yang, Y., Zhou, M., Wang, T., Yao, B., Han, P., Ji, D., Zhou, W., Sun, Y., Wang, G., and Wang, P.: Spatial and temporal variations of CO<sub>2</sub> mole fractions observed at Beijing, Xianghe, and Xinglong in North China, *Atmospheric Chemistry and Physics*, 21, 11741–11757, <https://doi.org/10.5194/acp-21-11741-2021>, 2021.

Yu, E., Bai, R., Chen, X., and Shao, L.: Impact of physical parameterizations on wind simulation with WRF V3.9.1.1 under stable conditions at planetary boundary layer gray-zone resolution: a case study over the coastal regions of North China, *Geoscientific Model Development*, 15, 8111–8134, <https://doi.org/10.5194/gmd-15-8111-2022>, 2022.

Zhao, X., Marshall, J., Hachinger, S., Gerbig, C., Frey, M., Hase, F., and Chen, J.: Analysis of total column CO<sub>2</sub> and CH<sub>4</sub> measurements in Berlin with WRF-GHG, *Atmospheric Chemistry and Physics*, 19, 11279–11302, <https://doi.org/10.5194/acp-19-11279-2019>, 2019.

Zhou, M., Wang, P., Kumps, N., Hermans, C., and Nan, W.: TCCON Data from Xianghe, China, Release GGG2020.R0 (Version R0), Caltech Library [data set], <https://doi.org/10.14291/tccon.ggg2020.xianghe01.R0>, 2022.